

GLIWICE RADIOCARBON DATES IX

MIECZYSŁAW F PAZDUR, ROMUALD AWSIUK,
ANDRZEJ BLUSZCZ, ANNA PAZDUR, ADAM WALANUS,
and ANDRZEJ ZASTAWNÝ

Institute of Physics, Silesian Technical University, Krzywoustego 2,
PL-44-100 Gliwice, Poland

The following list contains most of the measurements of geologic samples made during 1979 and 1980 using carbon-dioxide-filled proportional counters. Samples with laboratory numbers > 1000 were counted with counter No. 1 (L1), and those with numbers < 1000 , with counter No. 3 (L3), described by Mościcki and Zastawny (1977). Actual parameters of both counters were given in our previous date list (R, 1982, v 24, p 172).

Methods of sample pretreatment follow, with some minor modifications, those described by Olsson (1979). Some samples of peat, peaty mud, and gyttja were dated using both alkali-soluble fractions, precipitated by adding appropriate amounts of hydrochloric acid, and insoluble residue. The results obtained for series of such paired samples were discussed by Pazdur (1982) and conclusions are similar to those of Sheppard, Ali, and Mehringer (1979).

Combustion and purification lines were described by Pazdur *et al* (1979), the electronic system, designed and described by Bluszcz and Walanus (1980), is built of CAMAC units produced by POLON Enterprise, Warsaw. Measurement of any sample, including background and modern samples, consists of a series of 20 to 25 partial measurements of 100 minutes duration. Partial results obtained in such series were analyzed automatically at the computing center (Pazdur and Walanus, 1979a). Age calculations are based on contemporary value equal to 0.95 of the activity of NBS Oxalic Acid standard and on the Libby half-life of ^{14}C . Ages are reported as conventional ^{14}C dates in years before AD 1950. No corrections for isotopic fractionation in nature are made for samples presented in this list. Errors quoted ($\pm 1\sigma$) include estimated overall standard deviations of count rates of the unknown sample, contemporary standard, and background (Pazdur & Walanus, 1979a). In spite of our proposition (Walanus & Pazdur, 1980), infinite dates are reported according to 2σ criterion, as recommended by Stuiver and Polach (1977, p 362).

Most samples in this list were dated for investigations within the IGCP 158 Project "Paleohydrology of the temperate zone", Subproject A "Fluvial environment". A brief report on the activities of the Polish Group of the IGCP 158 Project was given by Starkel (1981a), the results obtained up to 1981 were presented at the Symposium "Paleohydrology of the temperate zone", Poznań, Poland, 1981, Sept 22-28 (Kozarski & Tobolski, 1981; Starkel, 1981b,c). All dates made for studies connected with Subproject 158B "Lake and mire environments" (Berglund, 1979) will be included in our next date list.

ACKNOWLEDGMENTS

The authors wish to thank Helena Skorupka for her excellent technical assistance in laboratory.

SAMPLE DESCRIPTIONS

I. GEOLOGIC SAMPLES

A. Poland

Gardno series

Peat from layers of varying thickness overlying sandy deposits and underlying gyttja in basin of Gardno Lake, Slovinian coastline. Coll and subm 1980 by Adam Wojciechowski, Inst Geog, Adam Mickiewicz Univ, Poznań. Samples dated for geomorphol and strat studies of basal deposits of Gardno Lake. Deposits underwent pollen (Zachowicz, 1977) and diatom analysis (Zaborowska, 1977). Profile Ga-A3 is 1200m W of Łupawa R mouth to Gardno Lake ($54^{\circ} 39' N$, $17^{\circ} 09' E$); coll Jan 1980. Profile Ga-D2 taken at S shore of lake ca 600m N of Retowo village ($54^{\circ} 38' 20'' N$, $17^{\circ} 07' 30'' E$); coll Feb 1980.

Gd-1286. Gardno Ga-A3a **9090 ± 70**

Carex peat from depth 255 to 260cm below water level, top of peat layer ca 1m thick, overlain by shell gyttja.

Gd-1288. Gardno Ga-A3b **$10,280 \pm 120$**

Carex-Phragmites peat from base of peat layer at depth 350 to 356cm below water level, at contact of medium-grained sands.

Gd-1285. Gardno Ga-D2a **5380 ± 50**

Carex-Phragmites peat from depth 135 to 140cm below water level, top of peat layer ca 1.2m thick, overlain by brown gyttja.

Gd-827. Gardno Ga-D2b **$10,140 \pm 150$**

Carex-Phragmites peat from base of peat layer at contact with gray humus sand, depth 245 to 252cm below water level.

Lubiatowo series

Peat from layer ranging from 1.5 to 3m underlain by medium and coarse-grained sands and overlain by fine-grained eolian (?) sands and peaty soil on coastal river plain near Baltic shore 1km W of Lubiatowo village ($54^{\circ} 48' N$, $17^{\circ} 51' E$), E of Łeba, Gardno-Łeba Plain, Slovinian coastline. Coll Aug 1979 and subm by Sylwester Skompski, Geol Inst, Warsaw.

Gd-749. Lubiatowo L-8:sol **8640 ± 200**

From lowermost part of peat layer at contact with sands, depth 2.9 to 3m, alkali-soluble fraction. *Comment:* undersized, diluted.

Gd-1216. Lubiatowo L-8:ins **8200 ± 100**

Same sample, insoluble fraction.

Gd-748. Lubiatowo L-1:sol **1350 ± 70**

From uppermost part of peat layer at contact with sands, depth 1.5 to 1.7m, alkali-soluble fraction.

Gd-1215. Lubiatowo L-1:ins **1030 ± 60**

Same sample, insoluble fraction.

Białogóra series

Peat from layer in fine-grained sands ranging from 0.7 to 2.4m on coastal river plain, 1.2km ENE of Białogóra village ($54^\circ 49' N$, $17^\circ 59' E$), NW of Zarnowieckie Lake, W part of Karwieńska Plain, 5m asl. Coll July 1979 and subm by Sylwester Skompski.

Gd-746. Białogóra B-7:sol **4470 ± 220**

From lowermost part of peat layer, depth 2.3 to 2.4m, alkali-soluble fraction. *Comment:* undersized, diluted.

Gd-745. Białogóra B-7:ins **4820 ± 100**

Same sample, insoluble fraction.

Gd-747. Białogóra B-1:sol **1750 ± 80**

From uppermost part of peat layer, depth 0.8 to 1m, alkali-soluble fraction.

Gd-1211. Białogóra B-1:ins **1620 ± 80**

Same sample, insoluble fraction.

Zalew Wiślany series

Organogenic sediments from Zalew Wiślany ($54^\circ 23' 07'' N$, $19^\circ 42' 47'' E$). Core 3a, 10m long taken at water depth 3m, reaching Pleistocene gray morainic till at depth 8.8m. Coll April 1979 and subm by Bogusław Rosa, Inst Geog, Dept Geomorphol and Quaternary Geol, Gdańsk Univ, Gdynia.

Gd-1238. Zalew Wiślany 1 **6730 ± 60**

Highly-decomposed homogeneous peaty slime without visible faunal remains, depth 5.5 to 5.75m, central part of peaty mud horizon ranging from 5.26 to 5.91m, overlain by poludine horizon.

Gd-1237. Zalew Wiślany 2 **7120 ± 100**

Highly-decomposed peat, brown at top and dark gray at base, depth 6 to 6.27m.

Gd-1239. Zalew Wiślany 3 **7600 ± 90**

Gray clotty peat, depth 6.9 to 7.15m. *Comment (BR):* characteristic consistency of this peat layer indicates temporal drying of peat-bog surface which means lowering of groundwater level and, at depth 6.88m, contact with local transgression or beginning of local flood.

Gd-1240. Zalew Wislany 4 **9390 ± 110**

Highly-decomposed peat, depth 7.3 to 7.5m.

Gd-772. Zalew Wislany 5 **11,090 ± 150**

Highly-decomposed peat, depth 7.5 to 7.7m.

Gd-773. Zalew Wislany 6 **11,240 ± 110**

Peaty slime rich in organic matter, brown at top and brown-gray at base, depth 7.7 to 7.95m.

Pomorsko station series

Sandy peat from top of peat layer filling up Odra R paleomeander and overlain by sandy sediments forming alluvial cones, 1km SE of Pomorsko RR sta ($52^{\circ} 02' 40''$ N, $15^{\circ} 32' 40''$ E), 7.5km SW of Sulechów, Cigacice-Krośno sec of Warsaw-Berlin Pradolina. Coll and subm 1979 by Bolesław Nowaczyk, Inst Geog, Adam Mickiewicz Univ, Poznań.

Gd-1195. Pomorsko sta 15/79/BN:sol **1460 ± 50**

Below older alluvial cone, depth 351 to 361cm, coll Aug 1979, alkali-soluble fraction dated.

Gd-726. Pomorsko sta 15/79/BN:ins **1720 ± 70**

Same sample, insoluble fraction.

Gd-1189. Pomorsko sta 16/79/BN:ins **1520 ± 80**

Below younger alluvial cone, depth 92 to 100cm, coll Oct 1979, insoluble fraction dated. *Comment* (BN&MFP): samples dated for systematic geomorphol and strat studies of dunes and lacustrine sediments in vicinity of Pomorsko (Nowaczyk, 1974; 1976; 1979; Nowaczyk & Tobolski, 1979; 1981); for other dates from this area, see Nowaczyk *et al* (1982) and next sec.

Czmoniec series

Organic sediments from peaty paleomeander of younger generation, Warta R valley near Czmoniec, ca 10km N of Srem, Great Poland Lowland ($52^{\circ} 11' N$, $17^{\circ} 00' E$). Coll July 1978 by Stefan Kozarski and Kazimierz Tobolski; subm 1980 by KT to supplement earlier dates from this site (R, 1982, v 24, p 187). Dated for systematic studies of Warta R paleomeanders initiated by Stefan Kozarski in 1967 and now included into IGCP 158A Project (Kozarski, 1981b,c; Okuniewska & Tobolski, 1981; Tobolski, 1981).

Gd-818. Czmoniec Cz I/78A/110-115 **3300 ± 120**

Wood peat with sand, top of peat layer at contact with detritus gyttja, depth 110 to 115cm. *Comment*: undersized, diluted.

Gd-817. Czmoniec Cz I/78A/144-150 **3550 ± 80**

Coarse detritus gyttja, with sand, top of gyttja layer at contact with wood peat layer, depth 144 to 150cm.

Zabno series

Channel deposits from paleochannels on terrace III of Warta R valley (Antczak, 1981; Kozarski, 1981b; Kozarski & Rotnicki, 1977) with traces of braided river flow, ca 3km W of recent Warta R channel at Zabno village ($52^{\circ} 11' 32''$ N, $16^{\circ} 54' 09''$ E), 30km S of Poznań, Srem sec of Warsaw-Berlin Pradolina. Coll March 1980 by Barbara Antczak; subm 1980 by Stefan Kozarski, Inst Geog, Adam Mickiewicz Univ, Poznań. Dated for investigations in IGCP 158A Project.

Gd-812. Zabno IM **6300 ± 790**

Organogenic silt from layer 30cm thick between two series of coarse and medium-grained sands, depth 1.06m. *Comment:* very small sample, undersized, diluted.

Gd-819. Zabno III Da **2200 ± 270**

Wood remnants, coarse fraction, from series of coarse- and medium-grained sands, depth 1.1m. *Comment:* undersized, diluted.

Gd-810. Zabno III Db **4560 ± 120**

Wood remnants, fine fraction from same series of sands.

General Comment (BA): samples dated to estimate end of period of water outflow on terrace III. Dates younger than expected, other dates from organic deposits filling up paleomeanders on terrace II are Gd-387, $11,430 \pm 630$; Gd-381, 9780 ± 340 ; Gd-380, 9770 ± 230 (R, 1979, v 21, p 167) and Gd-239, 9650 ± 240 (R, 1978, v 20, p 406); for general discussion of problem, see Kozarski (1981b) and Kozarski and Rotnicki (1977).

Gd-851. Sławie **7320 ± 170**

Wood, single fragments found at depth 2m in layer of sedge-reed peat in fossil peat bog at SW part of Sławskie Lake ($51^{\circ} 52' 40''$ N, $16^{\circ} 02' 22''$ E), ca 1km NW of Radzyń village, Sławskie Lakeland, Great Poland Lowland. Coll May 1979 and subm by Jerzy Janczak, Inst Meteorol and Water Economy, Poznań.

Wieruszów series

Samples from base layers of organic deposits filling up cut-off paleomeander channels on valley floor of Prosna R near Wieruszów, SE part of Great Poland Lowland. Coll March 1980 and subm by Karol Rotnicki, Inst Geog, Adam Mickiewicz Univ, Poznań. Samples dated for reconstruction of paleohydrological conditions in middle course of Prosna R during Holocene in connection with IGCP 158A Project (Kozarski & Rotnicki, 1977).

Gd-1274. Szpot 1/80KR **6390 ± 60**

Silt with admixture of plant remnants, depth 115 to 125cm, overlying coarse-grained sands and covered with peat, 850m N of small village Szpot, 6.5km S of Wieruszów ($51^{\circ} 14' 20''$ N, $18^{\circ} 10' 17''$ E).

Gd-1275. Szpot 4/80KR 5660 ± 60

Silt with plant remnants, depth ca 1m, 550m W of Prosna R, 7.5km S of Wieruszów ($51^{\circ} 13' 53''$ N, $18^{\circ} 10' 13''$ E).

Gd-1280. Dobrygosć 9/80KR 5600 ± 60

Peaty gyttja from base of layer separating two peat layers, depth 78 to 85cm, 300m SE of forester lodge Dobrygość, 300m W of Prosna R, 5km S of Wieruszów ($51^{\circ} 15' 00''$ N, $18^{\circ} 10' 02''$ E).

Gd-1283. Dobrygosć 11/80KR 8590 ± 70

Peat from base of lower peat layer, depth 142 to 150cm, same loc.

Gd-811. Dobrygosć 17/80KR 6320 ± 80

Peat from base of peat layer overlying thin layers of organic silt and silt with plant remnants, depth 170 to 180cm, 1km S of forester lodge Dobrygość, 5.5km S of Wieruszów ($51^{\circ} 14' 46''$ N, $18^{\circ} 09' 52''$ E).

Gd-1284. Dobrygosć 18/80KR 9050 ± 80

Silt with admixture of plant remnants, depth 200 to 210cm, same loc.

Sobieseki series

Organic deposits in form of layers in Würmian flood deposits on terrace 2 of Prosna R in N part of Grabów Basin, built of lacustrine silts, flood silts, and fine-grained sands and thin layers of peat, peaty silt, and silty peat with sand admixture, from 20 to 30cm thickness, accumulated from beginning of Würm till first phase of Upper Pleni-Würm. Total thickness of deposits on terrace 2 exceeds 35m. Two profiles were taken near Sobieseki village ($51^{\circ} 38'$ N, $18^{\circ} 18'$ E), ca 19km SE of Kalisz, profile I; coll June 1979 by Józef Wiśniewski; profile II coll Nov 1979 by Karol Rotnicki; subm 1980 by Karol Rotnicki. Samples dated for systematic studies of paleogeog and stratigraphy of Würmian deposits in Great Poland Lowland (Rotnicki, 1966).

Gd-758. Sobieseki SO-IA/79:sol 26,070 ± 900

Peaty silt with sand admixture, depth 835 to 840cm, alkali-soluble fraction.

Gd-1219. Sobieseki SO-IA/79:ins 26,080 ± 550

Same sample, insoluble fraction.

Gd-755. Sobieseki SO-IB/79:sol 26,070 ± 910

Peaty silt with sand admixture, depth 840 to 845cm, alkali-soluble fraction.

Gd-1220. Sobieseki SO-IB/79:ins 25,460 ± 450

Same sample, insoluble fraction.

Gd-752. Sobieseki SO-IIA/79:sol 26,300 ± 1200

Peaty silt with sand admixture, depth 535 to 550cm, alkali-soluble fraction.

Gd-753. Sobieseki SO-IIA/79:ins **$27,000 \pm 1100$**

Same sample, insoluble fraction.

Gd-756. Sobieseki SO-IIB/79:sol **$27,960 \pm 680$**

Sandy peat, depth 705 to 725cm, alkali-soluble fraction.

Gd-757. Sobieseki SO-IIB/79:ins **$27,780 \pm 590$**

Same sample, insoluble fraction.

General Comment (KR): organic silt from continuous layer at depth 520 to 550cm in Weglewice, 30km S of Sobieseki, from similar strat position, yielded dates $35,100 \pm 1000$ and $36,400 \pm 1200$; Lu-1632 and -1632A, (R, 1980, v 22, p 1055-1056).

Konin-Maliniec series

Organogenic interbeddings in series of sediments of stagnant water overlain by series of cross-stratified fluvioglacial sandy gravel sediments and morainic sediments in form of two-partite flow till of Leszno phase of Vistulian Glaciation in Konin-Maliniec site, ca 5km N of Konin ($52^{\circ} 16' N$, $18^{\circ} 15' E$); coll and subm 1979 by Wojciech Stankowski, Inst Geog, Adam Mickiewicz Univ, Poznań. Konin-Malinies site was presented during field excursion of Symposium on Vistulian Stratigraphy as one of ref sites of Vistulian in Poland (Kozarski, 1980; 1981a). Morphology, stratigraphy, and chronology of site in morainic upland in hinterland of marginal zone of Poznań phase was studied by Borówko-Dłuzakowa (1967; 1979), Stankowski (1979; 1980), Stankowska and Stankowski (1979), Tobolski (1979a,b) and Pazdur and Walanus (1979b), whose results were summarized briefly by Pazdur, Stankowski, and Tobolski (1980). Upper fossil flora level, 5 to 10cm thick, Maliniec II, found at depth 431 to 439cm in fine-grained sands with load deformation structures which penetrate to half thickness of fluvioglacial series, is composed of strongly sanded silts, lime silts with abundant plant detritus, and sanded and silted mossy peats with dominant macrofossils of *Drepanocladus resolvens* (main component), *Calliergon turgescens*, and *C trifarium* (id. by K Karczmarz) and represents accumulation in shallow, very wet tundra peat bog with calciphile mosses growing together with sedges in area of dwarf shrub willows. Lower fossil flora level, Maliniec I, at depth 678 to 753cm, is composed of 8 layers different in lithology and containing genetically different organogenic sediments with no traces of post-sedimentary deformations and representing complete sequence of climatic and vegetational changes of cool interstadial period with 4 distinct pollen zones (Tobolski, 1979a,b). Maliniec I level could be provisionally assigned to Moershoofd interstadial complex (Tobolski, 1979a; Kozarski, 1980; 1981a).

Gd-646. Maliniec II; KM-XII-B **$22,230 \pm 480$**

From Maliniec II level, layer XII, depth 431 to 439cm, alkali-soluble fraction.

Gd-645. Maliniec II; KM-XII-C **$22,050 \pm 450$**

Same sample, insoluble fraction.

+ 2300

Gd-647. Maliniec I; KM-VII-A **$34,000$**

- 1800

From top of Maliniec I level, layer VIII, depth 678 to 690cm, pollen zone D with *Betula nana*, *Empetrum*, and *Arctostaphylos uva-ursi*. Comment (MFP): acid-soluble fraction dated, undersized, diluted.

Gd-1077. Maliniec I; KM-VIII-B **$>42,900$**

Same sample, alkali-soluble fraction.

Gd-1076. Maliniec I; KM-VIII-C **$>42,500$**

Same sample, insoluble fraction.

Gd-668. Maliniec I; KM-I-A **$25,000 \pm 1000$**

From base of Maliniec I level, layer I, depth 740 to 748cm, pollen zone A with *Salix* and *Equisetum*. Comment (MFP): acid-soluble fraction dated, undersized, diluted.

Gd-1105. Maliniec I; KM-I-B **$>41,200$**

Same sample, alkali-soluble fraction.

Gd-671. Maliniec I; KM-I-C **$>40,700$**

Same sample, insoluble fraction.

General Comment (MFP): for general discussion of significance of Konin-Maliniec site for chronology and strat of Vistulian in Poland, see Kozarski (1980; 1981a). Organic horizons with ^{14}C dates close to those from Maliniec II level from profiles in German Democratic Republic and European part of USSR were described by Cepek (1965) and Krasnov (1978), respectively.

Gd-769. Młodocin 5 **1150 ± 90**

Black earth soil in form of thin layer at depth 60cm overlain by series of fluvial and lacustrine fine-grained sands in Noteć R valley in Młodocin village ($52^\circ 53' 29''$ N, $17^\circ 52'$ E), 750m NW of Wolickie Lake. Coll Aug 1977 by M Sinkiewicz; subm 1979 by Władysław Niewiarowski, Inst Geog, Mikołaj Kopernik Univ, Toruń.

Tuliszków series

Organic deposits of Vistulian age from vicinity of Tuliszków, Turek Uplands, Great Poland Lowland, S of Warta Pradolina. Coll Oct 1979 and subm by Alicja Mańkowska, Geol Inst, Warsaw. Dated to establish reach-line of Vistulian Glaciation in Konin-Tuliszków-Turek area and determine age and genesis of Złote Góry Mts glacial forms (Kawecki, 1969; Mańkowska, 1975).

Gd-785. Wymysłów s1193 **$31,200 \pm 800$**

Peaty mud in form of layer in fossil river channel at depth 1.9 to 2.1m in Wymysłów village ($52^\circ 04'$ N, $18^\circ 20'$ E) near Tuliszków, overlain by fossil soil, fine- and medium-grained sands, and boulder clay.

Gd-786. Zdzary s466**>27,000**

Peat from series of paludal sediments, depth 2.1 to 2.2m in Zdzary village ($52^{\circ} 08' N$, $18^{\circ} 16' E$), overlain by fine sands and boulder clay.
Comment: undersized, diluted.

Gd-1251. Zdzary s743**+ 3000****41,400****- 2200**

Peaty mud from series of lacustrine sediments at depth 1.6 to 1.7m overlain by clayey mud, silt, and boulder clay in Zdzary village ($52^{\circ} 08' 30'' N$, $18^{\circ} 16' E$). *Comment (AM):* pollen analysis by Zofia Borówko-Dłuzakowa indicates pollen spectrum representing cold phitophases of pine-birch forest of Pleistocene period.

General Comment (MFP): all dates of this series obtained on alkali-soluble fraction of sample, precipitated by adding appropriate amount of HCl.

Przyłubie series

Peat from thin fossil organic layer in substratum of flood plain of Wisła R valley, Toruń-Bydgoszcz Basin, ca 17km W of Toruń, near Przyłubie village ($53^{\circ} 03' 24'' N$, $18^{\circ} 19' 20'' E$); coll and subm 1979 by Anna Tomczak, Inst Geog, Mikołaj Kopernik Univ, Toruń (Galon, 1934; Niewiarowski & Tomczak, 1969; Tomczak, 1981).

Gd-1248. Przyłubie A**1870 ± 70**

From top part of organic layer, depth 60cm, at contact with overlying layer of clayey silt with admixture of organic matter.

Gd-1249. Przyłubie B**5480 ± 60**

From base of organic layer at contact with underlying fine-grained sands, depth 1.85m. *Comment (AT):* pollen analysis of base layer of peat indicates Atlantic or younger age.

Podgórz series

Peat from series of organic sediments filling former bed of Wisła R, flood plain on left bank of Wisła R valley, Toruń-Bydgoszcz Basin, Podgórz, Toruń ($52^{\circ} 59' 30'' N$, $18^{\circ} 35' E$); coll Nov 1979 and subm by Anna Tomczak.

Gd-1250. Podgórz A**710 ± 45**

Well-decomposed black peat from top of uppermost peat layer at contact with overlying series of clayey silt with plant fragments, depth 90 to 95cm.

Gd-784. Podgórz B**3190 ± 280**

Wood peat with undecomposed plant fragments from base of lowermost peat layer, depth 4.8 to 4.85m, at contact with underlying sands. *Comment (AT):* pollen analysis of lowermost peat layer by Bozena Noryśkiewicz indicate Atlantic or younger age.

Rzeczkowo series

Organic sediments filling up former bed of Wisła R, flood plain on right bank of Wisła R valley, Toruń-Bydgoszcz Basin, ca 17km W of Toruń, near Rzeczkowo village ($53^{\circ} 07' 06''$ N, $18^{\circ} 21' 27''$ E). Coll Oct 1979 by Anna Tomczak and Władysław Niewiarowski; subm 1980 by Anna Tomczak.

Gd-1246. Rzeczkowo A **940 ± 60**

Well-decomposed peat from top of peat layer, depth 30 to 35cm, overlain by clayey silt.

Gd-780. Rzeczkowo B **5580 ± 120**

Black peat from base of peat layer, depth 1.4 to 1.47m.

Gd-781. Rzeczkowo C **9760 ± 260**

Gray clayey gyttja from base of gyttja layer, depth 2.15 to 2.2m. *Comment:* undersized, diluted.

General Comment (AT): introductory pollen analysis of profile by Bozena Noryśkiewicz yielded following age estimates: A—younger than Atlantic, B—Atlantic, C—Younger Dryas. Samples from Przyłubie, Podgórz, and Rzeczkowo profiles dated for studies of changes of Wisła R valley between Toruń and Solec Kujawski during Late Glacial and Holocene, as part of IGCP 158A Project.

Warsaw-Płock Wisła River valley series

Peat and gyttja from base of organogenic layers filling paleochannels on Wisła R flood plain, Wisła R valley between Warsaw Depression and Płock Depression (Wiśniewski, 1981). Coll Sept 1979 and subm 1980 by Edward Wiśniewski, Inst Geog & Spatial Org, Dept Geomorphol & Hydrol of Lowland, Polish Acad Sci, Toruń.

Gd-741. Wąsosz 1:sol **5200 ± 150**

Gyttja from depth 3.95 to 4.25m, in Wąsosz near Piaski village ($52^{\circ} 26' N$, $19^{\circ} 49' E$), 6km NE of Gąbin, alkali-soluble fraction.

Gd-740. Wąsosz 1:ins **5250 ± 250**

Same sample, insoluble fraction. *Comment:* undersized, diluted.

Gd-733. Wykowo 5:sol **3360 ± 150**

Gyttja from depth 3.7 to 3.95m from Wykowo village ($52^{\circ} 29' N$, $19^{\circ} 52' E$), 13km SE of Płock, alkali-soluble fraction.

Gd-735. Wykowo 5:ins **3850 ± 100**

Same sample, insoluble fraction.

Gd-1199. Wola Ładowska 2:sol **8450 ± 100**

Peat from depth 1.52 to 1.72m in Wola Ładowska village ($52^{\circ} 02' N$, $20^{\circ} 02' E$), 2.5km N of Iłowo, alkali-soluble fraction.

Gd-1200. Wola Ładowska 2:ins **7520 ± 100**

Same sample, insoluble fraction.

Gd-1207. Podgórze 4:sol **5610 ± 70**

Peat from depth 0.8 to 1m in Podgórze village ($52^{\circ} 04' N$, $20^{\circ} 03' E$),
10km W of Wyszogród, alkali-soluble fraction.

Gd-1204. Podgórze 4:ins **6290 ± 60**

Same sample, insoluble fraction.

Gd-732. Nowa Wieś 3:sol **9540 ± 240**

Gyttja from depth 3.95 to 4.2m in Nowa Wieś ($52^{\circ} 21' N$, $20^{\circ} 08' E$),
6.5km ENE of Iłowo, alkali-soluble fraction. *Comment:* undersized,
diluted.

Gd-1202. Nowa Wieś 3:ins **9660 ± 110**

Same sample, insoluble fraction.

Rembertów series

Sandy peat from layer at depth 1.5 to 1.55m overlain by dune sands
on uppermost flood terrace of Wisła R, Mazowiecka Plain, 1km NE of
Rembertów ($51^{\circ} 16' N$, $21^{\circ} 13' E$). Coll Oct 1978 by Zdzisława Sarnacka;
subm 1980 by M D Baraniecka, Geol Inst, Warsaw.

Gd-1236. Rembertów:sol **410 ± 45**

Alkali-soluble fraction.

Gd-770. Rembertów:ins **440 ± 60**

Same sample, insoluble fraction.

Kobylarnia series

Peat from layer 1.8m thick underlain by silt, Wisła R valley, Kobylarnia
village, N part of Sandomierz Basin ($50^{\circ} 33' N$, $21^{\circ} 48' E$). Coll
Sept 1979 and subm 1980 by Elzbieta Mycielska-Dowgiałło, Dept Geog &
Regional Studies, Warsaw Univ (Mycielska-Dowgiałło, 1972).

Gd-1297. Kobylarnia 1a **6890 ± 70**

From depth 70cm.

Gd-1298. Kobylarnia 1b **8570 ± 100**

From depth 1.3 to 1.5m.

Gd-1299. Kobylarnia 1c **11,640 ± 100**

From lowermost layer, depth 1.8m.

Gd-808. Dobromierz S-79-73 **1150 ± 110**

Peat from thin layer between two series of gravels in alluvial fan,
depth 1m, Dobromierz ($50^{\circ} 50' N$, $16^{\circ} 20' E$), Sudety Mts. Coll Nov 1979
and subm 1980 by Alfred Jahn, Geog Inst, Wrocław Univ, Wrocław.

Gd-194. Smogornia **4900 ± 100**

Peat from base of peat-bog Smogornia in Karkonosze, Sudety Mts, S Poland ($50^{\circ} 41' N$, $15^{\circ} 40' E$), alt 1400m asl, depth 2m, in contact with thin layer of sandy clay overlying 45cm thick layer of granitic rubble (regolith). Coll and subm 1972 by Alfred Jahn. *Comment* (AJ): base of other peat bog in Jakuszyce, same region, was dated to $10,075 \pm 210$ BP (Dumanowski, Jahn, & Szczepankiewicz, 1962).

Jasiołka River valley series

Peat from base of peat layer at contact with lacustrine chalk, overlain by clays and ablation till in Jasiołka R valley, Jasło-Sanok Depression. Coll Aug 1979 and subm by Antoni Wójcik, Geol Inst, Carpathian Branch, Cracow.

Gd-1233. Roztoki 1:ins **9870 ± 110**

From fossil lake sediments on terrace of Jasiołka R valley, profile XVI, depth 3.95 to 4m, in Roztoki village ($49^{\circ} 44' 44'' N$, $21^{\circ} 32' 50'' E$) (Szafer & Jaroń, 1935; Szafer, 1948). *Comment*: insoluble fraction dated.

Gd-1232. Roztoki 1: 1st sol **9850 ± 110**

Same sample, 1st alkali-soluble extract dated.

Gd-766. Roztoki 1: 2nd sol **9920 ± 100**

Same sample, 2nd alkali-soluble extract dated.

General Comment (AW): both early studies (Szafer & Jaroń, 1935; Szafer, 1948) as well as results of actual pollen analysis indicate cold period from end of Pleistocene.

Gd-1235. Roztoki 3 **$10,950 \pm 120$**

From fossil lake sediments on terrace of Jasiołka R valley, profile XV (Szafer, 1948) in Roztoki village ($49^{\circ} 44' 36'' N$, $21^{\circ} 32' 52'' E$), depth 2.66 to 2.7m. *Comment* (AW): preliminary results of pollen analysis indicate cold period; expected age: Late Glacial.

Gd-767. Tarnowiec 5 **7930 ± 110**

From contact of peat layer with uppermost layer of lacustrine chalk, valley floor of left tributary of Jasiołka R, ca 1.1km from Tarnowiec ($49^{\circ} 44' 25'' N$, $21^{\circ} 35' 55'' E$), depth 1.4 to 1.45m. *Comment* (AW): expected age based on provisional pollen analysis by Krystyna Harmata—beginning of Holocene/end of Pleistocene.

Gd-1243. Jedlicze 8 **$36,700 \pm 2100$**

Wood (*Picea* or *Larix* sp), id. by Kazimierz Szczepanek, in lowest part of peat layer 0.8m thick at depth 4.3 to 4.4m in Jedlicze, Brzozowa St ($49^{\circ} 43' 30'' N$, $21^{\circ} 39' 30'' E$). Coll Oct 1979 by Kazimierz Szczepanek and Antoni Wójcik; subm by Antoni Wójcik. Peat layer overlain by clay, fossil soil, and 2 layers of till separated by thin layer of fossil soil. *Comment* (AW): results of provisional pollen analysis of peat layer indicate Brorup Interstadial.

San River series

Wood from San R valley, Sandomierz Basin. Coll May 1980 and subm by Andrzej Szumański, Dept Geol, Warsaw Univ, Warsaw. Dated for studies of evolution of San R valley during Late Glacial and Holocene in connection with IGCP 158A Project (Szumański, 1972; 1981).

Gd-838. Lezajsk S1**7080 ± 90**

From layer of gray-black silt with plant remains, depth 3m, overlain by gray-yellow loamy mud and yellow loess-like mud in buried oxbow of San R, rendzina terrace B. Coll from exposure at channel bank near Lezajsk ($50^{\circ} 30' N$, $22^{\circ} 40' E$).

Gd-1300. Rudnik S2**660 ± 50**

From top of series of gray fine-grained sands diagonally laminated with single trunks, overlain by gray-brown loamy mud and yellow loess-like mud, terrace 3 of San R valley at Rudnik ($50^{\circ} 50' N$, $22^{\circ} 22' E$), depth 3m.

Grabiny-Latoszyn series

Wood, mostly from individual oak trunks, and charcoal from alluvial sequence in gravel pit between Grabiny and Latoszyn, left bank of Wiślka R valley ($50^{\circ} 03' N$, $21^{\circ} 22' E$), ca 4km SW of Dębica, Carpathian Foreland. Gravel-pit Latoszyn, ca 1.5km long, belongs to series of ref sites in Wiślka R valley studied in IGCP 158A Project by Leszek Starkel and coworkers (Starkel, 1981d). S part of gravel-pit with profiles A and C is on terrace level IID, 189 to 190.5m asl (Mamakowa *et al*, 1981, p 64-67, fig 26), gently inclined to E and encompassing large mature paleochannel with cut meander neck resembling present channel, with outcrops revealing 4 parts of sediments of different age within exposed sequence, ca 7m, underlain by series of gravels ca 14m thick. N part of gravel pit with profile B is on terrace level IIA, ca 187m asl with sequence of sediments, 7 to 14m thick, consisting of top sandy alluvial loam, upper gravels, sands, and lower gravels. All samples but one coll 1978 and 1979 by Ewa Niedziałkowska, Leszek Starkel, and members of ^{14}C lab staff; subm by Leszek Starkel, Dept Geomorphol & Hydrol, Inst Geog, Polish Acad Sci, Cracow.

Gd-509. Grabiny-Latoszyn NII**2260 ± 120**

Oak wood, external fragment of trunk ca 1m diam, N part of gravel pit, provenience unknown. Coll and subm 1977 by Bolesław Kawalec, Rzeszów.

Gd-1011. Grabiny-Latoszyn 070878/1**2730 ± 70**

Wood, small trunk ca 3cm diam from silty organic detritus layer at depth ca 3.5m in profile B, N part of gravel-pit. Comment (MFP): pollen spectrum of silty detritus layer (Mamakowa *et al*, 1981, p 67, fig 27) reveals high frequency of *Abies*.

Gd-582. Grabiny-Latoszyn 180578/1 2420 ± 50

Oak trunk from gravels overlying sandy muds, profile B in N part of gravel pit, depth ca 2.5m.

Gd-581. Grabiny-Latoszyn 170878/4 5990 ± 80

Oak trunk, 50cm diam, from profile A1 in S part of gravel pit, from series of lower gravels at depth 5.3m. *Comment* (MFP): pollen analysis of clayey mud sample coll from environs of this trunk made by Kazimiera Mamakowa reveals *Abies* pollens, numerous culture indicators with different types of cereals (*Secale cereale*, *Triticum* type, *Hordeum* type, and high frequency of *Plantago lanceolata*) (Mamakowa *et al*, 1981, p 65, fig 27).

Gd-600. Grabiny-Latoszyn 170878/2 5950 ± 70

Oak trunk 50cm diam from profile A1bis, ca 5m E of profile A1, same series of lower gravels at depth ca 5m. *Comment* (MFP): sample of clayey mud from environs of this trunk reveals high frequency of *Abies* pollen.

Gd-580. Grabiny-Latoszyn 170878/1 5920 ± 60

Oak trunk 50cm diam from S part of gravel pit, same series of lower gravels at depth ca 5m, ca 25m N of samples Gd-581 and -600.

Gd-1009. Grabiny-Latoszyn 160878/3 960 ± 70

Oak trunk 60cm diam from series of upper gravels in S part of gravel pit at depth ca 2m, ca 10m E of sample Gd-600.

Gd-1012. Grabiny-Latoszyn 160878/1 890 ± 70

Oak trunk 25cm diam, same loc.

Gd-1014. Grabiny-Latoszyn 170878/LS 105 ± 45

Charcoal from series of unconsolidated covering deposits consisting of clayey, silty and sandy layers on E slope of terrace IID, S part of gravel pit near profile A4 at depth ca 1m. *Comment* (LS&MFP): top layer of this series was previously destroyed. Using Stuiver (1978) correction curve, calendar age of sample can be estimated as < 300 bp. This implies that migration of channel did not take place before 2nd half of 17th century AD. This agrees well with observations of Strzelecka (1958) from Wisłok R valley where old sinuous paleochannels were active to mid-18th century.

Gd-597. Grabiny-Latoszyn 101078/LS 7990 ± 110

Rounded piece of wood from same series near sample Gd-1014. *Comment* (LS): probably redeposited from right bank sites of early Holocene deposits in Podgrodzie area ca 4km upstream (Alexandrowicz *et al*, 1981; Mamakowa & Starkel, 1977).

Gd-1145. Grabiny-Latoszyn A-14 710 ± 80

Wood from tree-trunk with branches found in profile A-14, S part of gravel pit ca 120m N of profile A1, at base of upper series of sandy gravels, depth ca 2m.

Gd-707. Grabiny-Latoszyn A-15 2860 ± 150

Organic detritus from series of sands and silts of ob facies at top of gravel series, profile A-15 in S part of gravel pit, depth ca 3m. *Comment:* undersized, diluted.

Gd-1146. Grabiny-Latoszyn C-5 480 ± 60

Wood fragments from series of muds with organic detritus overlain by sands with some gravels, ca 1.5m below sandy base of paleochannel visible in relief, profile C-5 close to profile A-14, S part of gravel pit, depth 3.5m. *Comment (LS):* this paleochannel does not appear active on Mieg's map from AD 1780 (Mieg, 1779-1782).

Gd-1148. Grabiny-Latoszyn C-5a 1830 ± 70

Twigs from same layer, probably redeposited material.

Gd-583. Grabiny-Latoszyn 170878/4 5540 ± 80

Oak trunk 50cm diam from lower gravel series near profile A1 in S part of gravel-pit.

Gd-578. Grabiny-Latoszyn 080878/1 4540 ± 60

Oak trunk 55cm diam, S part of gravel pit. *Comment (MFP):* displaced, provenience unknown.

General Comment (LS): based on results of palynol investigations showing *Abies* pollens and numerous culture indicators, our previous view (Awsik *et al*, 1980) on age of gravels should be revised assuming that tree trunks of *Quercus* sp were reworked from lower bed at break of Sub-atlantic period. Only strongly consolidated gravels below 5.9m can be recognized as Atlantic.

*B. Norway***Gd-660. Elisebreen 1660 ± 130**

Tundra peat in form of layer 5cm thick underlain by brown loamy gravels and covered with fresh lodgement till, depth 60cm, in forefield of Elisebreen Glacier, Oscar II Land, NW Spitsbergen ($78^{\circ} 37' 45''$ N, $12^{\circ} 07'$ E). Coll July 1978 by Bozena Noryśkiewicz; subm by Władysław Niewiarowski. Dated to establish age of Elisebreen Glacier transgression (Niewiarowski, 1982).

Werenskiold series

Peat from surface of ice in marginal part of Werenskiold Glacier, SW Spitsbergen ($77^{\circ} 04' 55''$ N, $15^{\circ} 15' 20''$ E). Coll July 1979 by Henryk Chmal; subm 1980 by Alfred Jahn.

Gd-803. Werenskiold W-1 510 ± 50

Found 150m from ice front.

Gd-1267. Werenskiold W-2 750 ± 60

Found 200m from ice front.

Ytre Kjaes series

Peat from palsa hill in peat bog on coastal plain, Ytre Kjaes, Porsangen Fjord, Nord Norway ($70^{\circ} 30' N$, $25^{\circ} 30' E$). Coll June 1979 and subm 1980 by Alfred Jahn.

Gd-1266. Ytre Kjaes N-79-7 **970 ± 45**

From contact of unfrozen peat with permafrost, depth 30cm.

Gd-809. Ytre Kjaes N-79-6 **4760 ± 120**

From lowermost unfrozen peat layer 10cm thick at depth ca 1.2m.
Comment (MFP): for earlier dates from palsa and thufur-type hills and other permafrost structures, see R, 1980, v 22, p 63-64.

C. Mongolia

Gd-823. Tot 225/KR **$17,860 \pm 230$**

Remnants of rush plants with admixture of lake clay in continuous layer at depth 4.75m, overlain by layer, 2m thick, of lacustrine silt with plant admixture and lacustrine clay covered with stratified fluvial sands, in Tot-Nuurin Khot-nor Basin, 5km SW of Tot Lake, ca 20km S of small village Gallut ($46^{\circ} 40' N$, $100^{\circ} 5' E$), S Khangai Mts. Coll July 1975 and subm 1980 by Karol Rotnicki. Sampling point is in central part of tectonic basin filled with lacustrine sediments of great Pleistocene pluvial lake, alt ca 2000m asl. *Comment (KR):* upper layer of plant remnants yielded date $17,220 \pm 155$. For more detailed inf, see Klimek and Rotnicki (1978).

II. SOIL SAMPLES

Fractions of soil organic matter from fossil soil horizons in dune sands were dated to check suitability of simple pretreatment procedure, consisting of acid-alkali-acid treatment, for reliable dating of fossil soil horizons developed from dune sands. Pretreatment of soil samples, described by Pazdur (1982), includes enrichment of organic matter by sedimentation in distilled water, treatment with 2% HCl at $80^{\circ}C$ or $100^{\circ}C$ for 1 hr (or at room temp for 24 hr), evaporation of acid-soluble fraction and washing of acid-insoluble fraction, similar treatment with sodium hydroxide, precipitation of alkali-soluble fraction by adding appropriate amount of HCl, washing of alkali-insoluble residue, and final treatment of acid-insoluble, alkali-insoluble residue with 2% HCl at $80^{\circ}C$ for 30 min or with 1% HCl at room temp for ca 24 hr. Samples from soil horizons in Troszyn pretreated with 4% solutions of HCl and NaOH yielded amount of insoluble residue too small for accurate dating. For some soil samples age measurements were also made on total soil organic matter.

Troszyn series

Soil fraction samples from 3 fossil soil horizons in N part of parabolic dune, Troszyn, 7km E of Wolin, W Pomerania ($53^{\circ} 32' N$, $14^{\circ} 45' E$). Coll and subm Nov 1978 by Bolesław Nowaczyk and M F Pazdur.

Gd-1087. Troszyn 78/1: total **720 ± 50**

Gd-664. Troszyn 78/1: acid-sol **510 ± 110**

Gd-665. Troszyn 78/1: alkali-sol **1010 ± 60**

From upper soil horizon at depth ca 80cm. *Comment* (MFP): charcoal from this soil horizon yielded date 1580 ± 70 , Gd-546 (R, 1982, v 24, p 183).

Gd-657. Troszyn 78/2: total **2170 ± 70**

Gd-690. Troszyn 78/2: acid-sol **2020 ± 210**

Gd-1100. Troszyn 78/2: alkali-sol **2320 ± 50**

From middle soil horizon at depth 3.8m. *Comment* (MFP): 2 charcoal samples assoc with this horizon yielded dates, 2300 ± 170 and 2440 ± 60 , Gd-528 and -537 (R, 1982, v 24, p 183).

Gd-1088. Troszyn 78/3: total **3500 ± 60**

Gd-683. Troszyn 78/3: acid-sol **2790 ± 220**

Gd-666. Troszyn 78/3: alkali-sol **3600 ± 70**

From lowest soil horizon at depth 6.4m. *Comment* (MFP): charcoal from fire-layer at top of soil horizon yielded date, 3130 ± 70 , Gd-529 (R, 1982, v 24, p 183).

General Comment (MFP&BN): acid-soluble fractions of all samples did not produce enough CO₂ to fill counter to normal pressure. Dates on both total organic matter and alkali-soluble fraction agree well with corresponding dates on charcoal for middle and lowest soil horizons. For youngest soil level, rejuvenation due to rootlet penetration and infiltration of recent soluble organic matter is possible. More detailed discussion of dune structure and results of ¹⁴C dating is given by Karczewski and Nowaczyk (1978) and Nowaczyk and Pazdur (1982).

Pomorsko series

Soil and wood samples from 4 horizons of semihydromorphic and peaty soils in transverse dune close to glacial trough filled with lime gyttja and peat, Pomorsko ($52^{\circ} 05' 06''$ N, $15^{\circ} 31' 50''$ E), Cigacice-Krosno sec of Warsaw-Berlin Pradolina, Great Poland Lowland. Coll and subm Nov 1978 by Bolesław Nowaczyk and M F Pazdur. Geomorphol and geol investigations of dune and glacial trough were made by Nowaczyk (1974; 1976; 1979); pollen analysis of biogenic sediments by Nowaczyk and Tobolski (1979; 1981); malacol analysis by Alexandrowicz (1980) and Alexandrowicz and Nowaczyk (1982), and pedol studies by Kowalkowski (1977a,b). Numerous flint artifacts from top of dune were analyzed by Michał Kobusiewicz (Nowaczyk *et al*, 1982) indicating several phases of human occupation by Komorniki, Janisławice, Swiderian, and Lusatian cultures. Descriptions of organic horizons and content of organic matter are given by Kowalkowski (1977a, fig 3).

Gd-739. Pomorsko 78/1: acid-sol **1100 ± 210**

Gd-1064. Pomorsko 78/1: alkali-sol **710 ± 40**

Semihydromorphic muck soil developed postsedimentarily in upper part of younger eolian sands; coll from top of organic layer at depth 100 to 105cm. Organic matter content ca 4%.

Gd-642. Pomorsko 78/1H: acid-sol **1250 ± 70**

Gd-634. Pomorsko 78/1H: alkali-sol **1310 ± 60**

Same level, from base of soil layer at depth 105 to 115cm, organic matter content 2%.

Gd-636. Pomorsko 78/2: acid-sol **1850 ± 50**

Gd-1067. Pomorsko 78/2: alkali-sol **2440 ± 60**

Gd-1069. Pomorsko 78/2: res **2580 ± 60**

Semihydromorphic postsedimentary peaty soil in lower part of younger eolian sands, top of organic layer at depth ca 1.9m, organic matter content 5%.

Gd-637. Pomorsko 78/2H: acid-sol **1790 ± 100**

Gd-1068. Pomorsko 78/2H: alkali-sol **2270 ± 50**

Sample from top of same layer, ca 2m from previous sampling point.

Gd-641. Pomorsko 78/3H: acid-sol **2770 ± 60**

Gd-1070. Pomorsko 78/3H: alkali-sol **3070 ± 60**

Gd-648. Pomorsko 78/3H: res **2200 ± 70**

From semihydromorphic horizon of boggy soil with marked postsedimentary features, depth ca 2.5m, organic matter content ca 6%.

Gd-653. Pomorsko 78/4H: acid-sol **3650 ± 80**

Gd-1086. Pomorsko 78/4H: alkali-sol **4110 ± 60**

Gd-655. Pomorsko 78/4H: res **3480 ± 70**

From trisegmentary organic horizon 10cm thick, consisting of lower muck horizon formed postsedimentarily in stabilized boggy sands and upper synsedimentary muck horizon separated by thin peat layer, depth 2.8m, organic matter content from 10 to 20%.

Gd-1043. Pomorsko 78/5A **7090 ± 50**

Pinus and *Larix* cones from thin layer of eolian sands overlying layer of calcareous gyttja, 5m thick, underlying oldest soil, depth 3m.

Gd-643. Pomorsko 78/5B **10,200 ± 120**

Unid. wood fragments, same loc. Comment (BN&MFP): date of Gd-1043 determines end of accumulation of calcareous gyttja; date of Gd-643 on wood seems too old, probably wood fragments redeposited from older sediments. Oldest layer of calcareous gyttja from pine phase of Alleröd

dated palynologically by Kazimierz Tobolski (see Nowaczyk, 1976) has ^{14}C date, $11,380 \pm 275$, Gd-378 (R, 1979, v 21, p 166). Detailed discussion of validity of ^{14}C dates of soil fractions and comparison with results of other studies is given by Nowaczyk *et al* (1982).

Rabsztyn series

Soil and charcoal samples from fossil soil levels covered by dune sands in Rabsztyn, 4km NE of Olkus, NW part of Ojców Plateau ($50^\circ 18' 30''$ N, $19^\circ 36' 30''$ E). Coll and subm May 1979 by Bolesław Nowaczyk and Tadeusz Szczypek.

Gd-1206. Rabsztyn 79/1: alkali-sol	430 ± 50
Gd-736. Rabsztyn 79/1: res	260 ± 120
From uppermost soil horizon covered with series of well-stratified sands and recent soil; Site 1 in N part of dune, depth 0.6m.	
Gd-1118. Rabsztyn 79/2: total	1170 ± 60
Gd-680. Rabsztyn 79/2: alkali-sol	1360 ± 70
Gd-679. Rabsztyn 79/2: res	1320 ± 70
From middle soil horizon overlain by well-laminated sands, same loc, depth ca 2m.	
Gd-718. Rabsztyn 79/2W: charcoal	2640 ± 150
Fine charcoal from same soil horizon. <i>Comment:</i> undersized, diluted.	
Gd-1120. Rabsztyn 79/3: total	6460 ± 70
Gd-681. Rabsztyn 79/3: alkali-sol	6630 ± 80
From lowermost soil horizon between 2 series of unstratified sands, depth ca 3m, same loc.	
Gd-729. Rabsztyn 79/4: alkali-sol	800 ± 60
Gd-1197. Rabsztyn 79/4: res	750 ± 60
From soil horizon 20cm thick in central part of dune, Site 2, depth 2.5m.	
Gd-1122. Rabsztyn 79/5: total	1380 ± 60
Gd-1121. Rabsztyn 79/5: alkali-sol	1390 ± 60
Gd-1117. Rabsztyn 79/5: res	1480 ± 60
From soil horizon in S part of dune, Site 3, depth ca 2m.	
Gd-1116. Rabsztyn 79/6: total	840 ± 60
Gd-728. Rabsztyn 79/6: alkali-sol	900 ± 70
Gd-730. Rabsztyn 79/6: res	920 ± 130
From soil horizon in E part of dune, Site 4, depth 2m.	

Gd-1114. Rabsztyn 79/6W: charcoal **810 ± 60**

Large pieces of charcoal from same loc.

General Comment (MFP): in general, good agreement is seen between dates of alkali-soluble fraction and insoluble residue for all dated samples of soil humus; dates on total soil organic matter are in all cases younger than other corresponding dates from same sample. Detailed discussion of results of ^{14}C dating and their comparison with results of geomorphol, geol, and strat studies of dune is given by Nowaczyk, Pazdur, and Szczypek (1982); background inf on geomorphol and chronol of dune-forming processes in vicinity is given by Szczypek (1977; 1980).

Grodzewo series

Charcoal and sandy humus from fossil soil horizons in dune at Grodzewo ($52^\circ 34' \text{ N}$, $14^\circ 08' \text{ E}$), 25km W of Skwierzyna, Toruń-Eberswald Pradolina, W part of Gorzów Basin, Great Poland Lowland. Coll and subm Nov 1978 by Bolesław Nowaczyk and M F Pazdur.

Gd-632. Grodzewo 78/1W **2160 ± 60**

Fine charcoal from upper part of middle soil level, depth 1.6 to 1.73m.

Gd-629. Grodzewo 78/2W **2590 ± 60**

Fine charcoal from lower part of middle soil level, depth 1.86 to 2.03m.

Gd-779. Grodzewo 78/3H: alkali-sol **4120 ± 130** **Gd-783. Grodzewo 78/3H: res** **4000 ± 180**

Soil fractions from lowest soil horizon, depth 2.42 to 2.73m. *Comment*: undersized, diluted.

Budzyń series

Charcoal and sandy humus from fossil soil level in parabolic dune 5km W of Budzyń ($52^\circ 53' 30'' \text{ N}$, $16^\circ 56' 50'' \text{ E}$), Gniezno Uplands, Great Poland Lowland. Coll 1979 and subm 1980 by Bolesław Nowaczyk.

Gd-1208. Budzyń 17/79H: alkali-sol **970 ± 60** **Gd-1209. Budzyń 17/79H: res** **990 ± 50**

Soil fractions, depth 2m.

Gd-1201. Budzyń 17/79W **1120 ± 60**

Charcoal, same loc.

General Comment (BN): other dates on charcoal from soil horizons in dune are $11,400 \pm 320$ and 925 ± 125 , Gd-357 and -371 (R, 1979, v 21, p 166).

REFERENCES

Alexandrowicz, S W, 1980, Zespoły malakofauny w kredach jeziornych Ziemi Lubuskiej, Kreda jeziorna i gytje, konf naukowo-techn, materiały pozjazdowe: PTPNoZ, Oddział Ziemi Lubuskiej, Gorzów, p 24-32.

- Alexandrowicz, S W, Mamakowa, Kazimiera, Niedziałkowska, Ewa, and Starkel, Leszek, 1981, Locality Podgrodzie, in Starkel, Leszek, ed, The evolution of the Wiślka valley near Dębica during the Late Glacial and Holocene: *Folia Quaternaria*, v 53, p 38-55.
- Alexandrowicz, S W and Nowaczyk, Bolesław, 1982, Late-Glacial and Holocene lacustrine deposits at Pomorsko near Sulechów: *Quaestiones Geog*, v 9, p 000-000.
- Antczak, Barbara, 1981, Zabno. Morphology and deposits within the braided paleochannel pattern, in Kozarski, Stefan and Tobolski, Kazimierz, eds, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, p 26-28.
- Awsik, Romuald, Niedziałkowska, Ewa, Pazdur, Anna, Pazdur, M F, Starkel, Leszek, and Walanus, Adam, 1980, Preliminary results of studies on the age of the Holocene alluvia at the left bank of the Wiślka river near Dębica: *Studia geomorphologica Carpatho-Balcanica*, v 14, p 33-42.
- Berglund, B E, ed, 1979, Palaeohydrological changes in the temperature zone in the last 15,000 years. Subproject B. Lake and mire environments: Proj Guide I, Dept Quaternary Geol, Univ Lund, Sweden, 123 p.
- Bluszcz, Andrzej and Walanus, Adam, 1980, Zestaw do amplitudowej i czasowej analizy impulsów z zespołów licznikowych służących do pomiaru naturalnych aktywności ^{14}C : Informatyczne systemy pomiarowe o skupionej i rozłożonej inteligencji w systemie CAMAC, ZG SEP-ZZUJ Polon-IBJ, Warszawa, p 53-54.
- Borówko-Dłużakowa, Zofia, 1967, Badania paleobotaniczne osadów młodoplejstoceńskich (Brorup) w Koninie-Marantowie: Inst Geol, Prace, v 48, p 81-136.
- _____, 1979, Młodoplejstoceńskie i postglacialne stanowiska flor kopalnych w Koninie: *Kwartalnik Geol*, v 22, p 247-257.
- Cepék, A G, 1965, Geologische Ergebnisse der ersten Radiokarbondatierungen von Interstadien im Lausitzer Urstromtal: *Geologie*, v 14, p 625-657.
- Dumanowski, B, Jahn, A, and Szczepankiewicz, S, 1962, The Holocene of Lower Silesia in the light of the first radiocarbon dating: *Acad Pol Sci Bull*, v 10, no. 1.
- Galon, Rajmund, 1934, Dolina dolnej Wisły, jej kształt i rozwój na tle budowy dolnego Powiśla: Inst Geog, Prace, v 12/13.
- Håkansson, Soren, 1980, University of Lund radiocarbon dates XIII: Radiocarbon, v 22, p 1045-1063.
- Karczewski, Andrzej and Nowaczyk, Bolesław, 1978, Budowa geologiczna wydmy w Troszynie koło Wolina: *Pozn Tow Przyj Nauk sprawozdania*, no. 94, p 38-40.
- Kawecki, J, 1969, Geneza i wiek form krajobrazu między Tuliszkowem a Koninem, *Polskie Tow Geol*, 41 Zjazd, Przewodnik: Inst Geol, Warszawa, p 64-68.
- Klimek, Kazimierz and Rotnicki, Karol, 1978, Old shore-lines and deposits of pluvial lake in the Tot-Nuurin Khot-nor Basin: *Acad Pol Sci Bull, Ser Sci Terre*, v 25, no. 3-4, p 119-124.
- Kowalkowski, Alojzy, 1977a, A paleopedological investigations of dunes at Pomorsko: *Quaestiones Geog*, v 4, p 43-50.
- _____, 1977b, Dynamika rozwoju poznoplejstoceńskich i holocenowych gleb z piasków wydmowych w Pomorsku: *Roczniki Gleboznawcze*, v 28, no. 3-4, p 19-35.
- Kozarski, Stefan, 1980, An outline of Vistulian stratigraphy and chronology of the Great Poland Lowland: *Quaternary Studies in Poland*, v 2, p 21-35.
- _____, 1981a, Stratygrafia i chronologia Vistulianu Niziny Wielkopolskiej: PWN, Warszawa-Poznań, 54 p.
- _____, 1981b, River channel changes in the Warta valley to the south of Poznań, in Kozarski, Stefan and Tobolski, Kazimierz, eds, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, p 8-23.
- _____, 1981c, Czmoniec. Younger generation of paleomeanders, in Kozarski, Stefan and Tobolski, Kazimierz, eds, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, p 50-51.
- Kozarski, Stefan, Moczek, A A, Nowaczyk, Bolesław, and Tobolski, Kazimierz, 1982, Etapy i warunki rozwoju wydmy w Budzyniu w świetle analizy radiowęglowej, paleobotanicznej i pedologicznej: *Roczniki Gleboznawcze*, v 33, no. 2, p 159-174.
- Kozarski, Stefan, Nowaczyk, Bolesław, Rotnicki, Karol, and Tobolski, Kazimierz, 1969, The eolian phenomena in West-Central Poland with special reference to the chronology of phases of eolian activity: *Geog Polon*, v 17, p 231-248.
- Kozarski, Stefan and Rotnicki, Karol, 1977, Valley floors and changes of river channel patterns in the North Polish Plain during the Late Würm and Holocene: *Quaestiones Geog*, v 4, p 51-93.
- Kozarski, Stefan and Tobolski, Kazimierz, 1963, Wiek gleby kopalnej w wydmach w Pradolinie Noteći koło Czarnkowa: *Badania Fizjog nad Polską Zach*, v 11, p 213-229.

- Kozarski, S and Tobolski, K, 1968, Holoceński przeobrażenia wydm śródlądowych w Polsce w świetle badań geomorfologicznych i palynologicznych: *Folia Quaternaria*, v 29, p 127-134.
- _____, 1981, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, 120 p.
- Krasnov, I I, 1978, Stratigraphische Korrelation der Quartäralablagerungen im östlichen Gebiet der fennoskandische Vereisung: *Schriften geol Wiss*, v 9, p 69-79.
- Mamakowa, Kazimiera, Niedziałkowska, Ewa, Pazdur, M F, and Starkel, Leszek, 1981, Locality Grabiny-Latoszyn, in Starkel, Leszek, ed, The evolution of the Wiśłoka valley near Dębica during the Late Glacial and Holocene: *Folia Quaternaria*, v 53, p 64-68.
- Mamakowa, Kazimiera and Starkel, Leszek, 1977, Stratigraphy of the Eo- and Mesoholocene alluvia in Podgrodzie upon Wiśłoka river: *Studia geomorphol Carpatho-Balcanica*, v 11, p 101-110.
- Mańska, Alicja, 1975, Wybrane zagadnienia czwartorzędu rejonu Konina: *Kwart Geol*, v 19, p 464-465.
- Mieg, 1779-1782, Karte des Königreichs Galizien und Lodomerien (1:28,000): Wien.
- Mościcki, Włodzimierz, Pazdur, Anna, Pazdur, M F, and Zastawny, Andrzej, 1978, Gliwice radiocarbon dates IV: Radiocarbon, v 20, p 405-415.
- Mościcki, Włodzimierz and Zastawny, Andrzej, 1977, New proportional counter assembly in Gliwice ^{14}C laboratory, in Povinec, Pavel and Usačev, S, eds, Low-radioactivity measurements and applications, Internat'l Conf, Proc: Bratislava, Slov Pedagog Naklad, p 91-92.
- Mycielska-Dowgiążo, Elżbieta, 1972, Rozwój doliny środkowej Wisły w holocenie w świetle badań z okolic Tarnobrzega: *Przegląd geog*, v 44, p 73-83.
- Niewiarowski, Władysław, 1982, Morphology of the forefield of the Aavatsmark Glacier (Oscar II Land, Northwest Spitsbergen) and phases of its formation: *Acta Univ Nicolai Copernici, ser geog*, v 16, p 000-000.
- Niewiarowski, Władysław and Tomczak, Anna, 1969, Morfologia i rozwój rzezby obszaru miasta Torunia i jego okolic: Univ Mikołaja Kopernika, zesz nauk, geog, v 6.
- Nowaczyk, Bolesław, 1974, Geneza i rozwój wydm poprzecznej w Pomorsku, Rozwój den dolinnych, symp krajowe, przewodnik wycieczki: Poznań, Univ Adama Mickiewicza, p 40-43.
- _____, 1976, Geneza i rozwój wydm śródlądowych w zachodniej części Pradoliny Warszawsko-Berlińskiej w świetle badań struktury, uziarnienia i stratygrafii budujących je osadów: Pozn Tow Przyj Nauk, Kom geog-geol, prace, v 16, p 1-108.
- _____, 1979, Perspektywy powiększenia zasobów kredy jeziornej i gytji wapiennej na obszarze pomiędzy Pomorskiem a Kijami, Kreda jeziorna i gytje, konf naukowo-techni, materiały pojazdowe: PTPN0Z, Oddział Ziemi Lubuskiej, Gorzów, p 86-92.
- Nowaczyk, Bolesław, Pazdur, Anna, Pazdur, Mieczysław F, and Awsik, Romuald, 1982, Stratygrafia i warunki rozwoju wydm w Pomorsku koło Sulechowa w świetle nowych badań: Badania Fizjogr nad Polską Zach.
- Nowaczyk, Bolesław and Pazdur, M F, 1982, Próba datowania metoda ^{14}C gleb kopalnych z wydmą w Troszynie koto Wolina: *Roczniki Gleboznawcze*, v 33, no. 2, p 145-158.
- Nowaczyk, Bolesław, Pazdur, M F, and Szczępek, Tadeusz, 1982, Wiek przekształceń eolicznych wydm w północno-zachodniej części Płaskowyżu Ojcowskiego: *Geog Studia et Dissertationes*, v 6, p 34-49.
- Nowaczyk, Bolesław and Tobolski, Kazimierz, 1979, Geneza i wiek rynien glacjalnych i wypełniających je osadów w Wilczej i Pomorsku, Kreda jeziorna i gytje, konf naukowo-techni, materiały pojazdowe: PTPN0Z, Oddział Ziemi Lubuskiej, Gorzów, p 79-85.
- _____, 1981, W sprawie późnoglacialnych osadów wapiennych akumulowanych w środowisku wodnym: Badania Fizjogr nad Polską Zach, v 33, p 000-000.
- Okuniewska, Iwona and Tobolski, Kazimierz, 1981, Paleobotanical investigations of deposits from paleomeander of the younger generation at Czmoniec, in Kozarski, Stefan and Tobolski, Kazimierz, eds, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, p 52-56.
- Olsson, I U, 1979, Radiometric dating, in Berglund, B E, ed, Palaeohydrological changes in the temperate zone during the last 15,000 years. Subproject B. Lake and mire environments: Project Guide I, Dept Quaternary Geol, Univ Lund, Sweden, p 1-38.
- Pazdur, Anna, Awsik, Romuald, Bluszcz, Andrzej, Pazdur, M F, Walanus, Adam, and Zastawny, Andrzej, 1982a, Gliwice radiocarbon dates VII: Radiocarbon, v 24, p 171-181.
- Pazdur, Anna, Pazdur, M F, and Zastawny, Andrzej, 1979, Gliwice radiocarbon dates V: Radiocarbon, v 21, p 165-170.

- Pazdur, M F, 1982, Badanie dokładności datowania metoda ^{14}C północnołęściańskich i holocenickich osadów organogenicznych: Politechnika Śląska, Zeszyty nauk, ser mat fiz, No. 49, 77 p.
- Pazdur, M F, Awsik, Romuald, Bluszcz, Andrzej, Hałaś, Stanisław, Pazdur, Anna, Walanus, Adam, and Zastawny, Andrzej, 1979, Preliminary results of the study of isotopic fractionation during chemical purification of carbon dioxide for radiocarbon dating: Radiochem Radioanal Letters, v 39, p 157-168.
- Pazdur, M F, Awsik, Romuald, Bluszcz, Andrzej, Pazdur, Anna, Walanus, Adam, and Zastawny, Andrzej, 1982b, Gliwice radiocarbon dates VIII: Radiocarbon, v 24, p 182-193.
- Pazdur, M F, Pazdur, Anna, and Zastawny, Andrzej, 1980, Gliwice radiocarbon dates VI: Radiocarbon, v 22, p 61-67.
- Pazdur, M F, Stankowski, Wojciech, and Tobolski, Kazimierz, 1980, Litologiczna i stratygraficzna charakterystyka profilu z kopalnymi utworami organogenicznymi w Małiniu koło Konina: Badania Fizjog nad Polską Zach, v 33, p 79-88.
- Pazdur, M F and Walanus, Adam, 1979a, Statistical analysis of data and age calculations in Gliwice Radiocarbon Laboratory: Muz Archeol Etnog Łodzi, Prace Materiały, v 26, p 283-289.
- _____, 1979b, The Konin-Maliniec site—age assessment by radiocarbon method, in Mojski, J E, ed, Vistulian stratigraphy, Symposium, Guide-book of excursion: Warszawa, Wydawnictwa Geol, p 4.5-4.6.
- Rotnicki, Karol, 1966, Rzeźba Wzgórz Ostrzeszowskich jako rezultat rozwoju stoku podczas Würm: Pozn Tow Przyj Nauk, Kom geog geol, prace, v 5, pt 2, 260 p.
- Sheppard, J C, Ali, Syead Y, and Mehringer, P J Jr, 1979, Radiocarbon dating of organic components of sediments and peats, in Berger, Rainer, and Suess, H E, eds, Radiocarbon dating, Internatl radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 284-305.
- Stankowska, Anna and Stankowski, Wojciech, 1979, The Vistulian till covering stagnant water sediments with organic sediments, in Mojski, J E, ed, Vistulian stratigraphy, Symposium, Guide-book of excursion: Warszawa, Wydawnictwa Geol, p 4.2.
- Stankowski, Wojciech, 1979, The outline of geological structure and geomorphology of the Konin surroundings, in Mojski, J E, ed, Vistulian stratigraphy, Symposium, Guide-book of excursion: Warszawa, Wydawnictwa Geol, p 4.1.
- _____, 1980, The structural heterogeneity of Würm flow tills and melt-out tills exemplified by sets at Maliniec and Kleczew near Konin, in Stankowski, Wojciech, ed, Tills and glaciogene deposits: Poznań, Adam Mickiewicz Univ, p 67-69.
- Starkeł, Leszek, 1977, Paleogeografia holocenu: Warszawa, PWN, 362 p.
- _____, 1981a, Paleohydrologiczne zmiany w strefie umiarkowanej w ostatnich 15 000 lat—problem nr 158 Międzynarodowego Programu Korelacji Geologicznych (IGCP) i udział Polski w realizacji tego problemu: Przegląd geog, v 53, p 105-120.
- _____, 1981b, Report on research on the Vistula valley evolution during the last 15,000 years, in Paleohydrology of the temperate zone, Symposium, abs of papers: Poznań, Adam Mickiewicz Univ, p 51-52.
- _____, 1981c, Progress in the knowledge of the fluvial history in the temperate zone during last 15,000 years, in Paleohydrology of the temperate zone, Symposium, abs of papers: Poznań, Adam Mickiewicz Univ, p 50.
- _____, ed, 1981d, The evolution of the Wiślaka valley near Dębica during the Late Glacial and Holocene: Folia Quaternaria, v 53, 91 p.
- Strzelecka, Barbara, 1958, Historyczna dokumentacja niektórych młodszych zmian hydrograficznych na brzegu Karpat: Czas geog, v 29, p 455-472.
- Stuiver, Minze, 1978, Radiocarbon time scale tested against magnetic and other dating methods: Nature, v 273, p 271-274.
- Stuiver, Minze and Polach, H A, 1977, Discussion: Reporting of ^{14}C data: Radiocarbon, v 19, p 355-363.
- Szafer, Władysław, 1948, Późny glacjal w Roztokach pod Jasłem: Starunia, no. 26.
- Szafer, Władysław and Jaroń, B, 1935, Plejstoceńskie jezioro pod Jasłem: Starunia, no. 8.
- Szczypek, Tadeusz, 1977, Utwory i procesy eoliczne w północnej części Wyżyny Śląskiej: Uniwersytet Śląski, prace nauk, no. 190, 1-116.
- _____, 1980, Piaski eoliczne okolic Siewierza: Geog Studia et Dissert, v 3, p 44-68.
- Szumański, Andrzej, 1972, Changes in the development of the lower San channel pattern in the Late Pleistocene and Holocene, in Palaeogeographical changes of valley floors in the Holocene, Symposium, Guide-book of excursion: Polish Acad Sci, Warsaw, p 55-68.

- Szumański, Andrzej, 1981, The evolution of Lower San river during the Late Glacial and Holocene, *in* Paleohydrology of the temperate zone, Symposium, abs of papers: Poznań, Adam Mickiewicz Univ, p 81-82.
- Tobolski, Kazimierz, 1979a, Fossil flora in the Maliniec site, *in* Mojski, J E, ed, Vistulian stratigraphy, Symposium, Guide-book of excursion: Warszawa, Wydawnictwa Geol, p 4.3-4.5.
- 1979b, Biostratigraphy of the Late Pleistocene (Neo-Pleistocene) in the Konin neighbourhood, *in* Mojski, J E, ed, Vistulian stratigraphy, Symposium, Guide-book of excursion: Warszawa, Wydawnictwa Geol, p 4.6-4.9.
- 1981, Preliminary results of macrofossil investigations in the paleomeander of the younger generation at Czmoniec, *in* Kozarski, Stefan and Tobolski, Kazimierz, eds, Paleohydrology of the temperate zone, Symposium, Guide-book of excursions: Poznań, Adam Mickiewicz Univ, p 57.
- Tomczak, Anna, 1981, Evolution of the Vistula river valley between Toruń and Solec Kuj during the Late Glacial and Holocene, *in* Paleohydrology of the temperate zone, Symposium, abs of papers: Adam Mickiewicz Univ, Poznań, p 83-85.
- Walanus, Adam and Pazdur, M F, 1980, Age reporting of very old samples: Radiocarbon, v 22, p 1021-1027.
- Wiśniewski, Edward, 1981, The geomorphological evolution of the Vistula valley between Włocławek and Ciechocinek during the last 15,000 years, *in* Paleohydrology of the temperate zone, Symposium, abs of papers: Poznań, Adam Mickiewicz Univ, p 86-87.
- Zaborowska, Krystyna, 1977, Analiza flory okrzemek osadów dennych Jeziora Gardno: Studia materiały oceanol, no. 19, p 304-306.
- Zachowicz, Joanna, 1977, Analiza palinologiczna osadów Jeziora Gardno: Studia materiały oceanol, no. 19, p 299-303.