



Strong Logarithmic Sobolev Inequalities for Log-Subharmonic Functions

Piotr Graczyk, Todd Kemp, and Jean-Jacques Loeb

Abstract. We prove an intrinsic equivalence between strong hypercontractivity and a strong logarithmic Sobolev inequality for the cone of logarithmically subharmonic (LSH) functions. We introduce a new large class of measures, Euclidean regular and exponential type, in addition to all compactly-supported measures, for which this equivalence holds. We prove a Sobolev density theorem through LSH functions and use it to prove the equivalence of strong hypercontractivity and the strong logarithmic Sobolev inequality for such log-subharmonic functions.

1 Introduction

In this paper we study strong versions of logarithmic Sobolev inequalities (sLSI) and strong hypercontractivity (sHC) in the real spaces \mathbb{R}^n and for logarithmically subharmonic (LSH) functions, continuing our research published in [13] and solving the conjecture on the equivalence between sHC and sLSI formulated in [13, Remark 5.11]. The main difficulty to overcome, as already noted by Gross and Grothaus in [17], was efficient approximating of (logarithmically) subharmonic functions.

If μ is a probability measure, the *entropy* functional Ent_μ relative to μ , defined on all sufficiently integrable positive test functions g , is

$$\text{Ent}_\mu(g) = \int g \ln\left(\frac{g}{\|g\|_1}\right) d\mu$$

where $\|g\|_1 = \|g\|_{L^1(\mu)}$. (When $\|g\|_1 = 1$, so that g is a probability density, this gives the relative entropy of the density g to the measure μ .) The *logarithmic Sobolev inequality* is an energy-entropy functional inequality: a measure μ on \mathbb{R}^n (or more generally on a Riemannian manifold) satisfies a log Sobolev inequality if, for some constant $c > 0$ and for all sufficiently smooth positive test functions f ,

$$\text{(LSI)} \quad \text{Ent}_\mu(f^2) \leq c \int |\nabla f|^2 d\mu.$$

Making the substitution $g = f^2$ gives the equivalent form $\text{Ent}_\mu(g) \leq \frac{c}{4} \int |\nabla g|^2 / g d\mu$, the integral on the right defining the *Fisher information* of g relative to μ . In this form, the inequality was first discovered for the standard normal law μ on \mathbb{R} by Stam in [30]. It was rediscovered and named by Gross [15], who proved it for standard Gaussian measures on \mathbb{R}^n with sharp constant $c = 2$. Over the past four decades, it has become an enormously powerful tool making fundamental contributions to geometry and

Received by the editors November 18, 2013; revised June 25, 2014.

Published electronically September 15, 2015.

AMS subject classification: 47D06.

Keywords: logarithmic Sobolev inequalities.

global analysis [2–6, 8–10, 22, 24, 27], statistical physics [19, 32–34], mixing times of Markov chains [7, 11, 18], concentration of measure and optimal transport [23, 25, 31], random matrix theory [1, 26, 35], and many others.

Gross discovered the log Sobolev inequality through his work in constructive quantum field theory, particularly relating to Nelson's hypercontractivity estimates [29]. In fact, Gross showed in [15] that the log Sobolev inequality (LSI) is equivalent to hypercontractivity. Later, in [20, 21], Janson discovered a stronger form of hypercontractivity that holds for *holomorphic* test functions.

Theorem 1.1 (Janson [20]) *If μ is the standard Gaussian measure on \mathbb{C}^n , and $0 < p \leq q < \infty$, then for all holomorphic functions $f \in L^p(\mathbb{C}^n, \mu)$, we have $\|f(e^{-t} \cdot)\|_q \leq \|f\|_p$ for $t \geq \frac{1}{2} \ln \frac{q}{p}$. For $t < \frac{1}{2} \ln \frac{q}{p}$, the dilated function $f(e^{-t} \cdot)$ is not in $L^q(\mathbb{C}^n, \mu)$ in general.*

Remark 1.2 Nelson's hypercontractivity estimates ([29]) involve the semigroup e^{-tA_μ} , where A_μ is the Dirichlet form operator for the measure μ :

$$\int |\nabla f|^2 d\mu = \int \bar{f} A_\mu f d\mu.$$

If $d\mu = \rho dx$ has a smooth density ρ , integration by parts shows that $A_\mu = -\Delta - (\nabla \rho / \rho) \cdot \nabla$, and so when applied to holomorphic (hence harmonic) functions, e^{-tA_μ} is the flow of the vector field $\nabla \rho / \rho$. For the standard Gaussian measure, this is just the coordinate vector field x , the infinitesimal generator of dilations $Ef(x) = x \cdot \nabla f(x)$, also known as the *Euler operator*. The perspective of this paper, like its predecessor [13], is that the strong hypercontractivity theorem is essentially about the dilation semigroup $f \mapsto f(e^{-t} \cdot)$, independent of the underlying measure.

Janson's strong hypercontractivity differs from Nelson's hypercontractivity in two important ways: first, the time-to-contraction is smaller, $\frac{1}{2} \ln \frac{q}{p}$ as opposed to the larger Nelson time $\frac{1}{2} \ln \frac{q-1}{p-1}$, and second, the theorem applies even in the regime $0 < p, q < 1$ where the L^p "norms" are badly-behaved. Nevertheless, in [16], Gross showed that Janson's theorem is also a consequence of the same log Sobolev inequality (LSI); moreover, he generalized this implication considerably to complex manifolds (equipped with sufficiently nice measures). The reverse implication, however, was not established. The proof requires (LSI) to hold for non-holomorphic functions (in particular of the form $|f|^{p/2}$). We refer the reader to [13] for an extensive list of recent literature on strong hypercontractivity in the holomorphic category and related ideas (notably reverse hypercontractivity) in the subharmonic category.

The aim of this paper is to prove an *intrinsic* equivalence of strong hypercontractivity and a log Sobolev inequality. The starting point is a generalization of Theorem 1.1 beyond the holomorphic category. A function on \mathbb{R}^n is *log-subharmonic* (LSH) if $\ln |f|$ is subharmonic; holomorphic functions are prime examples. In [13], we proved that Theorem 1.1 holds in the larger class LSH, for the Gaussian measure, and several others. We also established a weak connection to a *strong log Sobolev inequality*.

Definition 1.3 A measure μ on \mathbb{R}^n satisfies a *strong logarithmic Sobolev inequality* if there is a constant $c > 0$ so that, for non-negative $g \in \text{LSH}$ sufficiently smooth and integrable,

$$(s\text{LSI}) \quad \text{Ent}_\mu(g) \leq \frac{c}{2} \int E g d\mu.$$

Inequality (sLSI) could be written equivalently in the form $\text{Ent}_\mu(f^2) \leq c \int f E f d\mu$; we will use it in L^1 -form throughout. In [13], we showed the strong log Sobolev inequality holds for the standard Gaussian measure on \mathbb{R}^n , with constant $c = 1$ (half the constant from (LSI)), and conjectured that (sLSI) is equivalent in greater generality to the following form of Janson's strong hypercontractivity.

Definition 1.4 A measure μ on \mathbb{R}^n satisfies the property of *strong hypercontractivity* if there is a constant $c > 0$ so that, for $0 < p \leq q < \infty$ and for every $f \in L^p(\mu) \cap \text{LSH}$, we have

$$(s\text{HC}) \quad \|f(r \cdot)\|_{L^q(\mu)} \leq \|f\|_{L^p(\mu)} \quad \text{if } 0 < r \leq (p/q)^{c/2}.$$

Remark 1.5 The statement in Definition 1.4 is given in multiplicative notation rather than additive, with $r = e^{-t}$ scaling the variable. It would appear more convenient to use the constant c instead of $\frac{c}{2}$ in (sLSI) and (sHC). We choose to normalize with $\frac{c}{2}$ for historical reasons. Gross's equivalence of the log Sobolev inequality and Nelson's hypercontractivity equates c in (LSI) to $\frac{c}{2}$ scaling the time to contraction.

Notation 1.6 For a function f on \mathbb{R}^n and $r \in [0, 1]$, f_r denotes the function $f_r(x) = f(rx)$.

1.1 Main Results

In [13], we showed that (sHC) implies (sLSI) in the special case that the measure μ is compactly supported. Our first result is the converse.

Theorem 1.7 Let μ be a compactly supported measure on \mathbb{R}^n . Suppose that μ satisfies (sLSI) for all sufficiently smooth functions $g \in \text{LSH}(\mathbb{R}^n)$. Then μ satisfies (sHC) for all functions $f \in \text{LSH}(\mathbb{R}^n)$.

Remark 1.8 We emphasize here that the domains in the equivalence consist of log-subharmonic functions a priori defined on all of \mathbb{R}^n , not just on the support of μ . Indeed, the dilation semigroup is not well defined if this is not satisfied. In fact, it is not hard to see that this result extends to log-subharmonic functions defined on any star-shaped open region containing the support of μ .

Theorem 1.7 and its converse have non-trivial applications. For example, [13, Proposition 4.2] implies that (sLSI) holds true for any compactly supported symmetric measure on \mathbb{R} , with constant $c \leq 2$. Nevertheless, it excludes the standard players in log Sobolev inequalities, most notably Gaussian measures. In [13, Theorem 5.8], we proved directly that (sLSI) holds true for the standard Gaussian measure on \mathbb{R}^n , with

best constant $c = 1$. This was proved directly from (LSI), and relied heavily on the precise form of the Gaussian measure; a direct connection to strong hypercontractivity (also proved for the Gaussian measure in [13, Theorem 3.2]) was not provided. That connection, for a wide class of measures, is the present goal.

The technicalities involved in establishing the equivalence of (sLSI) and (sHC) are challenging because of the rigidity of the class LSH. Standard cut-off approximations needed to use integrability arguments in the proof are unavailable for subharmonic functions. To amend this, we use a fundamentally different approximation technique: the *dilated convolution* introduced in [12,17] and developed in Section 2.1. In [12], the authors provided a local condition on the density of μ under which this operation is bounded on $L^p(\mu)$ (amounting to a bound on the Jacobian derivative of the translation and dilation). Here we present alternative conditions, which require little in terms of the local behavior of the measure (they are essentially growth conditions near infinity) and achieve the same effect.

Definition 1.9 Let $p > 0$ and let μ be a positive measure on \mathbb{R}^n with density ρ . Say that μ (or ρ) is *Euclidean exponential type p* if $\rho(x) > 0$ for all x and if the following two conditions hold:

$$(1.1) \quad \sup_x \sup_{|y| \leq s} |x|^p \frac{\rho(ax + y)}{\rho(x)} < \infty \quad \text{for any } a > 1, s \geq 0$$

$$(1.2) \quad \sup_x \sup_{1 < a < 1 + \epsilon} \frac{\rho(ax)}{\rho(x)} < \infty \quad \text{for some } \epsilon > 0.$$

If μ is Euclidean exponential type 0, we say it is *Euclidean regular*.

The terminology derives from the fact that conditions (1.1) and (1.2) insist that the Euclidean group acts on ρ in a controlled manner; exponential type refers to the growth condition involving $|x|^p$ (indeed, for $p > 0$ the measure must have tails that decay faster than any polynomial to be Euclidean exponential type p). For any probability measure μ with strictly positive density ρ , denote for $a \geq 1$ and $p, s \geq 0$,

$$(1.3) \quad C_\mu^p(a, s) = C_\rho^p(a, s) \equiv \sup_x \sup_{|y| \leq s} |x|^p \frac{\rho(ax + y)}{\rho(x)}.$$

Then the condition that μ is Euclidean exponential type p is precisely that $C_\mu^p(a, s) < \infty$ for each $a > 1$ and $s \geq 0$, and $C_\mu^0(a, 0)$ is uniformly bounded for a close to 1. It is clear from the definition that $C_\mu^p(a, s)$ is an increasing function of s . Moreover, if μ is Euclidean exponential type q , then it is Euclidean exponential type p for any $p < q$. For convenience, we will often write C_μ for C_μ^0 .

Example 1.10 On \mathbb{R} , the densities $(1 + x^2)^{-\alpha}$ for $\alpha > \frac{1}{2}$ are Euclidean regular. On \mathbb{R}^n the densities $e^{-c|x|^a}$ with $a, c > 0$ are Euclidean exponential type p for all $p > 0$.

More examples and properties that prove the Euclidean regular measures form a rich class are given in Appendix A. In order to justify the implication (sLSI) \implies (sHC) in the fully general (non-compactly-supported) case, we will insist on a further regularity property of the density ρ .

Definition 1.11 We say that μ (or ρ) is *exponentially sub-additive* if for all $x \in \mathbb{R}^n$ there exists $A \in \mathbb{R}^n$ such that for all $t \in \mathbb{R}^n$

$$\rho(x)\rho(t) \leq e^{\langle A, t \rangle} \rho(0)\rho(x+t).$$

Let $\alpha > 0$. We say that ρ is α -*subhomogeneous* if for all $c > 0$,

$$(1.4) \quad \rho(cy) \leq \rho(y)^{c^\alpha}.$$

Example 1.12 It is easy to check that the densities $e^{-c|x|^a}$ are exponentially sub-additive for $0 < a \leq 1$ (then $A = 0$) and for $a = 2$ (then $A = -2x$). All the densities $e^{-c|x|^a}$, $a, c > 0$ are evidently a -subhomogeneous, with equality in (1.4).

The purpose of introducing these classes of measures (notably Euclidean regularity) at present is its utility in proving a density theorem for an appropriate class of Sobolev-type spaces that we now proceed to define. These spaces, denoted $L_E^p(\mu)$, are exactly the domains of functions for which the strong log Sobolev inequality makes sense.

Definition 1.13 Let μ be a measure on \mathbb{R}^n , and let $p > 0$. Define the Sobolev space $L_E^p(\mu)$ to consist of those continuously differentiable (C^1) functions $f \in L^p(\mu)$ for which $Ef \in L^p(\mu)$. It is a normed space in the norm $f \mapsto \|f\|_p + \|Ef\|_p$.

Remark 1.14 The space $L_E^p(\mu)$ is generally not complete; its completion (for smooth μ) is the space of weakly differentiable functions $f \in L^p(\mu)$ satisfying $Ef \in L^p(\mu)$. To be precise, $Ef(x) = \sum_{j=1}^n x_j u_j(x)$, where u_j is the function (posited to exist) satisfying

$$-\int \partial_j \phi f \, dx = \int \phi u_j \, dx$$

for any $\phi \in C_c^\infty(\mathbb{R}^n)$, where dx denotes Lebesgue measure. We will not have occasion to need the completeness of this space in its norm; it will be more convenient to have Sobolev functions that are already at least C^1 , and so we restrict the definition thus.

Standard techniques, involving approximation by C_c^∞ functions, show that L_E^p is dense in L^p for reasonable measures. However, our goals here involve approximation of log-subharmonic functions, and the usual cut-off approximations fail to preserve subharmonicity. An alternative approach is to use a convolution approximate identity procedure, as is readily available for Lebesgue measure. The problem is that, for a given bump function ϕ , the operation $f \mapsto f * \phi$ is typically unbounded on $L^p(\mu)$ when μ is not Lebesgue measure. Indeed, for L^p of Gaussian measure, even the translation $f \mapsto f(\cdot + y)$ is unbounded if $y \neq 0$. The problem is that the convolution can shift mass in from near infinity. One might hope to dilate this extra mass back out near infinity, to preserve p -integrability; thus, we consider the dilated convolution $f \mapsto (f * \phi)_r$. Section 2.1 shows that this operation behaves well in L^p spaces of Euclidean regular measures; it also preserves the cone LSH.

The main technical theorem of this paper is the following smoothing procedure for LSH functions, *i.e.*, Sobolev density theorem, which is of its own independent interest.

Theorem 1.15 Let $p \in (0, \infty)$, and let μ be a Euclidean exponential type p probability measure on \mathbb{R}^n . Then the cone $C^\infty \cap \text{LSH} \cap L_E^p(\mu)$ is dense in the cone $\text{LSH} \cap L^p(\mu)$. More precisely, let $f \in \text{LSH} \cap L^p(\mu)$. Then there exists a sequence of functions $f_n \in C^\infty \cap \text{LSH} \cap L_E^p(\mu)$ that converges to f in $L^p(\mu)$.

Using Theorem 1.15, we will prove the equivalence of (sLSI) and (sHC), the first implication in a nominally weaker form that we now explain.

Definition 1.16 Let μ be a probability measure on \mathbb{R}^n , and let $0 < p < q < \infty$. Denote by $\text{LSH}_E^{p < q}(\mu)$ the closure of $L_E^q(\mu) \cap \text{LSH}$ in $L_E^p(\mu) \cap \text{LSH}$, with respect to the Sobolev norm $f \mapsto \|f\|_p + \|Ef\|_p$ of the space $L_E^p(\mu)$. Define the space

$$\text{LSH}_E^{p <}(\mu) = \overline{\bigcup_{q > p} \text{LSH}_E^{p < q}(\mu)}$$

with the closure being taken in $L_E^p(\mu)$.

For any probability measure, there is a common dense subspace (L^∞) for all the full L^q -spaces, $q > 0$, and so the closure of L^q in L^p is all of L^p for $p < q$; the standard proof uses cut-offs that do not respect subharmonicity, and indeed, there are no non-constant bounded subharmonic functions. In [16], Gross showed that, under certain conditions on a measure μ on a complex manifold (in terms of its Dirichlet form operator d^*d), in the presence of a full log Sobolev inequality (LSI), there is a common dense subspace for all holomorphic L^q spaces of μ . In the present context of logarithmically-subharmonic functions, no such technology is known, and we will content ourselves with the spaces $\text{LSH}_E^{p <}(\mu)$. We will consider the nature of these spaces in a future publication. A natural conjecture is that, for sufficiently nice measures μ , $\text{LSH}_E^{p <}(\mu) = \text{LSH} \cap L_E^p(\mu)$.

This brings us to our main theorem: the equivalence of (sLSI) and (sHC) for logarithmically subharmonic functions. Since slightly different hypotheses on the involved measures are required for the two directions of the equivalence, we state them separately. Moreover, because of some delicate issues with the (LSI) \implies (sHC) implication, we give two versions: one that requires the same conditions as the reverse implication but gives a slightly weaker form of strong hypercontractivity (b), and one that proves full strong hypercontractivity for exponentially subadditive and subhomogeneous measures (a).

Theorem 1.17 Let μ be an $O(n)$ -invariant probability measure on \mathbb{R}^n .

- (i) (a) Let μ be Euclidean exponential type p for all $p > 1$, exponentially subadditive, and α -subhomogeneous for some $\alpha > 0$. If μ satisfies the strong log Sobolev inequality (sLSI) for all functions in $\text{LSH} \cap L_E^1(\mu)$, with constant $c = \frac{2}{\alpha}$, then μ satisfies strong hypercontractivity (sHC): for $0 < p \leq q < \infty$ and $f \in \text{LSH} \cap L^p(\mu)$, $\|f_r\|_q \leq \|f\|_p$ for $0 < r \leq (p/q)^{c/2}$.

- (b) If μ is of Euclidean exponential type p for all $p > 1$ and (sLSI) holds for all functions in $\cup_{q>1} \text{LSH} \cap L_E^q(\mu)$, then μ satisfies partial strong hypercontractivity on each space $\text{LSH} \cap L^{q_0}(\mu)$, $q_0 > 1$, i.e., the inequality

$$\|f_r\|_{q(r)} \leq \|f\|_1, \quad q(r) = r^{-2/c}$$

from Proposition 1.19 holds for all functions $f \in \text{LSH} \cap L^{q_0}(\mu)$ and $r \in [1/q_0^{2/c}, 1]$.

- (ii) If μ is Euclidean exponential type p for some $p > 1$, and if μ satisfies (sHC) in the above sense, then μ satisfies the strong log Sobolev inequality (sLSI):

$$\text{Ent}_\mu(g) \leq \frac{c}{2} \int E g \, d\mu$$

for all $g \in \text{LSH}_E^{1<}$.

Remark 1.18 (i) The global assumption of rotational invariance in Theorem 1.17 is actually quite natural in this situation. The functional $g \mapsto \int E g \, d\mu$ on the right-hand-side of our strong log Sobolev inequality is not generally positive, since the operator E is not generally self-adjoint in $L^2(\mu)$; however, when μ is rotationally invariant, this functional is positive on the cone LSH, as pointed out in [13, Proposition 5.1].

(ii) In Theorem 1.17(i) we state the implication (sLSI) \implies (sHC) assuming the strong log Sobolev inequality (sLSI) holds for all functions in $\text{LSH} \cap L_E^1(\mu)$, which is the natural domain for which this inequality makes sense. In fact, our proof below actually shows the implication supposing (sLSI) holds on the nominally smaller space $\text{LSH}_E^{1<}$, and then the domains for (sLSI) are the same in both parts of Theorem 1.17.

We emphasize that Theorem 1.17 is *intrinsic*. While the two directions of the theorem require slightly different assumptions on the applicable measures, the implications between (sLSI) and (sHC) both stay within the cone LSH of log-subharmonic functions. This is the main benefit of extending Janson’s strong hypercontractivity theorem from holomorphic functions to this larger class, and restricting the log-Sobolev inequality to it: here, the two are precisely equivalent.

1.2 Alternative Formulation of sHC

The following equivalent characterization of strong hypercontractivity will be useful in what follows.

Proposition 1.19 Fix $c > 0$ and let $q(r)$ denote the function $q(r) = r^{-2/c}$. A measure μ satisfies strong hypercontractivity (sHC) if and only if for each function $f \in L^1(\mu) \cap \text{LSH}$,

$$\|f_r\|_{q(r)} \leq \|f\|_1 \quad \text{and} \quad \|f_r\|_1 \leq \|f\|_1, \quad \text{for } r \in (0, 1].$$

For the proof it is useful to note that the class LSH is closed under $f \mapsto f^p$ for any $p > 0$.

Proof First, suppose (sHC) holds with constant c . The case $p = q = 1$ yields $\|f_r\|_1 \leq \|f\|_1$ for $0 < r \leq (p/q)^{c/2} = 1$. More generally, by (sHC), $\|f_r\|_q \leq \|f\|_1$ whenever $0 < r \leq (1/q)^{c/2}$; i.e., whenever $q \leq r^{-2/c} = q(r)$. In particular, it follows that $\|f_r\|_{q(r)} \leq \|f\|_1$ as claimed.

Conversely, suppose the above conditions hold true. Fix $q \geq p > 0$ and let $f \in L^p(\mu) \cap \text{LSH}$. Then $f^p \in L^1(\mu) \cap \text{LSH}$, and so by assumption we have $\|(f^p)_r\|_{q(r)} \leq \|f^p\|_1$ for $0 < r \leq 1$. Since $(f^p)_r = (f_r)^p$, it follows immediately that $\|f_r\|_{p \cdot q(r)}^p \leq \|f\|_1^p$. Setting $q = p \cdot q(r)$ and solving for r , we have $r = r(p, q) \equiv (p/q)^{c/2}$, and so we have proved the equality case of (sHC). Finally, suppose that $r' \leq r(p, q) = (p/q)^{c/2}$; then there is $s \in (0, 1]$ so that $r' = s \cdot r(p, q)$. Dilations form a multiplicative semigroup, so $f_{r'} = (f_{r(p,q)})_s$. We have just proved that $f_{r(p,q)} \in L^q$, and hence $(f_{r(p,q)})^q$ is in $L^1(\mu)$. Therefore, by assumption, $\|[(f_{r(p,q)})^q]_s\|_1 \leq \|(f_{r(p,q)})^q\|_1$; unwinding this yields

$$\begin{aligned} \|f_{r'}\|_q^q &= \|(f_{r(p,q)})_s\|_q^q = \|[(f_{r(p,q)})^q]_s\|_1 = \|[(f_{r(p,q)})^q]\|_1 \leq \|(f_{r(p,q)})^q\|_1 \\ &= \|f_{r(p,q)}\|_q^q \leq \|f\|_p^q \end{aligned}$$

by the equality case, thus proving (sHC). ■

Remark 1.20 In fact, (sHC) implies the putatively stronger statement that $r \mapsto \|f_r\|_{q(r)}$ is non-decreasing on $[0, 1]$; however, the weaker form presented above is generally easier to work with.

1.3 Convolution Property

We will use the convolution operation to prove the Sobolev density theorem at the heart of this paper, as well as Theorem 1.7. We begin by showing that this operation preserves the cone LSH.

Lemma 1.21 *Let $f \in \text{LSH}$. Let $\phi \geq 0$ be a C^∞ test function. Then $f * \phi \in \text{LSH} \cap C^\infty$.*

Proof Since $f \in \text{LSH}$, $f \geq 0$ and $\ln f$ is subharmonic. In particular, $\ln f$ is upper semi-continuous and locally bounded above, and so the same holds for f . Thus f is locally bounded and measurable; thus, $f * \phi$ defines an $L^1_{\text{loc}} \cap C^\infty$ function. We must show that it is LSH.

Any subharmonic function is the decreasing limit of a sequence of C^∞ subharmonic functions; cf. [28, Appendix 1, Proposition 1.15]. Applying this to $\ln f$, there is a sequence $f_n \in \text{LSH} \cap C^\infty$ such that $f_n \downarrow f$. Let $g_n = f_n + \frac{1}{n}$, so g_n is strictly positive, and $g_n \downarrow f$. Since ϕ is ≥ 0 , it follows from the Monotone Convergence Theorem that $g_n * \phi \downarrow f * \phi$ pointwise.

Now, $(g_n * \phi)(x) = \int_{\mathbb{R}^n} g_n(x - \omega) \phi(\omega) d\omega$. Since translation and positive dilation preserve the cone LSH, the function $x \mapsto g_n(x - \omega) \phi(\omega)$ is continuous and LSH for each ω . Moreover, the function $\omega \mapsto g_n(x - \omega) \phi(\omega)$ is continuous and bounded. Finally, for small r , $\sup_{|t-x| \leq r} g_n(t - \omega) \phi(\omega) \leq \|\phi\|_\infty \sup_{|t| \leq |x| + r + s} g_n(t)$, where $s = \sup\{|\eta|: \eta \in \text{supp } \phi\}$, and this is bounded uniformly in ω . It follows from [13, Lemma 2.4] that $g_n * \phi$ is LSH. (The statement of that lemma apparently requires

the supremum to be uniform in x as well, but this is an overstatement; as the proof of the lemma clearly shows, only uniformity in ω is required).

Thus, $f * \phi$ is the decreasing limit of strictly positive LSH functions $g_n * \phi$. Applying the Monotone Convergence Theorem to integrals of $\ln(g_n * \phi)$ about spheres now shows that $\ln(f * \phi)$ is subharmonic, so $f * \phi \in \text{LSH}$ as claimed. ■

1.4 Compactly Supported Measures

This section is devoted to the proof of Theorem 1.7. It follows the now standard Gross proof of such equivalence: differentiating hypercontractivity at the critical time yields the log Sobolev inequality, and vice versa. The technical issues related to differentiating under the integral can be dealt with fairly easily in the case of a compactly supported measure; the remainder of this paper develops techniques for handling measures with non-compact support. The forward direction of the theorem, that (sHC) implies (sLSI) for compactly supported measures, is [13, Theorem 5.2], so we will only include the proof of the reverse direction here.

Proof of Theorem 1.7 By assumption, (sLSI) holds for sufficiently smooth and integrable functions; here we interpret that precisely to mean $\text{Ent}_\mu(g) \leq \frac{\epsilon}{2} \int E g d\mu$ for all $g \in C^1(\mathbb{R}^n)$ for which both sides are finite. Fix $f \in L^1(\mu) \cap \text{LSH} \cap C^1$. Utilizing Proposition 1.19, we must consider the function $\alpha(r) = \|f_r\|_{q(r)}$, where $q(r) = r^{-2/c}$. Let $\beta(r) = \alpha(r)^{q(r)} = \int f(rx)^{q(r)} \mu(dx)$ and set $\beta_x(r) = f(rx)^{q(r)}$ so that $\beta(r) = \int \beta_x(r) \mu(dx)$. Then

$$\frac{\partial}{\partial r} \ln \beta_x(r) = q'(r) \ln f(rx) + \frac{q(r)}{f(rx)} x \cdot \nabla f(rx).$$

Since $q'(r) = -\frac{2}{rc} q(r)$, and since $x \cdot \nabla f(rx) = \frac{1}{r} (Ef)_r(x) = \frac{1}{r} E(f_r)(x)$, we have

$$(1.5) \quad \frac{\partial}{\partial r} \beta_x(r) = -\frac{2}{rc} f_r(x)^{q(r)} \ln f_r(x)^{q(r)} + \frac{1}{r} q(r) f_r(x)^{q(r)-1} (Ef_r)(x).$$

Fix $0 < \epsilon < 1$. As f is C^1 , the function (of x) on the right-hand-side of (1.5) is uniformly bounded for $r \in (\epsilon, 1]$ and $x \in \text{supp } \mu$ (due to compactness). The Dominated Convergence Theorem thus allows differentiation under the integral, and so

$$(1.6) \quad \beta'(r) = \int \frac{\partial}{\partial r} \beta_x(r) \mu(dx).$$

Thus, since $\alpha(r) = \beta(r)^{1/q(r)}$ and $\beta(r) > 0$, it follows that α is C^1 on $(\epsilon, 1]$ and the chain rule yields

$$(1.7) \quad \alpha'(r) = \frac{\alpha(r)}{q(r)\beta(r)} \frac{2}{rc} \left[\beta(r) \ln \beta(r) + \frac{rc}{2} \beta'(r) \right].$$

From (1.5) and (1.6), the quantity in brackets is

$$\begin{aligned}
 (1.8) \quad & \int f_r^{q(r)} d\mu \cdot \ln \int f_r^{q(r)} d\mu + \frac{rc}{2} \int \left(-\frac{2}{rc} f_r^{q(r)} \ln f_r^{q(r)} + \frac{1}{r} q(r) f_r^{q(r)-1} E f_r \right) d\mu \\
 & = \int f_r^{q(r)} d\mu \cdot \ln \int f_r^{q(r)} d\mu - \int f_r^{q(r)} \ln f_r^{q(r)} d\mu + q(r) \frac{c}{2} \int f_r^{q(r)-1} E f_r d\mu \\
 & = -\text{Ent}_\mu(f_r^{q(r)}) + \frac{c}{2} \int E(f_r^{q(r)}) d\mu,
 \end{aligned}$$

where the equality in the last term follows from the chain rule.

Since $f \in C^1$, it is bounded on the compact set $\text{supp}\mu$, and so are all of its dilations f_r . Hence, both terms in (1.8) are finite, and so by the assumption of the theorem, this term is ≥ 0 . From (1.7), we therefore have $\alpha'(r) \geq 0$ for all $r > \epsilon$. Since this is true for each $\epsilon > 0$, it holds true for $r \in (0, 1]$. This verifies the first inequality in Proposition 1.19. For the second, we use precisely the same argument to justify differentiating under the integral to find

$$\frac{\partial}{\partial r} \|f_r\|_1 = \int \frac{\partial}{\partial r} f_r(x) \mu(dx) = \frac{1}{r} \int E f_r(x) \mu(dx) \geq \frac{2}{cr} \text{Ent}_\mu(f_r) \geq 0$$

by the assumption of (sLSI). This concludes the proof for $f \in C^1$.

Now, if $f \in L^1(\mu) \cap \text{LSH}$, we consider a smooth approximate identity sequence ϕ_k . The inequalities in Proposition 1.19 hold for $f * \phi_k$ by the first part of the proof and Lemma 1.21. Note by simple change of variables that $(f * \phi_k)_r = f_r * (r^n \phi_k)_r$, and that $(r^n \phi_k)_r$ is also an approximate identity sequence. The function f_r is LSH, so it is upper semi-continuous and consequently locally bounded. Thus, $f_r \in L^{q(r)}$ and $(f * \phi_k)_r$ converges to f_r in $L^{q(r)}$. This concludes the proof. ■

2 Density Results Through LSH Functions

This section is devoted to the approximation procedures we develop for smoothing LSH functions in L^p space of Euclidean regular measures, and in particular to the proof of Theorem 1.15. For a companion discussion of various closure properties of the class of Euclidean regular measures (testifying to the reasonably large size of this class), see the Appendix.

2.1 Continuity of the Dilated Convolution

One easy consequence of Definition 1.9 is that the operation $f \mapsto f_r$ is bounded on L^p .

Lemma 2.1 *Let μ be a Euclidean regular probability measure on \mathbb{R}^n , let $p > 0$, and let $r \in (0, 1)$. Then*

$$\|f_r\|_{L^p(\mu)} \leq r^{-n/p} C_\mu\left(\frac{1}{r}, 0\right)^{1/p} \|f\|_{L^p(\mu)}.$$

Proof We simply change variables $u = rx$ and use Definition 1.9:

$$\begin{aligned} \int |f_r(x)|^p \mu(dx) &= \int |f(rx)|^p \rho(x) dx = r^{-n} \int |f(u)|^p \rho(x/r) dx \\ &\leq r^{-n} C_\mu\left(\frac{1}{r}, 0\right) \int |f(u)|^p \rho(x) dx. \end{aligned} \quad \blacksquare$$

Remark 2.2 By condition (1.2) of Definition 1.9, the constant in Lemma 2.1 is uniformly bounded for $r \in (\epsilon, 1]$ for any $\epsilon > 0$; that is, there is a uniform (independent of r) constant C_ϵ so that, for $r \in (\epsilon, 1]$, $\|f_r\|_{L^p(\mu)} \leq C_\epsilon \|f\|_{L^p(\mu)}$.

The next proposition shows that, under the assumptions of Definition 1.9, the dilated convolution operation is indeed bounded on L^p . As usual, the conjugate exponent p' to $p \in [1, \infty)$ is defined by $\frac{1}{p} + \frac{1}{p'} = 1$.

Proposition 2.3 Let μ be a Euclidean regular probability measure on \mathbb{R}^n . Let $p \in [1, \infty)$, and let $\phi \in C_c^\infty$ be a test function. Then the dilated convolution operation $f \mapsto (f * \phi)_r$ is bounded on $L^p(\mu)$ for each $r \in (0, 1)$. Precisely, if $K = \text{supp } \phi$ and $s = \sup\{|w|; w \in K\}$, then

$$\|(f * \phi)_r\|_{L^p(\mu)} \leq r^{-n/p} C_\mu\left(\frac{1}{r}, \frac{s}{r}\right)^{1/p} \text{Vol}(K)^{1/p} \|\phi\|_{L^{p'}(K)} \|f\|_{L^p(\mu)},$$

where C_μ is the constant defined in (1.3).

Proof Denote by K the support of ϕ . By definition,

$$\|(f * \phi)_r\|_{L^p(\mu)}^p = \int_{\mathbb{R}^n} \left| \int_K f(rx - y)\phi(y) dy \right|^p \rho(x) dx.$$

We immediately estimate the internal integral using Hölder’s inequality:

$$\left| \int_K f(rx - y)\phi(y) dy \right|^p \leq \int_K |f(rx - y)|^p dy \cdot \|\phi\|_{L^{p'}(K)}^p.$$

This is finite, since the first integral is the p -th power of the L^p -norm of f restricted to the compact set $rx - K$. (Note that Euclidean regularity of μ implies that μ is equivalent to Lebesgue measure on compact sets.) Hence,

$$(2.1) \quad \|(f * \phi)_r\|_{L^p(\mu)}^p \leq \|\phi\|_{L^{p'}(K)}^p \int_{\mathbb{R}^n} \int_K |f(rx - y)|^p dy \rho(x) dx.$$

We apply Fubini’s theorem to the double integral, which is therefore equal to

$$(2.2) \quad \int_K \int_{\mathbb{R}^n} |f(rx - y)|^p \rho(x) dx dy = \int_K r^{-n} \int_{\mathbb{R}^n} |f(u)|^p \rho\left(\frac{u + y}{r}\right) du dy,$$

where we have made the change of variables $u = rx - y$ in the internal integral. By assumption, ρ is Euclidean regular, and so we have

$$(2.3) \quad \rho\left(\frac{1}{r}u + \frac{1}{r}y\right) \leq C_\mu\left(\frac{1}{r}, \frac{s}{r}\right) \rho(u), \quad y \in K.$$

where $s = \sup\{|w|; w \in K\}$. Substituting (2.3) into (2.2), we see that (2.1) yields

$$\|(f * \phi)_r\|_{L^p(\mu)}^p \leq r^{-n} C_\mu\left(\frac{1}{r}, \frac{s}{r}\right) \text{Vol}(K) \|\phi\|_{L^{p'}(K)}^p \int |f(u)|^p \rho(u) du.$$

This completes the proof. \blacksquare

Remark 2.4 The explicit constant in Proposition 2.3 appears to depend strongly on the support set of ϕ , but, in fact, it does not. Indeed, it is easy to check that the standard rescaling of a test function, $\phi^s(x) = s^{-n} \phi(x/s)$, which preserves total mass, also preserves the ϕ -dependent quantity above; to be precise, $\text{Vol}(\text{supp} \phi^s) \|\phi^s\|_{L^{p'}(\mathbb{R}^n)}^p$ does not vary with s . In addition, the constant $C_\mu(1/r, s/r)$ is well-behaved as s shrinks (indeed, it only decreases). It is for this reason that the proposition allows us to use the dilated convolution operation with an approximate identity sequence in what follows.

The use of Proposition 2.3 is that it allows us to approximate an L^p function by smoother L^p functions, along a path through LSH functions. To prove this, we first require the following continuity lemma.

Lemma 2.5 *Let μ be a Euclidean regular probability measure, and let $r \in (0, 1)$. Then for any $f \in L^p(\mu)$, the map $T_f: \mathbb{R}^n \rightarrow L^p(\mu)$ given by $[T_f(y)](x) = f_r(x - y)$ is continuous.*

Proof First note that, by the change of variables $u = rx - ry$,

$$\|T_f(y)\|_{L^p(\mu)}^p = \int |f(rx - ry)|^p \rho(x) dx = r^{-n} \int |f(u)|^p \rho\left(\frac{1}{r}u + y\right) du,$$

and the latter is bounded above by $r^{-n} C_\mu(\frac{1}{r}, |y|) \|f\|_{L^p(\mu)}^p$, showing that the range of T_f is truly in $L^p(\mu)$ for $y \in \mathbb{R}^n$. Now, fix $\epsilon > 0$ and let $\psi \in C_c(\mathbb{R}^n)$ be such that $\|f - \psi\|_{L^p(\mu)} < \epsilon$. Let $(y_k)_{k=1}^\infty$ be a sequence in \mathbb{R}^n with limit y_0 . Then

$$\begin{aligned} \|T_f(y_k) - T_f(y_0)\|_{L^p(\mu)} &\leq \|T_f(y_k) - T_\psi(y_k)\|_{L^p(\mu)} + \|T_\psi(y_k) - T_\psi(y_0)\|_{L^p(\mu)} \\ &\quad + \|T_\psi(y_0) - T_f(y_0)\|_{L^p(\mu)}. \end{aligned}$$

The first and last terms are simply $T_{\psi-f}(y_k)$ (with $k = 0$ for the last term), and so we have just proved that

$$\|T_{\psi-f}(y_k)\|_{L^p(\mu)} \leq r^{-n/p} C_\mu\left(\frac{1}{r}, |y_k|\right)^{1/p} \|\psi - f\|_{L^p(\mu)} < r^{-n/p} C_\mu\left(\frac{1}{r}, |y_k|\right)^{1/p} \epsilon.$$

Moreover, there is a constant s so that $|y_k| \leq s$ for all k , and since $C_\mu(a, s)$ is an increasing function of s , it follows that

$$\|T_f(y_k) - T_f(y_0)\|_{L^p(\mu)} \leq \|T_\psi(y_k) - T_\psi(y_0)\|_{L^p(\mu)} + 2r^{-n/p} C_\mu\left(\frac{1}{r}, s\right)^{1/p} \epsilon.$$

For each x , $(T_\psi(y_k)(x) - T_\psi(y_0)(x) = \psi(rx - ry_k) - \psi(rx - ry_0))$ converges to 0, since $ry_k \rightarrow ry_0$ and ψ is continuous. In addition, ψ_r is compactly supported and continuous, so it is uniformly bounded. Since μ is a probability measure, it now follows that $\|T_\psi(y_k) - T_\psi(y_0)\|_{L^p(\mu)} \rightarrow 0$ as $y_k \rightarrow y_0$, and the lemma follows by letting $\epsilon \downarrow 0$. ■

Corollary 2.6 *Let μ be a Euclidean regular probability measure, and let $r \in (0, 1)$. Then for any $f \in L^p(\mu)$, and ϕ_k an approximate identity sequence ($\phi_k \in C_c^\infty(\mathbb{R}^n)$ with $\int \phi_k(x) dx = 1$ and $\text{supp} \phi_k \downarrow \{0\}$),*

$$\|f_r * \phi_k - f_r\|_{L^p(\mu)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Proof Fix $\epsilon > 0$ and let $\psi \in C_c(\mathbb{R}^n)$ be such that $\|f - \psi\|_{L^p(\mu)} < \epsilon$. We estimate this in the following (standard) manner:

$$(2.4) \quad \|f_r * \phi_k - f_r\|_{L^p(\mu)} \leq \|(f_r - \psi_r) * \phi_k\|_{L^p(\mu)} + \|\psi_r * \phi_k - \psi_r\|_{L^p(\mu)} + \|\psi_r - f_r\|_{L^p(\mu)}.$$

By Lemma 2.1 applied to $f - \psi$, we have $\|f_r - \psi_r\|_{L^p(\mu)} \leq r^{-n/p} C_\mu(1/r, 0)^{1/p} \epsilon$, and from condition (1.2) of Definition 1.9, this is a uniformly bounded constant times ϵ for r away from 0. Also, note that

$$\begin{aligned} f_r * \phi_k(x) &= \int f_r(x - y)\phi_k(y) dy = \int f(rx - ry)\phi_k(y) dy \\ &= r^{-n} \int f(rx - u)\phi_k(u/r) du; \end{aligned}$$

that is to say, $f_r * \phi_k = r^{-n}(f * \tilde{\phi}_k)_r$, where we set $\tilde{\phi}_k = (\phi_k)_{1/r}$. Hence,

$$\begin{aligned} &\|(f - \psi)_r * \phi_k\|_{L^p(\mu)} \\ &= r^{-n} \|((f - \psi) * \tilde{\phi}_k)_r\|_{L^p(\mu)} \\ &\leq r^{-n} r^{-n/p} C_\mu\left(\frac{1}{r}, \frac{s_k}{r}\right)^{1/p} \text{Vol}(\text{supp}\tilde{\phi}_k)^{1/p} \|\tilde{\phi}_k\|_{L^{p'}(\mathbb{R}^n)} \cdot \|f - \psi\|_{L^p(\mu)} \end{aligned}$$

by Proposition 2.3, where $s_k = \sup\{|w|; w \in \text{supp}\phi_k\}$. Since $C_\mu(\frac{1}{r}, \frac{s}{r})$ is increasing in s , this constant is uniformly bounded as $k \rightarrow \infty$. In addition (cf. Remark 2.4), the product $\text{Vol}(\text{supp}\tilde{\phi}_k)^{1/p} \|\tilde{\phi}_k\|_{L^{p'}(\mathbb{R}^n)}$ can also be made constant with k (for example by choosing $\phi_k(x) = k^n \phi(kx)$ for some fixed unit mass C_c^∞ test-function ϕ). The result is that both the first and last terms in (2.4) are uniformly small as $k \rightarrow \infty$. Thus, we need only show that $\psi_r * \phi_k \rightarrow \psi_r$ in $L^p(\mu)$. The quantity in question is the p -th root of

$$(2.5) \quad \int \left| \int \psi_r(x - y)\phi_k(y) dy - \psi_r(x) \right|^p \mu(dx) = \int \left| \int_{K_k} [\psi_r(x - y) - \psi_r(x)] \phi_k(y) dy \right|^p \mu(dx),$$

where we have used the fact that ϕ_k is a probability density; here K_k denotes the support of ϕ_k . Since ψ_r is bounded, we may make the blunt estimate that the quantity in (2.5) is

$$\begin{aligned} &\leq \int \sup_{y \in K_k} |\psi_r(x - y) - \psi_r(x)|^p \left| \int_{K_k} \phi_k(y) dy \right|^p \mu(dx) \\ &= \int \sup_{y \in K_k} |\psi_r(x - y) - \psi_r(x)|^p \mu(dx). \end{aligned}$$

Since ψ_r is continuous and K_k is compact, there is a point $y_k \in K_k$ such that the supremum is achieved at y_k : $\sup_{y \in K_k} |\psi_r(x - y) - \psi_r(x)|^p = |\psi_r(x - y_k) - \psi_r(x)|^p$. As $k \rightarrow \infty$, the support K_k of ϕ_k shrinks to $\{0\}$, and so $y_k \rightarrow 0$. The function $|\psi_r(x - y_k) - \psi_r(x)|^p$ is continuous in x , and so converges to 0 pointwise as $y_k \rightarrow 0$. It therefore follows from the dominated convergence theorem that $\|\psi_r * \phi_k - \psi_r\|_{L^p(\mu)} \rightarrow 0$, completing the proof. ■

We will now use Proposition 2.3 and Corollary 2.6 to prove our main approximation theorem: that $L_E^p(\mu)$ is dense in $L^p(\mu)$ through log-subharmonic functions.

2.2 The Proof of Theorem 1.15

The basic idea of the proof is as follows: approximate a function $f \in \text{LSH} \cap L^p(\mu)$ by $(f * \phi)_r$, and let ϕ run through an approximate identity sequence and r tend to 1. We show that the dilated convolution $(f * \phi)_r$ is in $C^\infty \cap \text{LSH} \cap L^p_E(\mu)$, and that these may be used to approximate f in the L^p -sense.

Part 1: $(f * \phi)_r$ is in $C^\infty \cap \text{LSH} \cap L^p_E(\mu)$. Let $\phi \in C^\infty_c(\mathbb{R}^n)$ be a non-negative test function. Lemma 1.21 shows that $f * \phi$ is C^∞ and LSH. It is elementary to verify that the cone $C^\infty \cap \text{LSH}$ is invariant under dilations $g \mapsto g_r$; hence the dilated convolution $(f * \phi)_r$ is C^∞ and LSH. For fixed $r < 1$, Proposition 2.3 shows that $(f * \phi)_r$ is in $L^p(\mu)$, since $f \in L^p(\mu)$. We must now apply the differential operator E . Note that $(f * \phi)_r$ is C^∞ , and so

$$E[(f * \phi)_r](x) = x \cdot \nabla[(f * \phi)_r](x) = \int rx \cdot \nabla\phi(rx - y)f(y) dy.$$

Decomposing $rx = (rx - y) + y$, we break this up as two terms

$$(2.6) \quad E[(f * \phi)_r](x) = \int (rx - y) \cdot \nabla\phi(rx - y)f(y) dy + \int y \cdot \nabla\phi(rx - y)f(y) dy.$$

The first term is just $(f * E\phi)_r(x)$, and since $E\phi$ is also $C^\infty_c(\mathbb{R}^n)$, Proposition 2.3 bounds the L^p -norm of this term by the L^p -norm of f . Hence, it suffices to show that the second term in (2.6) defines an $L^p(\mu)$ -function of x . We now proceed analogously to the proof of Proposition 2.3. Changing variables $u = rx - y$ for fixed x in the internal integral and then using Hölder’s inequality

$$\begin{aligned} & \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} y \cdot \nabla\phi(rx - y)f(y) dy \right|^p \rho(x) dx \\ &= \int_{\mathbb{R}^n} \left| \int_K (rx - u) \cdot \nabla\phi(u)f(rx - u) du \right|^p \rho(x) dx \\ &\leq \int_{\mathbb{R}^n} \left(\int_K |rx - u|^p |f(rx - u)|^p du \right) \left(\int_K |\nabla\phi(u)|^{p'} dy \right)^{p/p'} \rho(x) dx, \end{aligned}$$

where $K = \text{supp}\phi$. Note that $\|\nabla\phi\|_{p'} < \infty$ is a constant independent of f . So we must consider the double integral, to which we apply Fubini’s theorem,

$$\int_{\mathbb{R}^n} \left(\int_K |rx - u|^p |f(rx - u)|^p du \right) \rho(x) dx = \int_K \left(\int_{\mathbb{R}^n} |rx - u|^p |f(rx - u)|^p \rho(x) dx \right) du.$$

Now we change variables $v = rx - u$ for fixed u in the internal integral, to achieve

$$(2.7) \quad \int_K \left(\int_{\mathbb{R}^n} |v|^p |f(v)|^p \rho\left(\frac{v + u}{r}\right) r^{-n} dv \right) du.$$

Finally, we utilize the assumption that ρ is exponential type p , and so there is a constant $C(p, r, K)$ so that $|v|^p \rho\left(\frac{v + u}{r}\right) \leq C(p, r, K)\rho(u)$ for $u \in K$. Hence the integral in (2.7) is bounded above by $C(p, r, K)r^{-n}\text{Vol}(K)$ times the finite norm $\int |f|^p d\mu$, which demonstrates that $E[(f * \phi)_r]$ is in $L^p(\mu)$.

Part 2: $(f * \phi)_r$ approximates f in $L^p(\mu)$. Let ϕ_k be an approximate identity sequence. Note by simple change of variables that $(f * \phi_k)_r = f_r * (r^n \phi_k)_r$, and that $(r^n \phi_k)_r$ is also an approximate identity sequence. Since $f_r \in L^p(\mu)$, by Lemma 2.1, it follows from Corollary 2.6 that $(f * \phi_k)_r \rightarrow f_r, k \rightarrow \infty$, in $L^p(\mu)$. We must now show

that $f_r \rightarrow f$ in $L^p(\mu)$ as $r \uparrow 1$. For this purpose, once again fix $\epsilon > 0$ and choose a $\psi \in C_c(\mathbb{R}^n)$ so that $\|f - \psi\|_{L^p(\mu)} < \epsilon$. Then

$$(2.8) \quad \|f - f_r\|_{L^p(\mu)} \leq \|f - \psi\|_{L^p(\mu)} + \|\psi - \psi_r\|_{L^p(\mu)} + \|\psi_r - f_r\|_{L^p(\mu)}.$$

The first term is $< \epsilon$, and changing variables the last term is

$$\begin{aligned} \|\psi_r - f_r\|_{L^p(\mu)}^p &= \int |\psi(rx) - f(rx)|^p \rho(x) \, dx = r^{-n} \int |\psi(u) - f(u)|^p \rho(u/r) \, du \\ &\leq r^{-n} C_\mu\left(\frac{1}{r}, 0\right) \int |\psi - f|^p \, d\mu. \end{aligned}$$

Here we have used the fact that μ is Euclidean regular. Note that, by condition (1.2) of Definition 1.9, the constant appearing here is uniformly bounded by, say, C , for $r \in (\frac{1}{2}, 1]$. Thence, the last term in (2.8) is bounded above by $C^{1/p}\epsilon$ and is also uniformly small. Finally, the middle term tends to 0 as $r \uparrow 1$, since $\psi_r \rightarrow \psi$ pointwise and the integrand is uniformly bounded. Letting ϵ tend to 0 completes the proof. ■

3 The Intrinsic Equivalence of (sLSI) and (sHC)

In this section, we prove Theorem 1.17, namely, if a measure μ is sufficiently Euclidean regular (satisfying the conditions of Definition 1.9), and if μ is invariant under rotations, then μ satisfies a strong log-Sobolev inequality precisely when it satisfies strong hypercontractivity. It will be useful to fix the following notation.

Notation 3.1 Let $c > 0$ be a fixed constant, let μ be a measure on \mathbb{R}^n , and let f be a function on \mathbb{R}^n .

(i) For $r \in (0, 1]$, let $q = q(r)$ denote the function $q(r) = r^{-2/c}$. Note that $q \in C^\infty(0, 1]$, is decreasing, and $q(1) = 1$.

(ii) Define a function $\alpha_{f,\mu}: (0, 1] \rightarrow [0, \infty)$ by

$$\alpha_{f,\mu}(r) \equiv \|f_r\|_{L^{q(r)}(\mu)} = \left(\int |f(rx)|^{q(r)} \mu(dx) \right)^{1/q(r)}.$$

When the function f and measure μ are clear from context, we denote $\alpha_{f,\mu} = \alpha$.

First we deal with the implication (sHC) \implies (sLSI).

3.1 (sHC) \implies (sLSI)

We begin with the following general statement.

Lemma 3.2 Suppose μ is a Euclidean regular probability measure. Let $q_0 > 1$, and let $f > 0$ be in $L^{q_0}(\mu) \cap C^1(\mathbb{R}^n)$. Let $\epsilon \in (0, 1)$, and suppose there are $q_\epsilon \geq 1$ and functions $h_1, h_2 \in L^{q_\epsilon}(\mu)$ such that for all $r \in (\epsilon, 1]$,

$$(3.1) \quad |f(rx)^{q(r)} \log f(rx)| \leq h_1(x), \quad |f(rx)^{q(r)-1} E f(rx)| \leq h_2(x) \quad \text{a.s.}[x].$$

Then for $r \in (\epsilon, 1]$, the functions $f_r^{q(r)}$ are in $L^{q_\epsilon}(\mu)$, and the function $\alpha = \alpha_{f, \mu}$ is differentiable on $(\epsilon, 1]$ with

(3.2)

$$\alpha'(r) = \frac{2}{cq(r)} \|f_r\|_{q(r)}^{1-q(r)} \left[\|f_r\|_{q(r)}^{q(r)} \log \|f_r\|_{q(r)}^{q(r)} - \int f(rx)^{q(r)} \log f(rx)^{q(r)} \mu(dx) + \frac{cq(r)}{2} \int f(rx)^{q(r)-1} Ef(rx) \mu(dx) \right].$$

Remark 3.3 Note that $(1/q(r))^{c/2} = r$. Hence, if $f \in \text{LSH}$ and μ satisfies the strong hypercontractivity property of (sHC) (with $p = 1$) we have $\alpha(r) \leq \|f\|_1 = \alpha(1)$ for $r \in (0, 1]$. The conditions of Lemma 3.2 guarantee that α is differentiable; hence, we essentially have that $\alpha'(1) \geq 0$. Equation (3.2) shows that $\alpha'(r)$ is closely related to the expression in (sLSI) for the function $f_r^{q(r)}$, and indeed this is our method for proving the equivalence of the logarithmic Sobolev inequality and strong hypercontractivity in what follows.

Proof Set $\beta(r, x) = f(rx)^{q(r)}$ so that $\alpha(r)^{q(r)} = \int \beta(r, x) \mu(dx)$. Note, $\beta(r, x) = f_r(x)^{q(r)}$.

First we show that if $f(rx)^{q(r)} \log f(rx) \in L^{q_\epsilon}(\mu)$, then, in fact, $f_r^{q(r)}$ is also in $L^{q_\epsilon}(\mu)$, and so $\beta(r, \cdot) \in L^1(\mu)$ for all $r \in (\epsilon, 1)$. The idea is simple: the logarithm cannot significantly improve the function $f(rx)^{q(r)}$. Rigorously, we fix $r \in (\epsilon, 1]$, we choose $0 < \delta < 1$, and we define

$$D = \{x \in \mathbb{R}^n : |f_r(x) - 1| < \delta\}.$$

The logarithm $\log f(rx)$ is bounded away from 0 on D^c , while the function $\mathbf{1}_D(x) f(rx)^{q(r)}$ is bounded, so it is in $L^{q_\epsilon}(\mu)$ on D . We have

$$f_r^{q(r)} = \mathbf{1}_D f_r^{q(r)} + \mathbf{1}_{D^c} f_r^{q(r)},$$

and there exists $c > 0$ such that $c \mathbf{1}_{D^c} f_r^{q(r)} \leq |f_r^{q(r)} \log f_r| \leq h_1$. Thus, $f_r^{q(r)} \in L^{q_\epsilon}(\mu)$.

Since $f \in C^1$ and strictly positive, we can check quickly that $\beta(\cdot, x)$ is as well. Using the fact that $q'(r) = -\frac{2}{c} r^{-2/c-1} = -\frac{2}{cr} q(r)$, and that $\frac{\partial}{\partial r} f(rx) = \frac{1}{r} Ef(rx)$, logarithmic differentiation yields

$$(3.3) \quad \frac{\partial}{\partial r} \beta(r, x) = q(r) \left[-\frac{2}{cr} f(rx)^{q(r)} \log f(rx) + \frac{1}{r} f(rx)^{q(r)-1} Ef(rx) \right].$$

From the hypotheses of the lemma, we therefore have

$$\left| \frac{\partial}{\partial r} \beta(r, x) \right| \leq \frac{q(r)}{r} \left[\frac{2}{c} h_1(x) + h_2(x) \right]$$

for almost every $x \in \mathbb{R}^n$, for $r \in (\epsilon', 1]$. As $q(r)/r$ is uniformly bounded on $(\epsilon', 1]$, we see that $|\frac{\partial}{\partial r} \beta(r, x)|$ is uniformly bounded above by an $L^1(\mu)$ function. It now follows from the Lebesgue differentiation theorem that $\alpha(r)^{q(r)} = \int \beta(r, x) \mu(dx)$ is

differentiable on a neighborhood of 1, and

$$\begin{aligned}
 \frac{d}{dr}[\alpha(r)^{q(r)}] &= \int \frac{\partial}{\partial r} \beta(r, x) \mu(dx) \\
 (3.4) \qquad \qquad \qquad &= -\frac{2}{cr} q(r) \int f(rx)^{q(r)} \log f(rx) \mu(dx) \\
 &\quad + \frac{1}{r} q(r) \int f(rx)^{q(r)-1} E f(rx) \mu(dx).
 \end{aligned}$$

Consequently $\alpha(r)$ is differentiable in a neighborhood of 1. Again using logarithmic differentiation,

$$\alpha'(r) = \alpha(r) \frac{d}{dr} \log \alpha(r) = \alpha(r) \frac{d}{dr} \left[\frac{1}{q(r)} \log \alpha(r)^{q(r)} \right],$$

and again using the fact that $q'(r) = -\frac{2}{cr} q(r)$,

$$\begin{aligned}
 \frac{d}{dr} \left[\frac{1}{q(r)} \log \alpha(r)^{q(r)} \right] &= \frac{2}{crq(r)} \log \alpha(r)^{q(r)} + \frac{1}{q(r)} \alpha(r)^{-q(r)} \frac{d}{dr} [\alpha(r)^{q(r)}] \\
 &= \frac{\alpha(r)^{-q(r)}}{q(r)} \left(\frac{2}{cr} \alpha(r)^{q(r)} \log \alpha(r)^{q(r)} + \frac{d}{dr} [\alpha(r)^{q(r)}] \right).
 \end{aligned}$$

Combining this with (3.4), we therefore have

$$\begin{aligned}
 (3.5) \qquad \alpha'(r) &= \frac{\alpha(r)^{1-q(r)}}{q(r)} \left[\frac{2}{cr} \alpha(r)^{q(r)} \log \alpha(r)^{q(r)} - \frac{2}{cr} q(r) \int f(rx)^{q(r)} \log f(rx) \mu(dx) \right. \\
 &\quad \left. + \frac{1}{r} q(r) \int f(rx)^{q(r)-1} E f(rx) \mu(dx) \right].
 \end{aligned}$$

Simplifying (3.5) and using the definition $\alpha(r) = \|f_r\|_{q(r)}$ yields (3.2), proving the lemma. ■

We therefore seek conditions on a function f (and on the measure μ) that guarantee the hypotheses of Lemma 3.2 (specifically the existence of the Lebesgue dominating functions h_1 and h_2). Naturally, we will work with LSH functions f . We will also make the fairly strong assumption that μ is rotationally invariant.

Notation 3.4 Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be locally-bounded. Denote by \tilde{f} the spherical average of f . That is, with ϑ denoting Haar measure on the group $O(n)$ of rotations of \mathbb{R}^n ,

$$\tilde{f}(x) = \int_{O(n)} f(ux) \vartheta(du).$$

If μ is rotationally invariant, then $\int f d\mu = \int \tilde{f} d\mu$ for any $f \in L^1(\mu)$. As such, we can immediately weaken the integrability conditions of Lemma 3.2 as follows.

Lemma 3.5 Suppose μ is a Euclidean regular probability measure that is invariant under rotations of \mathbb{R}^n . Let $q_0 > 1$ and let $f > 0$ be in $L^{q_0}_E(\mu)$. Denote by $f_1, f_2: (0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}$ the functions

$$(3.6) \qquad f_1(r, x) = f(rx)^{q(r)} \log f(rx), \quad f_2(r, x) = f(rx)^{q(r)-1} E f(rx).$$

Fix $\epsilon \in (0, 1)$, and suppose that there exist functions $h_1, h_2 \in L^1(\mu)$ such that, for $r \in (\epsilon, 1]$, $|\tilde{f}_j(r, x)| \leq h_j(x)$ for almost every x , $j = 1, 2$. (Here $\tilde{f}_j(r, \cdot)$ refers to the rotational average of $f_j(r, \cdot)$, as per Notation 3.4.) Then the conclusion of Lemma 3.2 stands: the function $\alpha = \alpha_{f, \mu}$ is differentiable on $(\epsilon, 1]$, and its derivative is given by (3.2).

Proof Following the proof of Lemma 3.2, only a few modifications are required. Defining $\beta(r, x)$ as above, $\alpha(r)^{q(r)} = \int \beta(r, x) \mu(dx)$; since μ is rotationally invariant, this is equal to $\int \tilde{\beta}(r, x) \mu(dx)$, where $\tilde{\beta}$ refers to the rotational average of β in the variable x . Evidently, $\tilde{\beta}(r, \cdot)$ is μ -integrable for sufficiently large $r < 1$ (since β is). To use the Lebesgue differentiation technique, we must verify that $\frac{\partial}{\partial r} \tilde{\beta}(r, x)$ exists for almost every x and is uniformly bounded by an $L^1(\mu)$ dominator. Note that $\beta(r, x)$ is locally-bounded in x for each r , and so for fixed x it is easy to verify that indeed

$$\frac{\partial}{\partial r} \tilde{\beta}(r, x) = \int_{O(n)} \frac{\partial}{\partial r} \beta(r, ux) \vartheta(du).$$

Using (3.3), we then have

$$\begin{aligned} \frac{\partial}{\partial r} \tilde{\beta}(r, x) = & \\ & q(r) \int_{O(n)} \left(-\frac{2}{cr} f(rux)^{q(r)} \log f(rux) + \frac{1}{r} f(rux)^{q(r)-1} E f(rux) \right) \vartheta(du). \end{aligned}$$

That is, using (3.6), $\frac{\partial}{\partial r} \tilde{\beta}(r, x) = q(r) \left[-\frac{2}{cr} \tilde{f}_1(r, x) + \frac{1}{r} \tilde{f}_2(r, x) \right]$. Hence, from the assumptions of this lemma,

$$\left| \frac{\partial}{\partial r} \tilde{\beta}(r, x) \right| \leq \frac{q(r)}{r} \left[\frac{2}{c} h_1(x) + h_2(x) \right]$$

and so, since $q(r)/r$ is uniformly bounded for $r \in (\frac{1}{2}, 1]$, it follows that $\alpha(r)^{q(r)} = \int \tilde{\beta}(r, x) \mu(dx)$ is differentiable near 1, with the derivative given by

$$\int \frac{\partial}{\partial r} \tilde{\beta}(r, x) \mu(dx) = q(r) \left[-\frac{2}{rc} \int \tilde{f}_1(r, x) \mu(dx) + \frac{1}{r} \int \tilde{f}_2(r, x) \mu(dx) \right].$$

Now using the rotational invariance of μ again, these integrals are the same as the corresponding non-rotated integrands $\int f_j(r, x) \mu(dx)$, yielding the same result as (3.4). The remainder of the proof follows the proof of Lemma 3.2 identically. ■

Remark 3.6 The point of Lemma 3.5 – that it is sufficient to find uniform Lebesgue dominators for the rotational averages of the terms in (3.1) – is actually quite powerful for us. While a generic subharmonic function in dimension ≥ 2 may not have good global properties, a rotationally invariant subharmonic function does, as the next proposition demonstrates. We will exploit this kind of behavior to produce the necessary bounds to verify the conditions of Lemma 3.5 and prove the differentiability of the norm.

Proposition 3.7 Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be subharmonic and locally-bounded. Then \tilde{f} is also subharmonic; moreover, for fixed $x \in \mathbb{R}^n$, $r \mapsto \tilde{f}(rx)$ is an increasing function of $r \in [0, 1]$.

Proof Fix $u \in O(n)$. Since f is locally-bounded, subharmonicity means that

$$\int_{B(x,r)} f(t) dt \geq f(x) \quad \text{for every } x \in \mathbb{R}^n, r \in (0, \infty).$$

Changing variables, we have

$$\int_{B(x,r)} f(ut) dt = \int_{u \cdot B(x,r)} f(t) dt = \int_{B(ux,r)} f(t) dt \geq f(ux).$$

Hence, $f \circ u$ is subharmonic for each $u \in O(n)$. The local-boundedness of f means that the function $u \mapsto f(ux)$ is uniformly bounded in $L^1(O(n), \vartheta)$ for x in a compact set, and hence it follows that \tilde{f} is subharmonic.

Hence \tilde{f} is a rotationally invariant subharmonic function. Fix $x \in \mathbb{R}^n$ and $r \in [0, 1]$. Then rx is in the ball $B(0, |x|)$, and since \tilde{f} is subharmonic, the maximum principle (cf. [14, Prop. 7.7.7]) asserts that $\tilde{f}(rx)$ is no larger than the maximum of \tilde{f} on $\partial B(0, |x|)$. But \tilde{f} is constantly equal to $\tilde{f}(x)$ on $\partial B(0, |x|)$ by rotational invariance, and so $\tilde{f}(rx) \leq \tilde{f}(x)$, proving the proposition. ■

Proposition 3.7 makes it quite easy to provide a uniform Lebesgue dominating function for the function f_1 in Lemma 3.5.

Proposition 3.8 *Suppose μ is a rotationally invariant probability measure on \mathbb{R}^n . Let $q_0 > 1$, and let $f \geq 0$ be subharmonic and in $L^{q_0}(\mu)$. Define f_1 as in (3.6): $f_1(r, x) = f(rx)^{q(r)} \log f(rx)$. Set $g_1(x) = f(x)^{q_0}$, and set $h_1 = \tilde{g}_1 + 1$; i.e., $h_1(x) = 1 + \int_{O(n)} f(ux)^{q_0} \vartheta(du)$. Then $h_1 \in L^1(\mu)$ and there is an $\epsilon \in (0, 1)$ and a constant $C > 0$ so that for all $r \in (\epsilon, 1]$, $|\tilde{f}_1(r, x)| \leq Ch_1(x)$ for almost every x .*

Remark 3.9 By the rotational invariance of μ , $\int h_1 d\mu = \int \tilde{g}_1 d\mu + 1 = \int g_1 d\mu + 1 = \int f^{q_0} d\mu + 1 < \infty$, and so h_1 is a uniform $L^1(\mu)$ dominator verifying the first condition of Lemma 3.5.

Proof Choose some small $\delta \in (0, 1)$. First note from simple calculus that, for $u \geq 1$, $u^{-\delta} \log u \leq \frac{1}{e\delta}$. Now, choose $\epsilon \in (0, 1)$ so that $q(\epsilon) < q_0 - \delta$; then $q(r) < q_0 - \delta$ for $r \in (\epsilon, 1]$. Consequently, if $f(y) \geq 1$, we have

$$0 \leq f(y)^{q(r)} \log f(y) \leq f(y)^{q_0 - \delta} \log f(y) \leq \frac{1}{e\delta} f(y)^{q_0}.$$

On the other hand, for $0 \leq u \leq 1$, $|u^{q(r)} \log u| \leq \frac{1}{eq(r)} \leq \frac{1}{e}$ (again by simple calculus). Thus, since $f \geq 0$, in total we have

$$(3.7) \quad |f(y)^{q(r)} \log f(y)| \leq \frac{1}{e} \max\left\{\frac{1}{\delta} f(y)^{q_0}, 1\right\} \leq \frac{1}{e\delta} [f(y)^{q_0} + 1].$$

Set $C = \frac{1}{e\delta}$. With $y = rx$, the left-hand side of (3.7) is precisely $\tilde{f}_1(r, x)$. Averaging (3.7) over $O(n)$ and recalling that $g_1(y) = f(y)^{q_0}$, we have

$$|\tilde{f}_1(r, x)| \leq C[\tilde{g}_1(rx) + 1].$$

Recall that if ϕ is convex and f is subharmonic, then $\phi \circ f$ is also subharmonic. Thus, since $q_0 > 1$ and f is subharmonic, g_1 is also subharmonic, and hence from Proposition 3.7, $\tilde{g}_1(rx) \leq \tilde{g}_1(x)$. This proves the proposition. ■

We must now bound the second term $\tilde{f}_2(r, \cdot)$ uniformly for r in a neighborhood of 1. The following Lemma is useful in this regard.

Lemma 3.10 *Let \tilde{k} be a C^1 non-negative subharmonic rotationally invariant function. Then for $x \in \mathbb{R}^n$ and $r \in (0, 1]$,*

$$(3.8) \quad E\tilde{k}(rx) \leq r^{2-n} E\tilde{k}(x).$$

Proof First, note that it suffices to assume \tilde{k} is in fact C^∞ . Indeed, for more general \tilde{k} , let ϕ be a rotationally invariant non-negative compactly-supported bump function, and replace \tilde{k} with $\tilde{k} * \phi$. By Lemma 1.21, this function is subharmonic and C^∞ ; it is also rotationally invariant. If we proceed to prove (3.8) for this mollified function, we may then take an approximate identity sequence of ϕ . Now, since $\tilde{k} \in C^1$, $\partial_j(\tilde{k} * \phi) = (\partial_j\tilde{k}) * \phi$ for $j = 1, \dots, n$, and the functions $\partial_j\tilde{k}$ are continuous and locally bounded. Hence, we may choose the approximate identity sequence so that the derivatives converge pointwise (or even uniformly on compact sets), which shows that both sides of (3.8) converge appropriately. Henceforth, we assume \tilde{k} is C^∞ .

Since \tilde{k} is rotationally invariant, there is a function $h: [0, \infty) \rightarrow \mathbb{R}$ so that $\tilde{k}(x) = h(|x|)$. The Laplacian of \tilde{k} can then be expressed in terms of derivatives of h ; the result is

$$\Delta\tilde{k}(x) = h''(|x|) + (n-1)\frac{1}{|x|}h'(|x|).$$

Hence, since \tilde{k} is subharmonic and smooth, it follows that for $t > 0$,

$$(3.9) \quad t h''(t) + (n-1)h'(t) \geq 0.$$

One can also check that, in this case, $E\tilde{k}(x) = |x|h'(|x|)$. Now define $F(r) = r^{n-2}E\tilde{k}(rx) = r^{n-2}r|x|h'(r|x|)$. Then F is smooth on $(0, \infty)$ and $F(1) = |x|h'(|x|) = E\tilde{k}(x)$. We differentiate, yielding

$$\begin{aligned} F'(r) &= |x|\frac{d}{dr}r^{n-1}h'(r|x|) = |x|(n-1)r^{n-2}h'(r|x|) + |x|r^{n-1}h''(r|x|)|x| \\ &= |x|r^{n-2}[r|x|h''(r|x|) + (n-1)h'(r|x|)]. \end{aligned}$$

Equation (3.9) with $t = r|x|$ now yields that $F'(r) \geq 0$ for $r > 0$. Hence, $F(r) \leq F(1)$ for $r \leq 1$. This is precisely the statement of the lemma. ■

Proposition 3.11 *Let $q_0 > 1$ and let μ be a rotationally invariant probability measure on \mathbb{R}^n . Let $f > 0$ be subharmonic, C^1 , and in $L_E^{q_0}(\mu)$. Define f_2 as in (3.6): $f_2(r, x) = f(rx)^{q(r)-1}Ef(rx)$. Set $g_3(x) = (f(x)^{q_0-1} + 1)|Ef(x)|$, and set $h_2 = \tilde{g}_3$. Then there is an $\epsilon \in (0, 1)$ and a constant $C > 0$ so that for all $r \in (\epsilon, 1]$, $|\tilde{f}_2(r, x)| \leq Ch_2(x)$ for almost every x ; moreover, $h_2 \in L^1(\mu)$.*

Proof Fix $\epsilon \in (0, 1)$ small enough that $q(r) < q_0$ for all $r \in (\epsilon, 1]$. Define $g_2(r, y) = f(y)^{q(r)-1}Ef(y)$, and note that $f_2(r, x)$ is given by the dilation $f_2(r, x) = g_2(r, rx)$.

Since E is a first-order differential operator, we can quickly check that

$$g_2(r, y) = \frac{1}{q(r)} E(f^{q(r)})(y).$$

We now average both sides over $O(n)$. Set $k = f^{q(r)}$, which is C^1 , and let $u \in O(n)$. Then we have the following calculus identity:

$$E(k \circ u)(y) = y \cdot \nabla(k \circ u)(y) = y \cdot u^\top \nabla k(uy) = (uy) \cdot \nabla k(uy) = (Ek)(uy).$$

For fixed y , the function $u \mapsto (Ek)(uy)$ is uniformly bounded and so we integrate both sides to yield

$$\begin{aligned} \widetilde{Ek}(y) &= \int_{O(n)} (Ek)(uy) \vartheta(du) = \int E(k \circ u)(y) \vartheta(du) \\ &= E \int k \circ u(y) \vartheta(du) = E(\widetilde{k})(y). \end{aligned}$$

In other words, $\widetilde{g}_2(r, y) = \frac{1}{q(r)} E(\widetilde{f^{q(r)}})(y)$. As in the proof of Proposition 3.8, the function $\widetilde{k} = \widetilde{f^{q(r)}}$ is subharmonic and rotationally invariant. Hence, we employ Lemma 3.10 and have

$$\widetilde{g}_2(r, rx) = \frac{1}{q(r)} E\widetilde{k}(rx) \leq \frac{1}{q(r)} r^{2-n} E\widetilde{k}(x) = r^{2-n} \widetilde{g}_2(r, x).$$

Since r^{2-n} is uniformly bounded for $r \in (\epsilon, 1]$, it now suffices to find a uniform dominator for $\widetilde{g}_2(r, x)$.

We therefore make the following estimates. Since $q(r) < q_0$, we have

$$\begin{aligned} |g_2(r, x)| &= f(x)^{q(r)-1} |Ef(x)| \leq \max\{f(x)^{q(r)-1}, 1\} |Ef(x)| \\ &\leq \max\{f(x)^{q_0-1}, 1\} |Ef(x)| \leq (f(x)^{q_0-1} + 1) |Ef(x)|. \end{aligned}$$

That is to say, $|g_2(r, x)| \leq g_3(x)$ for $r \in (\epsilon, 1]$. Hence,

$$\begin{aligned} |\widetilde{g}_2(r, x)| &= \left| \int_{O(n)} g_2(r, ux) \vartheta(du) \right| \leq \int_{O(n)} |g_2(r, ux)| \vartheta(du) \\ &\leq \int_{O(n)} g_3(ux) \vartheta(du) = \widetilde{g}_3(x) = h_2(x), \end{aligned}$$

thus proving the estimate.

As usual, by rotational invariance of μ , $\int \widetilde{g}_3 d\mu = \int g_3 d\mu$, and so to show $h_2 \in L^1(\mu)$ we need only verify that $g_3 \in L^1(\mu)$. To that end, we break up $g_3(x) = f(x)^{q_0-1} |Ef(x)| + |Ef(x)|$. By assumption, $f \in L^{q_0}_E(\mu)$ and so $|Ef| \in L^{q_0}(\mu)$; as μ is a finite measure, this means that $|Ef| \in L^1(\mu)$ and hence the second term is integrable. For the first term, we use Hölder's inequality:

$$\int f^{q_0-1} |Ef| d\mu \leq \|f^{q_0-1}\|_{q'_0} \|Ef\|_{q_0} = \|f\|^{q_0-1}_{q_0} \|Ef\|_{q_0}.$$

Both terms are finite, since $f \in L^{q_0}_E(\mu)$, and hence $g_3 \in L^1(\mu)$, proving the proposition. ■

Combining Lemma 3.5 and Propositions 3.8 and 3.11, we therefore have the following theorem.

Theorem 3.12 *Let $q_0 > 1$ and let μ be a probability measure of Euclidean type q_0 that is invariant under rotations of \mathbb{R}^n . Suppose that μ satisfies strong hypercontractivity of (sHC) with constant $c > 0$. Let $f \in L_E^{q_0}(\mu) \cap \text{LSH}$. Then the strong log-Sobolev inequality, (sLSI), holds for f :*

$$\int f \log f \, d\mu - \int f \, d\mu \log \int f \, d\mu \leq \frac{c}{2} \int E f \, d\mu.$$

Proof Under the conditions stated above, the results of the preceding section show that the function $\alpha = \alpha_{f,\mu}$ is differentiable on $(\epsilon', 1]$ for some $\epsilon' \in (0, 1)$. Since μ satisfies strong hypercontractivity, Proposition 1.19 shows that the function α is non-decreasing on $(0, 1]$. It therefore follows that $\alpha'(r) \geq 0$ for $r \in (\epsilon', 1]$ (here $\alpha'(1)$ denotes the left derivative). Hence, from (3.2) we have, for $r \in (\epsilon', 1]$,

$$\begin{aligned} \|f_r\|_{q(r)}^{q(r)} \log \|f_r\|_{q(r)}^{q(r)} - \int f(rx)^{q(r)} \log f(rx)^{q(r)} \mu(dx) \\ + \frac{cq(r)}{2} \int f(rx)^{q(r)-1} E f(rx) \mu(dx) \geq 0. \end{aligned}$$

At $r = 1$, this reduces precisely to (sLSI), proving the result. ■

Theorem 3.12 implies Theorem 1.17(ii). Indeed, let $g \in \text{LSH}_E^{1<}$ and let (f_k) be a sequence of functions converging to g in $L_E^1(\mu)$ and such that (sLSI) holds for each f_k . Then, by the definition of the norm of the Sobolev space $L_E^1(\mu)$, we have

$$\int f_k \, d\mu \rightarrow \int f \, d\mu \quad \text{and} \quad \int E f_k \, d\mu \rightarrow \int E f \, d\mu.$$

There exists a subsequence $f_{k'}$ tending to f almost surely. We apply the Dominated Convergence Theorem to the sequence $\int f_{k'} \log f_{k'} 1_{\{f_{k'} \leq 1\}} \, d\mu$ and Fatou's Lemma to the sequence $\int f_{k'} \log f_{k'} 1_{\{f_{k'} > 1\}} \, d\mu$. The inequality (sLSI) for f follows.

We now turn to Theorem 1.17(i). We will need the following refinement of Propositions 3.8 and 3.11. The proofs are the same, paying more attention to L^q -integrability, $q > 1$, and to the precise value of ϵ .

Corollary 3.13 *Propositions 3.8 and 3.11 hold for any $\epsilon = 1/q_0^{2/c} + \delta < 1$, with $\delta > 0$. For any fixed ϵ of this form, the majorizing functions h_1, h_2 belong to $L^q(\mu)$ for some $q > 1$.*

3.2 (sLSI) \implies (sHC)

We utilize many of the results in the previous section in the same manner they were stated; we therefore outline this direction more briefly.

First we prove Theorem 1.17(i)(b). Fix some $q_0 > 1$, and let $g \in \text{LSH} \cap L_E^{q_0}(\mu)$. We proceed as in the proof of Theorem 1.7. In order to justify differentiating under the integral, we use Lemma 3.5 and Propositions 3.8 and 3.11 with Corollary 3.13. Using Fatou's Lemma, we obtain the strong hypercontractivity inequalities from Proposition 1.19 for $g \in \text{LSH} \cap L_E^{q_0}(\mu)$ and $r \in [1/(q_0^{2/c}), 1]$.

In the next step of the proof we show that the partial strong hypercontractivity inequalities from Proposition 1.19 hold for $h \in \text{LSH} \cap L^q(\mu)$ for any $q > 1$ and $r \in$

$[1/(q^{2/\epsilon}), 1]$. By Theorem 1.15, there exists a sequence $(g_k) \subset \text{LSH} \cap L_E^q(\mu)$ converging to h in $L^q(\mu)$, so also in $L^1(\mu)$. By passing to a subsequence, we may suppose that g_k converge to h almost surely; thus $(g_k)_r$ converge to h_r almost surely for any $r \in (0, 1]$. Fatou's Lemma then implies that $\|h_r\|_{q(r)} \leq \|h\|_1$. It is in this step of the proof that the hypothesis of p -Euclidean exponential type of μ for every $p > 1$ is essential. Thence, we obtain Theorem 1.17(i)(b).

In order to prove Theorem 1.17(i)(a), we first prove the following property of exponentially sub-additive and α -subhomogeneous measures μ :

$$(3.10) \quad f \in L^1(\mu) \cap \text{LSH} \Rightarrow f_r^{q(r)} \in L^1(\mu), \quad q(r) = r^{-\frac{2}{\alpha}}.$$

The property of exponential sub-additivity of μ allows us to show that if $f \in L^1(\mu) \cap \text{LSH}$, then the function $\rho(x)f(x)$ is bounded by a multiple of $\|f\|_1$. (We profit from the fact that the product $e^{-(A,t)f(x+t)}$ is LSH, so also subharmonic. We use the fact that $g(0) \leq \int g d\mu$ for any subharmonic function). Next, it is easy to show that the α -subhomogeneity of μ together with the boundedness of ρf implies that $f_r^{q(r)} \in L^1(\mu)$.

Now, as in the proof of (i)(a), we suppose that $g \in \text{LSH} \cap L_E^{q_0}(\mu)$ for some $q_0 > 1$. We obtain partial (sHC) inequalities for $r \in [q_0^{-2/\epsilon}, 1]$, but property (3.10) allows us to iterate the proof procedure and to get partial (sHC) inequalities for $r \in [(q_0^2)^{-2/\epsilon}, 1]$. By induction, the (sHC) inequalities hold on any segment $[(q_0^{2^n})^{-2/\epsilon}, 1]$, so on $(0, 1]$.

Finally, we eliminate the L_E hypothesis precisely as in the proof of (i)(a): consider $f \in L^1(\mu) \cap \text{LSH}$. Let $\alpha < 1$. Then $f^\alpha \in L^{\frac{1}{\alpha}}(\mu) \cap \text{LSH}$. By the previous step, the inequalities from Proposition 1.19 hold for f^α . Now let $\alpha \nearrow 1$. By the Monotone Convergence Theorem applied on the domain $\{f \geq 1\}$ and the Dominated Convergence Theorem applied on the domain $\{f < 1\}$ we get the same inequalities for f . This completes the proof of Theorem 1.17(i).

Appendix A Properties of Euclidean Regular Measures

In this brief appendix, we show several closure properties of the class of Euclidean regular measures (of any given exponential type $p \in [0, \infty)$): it is closed under bounded perturbations, convex combinations, product, and convolution. Throughout, we use μ_i ($i = 1, 2$) to stand for such measures, and ρ_i to stand for their densities.

Proposition A.1 *Let μ_1 and μ_2 be positive measures on \mathbb{R}^n , and suppose μ_1 is Euclidean exponential type $p \in [0, \infty)$. If there are constants $C, D > 0$ such that $C\mu_1 \leq \mu_2 \leq D\mu_1$, then μ_2 is also Euclidean exponential type p .*

Proof The assumption is that $C\rho_1 \leq \rho_2 \leq D\rho_1$. Let $\epsilon > 0$ be such that

$$\sup_{1 < a < 1+\epsilon} C_{\rho_1}^0(a, 0) < \infty.$$

Then for any such a ,

$$\frac{\rho_2(ax)}{\rho_2(x)} \leq \frac{D\rho_1(ax)}{C\rho_1(x)} \leq \frac{D}{C} C_{\rho_1}^0(a, 0)$$

for all x ; thus $C_{\rho_2}^0(a, 0) \leq \frac{D}{C} C_{\rho_1}^0(a, 0)$, and so $\sup_{1 < a < 1 + \epsilon} C_{\rho_2}^0(a, 0) < \infty$. Similarly, for $x, y \in \mathbb{R}^n$ and $a > 1$,

$$|x|^p \frac{\rho_2(ax + y)}{\rho_2(x)} \leq |x|^p \frac{D\rho_1(ax + y)}{C\rho_1(x)} \leq \frac{D}{C} C_{\rho_1}^p(a, |y|)$$

and so $C_{\rho_2}^p(a, s) \leq \frac{D}{C} C_{\rho_1}^p(a, s) < \infty$. ■

Proposition A.2 Let μ_1 and μ_2 be Euclidean regular measures of exponential type $p \in [0, \infty)$. For any $t \in [0, 1]$, $\mu = (1 - t)\mu_1 + t\mu_2$ is Euclidean exponential type p .

Proof Let $\epsilon > 0$ be such that $\sup_{1 < a < 1 + \epsilon} C_{\rho_i}^0(a, 0) < \infty$ for $i = 1, 2$. Let ρ be the density of μ . Then for any $x \in \mathbb{R}^n$,

$$\begin{aligned} \rho(ax) &= (1 - t)\rho_1(ax) + t\rho_2(ax) \leq (1 - t)C_{\rho_1}^0(a, 0)\rho_1(x) + tC_{\rho_2}^0(a, 0)\rho_2(x) \\ &\leq \max\{C_{\rho_1}^0(a, 0), C_{\rho_2}^0(a, 0)\}\rho(x) \end{aligned}$$

and so $C_{\rho}^0(a, 0) \leq \max\{C_{\rho_1}^0(a, 0), C_{\rho_2}^0(a, 0)\}$ is uniformly bounded for $1 < a < 1 + \epsilon$, as required. Similarly, for $x, y \in \mathbb{R}^n$ and $a > 1$,

$$\begin{aligned} |x|^p \rho(ax + y) &\leq (1 - t)|x|^p \rho_1(ax + y) + t|x|^p \rho_2(ax + y) \\ &\leq (1 - t)C_{\rho_1}^p(a, |y|)\rho_1(x) + tC_{\rho_2}^p(a, |y|)\rho_2(x) \\ &\leq \max\{C_{\rho_1}^p(a, |y|), C_{\rho_2}^p(a, |y|)\}\rho(x) \end{aligned}$$

which shows that $C_{\rho}^p(a, s) \leq \max\{C_{\rho_1}^p(a, s), C_{\rho_2}^p(a, s)\} < \infty$ for $a \geq 1$ and $s \geq 0$. ■

Proposition A.3 Let $p \in [0, \infty)$, let μ_1 be a Euclidean exponential type p measure on \mathbb{R}^{n_1} , and let μ_2 be a Euclidean exponential type p measure on \mathbb{R}^{n_2} . Then the product measure $\mu_1 \otimes \mu_2$ is Euclidean exponential type p on $\mathbb{R}^{n_1 + n_2}$.

Proof For $i = 1, 2$ let ρ_i be the density of μ_i ; then $\mu_1 \otimes \mu_2$ has density $\rho_1 \otimes \rho_2(x_1, x_2) = \rho_1(x_1)\rho_2(x_2)$. Fix $\epsilon > 0$ so that $\sup_{1 < a < 1 + \epsilon} C_{\rho_i}^p(0, a) < \infty$ for $i = 1, 2$. Then, letting $\mathbf{x} = (x_1, x_2)$,

$$\rho_1 \otimes \rho_2(a\mathbf{x}) = \rho_1(ax_1)\rho_2(ax_2) \leq C_{\rho_1}^0(a, 0)\rho_1(x_1) \cdot C_{\rho_2}^0(a, 0)\rho_2(x_2)$$

and so $C_{\rho_1 \otimes \rho_2}^0(a, 0) \leq C_{\rho_1}^0(a, 0) \cdot C_{\rho_2}^0(a, 0)$, meaning $\sup_{1 < a < 1 + \epsilon} C_{\rho_1 \otimes \rho_2}^0(a, 0) < \infty$. Similarly, for fixed $\mathbf{x}, \mathbf{y} \in \mathbb{R}^{n_1 + n_2}$ and $a > 1$,

$$|\mathbf{x}|^p \rho_1 \otimes \rho_2(a\mathbf{x} + \mathbf{y}) = (|x_1| + |x_2|)^p \rho_1(ax_1 + y_1)\rho_2(ax_2 + y_2).$$

By elementary calculus, $(|x_1| + |x_2|)^p \leq 2^{p-1}(|x_1|^p + |x_2|^p)$, and so we have

$$\begin{aligned} |\mathbf{x}|^p \rho_1 \otimes \rho_2(a\mathbf{x} + \mathbf{y}) &\leq 2^{p-1}|x_1|^p \rho_1(ax_1 + y_1) \cdot \rho_2(ax_2 + y_2) \\ &\quad + 2^{p-1}\rho_1(ax_1 + y_1) \cdot |x_2|^p \rho_2(ax_2 + y_2). \end{aligned}$$

For the first term, we have $|x_1|^p \rho_1(ax_1 + y_1) \leq C_{\rho_1}^p(a, |y_1|)\rho_1(x_1)$, while $\rho_2(ax_2 + y_2) \leq C_{\rho_2}^0(a, |y_2|)$; for the second term, we have $\rho_1(ax_1 + y_1) \leq C_{\rho_1}^0(a, |y_1|)\rho_1(x_1)$, while

$|x_2|^p \rho_2(ax_2 + y_2) \leq C_{\rho_2}^p(a, |y_2|)$. If $|y| \leq s$, then $|y_i| \leq s$ for $i = 1, 2$. All together, this shows that

$$C_{\rho_1 \otimes \rho_2}^p(a, s) \leq 2^{p-1} [C_{\rho_1}^p(a, s) C_{\rho_2}^0(a, s) + C_{\rho_1}^0(a, s) C_{\rho_2}^p(a, s)],$$

which is finite, since both ρ_1, ρ_2 are Euclidean exponential type p (and hence also Euclidean regular). This proves the proposition. ■

Proposition A.4 *Let μ_1 and μ_2 be positive measures on \mathbb{R}^n , each of Euclidean exponential type $p \in [0, \infty)$. Then $\mu_1 * \mu_2$ is Euclidean exponential type p .*

Proof Let ρ_j be the density of μ_j . By assumption, for $i = 1, 2$, $C_{\rho_i}^{p_i}(a, s) < \infty$ for $a > 1$ and $s \geq 0$, and there is $\epsilon > 0$ such that $\sup_{1 < a < 1+\epsilon} C_{\rho_i}^0(a, 0) < \infty$ (cf. (1.3)). Then for $a \geq 1$ and $x \in \mathbb{R}^n$,

$$\rho_1 * \rho_2(ax) = \int \rho_1(ax - u) \rho_2(u) du = a^n \int \rho_1(ax - av) \rho_2(av) dv.$$

By definition, $\rho_1(a(x - v)) \leq C_{\rho_1}^0(a, 0) \rho_1(x - v)$ and $\rho_2(av) \leq C_{\rho_2}^0(a, 0) \rho_2(v)$ for all x, v . Thus,

$$\begin{aligned} \rho_1 * \rho_2(ax) &\leq a^n C_{\rho_1}^0(a, 0) \cdot C_{\rho_2}^0(a, 0) \int \rho_1(x - v) \rho_2(v) dv \\ &= a^n C_{\rho_1}^0(a, 0) \cdot C_{\rho_2}^0(a, 0) \rho_1 * \rho_2(x). \end{aligned}$$

It follows that $C_{\rho_1 * \rho_2}^0(a, 0) \leq a^n C_{\rho_1}^0(a, 0) \cdot C_{\rho_2}^0(a, 0)$, and hence

$$(A.1) \quad \sup_{1 < a < 1+\epsilon} C_{\rho_1 * \rho_2}^0(a, 0) \leq (1 + \epsilon)^n \sup_{1 < a < 1+\epsilon} C_{\rho_1}^0(a, 0) \cdot \sup_{1 < a < 1+\epsilon} C_{\rho_2}^0(a, 0) < \infty,$$

as required. Similarly, for $x, y \in \mathbb{R}^n$ and $a > 1$,

$$\begin{aligned} |x|^p \rho_1 * \rho_2(ax + y) &= |x|^p \int \rho_1(ax + y - u) \rho_2(u) du \\ &= a^n \int |x|^p \rho_1(a(x - v) + y) \rho_2(av) dv. \end{aligned}$$

Note (by elementary calculus) that $|x|^p \leq 2^{p-1}(|x - v|^p + |v|^p)$, and so

$$\begin{aligned} |x|^p \rho_1 * \rho_2(ax + y) &\leq 2^{p-1} a^n \left[\int |x - v|^p \rho_1(a(x - v) + y) \rho_2(av) dv \right. \\ &\quad \left. + \int \rho_1(a(x - v) + y) |v|^p \rho_2(av) dv \right]. \end{aligned}$$

In the first term, we have $|x - v|^p \rho_1(a(x - v) + y) \leq C_{\rho_1}^p(a, |y|) \rho_1(x - v)$ and $\rho_2(av) \leq C_{\rho_2}^0(a, 0)$, and so

$$\int |x - v|^p \rho_1(a(x - v) + y) \rho_2(av) dv \leq C_{\rho_1}^p(a, |y|) \cdot C_{\rho_2}^0(a, 0) \rho_1 * \rho_2(x).$$

In the second term, we have $\rho_1(a(x - v) + y) \leq C_{\rho_1}^0(a, |y|) \rho_1(x - v)$ and $|v|^p \rho_2(av) \leq C_{\rho_2}^p(a, 0)$, and so

$$\int \rho_1(a(x - v) + y) |v|^p \rho_2(av) dv \leq C_{\rho_1}^0(a, |y|) \cdot C_{\rho_2}^p(a, 0) \rho_1 * \rho_2(x).$$

All together, for any $s \geq |y|$, this gives

$$(A.2) \quad C_{\rho_1 * \rho_2}^p(a, s) \leq 2^{p-1} a^n [C_{\rho_1}^p(a, s) \cdot C_{\rho_2}^0(a, 0) + C_{\rho_1}^0(a, s) \cdot C_{\rho_2}^p(a, 0)]$$

which is finite, since both ρ_1 and ρ_2 are Euclidean exponential type p (and thus also Euclidean regular). Equations (A.1) and (A.2) prove the proposition. ■

Acknowledgments The authors wish to express their gratitude to the anonymous referee, who not only helped us significantly improve the exposition of this paper, but whose comments also led us to strengthen the conclusions of our main theorems which, in original form, had included technical assumptions that were unnecessary.

References

- [1] G. Anderson, A. Guionnet, and O. Zeitouni, *An introduction to random matrices*. Cambridge Studies in Advanced Mathematics, 118, Cambridge University Press, Cambridge, 2010.
- [2] D. Bakry, *L'hypercontractivité et son utilisation en théorie des semigroupes*. In: Lectures on probability theory (Saint-Flour, 1992), Lecture Notes in Math., 1581, Springer, Berlin, 1994, pp. 1–114.
- [3] ———, *On Sobolev and logarithmic Sobolev inequalities true for Markov semigroups*. In: New trends in stochastic analysis (Charingworth, 1994), World Sci. Publ., River Edge, NJ, 1997, pp. 43–75.
- [4] D. Bakry and M. Emery, *Diffusions hypercontractives*. Séminaire de Probabilités, XIX, Lecture Notes in Math. 1123, Springer, Berlin, 1985, pp. 177–206.
- [5] S. Bobkov and C. Houdré, *Some connections between isoperimetric and Sobolev-type inequalities*. Mem. Amer. Math. Soc. 129(1997), no. 616.
- [6] S. Bobkov and M. Ledoux, *From Brunn-Minkowski to Brascamp-Lieb and to logarithmic Sobolev inequalities*. Geom. Funct. Anal. 10(2000), no. 5, 1028–1052.
<http://dx.doi.org/10.1007/PL00001645>
- [7] S. Bobkov and P. Tetali, *Modified logarithmic Sobolev inequalities in discrete settings*. J. Theoret. Probab. 19(2006), no. 2, 289–336. <http://dx.doi.org/10.1007/s10959-006-0016-3>
- [8] E. B. Davies, *Explicit constants for Gaussian upper bounds on heat kernels*. Amer. J. Math. 109(1987), no. 2, 319–333. <http://dx.doi.org/10.2307/2374577>
- [9] ———, *Heat kernels and spectral theory*. Cambridge Tracts in Mathematics, 92, Cambridge University Press, Cambridge, 1990.
- [10] E. B. Davies and B. Simon, *Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians*. J. Funct. Anal. 59(1984), no. 2, 335–395.
[http://dx.doi.org/10.1016/0022-1236\(84\)90076-4](http://dx.doi.org/10.1016/0022-1236(84)90076-4)
- [11] P. Diaconis and L. Saloff-Coste, *Logarithmic Sobolev inequalities for finite Markov chains*. Ann. Appl. Probab. 6(1996), no. 3, 695–750. <http://dx.doi.org/10.1214/aop/1034968224>
- [12] F. Galaz-Fontes, L. Gross, and S. B. Sontz, *Reverse hypercontractivity over manifolds*. Ark. Math. 39(2001), no. 2, 283–309. <http://dx.doi.org/10.1007/BF02384558>
- [13] P. Graczyk, T. Kemp, and J.-J. Loeb, *Hypercontractivity for log-subharmonic functions*. J. Funct. Anal. 258(2010), no. 6, 1785–1805. <http://dx.doi.org/10.1016/j.jfa.2009.08.014>
- [14] R. Greene and S. Krantz, *Function theory of one complex variable*. Third ed., Graduate Studies in Mathematics, 40, American Mathematical Society, Providence, RI, 2006.
- [15] L. Gross, *Logarithmic Sobolev inequalities*. Amer. J. Math. 97(1975), no. 4, 1061–1083.
<http://dx.doi.org/10.2307/2373688>
- [16] ———, *Hypercontractivity over complex manifolds*. Acta Math. 182(1999), no. 2, 159–206.
<http://dx.doi.org/10.1007/BF02392573>
- [17] L. Gross and M. Grothaus, *Reverse hypercontractivity for subharmonic functions*. Canad. J. Math. 57(2005), no. 3, 506–534. <http://dx.doi.org/10.4153/CJM-2005-022-2>
- [18] A. Guionnet and B. Zegarlinski, *Lectures on logarithmic Sobolev inequalities*. Séminaire de Probabilités, XXXVI, Lecture Notes in Math., 1801, Springer, Berlin, 2003, pp. 1–134.
- [19] R. Holley and D. Stroock, *Logarithmic Sobolev inequalities and stochastic Ising models*. J. Statist. Phys. 46(1987), Ino. 5–6, 159–1194. <http://dx.doi.org/10.1007/BF01011161>
- [20] S. Janson, *On hypercontractivity for multipliers of orthogonal polynomials*. Ark. Mat. 21(1983), no. 1, 97–110. <http://dx.doi.org/10.1007/BF02384302>
- [21] ———, *On complex hypercontractivity*. J. Funct. Anal. 151(1997), no. 1, 270–280.
<http://dx.doi.org/10.1006/jfan.1997.3144>
- [22] M. Ledoux, *Isoperimetry and Gaussian analysis*. In: Lectures on probability theory and statistics, Lecture Notes in Math 1648, Springer, Berlin, 1996, pp. 165–294.

- [23] ———, *Concentration of measure and logarithmic Sobolev inequalities*. Séminaire de Probabilités, XXXIII, Lecture Notes in Math., 1709, Springer, Berlin, 1999, pp. 120–216.
- [24] ———, *The geometry of Markov diffusion generators*. Ann. Fac. Sci. Toulouse Math. (6) 9(2000), no. 2, 305–366. <http://dx.doi.org/10.5802/afst.962>
- [25] ———, *The concentration of measure phenomenon*. Mathematical Surveys and Monographs, 89, American Mathematical Society, Providence, RI, 2001.
- [26] ———, *A remark on hypercontractivity and tail inequalities for the largest eigenvalues of random matrices*. Séminaire de Probabilités XXXVII, Lecture Notes in Math., 1832, Springer, Berlin, 2003, pp. 360–369.
- [27] ———, *Spectral gap, logarithmic Sobolev constant, and geometric bounds*. In: Surveys in differential geometry. Vol. IX, Surv. Differ. Geom., IX, Int. Press, Somerville, MA, 2004, pp. 219–240.
- [28] P. Lelong and L. Gruman, *Entire functions of several complex variables*. Grundlehren der Mathematischen Wissenschaften, 282, Springer-Verlag, Berlin, 1986.
- [29] E. Nelson, *The free Markov field*. J. Funct. Anal. 12(1973), 211–227. [http://dx.doi.org/10.1016/0022-1236\(73\)90025-6](http://dx.doi.org/10.1016/0022-1236(73)90025-6)
- [30] A. J. Stam, *Some inequalities satisfied by the quantities of information of Fisher and Shannon*. Information and Control 2(1959), 101–112. [http://dx.doi.org/10.1016/S0019-9958\(59\)90348-1](http://dx.doi.org/10.1016/S0019-9958(59)90348-1)
- [31] C. Villani, *Topics in optimal transportation*. Graduate Studies in Mathematics, 58, American Mathematical Society, Providence, RI, 2003.
- [32] H.-T. Yau, *Logarithmic Sobolev inequality for the lattice gases with mixing conditions*. Commun. Math. Phys. 181(1996), no. 2, 367–408. <http://dx.doi.org/10.1007/BF02101009>
- [33] ———, *Logarithmic Sobolev inequality for generalized simple exclusion processes*. Probab. Theory Related Fields 109(1997), no. 4, 507–538. <http://dx.doi.org/10.1007/s004400050140>
- [34] B. Zegarlinski, *Dobrushin uniqueness theorem and logarithmic Sobolev inequalities*. J. Funct. Anal. 105(1992), no. 1, 77–111. [http://dx.doi.org/10.1016/0022-1236\(92\)90073-R](http://dx.doi.org/10.1016/0022-1236(92)90073-R)
- [35] D. Zimmermann, *Logarithmic Sobolev inequalities for mollified compactly supported measures*. J. Funct. Anal. 265(2013), no. 6, 1064–1083. <http://dx.doi.org/10.1016/j.jfa.2013.05.029>

Université d'Angers, 2 Boulevard Lavoisier, 49045 Angers Cedex 01, France

e-mail: piotr.graczyk@univ-angers.fr

Department of Mathematics, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0112, USA

e-mail: tkemp@math.ucsd.edu

Université d'Angers, 2 Boulevard Lavoisier, 49045 Angers Cedex 01, France

e-mail: jean-jacques.loeb@univ-angers.fr