

July 1959 ✓

J. P. Thomsen

Vol. LXXIII No. 7

The Journal of Laryngology and Otology

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

Contents

THE INFLUENCE OF PRESSURE VARIATIONS ON THE IMPEDANCE OF THE HUMAN EAR DRUM	K. TERKILDSEN ✓ and K. A. THOMSEN
LIPOMATA OF THE UPPER AIR AND FOOD PASSAGES	ALEX R. HARPER ✓
THE TREATMENT OF RECURRENT NASAL POLYPI BY INTERO-EXTERNAL "WINDOW" ETHMOIDECTOMY	F. BOYES KORKIS ✓
UVULOTOMY AMONG EAST AFRICAN TRIBES	J. F. JARVIS and ✓ S. N. MWATHI
A NOTE ON THE DEVELOPMENT OF THE EXTERNAL EAR	T. G. WILSON ✓
CLINICAL RECORD— FATALITY RESULTING FROM INTRA-NASAL POLYPECTOMY	GILBERT FORBES and ✓ ANDREW BRADLEY
CLINICAL NOTE— TYMPANO DYNAMOMETRY	A. TUMARKIN ✓
SOCIETIES' PROCEEDINGS— ROYAL SOCIETY OF MEDICINE—SECTION OF OTOTOLOGY	✓
GENERAL NOTES	

London

Headley Brothers Ltd

109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13

Monthly, 10/- net post free

The Journal of Laryngology and Otology

CONTENTS

	PAGE
THE INFLUENCE OF PRESSURE VARIATIONS ON THE IMPEDANCE OF THE HUMAN EAR DRUM. K. Terkildsen and K. A. Thomsen (Copenhagen)	409
LIPOMATA OF THE UPPER AIR AND FOOD PASSAGES. Alex R. Harper (Glasgow)	419
THE TREATMENT OF RECURRENT NASAL POLYPI BY INTERO-EXTERNAL "WINDOW" ETHMOIDECTORY. F. Boyes Korkis (London)	424
UVULOTOMY AMONG EAST AFRICAN TRIBES. J. F. Jarvis and S. N. Mwathi (Nairobi)	436
A NOTE ON THE DEVELOPMENT OF THE EXTERNAL EAR. T. G. Wilson (Dublin)	439
CLINICAL RECORD—	
Fatality Resulting from Intra-Nasal Polypectomy. Gilbert Forbes and Andrew Bradley (Sheffield)	445
CLINICAL NOTE—	
Tympano Dynamometry. A. Tumarkin (Liverpool)	448
SOCIETIES' PROCEEDINGS—	
Royal Society of Medicine—Section of Otology	451
GENERAL NOTES	488

For Advertisement space in this Journal apply to:
HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) *J. Laryng.*, 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of £10 per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., 109 Kingsway London, WC2.

5. Orders for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free, and is payable in advance.


9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol. LXIII may be purchased at 4s. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2.

United States of America

Orders for this *Journal* may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2, England.

**When a
hearing
aid is
advised . .**



**. . you may confidently
recommend your patients
to S. C. INGRAM**

- ★ They can try the different makes of hearing aids
- ★ They get impartial advice on all makes of hearing aids
- ★ Their personal needs are given individual and continued attention
- ★ A full report on each fitting recommended is sent to the patient's ear specialist
- ★ They can compare at one consultation the different makes one with another

S. C. INGRAM Independent Hearing Aid Consultant
On the approved List of the National Institute for the Deaf
2 SHEPHERD STREET, SHEPHERD MARKET, LONDON, W.1
 Hyde Park 9042


Please mention *The Journal of Laryngology and Otology* when replying to advertisements

EAR WAX

Removed this easy way

The removal of wax from the external auditory meatus has, in the past, normally entailed attendance by the patient for diagnosis and for the prescription of a suitable loosening agent, and a second attendance a few days later for syringing.

Now, by the use of Cerumol Ear Drops, wax can be removed in most cases at one visit. A few drops of Cerumol can be instilled into the ear and, while another patient is being attended to, the soft cerumen dissolves and the harder wax disimpacts. The wax can then be removed by gentle syringing or with cotton wool. The wax may even be found to run out of the ear on its own accord, in which case patients themselves may instil Cerumol at home, obviating further attendances. Cerumol is anti-bacterial, non-irritating and harmless to the lining of the external auditory meatus or the tympanic membrane. Cerumol is included in Category No. 4 of the M.O.H. classified list and may be prescribed on N.H.S. Form E.C.10.



CERUMOL
REGD. TRADE MARK

EAR DROPS

for the easier removal of wax

If you wish to test for yourself and have not received recently a 10. c.c. vial please write or telephone direct to:

LABORATORIES FOR APPLIED BIOLOGY LTD.,
91, AMHURST PARK, LONDON, N.16 Tel.: STA 2252

*PACKS For Surgery Use :
 10 c.c. vial — separate
 dropper included
 (Basic N.H.S. price 2/8)
 for Hospital Use : 2 oz.
 and 10 oz. bottles.*

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

In schools for the DEAF— multitone Telesonic INDUCTION LOOP SYSTEM brings new advantages

- 1 Can be installed in adjoining classrooms without risk of 'spill over'.
- 2 Combines good hearing with complete freedom of movement. Unaffected by distance between pupil and teacher.
- 3 Clear speech at high amplification.
- 4 Any number of pupils may use the same installation.
- 5 The child can use the instrument as a normal hearing aid outside school.



The Kindergarten Class at the London Residential School for Jewish Deaf Children where all classrooms have been installed with the Multitone Telesonic Induction Loop System

Write for further details to:


multitone

ELECTRIC COMPANY LTD., 12-20 UNDERWOOD STREET, LONDON, N.1.
CRC M23

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

**THE
HALLPIKE—**

**BLACKMORE
EAR
MICROSCOPE***



USES

DIAGNOSIS

The excellent optical performance of the Hallpike Ear Microscope gives the user greatly increased confidence in the diagnosis of abnormal conditions; and many disorders may be recognized at an earlier stage than has hitherto been possible with "headlamp and speculum" techniques. Thus slight pulsation of swellings and vascular tumours—of high diagnostic significance and almost impossible to detect by conventional methods—can readily be detected under this microscope.

Use of the Siegle attachment permits observation of small bubbles present in fluid collections within the tympanum; it is also invaluable for demonstrating minute perforations. Where defects of the tympanic membrane are present, the middle-ear cavity can be scrutinized minutely; the detailed structure of the stapes, stapedius tendon, round window and Eustachian orifice—not normally visible by ordinary methods of examination—are seen with quite remarkable clarity when the Ear Microscope is used.

TREATMENT

Manipulative procedures may be undertaken with greater confidence and precision if the Ear Microscope and its special instruments are used.

The curettes enable adherent wax and debris to be removed with the greatest precision from the walls of the deep meatus or from the tympanic membrane itself. Cholesteatomatous masses or neoplasms may be examined critically and biopsy specimens obtained by means of the curettes. These may also be used as delicate probes for palpation of the incus and head of the stapes in certain cases of otosclerosis with retraction and atrophy of the tympanic membrane; in this way unmistakable evidence may be obtained of bony fixation of the footplate. Or again, it is sometimes possible to identify by palpation defects of the bony tympanic ring or meatal wall which may underlie granulations.

Myringotomy or removal of foreign bodies can be undertaken with surprisingly high precision because the optics of the Ear Microscope and the specially angled myringotome blade eliminate all shadows so that the surgeon can see clearly where and what he is cutting. The outfit is invaluable also for post-operative examinations and treatment of mastoid cavities. Granulations can be cauterized and exposed areas of dura mater carefully palpated.

**J. Laryng.* (1953) 67, 108.

**39 WIGMORE STREET
LONDON W.1**



**5241 WHITBY AVENUE
PHILADELPHIA 43, Pa.,
U.S.A.**

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

PERDILATAL

**in the treatment of sudden
perceptive deafness
and Ménière's syndrome.**

* Vascular spasm may well be responsible for the majority of these cases.¹

* Vascular changes in the labyrinth are an important factor in the physiopathology of Ménière's disease.²

* In the treatment of Ménière's disease emphasis must be put upon early restoration of blood flow to the inner ear.³

* Perdilatal in doses of at least 6 mg. 4 times a day is the most effective vasodilator for the treatment of these conditions.⁴

1 Fowler, E. P. *Ann. Otorhinolaryng.*, 1950, 59, 980.

2 Atkinson, M. *Proc. roy. Soc. Med.*, 1946, 39, 807

3 Wilmot, T. J. *Brit. med. J.*, 1957, 2, 1047.

4 Wilmot, T. J. Paper read on February 6th, 1959 at a Meeting of the Section of Otology, Roy. Soc. Med.

Perdilatal is available in tablets of 6 mg. as PERDILATAL FORTE in 50's, 250's and 1,000's.



Smith & Nephew Pharmaceuticals Limited

WELWYN GARDEN CITY · HERTFORDSHIRE

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

POST-TONSILLECTOMY COMFORT

Immediate pain relief—Speedier Convalescence

The pain of traumatized tissues following tonsillectomy, demands its own relief—and points the need for analgesia that quickly reaches the irritated area.

ASPERGUM provides 'salivary analgesia' through the simple act of chewing—it brings pain-relieving acetylsalicylic acid into *intimate* and *prolonged* contact with the tonsillar

region, seldom reached even intermittently by gargling. The rhythmic stimulation of muscular action also aids in relieving local spasticity & stiffness: more rapid tissue repair is promoted. Each pleasantly flavoured chewing gum tablet provides $3\frac{1}{2}$ grains acetylsalicylic acid, permitting frequent use. Particularly suitable for children.

Aspergum (Category 3)

for more than two decades a dependable
and welcome aid to patient-comfort

Ethically promoted in packages of 16 tablets and moisture proof bottles of 36 and 250

WHITE LABORATORIES LTD., 428, SOUTHCROFT ROAD, LONDON, S.W.16

THE LARYNGOSCOPE

A Monthly Journal
devoted to the disease of
EAR, NOSE AND THROAT

*Official organ for the American Laryngological
Rhinological and Otological Society*

Price \$14.00 per year

Canada \$13.00 per year

MAX A. GOLDSTEIN, M.D.
FOUNDER

THEODORE E. WALSH, M.D.
EDITOR


640 SOUTH KINGSHIGHWAY
SAINT LOUIS 10, MO.

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

An outstanding accomplishment
in hearing aid design

Amplivox

"C" SERIES HEARING AIDS



FACILITIES

Individual fitting related to the Audiogram by means of selective compensation. Tone control giving low frequency attenuation combined with an overall response extending to 4000 c/s provides excellent speech discrimination in cases of perceptive deafness.

Two-stage adjustable automatic volume compression.

SPECIFICATION

4 stage transistor circuit. Stable operation from 0° to 40°C.

Miniature earphone with extended response. Magnetic microphone. Telephone inductor. Mercury cell battery—life 100 hours. Size $2\frac{1}{8}$ " x $1\frac{1}{16}$ " x $\frac{1}{32}$ " Weight 1 oz.


Mirror gold finish.

AMPLIVOX LTD . LONDON . ENGLAND

80 New Bond Street, London, W.1. Hyde Park 9888

WORLD-WIDE DISTRIBUTION AND SERVICE

Please mention *The Journal of Laryngology and Otology* when replying to advertisements



'OTOSPORIN' clears infected ears

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' brand Drops.

'Otosporin' is an aqueous suspension containing both neomycin (5 mgm. per c.c.) and polymyxin B sulphate (10,000 units per c.c.) which are together effective against practically all the bacteria found in ear infections. They are unlikely to induce bacterial resistance or skin sensitisation, and neither of them gives rise to cross-resistance or cross-sensitisation to penicillin and other antibiotics. Their effectiveness is visibly enhanced by the hydrocortisone (5 mgm. per c.c.) in 'Otosporin'; this, by reducing inflammation, not only relieves pain, but provides greater access for the antibiotics.

'OTOSPORIN'^{BRAND} drops

issued in bottles of 5 c.c.



BURROUGHS WELLCOME & CO. (The Wellcome Foundation Ltd.) **LONDON**

Please mention *The Journal of Laryngology and Otology* when replying to advertisements