

## PHASE TRANSITIONS FOR RANDOM GEOMETRIC PREFERENTIAL ATTACHMENT GRAPHS

JONATHAN JORDAN,\* *University of Sheffield*

ANDREW R. WADE,\*\* *Durham University*

### Abstract

Vertices arrive sequentially in space and are joined to existing vertices at random according to a preferential rule combining degree and spatial proximity. We investigate phase transitions in the resulting graph as the relative strengths of these two components of the attachment rule are varied.

Previous work of one of the authors showed that when the geometric component is weak, the limiting degree sequence mimics the standard Barabási–Albert preferential attachment model. We show that at the other extreme, in the case of a sufficiently strong geometric component, the limiting degree sequence mimics a purely geometric model, the on-line nearest-neighbour graph, for which we prove some extensions of known results. We also show the presence of an intermediate regime, with behaviour distinct from both the on-line nearest-neighbour graph and the Barabási–Albert model; in this regime, we obtain a stretched exponential upper bound on the degree sequence.

*Keywords:* Random spatial network; preferential attachment; on-line nearest-neighbour graph; degree sequence

2010 Mathematics Subject Classification: Primary 60D05

Secondary 05C80; 90B15

### 1. Introduction

Stochastic models in which a network evolves via the sequential addition of new nodes, each connected by an edge to an existing node in the graph according to some probabilistic rule, have been the subject of an explosion of interest over the past decade or so, motivated by real-world graphs such as those associated with social networks or the internet. The subject of this paper is a model whose connectivity rule combines degree-based preferential attachment with a spatial component; we describe our model in detail below. This model, previously studied in [11], is a variant of the geometric preferential attachment model of Flaxman *et al.* [7], [8], which itself can be viewed as a generalization of an earlier model of Manna and Sen [12]. A continuous time model with a similar flavour has recently been studied by Jacob and Mörters [9].

In a sense to be explained in this paper, the behaviour of the model studied here interpolates between pure preferential attachment (essentially the well-known Barabási–Albert model) and a purely geometric model (the on-line nearest-neighbour graph). It was shown in [11] that for a sufficiently weak geometric component of the attachment rule, the limiting degree distribution coincides with that of the Barabási–Albert model, which famously has a ‘scale-free’ or ‘power-law’ degree distribution [2], [10].

---

Received 19 November 2013; revision received 4 June 2014.

\* Postal address: School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK. Email address: jonathan.jordan@sheffield.ac.uk

\*\* Postal address: Department of Mathematical Sciences, Durham University, South Road, Durham DH1 2LE, UK.

The focus of the present paper is the complementary setting, in which the geometric component has a significant impact. We show that in the extreme case of a dominant geometric effect, the model behaves similarly to the on-line nearest-neighbour graph, which by contrast has a degree distribution with exponential tails (cf. [1]). We also study an intermediate regime in which the model behaves differently from both of the extreme cases, and in which the degree distribution satisfies a stretched exponential tail bound. Thus, we demonstrate the existence of nontrivial phase transitions.

In the next section we describe our models precisely and state our main results.

## 2. Random spatial graph models and main results

### 2.1. Notation

Write  $\mathbb{N} := \{1, 2, \dots\}$ ,  $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$ , and  $\mathbb{R}_+ := [0, \infty)$ . The vertices of our graphs are associated with *sites* in  $S \subset \mathbb{R}^d$ , assumed to be compact, convex, and of positive  $d$ -dimensional Lebesgue measure. The locations of the sites are  $X_0, X_1, \dots$ , independent random variables with density  $f$  supported on  $S$ . For  $n \in \mathbb{N}$ , set  $\mathcal{X}_n := \{X_0, \dots, X_n\}$ . We assume throughout that  $f$  is bounded away from 0 and  $\infty$ :

$$0 < \lambda_0 := \inf_{x \in S} f(x) \leq \sup_{x \in S} f(x) =: \lambda_1 < \infty. \quad (2.1)$$

We write  $\|\cdot\|$  for the Euclidean norm on  $\mathbb{R}^d$ , and  $\rho(x, y) = \|x - y\|$  for the Euclidean distance between  $x$  and  $y$  in  $\mathbb{R}^d$ . Denote by  $B(x; r)$  the open Euclidean  $d$ -ball centred at  $x \in \mathbb{R}^d$  with radius  $r > 0$ . Throughout we understand  $\log x$  to stand for  $\max\{0, \log x\}$ . Let  $\#A$  denote the number of elements of a finite set  $A$ .

### 2.2. On-line nearest-neighbour graph

The on-line nearest-neighbour graph (ONG) is constructed on points arriving sequentially in  $\mathbb{R}^d$  by connecting each point after the first to its nearest predecessor. The ONG is a natural and basic model of evolving spatial networks. It is a special case (or limiting case) of several models that have appeared in the literature, including a version of the Fabrikant, Koutsoupias and Papadimitriou (FKP) network model [1], [5] and geometric preferential attachment models such as [7], [11], and [12] (specifically, it is the ‘ $\alpha = -\infty$ ’ case of the model of Manna and Sen [12]); one contribution of the present paper is to explore this latter connection. The name ‘on-line nearest-neighbour graph’ was apparently introduced by Penrose [16].

In the ONG on  $(X_0, \dots, X_n)$ , the  $n$ th edge ( $n \in \mathbb{N}$ ) is between  $X_n$  and its almost surely (a.s.) unique nearest neighbour among  $\mathcal{X}_{n-1}$ . In other words, writing

$$\eta_1(n) := \arg \min_{i \in \{0, \dots, n-1\}} \rho(X_n, X_i), \quad (2.2)$$

the ONG on  $(X_0, \dots, X_n)$  consists of the edges  $(i, \eta_1(i))$  for  $1 \leq i \leq n$ ; it is natural to view these as directed edges when constructing the graph, but we largely treat them as undirected. We call  $X_{\eta_1(n)}$  the *on-line nearest neighbour* of  $X_n$ .

Let  $\deg_n(i)$  denote the degree of vertex  $i$  in the ONG on  $(X_0, \dots, X_n)$ . Let  $N_n^{\text{ONG}}(k)$  denote the number of vertices with degree at least  $k$  in the ONG on  $(X_0, \dots, X_n)$ :

$$N_n^{\text{ONG}}(k) = \sum_{i=0}^n \mathbf{1}\{\deg_n(i) \geq k\}.$$

We study the *asymptotic degree sequence*, i.e. the asymptotic proportion of vertices with degree at least  $k$  (for each  $k$ ); for convenience we work with  $n^{-1}N_n^{\text{ONG}}(k)$ .

Part of the statement of our main result on the ONG, Theorem 2.1 below, is that, for each  $k$ ,  $\lim_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)]$  exists; this was stated, apparently without proof, in [1, Section 2], but can be justified for the ONG using stabilization arguments of Penrose [16], as we explain in Section 6 below. Stabilization also provides an explicit description of the limit in terms of a version of the ONG defined on an infinite Poisson point process, as we describe next; in particular, the limit depends only on  $d$  and not on  $S$  or  $f$ .

Let  $\mathcal{H}$  be a unit-rate homogeneous Poisson point process on  $\mathbb{R}^d \times [0, 1]$ ; the  $[0, 1]$ -valued marks play the role of time in the finite ONG. For  $u \in [0, 1]$ , let  $\mathcal{H}_u := \mathcal{H} \cap (\mathbb{R}^d \times [0, u])$ , those Poisson points with marks in  $[0, u]$ . For  $x, y \in \mathbb{R}^d$  let  $B_x(y)$  be the open ball with centre  $y$  whose boundary includes  $x$ . Given  $x \in \mathbb{R}^d$  and  $u \in [0, 1]$ , let

$$\xi(x, u; \mathcal{H}) := 1 + \sum_{(y,v) \in \mathcal{H}, v > u} \mathbf{1}\{\mathcal{H}_v \cap (B_x(y) \times [0, 1]) = \{(y, v)\}\}.$$

By stabilization for the ONG (see [16]),  $\xi(x, u; \mathcal{H}) < \infty$  a.s. for any  $x \in \mathbb{R}^d$  and any  $u \in (0, 1)$ . We call  $\xi(x, u; \mathcal{H})$  the *degree* of  $(x, u)$  in the *infinite Poisson on-line nearest-neighbour graph*, defined locally by joining each point to the nearest Poisson point with mark equal to or less than the mark of the given point; note that  $(x, u)$  itself need not be in  $\mathcal{H}$ . Let  $U$  denote a uniform  $[0, 1]$  random variable, independent of  $\mathcal{H}$ .

**Theorem 2.1.** *Let  $d \in \mathbb{N}$ . Then for any  $k \in \mathbb{N}$ ,*

$$\lim_{n \rightarrow \infty} n^{-1} N_n^{\text{ONG}}(k) = \lim_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] = \mathbb{P}[\xi(0, U; \mathcal{H}) \geq k] =: \rho_k, \tag{2.3}$$

the first limit equality holding a.s. and in  $L^1$ . Here  $\rho_k \in [0, 1]$  are nonincreasing with  $\rho_1 = 1$ ,  $\lim_{k \rightarrow \infty} \rho_k = 0$ , and  $\sum_{k \in \mathbb{N}} \rho_k = 2$ . Moreover, there exist finite positive constants  $A, A', C$ , and  $C'$  such that, for all  $k \in \mathbb{N}$ ,

$$A'e^{-C'k} \leq \rho_k \leq Ae^{-Ck}, \tag{2.4}$$

and, more precisely,

$$\frac{1}{2} \log(1 + (2^{2d} - 1)^{-1}) \leq \liminf_{k \rightarrow \infty} (-k^{-1} \log \rho_k) \leq \limsup_{k \rightarrow \infty} (-k^{-1} \log \rho_k) \leq 1. \tag{2.5}$$

Finally, there exists a constant  $D < \infty$  for which, a.s., for all sufficiently large  $n$ ,

$$\max_{0 \leq i \leq n} \deg_n(i) \leq D \log n. \tag{2.6}$$

This result extends a result of Berger *et al.* [1]. Specifically, [1, Theorem 3] showed that

$$A'e^{-C'k} \leq \liminf_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \leq \limsup_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \leq Ae^{-Ck},$$

in the special case where  $d = 2$  and  $f$  is the indicator of the unit square  $S = (0, 1)^2$ . Our proof of Theorem 2.1, which we give in Section 6 below, is based in part on the proof of the analogous result in [1], with additional arguments required to obtain the existence of the limit and the almost-sure convergence in (2.3). Some extra work is also needed to obtain the quantitative bounds in (2.5): the  $d = 2$  case of the lower bound,  $\frac{1}{2} \log \frac{16}{15}$ , is contained in the argument of [1]; the other bounds are new.

TABLE 1: Estimated  $\mathbb{P}[\xi(0, U; \mathcal{H}) = k]$  (4dp) for  $1 \leq k \leq 10$ , for  $d \in \{1, 2, 100\}$ . For each  $d$ , the estimates are based on 500 simulations with  $n = 10^5$  for  $f$  uniform on the  $d$ -dimensional torus, to avoid finite-sample boundary effects.

	Degree ( $k$ )									
	1	2	3	4	5	6	7	8	9	10
$d = 1$	0.4728	0.2675	0.1394	0.0670	0.0304	0.0132	0.0056	0.0024	0.0001	0.0000
$d = 2$	0.4777	0.2636	0.1369	0.0668	0.0308	0.0137	0.0060	0.0026	0.0001	0.0000
$d = 100$	0.4999	0.2501	0.1250	0.0625	0.0312	0.0156	0.0078	0.0039	0.0002	0.0001

**Remark 2.1.** In view of (2.5), it is natural to conjecture that, for each  $d \in \mathbb{N}$ ,

$$\lim_{k \rightarrow \infty} (-k^{-1} \log \rho_k) = \mu(d) \text{ exists in } (0, 1];$$

the upper bound of 1 comes from (2.5). In [18, Section 7.6.5] it was conjectured that one might have  $\mu(d) = \mu = 1$ . The simpler, nonspatial, *uniform attachment* model in which vertex  $n$  is connected uniformly at random to a vertex from  $\{0, 1, \dots, n - 1\}$  leads to an analogous result with  $\mu = \log 2$ , as follows from [2, Section 4]. We think it unlikely that  $\mu(d) \in \{1, \log 2\}$  for any  $d \in \mathbb{N}$ ; we conjecture, however, that  $\lim_{d \rightarrow \infty} \mu(d) = \log 2$ . Simulations suggest that  $\mu(1) \approx 0.79$ ,  $\mu(2) \approx 0.77$ , and  $\mu(100) \approx 0.69$ ; cf. Table 1.

### 2.3. Geometric preferential attachment graph

The geometric preferential attachment (GPA) model that we study is as follows; often our notation coincides with [11]. We define a (random) sequence of finite graphs  $G_n = (V_n, E_n)$ ,  $n \in \mathbb{N}$ . The vertex set of  $G_n$  is  $V_n = \{0, 1, \dots, n\}$ . For  $v \in V_n$ , we denote by  $\text{deg}_n(v)$  the degree of  $v$  in the GPA graph  $G_n$  (viewed as an undirected graph); this notation is the same as for degrees in the ONG, but the graph under consideration will be clear in context.

The construction uses an *attractiveness function*  $F: (0, \infty) \rightarrow (0, \infty)$ . Recall that  $X_0, X_1, \dots$  are random sites in  $S$ . For simplicity, we start with  $G_1 = (V_1, E_1)$  consisting of vertices with labels 0 and 1 joined by a single edge, so  $V_1 = \{0, 1\}$  and  $E_1 = \{(1, 0)\}$ . Vertices 0 and 1 are associated with sites  $X_0$  and  $X_1$  in  $S$ , respectively.

We proceed via iterated addition of vertices to construct  $G_{n+1} = (V_{n+1}, E_{n+1})$  from  $G_n = (V_n, E_n)$ ,  $n \in \mathbb{N}$ . Given  $G_n$ ,  $n \in \mathbb{N}$ , and the spatial locations  $\mathcal{X}_n$  of its vertices, we add a vertex with label  $n + 1$  at site  $X_{n+1} \in S$ , and we add a new edge  $(n + 1, v_{n+1})$  where  $v_{n+1}$  is chosen randomly from  $V_n$  with distribution specified by

$$\mathbb{P}[v_{n+1} = v \mid G_n, \mathcal{X}_{n+1}] = \frac{\text{deg}_n(v)F(\rho(X_v, X_{n+1}))}{D_n(X_{n+1})}, \quad v \in V_n, \tag{2.7}$$

where for  $n \in \mathbb{N}$  and  $x \in S$ ,  $D_n(x) := \sum_{v \in V_n} \text{deg}_n(v)F(\rho(X_v, x))$ .

We call  $G_n$  so constructed a GPA graph with attractiveness function  $F$ . In [11], it was assumed that  $\int_S F(\rho(x, y))f(y) dy < \infty$ , so  $F$  cannot blow up too rapidly at 0. In this paper, our primary interest is in  $F$  for which this condition is not satisfied.

### 2.4. The strong geometric regime

For  $\gamma > 1$ , define  $F_\gamma$  for  $r > 0$  by  $F_\gamma(r) := \exp\{(\log(1/r))^\gamma\}$ . Note that  $F_\gamma(r)$  blows up at 0 faster than  $r^{-s}$  for any power  $s$ . Recall that the convention  $\log x \equiv \max\{0, \log x\}$  is in force,

so  $F_\gamma(r) = 1$  for  $r \geq 1$ . Also,  $F_\gamma(r)$  is strictly decreasing for  $r \in (0, 1)$ , with  $F_\gamma(r) \rightarrow \infty$  as  $r \downarrow 0$ .

Our main result in this setting (i) gives an almost-sure degree bound analogous to (2.6) above for the ONG, and (ii) shows that the limiting degree sequence for the GPA graph is the same as for the ONG, for a strong enough geometric component to the interaction (under the condition  $\gamma > \frac{3}{2}$ ). Let  $N_n^{\text{GPA}}(k)$  denote the number of vertices with degree at least  $k$  in the GPA graph  $G_n$ .

**Theorem 2.2.** *Suppose that  $F = F_\gamma$  for some  $\gamma > 1$ .*

(i) *For any  $\nu \in (0, 1)$  with  $\nu > 2 - \gamma$ , a.s. for all sufficiently large  $n$ ,*

$$\max_{0 \leq i \leq n} \text{deg}_n(i) \leq \exp\{(\log n)^\nu\}. \tag{2.8}$$

(ii) *Suppose that  $\gamma > \frac{3}{2}$ . Then  $\lim_{n \rightarrow \infty} \mathbb{P}[v_n = \eta_1(n)] = 1$  and*

$$\lim_{n \rightarrow \infty} n^{-1} \mathbb{E} \sum_{i=1}^n \mathbf{1}\{v_i \neq \eta_1(i)\} = 0. \tag{2.9}$$

Moreover, for any  $k \in \mathbb{N}$ ,

$$\lim_{n \rightarrow \infty} n^{-1} N_n^{\text{GPA}}(k) = \lim_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{GPA}}(k)] = \rho_k, \tag{2.10}$$

the first limit equality holding in  $L^1$ , where  $\rho_k$  is as in Theorem 2.1.

We give the proof of Theorem 2.2 in Section 4.

**Conjecture 2.1.** *We suspect the conclusion of Theorem 2.2(ii) to hold for any  $\gamma > 1$ .*

**2.5. The intermediate regime: power-law attractiveness**

Take  $F(r) = r^{-s}$  for  $s \in (0, \infty)$ . The next result contrasts with (2.9) in the strong geometric attraction regime, and shows that in this case, in expectation, there is a nonnegligible proportion of vertices not connecting to their nearest neighbour.

**Theorem 2.3.** *Suppose that  $F(r) = r^{-s}$  for  $s \in (0, \infty)$ . Then  $\limsup_{n \rightarrow \infty} \mathbb{P}[v_n = \eta_1(n)] < 1$  and*

$$\liminf_{n \rightarrow \infty} n^{-1} \mathbb{E} \sum_{i=1}^n \mathbf{1}\{v_i \neq \eta_1(i)\} > 0. \tag{2.11}$$

It was proved in Theorem 2.1 of [11] that in the  $s \in (0, d)$  case, under certain conditions on  $S$  and  $f$ , the degree distribution of the GPA graph converges to a power-law distribution, as in the Barabási–Albert model:  $\lim_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{GPA}}(k)] = r_k$  where  $r_k \sim 2k^{-2}$  as  $k \rightarrow \infty$ . In contrast, the next result gives a stretched exponential upper bound for the tail of the degree distribution when  $s > d$ .

**Theorem 2.4.** *Suppose that  $F(r) = r^{-s}$  for  $s > d$ . For any  $\gamma \in (0, (s - d)/(2s - d))$ , there exists a constant  $C < \infty$  such that, for all  $k$ ,*

$$\limsup_{n \rightarrow \infty} n^{-1} N_n^{\text{GPA}}(k) \leq C e^{-k^\gamma}, \quad \text{a.s. and} \quad \limsup_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{GPA}}(k)] \leq C e^{-k^\gamma}.$$

This result confirms the presence of a phase transition in the character of the degree distribution at  $s = d$ , as intimated in [11, Section 5] and in line with  $d \in \{1, 2\}$  simulation results of [12] (who actually conjectured that the phase transition point was  $s = d - 1$ ) and [6] (who did suggest  $s = d$  for the transition). The stretched exponential for  $s > d$  is also consistent with the observations of [6] and [12]. We remark that as  $s \rightarrow \infty$ , Theorem 2.4 provides an upper bound of order almost  $e^{-\sqrt{k}}$ .

The rest of the paper is organized as follows. In Section 3 we collect some preparatory lemmas. In Section 4 we deal with the strong geometric regime, culminating in the proof of Theorem 2.2. In Section 5 we deal with the case of power-law attractiveness, and present the proofs of Theorems 2.3 and 2.4. Finally, in Section 6 we deal with the ONG and present the proof of Theorem 2.1.

### 3. Preliminaries to the proofs

First we state a basic property of the set  $S$ , under our standing assumptions. Let  $\omega_d$  be the volume of the unit-radius Euclidean  $d$ -ball, and set  $\text{diam}(S) := \sup_{x,y \in S} \rho(x, y)$ .

**Lemma 3.1.** *There exists  $\delta_S > 0$  such that, for all  $r \in [0, \text{diam}(S)]$ ,*

$$\inf_{x \in S} |B(x; r) \cap S| \geq \delta_S \omega_d r^d.$$

*Proof.* Since  $S$  is convex, compact, and of positive measure, there exist  $x_0 \in S$  and  $r_0 > 0$  such that  $B(x_0; r_0)$  is contained in the interior of  $S$ . It suffices to suppose that either (i)  $\rho(x, x_0) \geq 2r_0$ , or (ii)  $\rho(x, x_0) \leq r_0/2$ . To see this, suppose that  $r_0/2 < \rho(x, x_0) < 2r_0$ . Then we may carry out the argument for case (i) after having replaced  $r_0$  by  $r_0/4$ , introducing only a constant multiplicative factor into the argument.

So now suppose that (i) holds. For  $r \leq r_0$ , let  $C(x, r)$  denote the cone with apex  $x$ , axis passing through  $x_0$ , and half-angle  $\theta(x, r) = \sin^{-1}(r/\rho(x, x_0))$ . Since  $\rho(x, x_0) \leq \text{diam}(S)$ ,  $\theta(x, r) \geq \theta(r) := \sin^{-1}(r/\text{diam}(S))$ . By construction and convexity of  $S$ ,  $C(x, r) \cap S$  contains the cone segment  $\{y \in C(x, r) : \rho(x, y) \leq \rho(x, x_0) \cos \theta(x, r)\}$ . So, if  $\rho(x, x_0) \geq 2r_0$ , then  $B(x; r) \cap S$  contains the cone segment  $\{y \in C(x, r) : \rho(x, y) \leq r \wedge r_0\}$ , which has volume bounded below by  $c_d \theta(r)^{d-1} r$ , provided  $r \leq r_0$ , where  $c_d > 0$  is an absolute constant. Hence,  $|B(x; r) \cap S|$  is bounded below by a constant multiplied by  $r^d$ , for all  $r \leq r_0$ . The same conclusion follows if  $r \in (r_0, \text{diam}(S))$ , using the lower bound  $c_d \theta(r_0)^{d-1} r_0 \geq c'_d (r_0/\text{diam}(S))^d r^d$  for  $c'_d > 0$  not depending on  $r$ .

Finally, in case (ii), we have that  $B(x; r) \cap S$  contains the ball  $B(x; r \wedge (r_0/2))$ , and a similar argument to that for part (i) completes the proof.

We next present some basic results on nearest-neighbour distances. For  $n \in \mathbb{N}$ , let

$$Z_n := \rho(X_n; \mathcal{X}_{n-1}) := \min_{0 \leq i \leq n-1} \rho(X_n, X_i) = \rho(X_n, X_{\eta_1(n)}),$$

the distance from  $X_n$  to its on-line nearest neighbour. Write  $x^+ := x \mathbf{1}\{x > 0\}$ .

**Lemma 3.2.** *Let  $\delta_S > 0$  be the constant in Lemma 3.1. Then for  $r > 0$ ,*

$$\mathbb{P}[Z_n \geq r] \leq (1 - \delta_S \lambda_0 \omega_d r^d)^n \mathbf{1}\{r \leq \text{diam}(S)\}; \tag{3.1}$$

$$\mathbb{P}[Z_n \geq r] \geq ((1 - \lambda_1 \omega_d r^d)^+)^n \mathbf{1}\{r \leq \text{diam}(S)\}. \tag{3.2}$$

*Proof.* Conditional on  $X_n$ , we have, for any  $r > 0$ , a.s.,

$$\begin{aligned} \mathbb{P}[Z_n \geq r \mid X_n] &= \mathbb{P}[S \cap B(X_n; r) \cap \mathcal{X}_{n-1} = \emptyset \mid X_n] \\ &= \left(1 - \int_{S \cap B(X_n; r)} f(x) \, dx\right)^n. \end{aligned} \tag{3.3}$$

Note that  $\mathbb{P}[Z_n > \text{diam}(S)] = 0$ , so it suffices to suppose that  $r \leq \text{diam}(S)$ . Then,

$$\delta_S \omega_d r^d \leq |S \cap B(X_n; r)| \leq \omega_d r^d, \text{ a.s.}, \tag{3.4}$$

for all  $r \leq \text{diam}(S)$ , by Lemma 3.1. It follows from (3.3) and (2.1) that  $\mathbb{P}[Z_n \geq r \mid X_n] \leq (1 - \lambda_0 |S \cap B(X_n; r)|)^n$ , which, with the first inequality in (3.4), gives (3.1). Similarly, (3.2) follows from (3.3) and (2.1) with the second inequality in (3.4).

Next we state a simple but useful result on degrees, which is basically Markov’s inequality. Write  $N_n(k)$  for  $N_n^{\text{GPA}}(k)$  or  $N_n^{\text{ONG}}(k)$ . Either graph is a tree, so using the degree sum formula and interchanging the order of summation we obtain

$$2n = \sum_{i=0}^n \text{deg}_n(i) = \sum_{i=0}^n \sum_{k \geq 1} \mathbf{1}\{\text{deg}_n(i) \geq k\} = \sum_{k \geq 1} N_n(k).$$

So for any  $k_0 \in \mathbb{N}$ ,  $2n \geq \sum_{k=1}^{k_0} N_n(k) \geq k_0 N_n(k_0)$ , since  $N_n(\cdot)$  is nonincreasing. Thus,

$$N_n(k) \leq \frac{2n}{k}, \text{ a.s.}, \text{ for any } k \text{ and } n. \tag{3.5}$$

Finally, we introduce some notation for dealing with conditional probabilities. Let  $\mathcal{F}_n := \sigma(\mathcal{X}_n, v_2, v_3, \dots, v_{n-1})$ , the  $\sigma$ -algebra generated by the sites up to and including  $X_n$  and by the edge choices made on previous steps. Then  $\mathcal{F}_n$  contains all the information about  $G_{n-1}$  as well as  $X_0, \dots, X_n$ , and (2.7) can be expressed as

$$\mathbb{P}[v_n = v \mid \mathcal{F}_n] = \frac{\text{deg}_{n-1}(v)F(\rho(X_v, X_n))}{D_{n-1}(X_n)}, \quad v \in \{0, \dots, n-1\}. \tag{3.6}$$

Also set  $\tilde{\mathcal{F}}_n := \sigma(\mathcal{X}_n, v_2, v_3, \dots, v_{n-1}, v_n)$ , which adds to  $\mathcal{F}_n$  information about  $G_n$ .

### 4. Proofs for the strong geometric regime

In this section we prove our results from Section 2.4. First we outline the central idea of the proof of Theorem 2.2, to show that  $X_n$  joins to its on-line nearest neighbour  $X_{\eta_1(n)}$  with probability  $1 - o(1)$  (cf. Lemma 4.3). By (3.6), this probability satisfies

$$\mathbb{P}[v_n = \eta_1(n) \mid \mathcal{F}_n] = \frac{\text{deg}_{n-1}(\eta_1(n))F(Z_n)}{D_{n-1}(X_n)}.$$

For  $F = F_\gamma$ , the fact that  $F_\gamma$  is decreasing and the degree sum formula gives

$$D_{n-1}(X_n) = \sum_{i=0}^{n-1} \text{deg}_{n-1}(i)F_\gamma(\rho(X_i, X_n)) \leq 2nF_\gamma(W_n) + \text{deg}_{n-1}(\eta_1(n))F_\gamma(Z_n),$$

where  $W_n$  is the distance from  $X_n$  to its *second* nearest neighbour among  $\mathcal{X}_{n-1}$ , so

$$\mathbb{P}[v_n = \eta_1(n) \mid \mathcal{F}_n] \geq 1 - \frac{2nF_\gamma(W_n)}{F_\gamma(Z_n)}.$$

With probability  $1 - o(1)$ ,  $W_n > Z_n + \theta_n$  where  $\theta_n = o(n^{-1/d})$ , so to show  $\mathbb{P}[v_n = \eta_1(n)] = 1 - o(1)$  it suffices to show that, as  $n \rightarrow \infty$ ,

$$\frac{nF_\gamma(Z_n + \theta_n)}{F_\gamma(Z_n)} \rightarrow 0, \text{ in probability.}$$

Taylor’s formula shows that this holds provided  $\gamma > 2$ . Further progress requires control of both the vertex degrees and the number of ‘plausible alternatives’ for  $v_n$ .

Theorem 2.2(i) gives sufficient control of degrees to achieve the  $\gamma > 3/2$  case of Theorem 2.2(ii). To achieve  $\gamma > 1$  (cf. Conjecture 2.1) seems to need tighter control, and a technique that enables one to replace almost-sure upper bounds growing with  $n$  by ‘typical’ statistics, as might be available given some suitable stabilization property, such as that enjoyed by ONG. This seems to be a challenging problem.

For  $\nu \in (0, 1)$  and  $n \geq 2$  set  $\beta(n, \nu) := n^{-1/d} \exp\{(\log n)^\nu\}$ , and let

$$E(n, \nu) := \{\rho(X_{v_n}, X_n) \geq \beta(n, \nu)\},$$

the event that the edge from vertex  $n$  connects to any vertex outside  $B(X_n; \beta(n, \nu))$ .

**Lemma 4.1.** *Suppose that  $F = F_\gamma$  for some  $\gamma > 1$  and that  $\nu \in (0, 1)$  with  $\nu > 2 - \gamma$ . Then for any  $p < \infty$ , as  $n \rightarrow \infty$ ,*

$$\mathbb{P}[E(n, \nu)] = O(\exp\{-\gamma d^{1-\gamma} (1 + o(1))(\log n)^{\gamma+\nu-1}\}) = O(n^{-p}).$$

*Proof.* Note that for any  $\nu \in (0, 1)$ ,

$$F_\gamma(\beta(n, \nu)) = \exp\{d^{-\gamma} (\log n)^\gamma - \gamma d^{1-\gamma} (1 + o(1))(\log n)^{\gamma+\nu-1}\}. \tag{4.1}$$

We obtain from (3.1) and (2.1) that

$$\mathbb{P}[Z_n > \beta(n, \nu)] = O(\exp\{-\delta_S \lambda_0 \omega_d \exp\{d(\log n)^\nu\}\}) = O(\exp\{-(\log n)^K\}), \tag{4.2}$$

for any  $K < \infty$ . Fix  $\nu \in (0, 1)$  and choose  $\nu' \in (0, \nu)$ . Then

$$\mathbb{P}[E(n, \nu)] \leq \mathbb{P}[Z_n > \beta(n, \nu')] + \mathbb{P}[E(n, \nu) \mid Z_n \leq \beta(n, \nu')]. \tag{4.3}$$

Suppose that  $Z_n \leq \beta(n, \nu')$ . Then, since  $F_\gamma(r)$  is nonincreasing in  $r > 0$ ,

$$\deg_{n-1}(\eta_1(n))F_\gamma(\rho(X_{\eta_1(n)}, X_n)) \geq F_\gamma(Z_n) \geq F_\gamma(\beta(n, \nu')),$$

so that  $D_{n-1}(X_n) \geq F_\gamma(\beta(n, \nu'))$ , given  $Z_n \leq \beta(n, \nu')$ . On the other hand, any vertex  $j < n$  with  $X_j \notin B(X_n; \beta(n, \nu))$  has  $\deg_{n-1}(j)F_\gamma(\rho(X_j, X_n)) \leq nF_\gamma(\beta(n, \nu))$ , using the crude bound  $\deg_{n-1}(j) \leq n$ . Hence, by (2.7) and (4.1),

$$\begin{aligned} &\mathbb{P}[E(n, \nu) \mid Z_n \leq \beta(n, \nu')] \\ &= \sum_{j=0}^{n-1} \mathbb{P}[\{v_n = j\} \cap E(n, \nu) \mid Z_n \leq \beta(n, \nu')] \\ &\leq \frac{n^2 F_\gamma(\beta(n, \nu))}{F_\gamma(\beta(n, \nu'))} \end{aligned}$$

$$\begin{aligned}
 &= O(\exp\{2 \log n - \gamma d^{1-\gamma}(1 + o(1))((\log n)^{\gamma+\nu-1} - (\log n)^{\gamma+\nu'-1})\}) \\
 &= O(\exp\{-\gamma d^{1-\gamma}(1 + o(1))(\log n)^{\gamma+\nu-1}\}), \tag{4.4}
 \end{aligned}$$

provided that  $\gamma + \nu - 1 > 1$ , i.e.  $\nu > 2 - \gamma$ , which we can ensure by choosing  $\nu \in (0, 1)$  close enough to 1 since  $\gamma > 1$ . The result now follows from (4.2), (4.3), and (4.4).

The next result is a bound on degrees that amounts to Theorem 2.2(i), and which will also be an ingredient in our proof of Theorem 2.2(ii).

**Lemma 4.2.** *Suppose that  $F = F_\gamma$  for some  $\gamma > 1$ . Then for any  $\nu \in (0, 1)$  with  $\nu > 2 - \gamma$ , a.s., for all but finitely many  $n \in \mathbb{N}$ , (2.8) holds.*

*Proof.* Let  $\gamma > 1$  and  $\nu > 2 - \gamma$ . By Lemma 4.1,  $\mathbb{P}[E(j, \nu)] = O(j^{-2})$ . Hence, by the Borel–Cantelli lemma, for only finitely many  $j \in \mathbb{N}$  does the vertex  $j$  connect to a vertex  $i < j$  with  $\rho(X_i, X_j) \geq \beta(j, \nu)$ . It follows that there exists some finite random variable  $D_\nu = 1 + \sum_{j=1}^\infty \mathbf{1}(E(j, \nu))$  such that, for all  $n \in \mathbb{N}$  and all  $i \in \{0, 1, \dots, n\}$ ,

$$\deg_n(i) \leq D_\nu + \sum_{j=i+1}^n \xi_{i,j},$$

where we set  $\xi_{i,j} := \mathbf{1}\{\rho(X_j, X_i) \leq \beta(j, \nu)\}$  for  $i \neq j$  and  $\xi_{i,i} := 0$ . Hence,

$$\max_{0 \leq i \leq n} \deg_n(i) \leq D_\nu + \max_{0 \leq i \leq n} \sum_{j=1}^n \xi_{i,j}. \tag{4.5}$$

For fixed  $i$ , conditional on  $X_i$ , the  $n - 1$  terms  $\xi_{i,j}$  with  $j \neq i$  in the sum on the right-hand side of (4.5) are independent and  $\{0, 1\}$ -valued, and a suitable version of Talagrand’s inequality (see, e.g. [14, p. 81]) will show that their sum is concentrated around its mean (in fact, we only need an upper bound here). Specifically, for  $n \in \mathbb{N}$ ,

$$\mathbb{E} \sum_{j=1}^n \xi_{i,j} = \sum_{j=1}^n \mathbb{P}[X_i \in B(X_j; \beta(j, \nu))] = \Theta\left(\sum_{j=1}^n \beta(j, \nu)^d\right), \tag{4.6}$$

uniformly for  $i \in \{1, \dots, n\}$ , where the implicit constants depend on  $S$ ,  $\lambda_0$  and  $\lambda_1$  (we use Lemma 3.1 here). We claim that

$$\sum_{j=1}^n \beta(j, \nu)^d = \exp\{d(\log n)^\nu(1 + o(1))\}. \tag{4.7}$$

To verify (4.7), we combine the upper and lower bounds

$$\sum_{j=1}^n \beta(j, \nu)^d \leq \sum_{j=1}^n \frac{1}{j} \exp\{d(\log n)^\nu\} \leq (1 + \log n) \exp\{d(\log n)^\nu\},$$

and

$$\sum_{j=1}^n \beta(j, \nu)^d \geq \sum_{j=\lceil n/2 \rceil}^n \frac{1}{n} \exp\left\{d\left(\log\left(\frac{n}{2}\right)\right)^\nu\right\} \geq \frac{1}{2} \exp\left\{d\left(\log\left(\frac{n}{2}\right)\right)^\nu\right\}.$$

From (4.6) and (4.7), we have  $\mathbb{E} \sum_{j=1}^n \xi_{i,j} = \exp\{d(\log n)^\nu(1+o(1))\}$ . Talagrand’s inequality implies that for all  $n$ ,

$$\max_{0 \leq i \leq n} \mathbb{P} \left[ \sum_{j=1}^n \xi_{i,j} > \exp\{2d(\log n)^\nu\} \right] \leq O(\exp\{-e^{d(\log n)^\nu}\}),$$

which is  $O(n^{-3})$ , say, so that Boole’s inequality yields

$$\mathbb{P} \left[ \max_{0 \leq i \leq n} \sum_{j=1}^n \xi_{i,j} > \exp\{2d(\log n)^\nu\} \right] = O(n^{-2}).$$

Now, another application of the Borel–Cantelli lemma together with (4.5) completes the proof of the lemma, noting that  $\nu > 2 - \gamma$  was arbitrary.

The main step remaining in the proof of Theorem 2.2 is the following.

**Lemma 4.3.** *Suppose that  $F = F_\gamma$  for  $\gamma > \frac{3}{2}$ . Then  $\mathbb{P}[v_n \neq \eta_1(n)] \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* Take a sequence  $\theta_n > 0$  with  $\theta_n = o(n^{-1/d})$ , and, given  $X_n$  and  $Z_n$ , define the shells  $A_n := B(X_n; Z_n + \theta_n) \setminus B(X_n; Z_n)$ . Let  $a_n := \#(A_n \cap \mathcal{X}_{n-1} \setminus \{X_{\eta_1(n)}\})$ , the number of predecessors to  $X_n$ , other than its on-line nearest neighbour, inside  $A_n$ .

Conditional on  $X_n$  and  $Z_n$ , the points of  $\mathcal{X}_{n-1} \setminus \{X_{\eta_1(n)}\}$  are independent and identically distributed (i.i.d.) on  $S \setminus B(X_n; Z_n)$  with conditional distribution given for measurable  $\Gamma \subseteq S \setminus B(X_n; Z_n)$  by  $\mathbb{P}[\cdot \in \Gamma] = \int_\Gamma g_n(x) dx$ , where

$$g_n(x) = \frac{f(x)}{\mathbb{P}[X_0 \in S \setminus B(X_n; Z_n) \mid X_n, Z_n]}.$$

Note that, a.s.,

$$\mathbb{P}[X_0 \in S \setminus B(X_n; Z_n) \mid X_n, Z_n] = 1 - \int_{S \cap B(X_n; Z_n)} f(x) dx \geq 1 - \lambda_1 \omega_d Z_n^d \geq \frac{1}{2},$$

provided  $Z_n \leq (2\lambda_1 \omega_d)^{-1/d}$ . Moreover,  $S \cap A_n$  has volume bounded above by

$$\omega_d(Z_n + \theta_n)^d - \omega_d Z_n^d \leq C_d \theta_n (\theta_n^{d-1} + Z_n^{d-1})$$

for some  $C_d < \infty$  depending only on  $d$ . Hence, conditional on  $X_n$  and  $Z_n$ , each of the  $n - 1$  points  $X_0, \dots, X_{n-1}$ , excluding  $X_{\eta_1(n)}$ , lands in  $A_n$  with probability at most

$$\frac{\int_{S \cap A_n} f(x) dx}{\mathbb{P}[X_0 \in S \setminus B(X_n; Z_n) \mid X_n, Z_n]} \leq 2\lambda_1 C_d \theta_n (\theta_n^{d-1} + Z_n^{d-1}) + \mathbf{1}\{Z_n > (2\lambda_1 \omega_d)^{-1/d}\}.$$

It follows that

$$\mathbb{E}[a_n \mid Z_n] \leq 2\lambda_1 C_d n \theta_n (\theta_n^{d-1} + Z_n^{d-1}) + n \mathbf{1}\{Z_n > (2\lambda_1 \omega_d)^{-1/d}\}.$$

Taking expectations and using (3.1) we have  $n\mathbb{P}[Z_n > (2\lambda_1 \omega_d)^{-1/d}] = o(1)$ , while, for any  $\alpha > 0$ , by another application of (3.1), for some  $C < \infty$ ,

$$\mathbb{E}[Z_n^\alpha] = \int_0^\infty \mathbb{P}[Z_n > r^{1/\alpha}] dr \leq \int_0^\infty \exp\{-Cnr^{d/\alpha}\} dr,$$

which gives  $\mathbb{E}[Z_n^\alpha] = O(n^{-\alpha/d})$ . Hence,  $\mathbb{E}[a_n] = O(\theta_n^d n) + O(\theta_n n^{1/d}) + o(1)$ , which is  $o(1)$  provided  $\theta_n = o(n^{-1/d})$ , so that, by Markov's inequality,  $\mathbb{P}[a_n > 0] \leq \mathbb{E}[a_n] = o(1)$ .

Now we condition on the whole of  $\mathcal{F}_n$ . Again take  $\beta(n, \nu) = n^{-1/d} \exp\{(\log n)^\nu\}$ . Let  $E'_n$  denote the event that  $X_n$  is joined to a point outside  $B(X_n; Z_n + \theta_n)$ :

$$E'_n := \{\rho(X_{v_n}, X_n) \geq Z_n + \theta_n\}.$$

Also, for a constant  $b > 1$  (which we will later choose to be large), set

$$E''_n := \{Z_n \leq b^{-1}n^{-1/d}\} \cup \{Z_n \geq bn^{-1/d}\}.$$

Finally, define the event (for another constant  $C$  to be chosen later)

$$E'''_n := \{\#(\mathcal{X}_{n-1} \cap B(X_n; \beta(n, \nu))) \geq C \exp\{d(\log n)^\nu\}\}.$$

The ball  $B(X_n; \beta(n, \nu))$  has volume bounded above by  $\omega_d n^{-1} \exp\{d(\log n)^\nu\}$ . The events  $\{X_j \in B(X_n; \beta(n, \nu))\}$ ,  $0 \leq j \leq n - 1$  are independent each with probability at most  $\lambda_1 \omega_d n^{-1} \exp\{d(\log n)^\nu\}$ , so  $\#(\mathcal{X}_{n-1} \cap B(X_n; \beta(n, \nu)))$  is stochastically dominated by a binomial  $(n, \lambda_1 \omega_d n^{-1} \exp\{d(\log n)^\nu\})$  random variable. Standard binomial tail bounds show that, for an appropriate  $C < \infty$ ,  $\mathbb{P}[E'''_n] = o(1)$ .

Since  $\{a_n = 0\} \cap (E'_n)^c$  implies that  $X_n$  is joined to its on-line nearest neighbour,

$$\begin{aligned} \mathbb{P}[v_n \neq \eta_1(n) \mid \mathcal{F}_n] &\leq \mathbb{P}[E'_n \mid \mathcal{F}_n] \mathbf{1}\{a_n = 0\} \cap (E''_n)^c \cap (E'''_n)^c \\ &\quad + \mathbf{1}\{a_n > 0\} + \mathbf{1}(E''_n) + \mathbf{1}(E'''_n). \end{aligned} \tag{4.8}$$

For any  $\varepsilon > 0$ , Lemma 3.2 shows that we can choose  $b$  and  $n_0$  sufficiently large so that  $\mathbb{P}[E''_n] < \varepsilon$  for all  $n \geq n_0$ . We have already seen that  $\mathbb{P}[a_n > 0] = o(1)$  and  $\mathbb{P}[E'''_n] = o(1)$ . We also claim that

$$\mathbb{P}[E'_n \mid \mathcal{F}_n] \mathbf{1}\{a_n = 0\} \cap (E''_n)^c \cap (E'''_n)^c = o(1), \text{ a.s.} \tag{4.9}$$

The bounded convergence theorem implies that the expectation of this last quantity is also  $o(1)$ , so taking expectations in (4.8) we see that for any  $\varepsilon > 0$ , we may choose  $b$  such that  $\limsup_{n \rightarrow \infty} \mathbb{P}[v_n \neq \eta_1(n)] \leq \varepsilon$ . This gives the statement in the lemma.

It remains to prove the claim (4.9). First, we note that

$$D_{n-1}(X_n) \geq \deg_{n-1}(\eta_1(n)) F_\gamma(\rho(X_{\eta_1(n)}, X_n)) \geq F_\gamma(Z_n).$$

On the other hand, on  $\{a_n = 0\}$ , any alternative  $X_j$  to  $X_{\eta_1(n)}$  among  $\mathcal{X}_{n-1}$  is at distance at least  $Z_n + \theta_n$  from  $X_n$ , so that for  $j \neq \eta_1(n)$ ,

$$\deg_{n-1}(j) F_\gamma(\rho(X_j, X_n)) \leq \exp\{(\log n)^\nu\} F_\gamma(Z_n + \theta_n), \text{ a.s.,}$$

for all  $n$  large enough, by Lemma 4.2, provided  $\nu \in (0, 1)$  with  $\nu > 2 - \gamma$ .

On  $(E'''_n)^c \cap \{a_n = 0\}$ , the contribution of points inside  $B(X_n; \beta(n, \nu))$ , other than  $X_{\eta_1(n)}$ , to  $D_{n-1}(X_n)$  is bounded above by  $C \exp\{2d(\log n)^\nu\} F_\gamma(Z_n + \theta_n)$ , since there are at most  $O(\exp\{d(\log n)^\nu\})$  of these points, their degrees are at most  $O(\exp\{(\log n)^\nu\})$ , a.s., by Lemma 4.2, and they are all at distance at least  $Z_n + \theta_n$  from  $X_n$ . Moreover, similarly to as in the proof of Lemma 4.1, the contribution to  $D_{n-1}(X_n)$  from any points outside  $B(X_n; \beta(n, \nu))$  is at most  $n^2 F_\gamma(\beta(n, \nu))$ .

So from (3.6) we have, on  $\{a_n = 0\} \cap (E''_n)^c$ , for all  $n$  large enough,

$$\mathbb{P}[E'_n \mid \mathcal{F}_n] \leq \frac{C \exp\{2d(\log n)^\nu\} F_\gamma(Z_n + \theta_n) + n^2 F_\gamma(\beta(n, \nu))}{F_\gamma(Z_n)}.$$

Here, similarly to (4.4),

$$\frac{n^2 F_\gamma(\beta(n, \nu))}{F_\gamma(Z_n)} = O(\exp\{-c(\log n)^{\gamma+\nu-1}\}),$$

for some  $c > 0$ , as long as  $\nu > 2 - \gamma$ . Also, we have that, on  $(E''_n)^c$ ,

$$\begin{aligned} \frac{F_\gamma(Z_n + \theta_n)}{F_\gamma(Z_n)} &= \exp\left\{\left(\log\left(\frac{1}{Z_n}\right)\right)^\gamma \left(\left(1 + \frac{\log(1 + (\theta_n/Z_n))}{\log Z_n}\right)^\gamma - 1\right)\right\} \\ &= \exp\{-c'(\log n)^{\gamma-1} n^{1/d} \theta_n(1 + o(1))\}, \end{aligned}$$

for some  $c' > 0$ , if  $\theta_n = o(n^{-1/d})$ . In particular, for  $\gamma - 1 > \nu$ , we can choose  $\theta_n = n^{1/d}(\log n)^{1-\gamma+\nu+\varepsilon}$  for some  $\varepsilon > 0$  and  $1 - \gamma + \nu + \varepsilon < 0$ . The constraints  $\gamma - 1 > \nu$  and  $\nu > 2 - \gamma$  entail  $\gamma > \frac{3}{2}$ . With this choice of  $\theta_n$ , we thus verify (4.9).

Now we can complete the proof of Theorem 2.2.

*Proof of Theorem 2.2.* Part (i) is Lemma 4.2. It remains to prove part (ii). Let  $R_n = \sum_{i=1}^n \mathbf{1}\{v_i \neq \eta_1(i)\}$ . Then, by Lemma 4.3,  $\mathbb{E}R_n = o(n)$ , which gives (2.9). Moreover, the obvious coupling is such that, given  $\mathcal{X}_n$  and  $G_n$ , one can transform the GPA graph  $G_n$  into the ONG on the same vertex sequence by the reassignment of the endpoint with smaller index of  $R_n$  edges, so affecting the degrees of at most  $2R_n$  vertices. Hence, with this coupling, for any  $k \in \mathbb{N}$ ,  $n^{-1}|N_n^{\text{GPA}}(k) - N_n^{\text{ONG}}(k)| \leq 2n^{-1}R_n$ , which tends to 0 in  $L^1$ . Now the  $L^1$  limit statement in (2.3) yields (2.10).

### 5. Proofs for power-law attractiveness

#### 5.1. Rejecting on-line nearest-neighbours

We introduce some notation on Voronoi cells that will also be used in Section 6. Let  $\mathcal{V}_n(i)$  denote the (bounded) Voronoi cell of  $X_i$  with respect to  $\mathcal{X}_n$  in  $S$ , i.e.

$$\mathcal{V}_n(i) := \{x \in S : \rho(x, X_i) < \min\{\rho(x, X_j) : 0 \leq j \leq n, j \neq i\}\}. \tag{5.1}$$

We need an elementary result showing that Voronoi cells are unlikely to be very small.

**Lemma 5.1.** *For any  $z > 0$ , with  $\delta_S > 0$  the constant in Lemma 3.1,*

$$\mathbb{P}[|\mathcal{V}_n(i)| < z] \leq 2^d \lambda_1 \delta_S^{-1} n z. \tag{5.2}$$

*Proof.* We follow the idea from [1, p. 311]. If no  $X_j$  with  $0 \leq j \leq n, j \neq i$  lies in  $B(X_i; r)$ , then  $S \cap B(X_i; r/2) \subseteq \mathcal{V}_n(i)$  and, hence,  $|\mathcal{V}_n(i)| \geq \delta_S \omega_d (r/2)^d$  by Lemma 3.1. That is,  $\mathbb{P}[|\mathcal{V}_n(i)| \geq \delta_S \omega_d (r/2)^d] \geq \mathbb{P}[\mathcal{X}_n \cap B(X_i; r) = \{X_i\}]$ . Complementation then shows that  $|\mathcal{V}_n(i)| < z$  ( $z > 0$ ) implies that at least one of  $n$  points  $X_j$  falls in  $B(X_i; 2z^{1/d}/(\omega_d \delta_S)^{1/d})$ . Now (5.2) follows from an application of Boole’s inequality:

$$\mathbb{P}[|\mathcal{V}_n(i)| < z] \leq n \mathbb{P}[X_j \in B(X_i; 2z^{1/d}/(\omega_d \delta_S)^{1/d})] \leq 2^d \delta_S^{-1} \lambda_1 n z.$$

Take  $F(r) = r^{-s}$  for  $s \in (0, \infty)$ . To prove Theorem 2.3, we consider the event  $\{v_n \neq \eta_1(n)\}$  that  $X_n$  is joined to a point *other than* its nearest neighbour.

*Proof of Theorem 2.3.* Extending the notation of (2.2), for  $\ell \in \mathbb{N}$  we let  $\eta_\ell(n)$  be the index of the  $\ell$ th nearest neighbour of  $X_n$  among  $\mathcal{X}_{n-1}$ . Again set  $Z_n = \rho(X_n, X_{\eta_1(n)})$  and  $W_n = \rho(X_n, X_{\eta_2(n)})$ . Then by (3.6),

$$\frac{\mathbb{P}[v_n \neq \eta_1(n) \mid \mathcal{F}_n]}{\mathbb{P}[v_n = \eta_1(n) \mid \mathcal{F}_n]} \geq \frac{\mathbb{P}[v_n = \eta_2(n) \mid \mathcal{F}_n]}{\mathbb{P}[v_n = \eta_1(n) \mid \mathcal{F}_n]} \geq \frac{F(W_n)}{\deg_{n-1}(\eta_1(n))F(Z_n)}.$$

Rearranging and using the fact that  $F(r) = r^{-s}$ , we obtain

$$\mathbb{P}[v_n \neq \eta_1(n) \mid \mathcal{F}_n] \geq \frac{1}{1 + \deg_{n-1}(\eta_1(n))(W_n/Z_n)^s} \geq \frac{(Z_n/W_n)^s}{2 \deg_{n-1}(\eta_1(n))}. \tag{5.3}$$

Then (2.11) will follow from (5.3) together with the following two claims: first, there exist constants  $k_0 \in \mathbb{N}$  and  $\theta_0 \in (0, 1)$  such that

$$\liminf_{n \rightarrow \infty} \mathbb{P}[\deg_n(\eta_1(n+1)) \leq k_0] \geq 2\theta_0, \tag{5.4}$$

and second, that for any  $\theta > 0$  there exist constants  $c, C \in (0, \infty)$  such that,

$$\mathbb{P}[Z_n \geq cn^{-1/d}] \geq 1 - \left(\frac{\theta}{3}\right), \quad \text{and} \quad \mathbb{P}[W_n \leq Cn^{-1/d}] \geq 1 - \left(\frac{\theta}{3}\right) \tag{5.5}$$

for all sufficiently large  $n$ . Indeed, it follows from (5.5) that  $\mathbb{P}[Z_n/W_n \geq c/C] \geq 1 - (2\theta_0/3)$  for suitable choice of  $c$  and  $C$ , so that, by (5.3) and (5.4),  $\mathbb{P}[v_n \neq \eta_1(n) \mid \mathcal{F}_n] \geq (1/2k_0)(c/C)^s$  with probability at least  $\theta_0/3$  for all sufficiently large  $n$ . Then, taking expectations, we obtain (2.11). Thus, it remains to prove the claims (5.4) and (5.5).

The idea behind (5.4) is that a large proportion of vertices have degrees bounded above by some  $k_0$ , and the union of the corresponding Voronoi cells will have volume bounded uniformly below in expectation, so that  $X_{n+1}$  will have such a vertex as its nearest neighbour with strictly positive probability. We formalize this idea.

With  $I_n(k) := \{i \in \{0, \dots, n\} : \deg_n(i) \leq k\}$ , we have  $\#I_n(k) = n + 1 - N_n^{\text{GPA}}(k + 1)$ . Taking  $k_0 = 9$ , we obtain from (3.5) that  $\#I_n(k_0) \geq 4n/5$  for all  $n$ . Each vertex  $i \in I_n(k_0)$  is associated with a Voronoi cell  $\mathcal{V}_n(i)$ .

Let  $\Lambda_n(r) = \#\{i \in \{0, \dots, n\} : |\mathcal{V}_n(i)| \geq r/n\}$ . Then

$$\mathbb{E}[\Lambda_n(r)] = \sum_{i=0}^n \mathbb{P}\left[|\mathcal{V}_n(i)| \geq \frac{r}{n}\right] = (n + 1)\mathbb{P}\left[|\mathcal{V}_n(i)| \geq \frac{r}{n}\right],$$

by exchangeability. Here, by (5.2),  $\mathbb{P}[|\mathcal{V}_n(i)| \geq r/n] \geq 1 - 2^d \lambda_1 \delta_S^{-1} r$ . Hence, we can (and do) choose  $r = r_0$  sufficiently small so that  $\mathbb{E}[\Lambda_n(r_0)] \geq 9n/10$ , say. Then, by Markov's inequality and the fact that  $\Lambda_n(r_0) \leq 1 + n$ ,

$$\mathbb{P}\left[\Lambda_n(r_0) \leq \frac{n}{2}\right] \leq \mathbb{P}\left[n + 1 - \Lambda_n(r_0) \geq \frac{n}{2}\right] \leq \frac{1 + (n/10)}{n/2} \leq \frac{1}{4} \quad \text{for all } n \geq 40.$$

So  $\mathbb{P}[\Lambda_n(r_0) \geq n/2] \geq \frac{3}{4}$  for all  $n \geq 40$ . On  $\{\Lambda_n(r_0) \geq n/2\}$ , since  $\#I_n(k_0) \geq 4n/5$ , at least  $3n/10$  vertices in  $I_n(k_0)$  have Voronoi cells of volume at least  $r_0/n$ , so

$$\mathbb{P}\left[\left|\bigcup_{i \in I_n(k_0)} \mathcal{V}_n(i)\right| \geq \frac{3r_0}{10}\right] \geq \frac{3}{4}, \tag{5.6}$$

for all sufficiently large  $n$ . Hence,

$$\mathbb{P}[\text{deg}_n(\eta_1(n + 1)) \leq k_0] \geq \mathbb{P}\left[X_{n+1} \in \bigcup_{i \in I_n(k_0)} \mathcal{V}_n(i)\right] \geq \lambda_0 \mathbb{E}\left[\left|\bigcup_{i \in I_n(k_0)} \mathcal{V}_n(i)\right|\right],$$

which with (5.6) gives (5.4), for  $2\theta_0 = 9r_0\lambda_0/40 > 0$ .

Finally, (5.5) can be verified by a similar argument to Lemma 3.2.

**5.2. Stretched exponential degree estimates**

Throughout this section we take  $F(r) = r^{-s}$  for  $s > d$ . By (3.6), for  $0 \leq i \leq n - 1$ ,

$$\mathbb{P}[v_n = i \mid \mathcal{F}_n] = \frac{\text{deg}_{n-1}(i)\rho(X_i, X_n)^{-s}}{\sum_{j=0}^{n-1} \text{deg}_{n-1}(j)\rho(X_j, X_n)^{-s}}.$$

Define, for any  $x \in S$ ,

$$\zeta_{n-1}(x) := n^{-s/d} \sum_{j=0}^{n-1} \rho(X_j, x)^{-s}. \tag{5.7}$$

Then we can write

$$\mathbb{P}[v_n = i \mid \mathcal{F}_n] \leq \frac{\text{deg}_{n-1}(i)\rho(X_i, X_n)^{-s}}{n^{s/d}\zeta_{n-1}(X_n)}. \tag{5.8}$$

The next result gives an estimate for the probability that  $\zeta_{n-1}(X_n)$  is small.

**Lemma 5.2.** *There exist constants  $C_0 < \infty$  and  $u_0 > 0$  such that, for all  $t > 0$ ,*

$$\limsup_{n \rightarrow \infty} \mathbb{P}[\zeta_{n-1}(X_n) \leq t] \leq C_0 \exp\{-u_0 t^{-d/(s-d)}\}. \tag{5.9}$$

*Proof.* First, for fixed  $x \in S$ , we have the tail estimate, for  $r > 0$ ,

$$\mathbb{P}[\rho(X_j, x)^{-s} > r] = \mathbb{P}[X_j \in B(x; r^{-1/s})] \geq \lambda_0 \delta_S \omega_d r^{-d/s},$$

using (2.1) and the lower bound in (3.4). Hence,  $\zeta_{n-1}(x)$  stochastically dominates  $\zeta_{n-1} := n^{-s/d} \sum_{j=0}^{n-1} \xi_j$ , where  $\xi_j \in \mathbb{R}_+$  are i.i.d. with  $\mathbb{P}[\xi_j > r] = \lambda_0 \delta_S \omega_d r^{-d/s}$ . Here,  $\zeta_{n-1}$  converges in distribution as  $n \rightarrow \infty$  to a random variable  $\zeta$  with a positive stable law of index  $d/s \in (0, 1)$ . Hence, for all  $x \in S$  and any  $t > 0$ ,

$$\limsup_{n \rightarrow \infty} \mathbb{P}[\zeta_{n-1}(x) \leq t] \leq \lim_{n \rightarrow \infty} \mathbb{P}[\zeta_{n-1} \leq t] = \mathbb{P}[\zeta \leq t].$$

Given that  $\zeta$  is a random variable with a positive stable law with index  $\alpha \in (0, 1)$ , for  $p > 0$  the random variable  $\zeta^{-p}$  satisfies  $\mathbb{E}[\exp(u\zeta^{-p})] < \infty$  for  $u \geq 0$  in a neighbourhood of zero, provided  $p \leq \alpha/(1 - \alpha)$ ; see, e.g. the proof of Lemma 1 in [3]. Hence, there exist  $u_0 > 0$  and  $C_0 < \infty$  such that, for  $p = d/(s - d) > 0$ ,  $\mathbb{E}[\exp(u_0\zeta^{-p})] \leq C_0$ . Thus,  $\mathbb{P}[\zeta \leq t] = \mathbb{P}[\exp(u_0\zeta^{-p}) \geq \exp(u_0t^{-p})]$ , and the result now follows from Markov’s inequality.

The next result is a conditional version of (5.9), given  $\mathcal{X}_{n-1}$ . The proof uses a concentration argument based on independently ‘resampling’ sites. Let  $X'_0, X'_1, \dots$  be an independent copy of the sequence  $X_0, X_1, \dots$ . For  $0 \leq i \leq n$ , let  $\mathcal{X}_n^i = (X_0, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n)$ , the sites  $\mathcal{X}_n$  but with  $X_i$  replaced by  $X'_i$ .

**Lemma 5.3.** *There exist constants  $C_1 < \infty$  and  $u_1 > 0$  such that, for any  $t > 0$ , a.s.,*

$$\limsup_{n \rightarrow \infty} \mathbb{P}[\zeta_{n-1}(X_n) \leq t \mid \mathcal{X}_{n-1}] \leq C_1 \exp\{-u_1 t^{-d/(s-d)}\}.$$

*Proof.* We approximate the indicator function  $\mathbf{1}_{[0,t]}$  by  $\chi_t^n: \mathbb{R}_+ \rightarrow [0, 1]$  defined by

$$\chi_t^n(x) := \begin{cases} 1 & \text{if } x \leq t, \\ 1 - (x - t)n^\delta & \text{if } t \leq x \leq t + n^{-\delta}, \\ 0 & \text{if } x \geq t + n^{-\delta}, \end{cases}$$

where  $\delta > 0$  is a constant to be specified later. Then

$$\mathbb{P}[\zeta_{n-1}(X_n) \leq t \mid \mathcal{X}_{n-1}] = \mathbb{E}[\mathbf{1}_{[0,t]}(\zeta_{n-1}(X_n)) \mid \mathcal{X}_{n-1}] \leq \mathbb{E}[\chi_t^n(\zeta_{n-1}(X_n)) \mid \mathcal{X}_{n-1}],$$

and  $\chi_t^n$  has the one-sided Lipschitz property  $\chi_t^n(r) - \chi_t^n(s) \leq n^\delta(s - r)^+$ . Now

$$\mathbb{E}[\chi_t^n(\zeta_{n-1}(X_n)) \mid \mathcal{X}_{n-1}] = \int_S f(x)\chi_t^n(\zeta_{n-1}(x)) \, dx = \phi(\mathcal{X}_{n-1})$$

for some measurable  $\phi: S^n \rightarrow [0, 1]$ . To obtain a concentration result for  $\phi(\mathcal{X}_{n-1})$ , we estimate  $\phi(\mathcal{X}_{n-1}^i) - \phi(\mathcal{X}_{n-1})$ . We introduce the notation

$$\zeta_{n-1}^i(x) = \zeta_{n-1}(x) + n^{-s/d}(\rho(X_i', x)^{-s} - \rho(X_i, x)^{-s}), \tag{5.10}$$

the change in the quantity given by (5.7) on resampling  $X_i$ . Then, for  $r_n > 0$ ,

$$\begin{aligned} \phi(\mathcal{X}_{n-1}^i) - \phi(\mathcal{X}_{n-1}) &\leq \int_{B(X_i; r_n)} f(x) \, dx \\ &\quad + \int_{S \setminus B(X_i; r_n)} f(x)(\chi_t^n(\zeta_{n-1}^i(x)) - \chi_t^n(\zeta_{n-1}(x))) \, dx \\ &\leq \lambda_1 \omega_d r_n^d + \int_{S \setminus B(X_i; r_n)} n^\delta f(x)(\zeta_{n-1}(x) - \zeta_{n-1}^i(x))^+ \, dx \end{aligned}$$

using the one-sided Lipschitz property of  $\chi_t^n$ . Now, by (5.10),

$$(\zeta_{n-1}(x) - \zeta_{n-1}^i(x))^+ \leq n^{-s/d} \rho(x, X_i)^{-s} \leq n^{-s/d} r_n^{-s}$$

provided  $x \notin B(X_i; r_n)$ . So, we obtain  $\phi(\mathcal{X}_{n-1}^i) - \phi(\mathcal{X}_{n-1}) \leq \lambda_1 \omega_d r_n^d + n^\delta n^{-s/d} r_n^{-s}$ . Since  $s > d$ , we may choose  $\delta > 0$  such that  $(s/d) - \delta > 1$ . Take  $r_n = n^{-\nu}$  where  $\nu = ((s/d) - \delta)/(s + d) > 0$ . Then we have that, for some constant  $C < \infty$ ,

$$\phi(\mathcal{X}_{n-1}^i) - \phi(\mathcal{X}_{n-1}) \leq C n^{-d((s/d)-\delta)/(s+d)} \leq C n^{-d/(s+d)}.$$

A version of Talagrand’s inequality [13, Theorem 4.5] yields, for some  $c_1 > 0$ ,

$$\mathbb{P}[|\phi(\mathcal{X}_{n-1}) - m_{n-1}| \geq r] \leq 4 \exp\{-c_1 n^{2d/(s+d)} r^2\}, \quad \text{for all } r > 0, \tag{5.11}$$

where  $m_{n-1}$  is a median of  $\phi(\mathcal{X}_{n-1})$ . In turn, (5.11) implies, by Lemma 4.6 of [13], that  $|m_{n-1} - \mathbb{E}\phi(\mathcal{X}_{n-1})| \leq c_2 n^{-d/(s+d)}$  for some  $c_2 < \infty$ . Here,  $\mathbb{E}\phi(\mathcal{X}_{n-1}) = \mathbb{E}[\chi_t^n(\zeta_{n-1}(X_n))]$   $\geq \mathbb{P}[\zeta_{n-1}(X_n) \leq t]$ , which for a fixed  $t > 0$  is bounded below uniformly in  $n$ , as can be proved

using an analogous argument to the proof of Lemma 5.2, this time using the upper bound in (3.4). It follows that, for some  $c_3 > 0$ ,

$$\mathbb{P}[\phi(\mathcal{X}_{n-1}) \geq 2\mathbb{E}\phi(\mathcal{X}_{n-1})] \leq 4 \exp\{-c_3 n^{2d/(s+d)}\}. \tag{5.12}$$

The right-hand side of (5.12) is summable in  $n$ , so the Borel–Cantelli lemma shows

$$\mathbb{P}[\zeta_{n-1}(X_n) \leq t \mid \mathcal{X}_{n-1}] \leq \phi(\mathcal{X}_{n-1}) \leq 2\mathbb{E}\phi(\mathcal{X}_{n-1}), \text{ a.s.,}$$

for all but finitely many  $n$ . Here, for  $t > 0$ ,

$$\mathbb{E}\phi(\mathcal{X}_{n-1}) \leq \mathbb{P}[\zeta_{n-1}(X_n) \leq t + n^{-\delta}] \leq \mathbb{P}[\zeta_{n-1}(X_n) \leq 2t]$$

for all large enough  $n$ . Now the statement follows from (5.9).

Choosing  $t = k^{-\gamma(s-d)/d}$  with  $\gamma \in (0, 1)$  in Lemma 5.3, we obtain the key estimate

$$\limsup_{n \rightarrow \infty} \mathbb{P}[\zeta_{n-1}(X_n) \leq k^{-\gamma(s-d)/d} \mid \mathcal{X}_{n-1}] \leq C_1 \exp\{-u_1 k^\gamma\}, \text{ a.s.} \tag{5.13}$$

In what follows,  $C_2, C_3, \dots$  represent constants not depending on  $n$  or  $k$ . We have

$$\begin{aligned} &\mathbb{P}[v_n = i, \zeta_{n-1}(X_n) > t \mid \tilde{\mathcal{F}}_{n-1}] \\ &\leq \mathbb{P}[\rho(X_i, X_n) \leq Bn^{-1/d} \mid \tilde{\mathcal{F}}_{n-1}] \\ &\quad + \mathbb{P}[v_n = i, \rho(X_i, X_n) > Bn^{-1/d}, \zeta_{n-1}(X_n) > t \mid \tilde{\mathcal{F}}_{n-1}] \end{aligned} \tag{5.14}$$

for any  $B > 0$  and any  $t > 0$ . The first term on the right-hand side of (5.14) is at most  $C_2 B^d n^{-1}$ , and the second term, by (5.8), is bounded above by

$$\frac{\deg_{n-1}(i)}{tn^{s/d}} \int_S f(x) \rho(X_i, x)^{-s} \mathbf{1}\{\rho(X_i, x) > Bn^{-1/d}\} dx.$$

For  $s > d$ , the latter integral is bounded above by

$$C_3 \int_{Bn^{-1/d}}^\infty \rho^{-s} \rho^{d-1} d\rho = C_4 B^{d-s} n^{(s/d)-1}.$$

Hence, we obtain from (5.14) that

$$\mathbb{P}[v_n = i, \zeta_{n-1}(X_n) > t \mid \tilde{\mathcal{F}}_{n-1}] \leq n^{-1} \left( C_2 B^d + \frac{C_4}{t} B^{d-s} \deg_{n-1}(i) \right). \tag{5.15}$$

For ease of notation, let  $q_k^{(n)}$  be the proportion of vertices of  $G_n$  with degree at least  $k$ , so that  $q_k^{(n)} := (n + 1)^{-1} N_n^{\text{GPA}}(k)$ . Then the proportion of vertices of  $G_n$  with degree  $k$  is equal to  $q_k^{(n)} - q_{k+1}^{(n)}$ , so that (5.15) yields

$$\begin{aligned} \mathbb{P}[\deg_{n-1}(v_n) = k, \zeta_{n-1}(X_n) > t \mid \tilde{\mathcal{F}}_{n-1}] &= \sum_{i: \deg_{n-1}(i)=k} \mathbb{P}[v_n = i, \zeta_{n-1}(X_n) > t \mid \tilde{\mathcal{F}}_{n-1}] \\ &\leq (q_k^{(n-1)} - q_{k+1}^{(n-1)}) \left( C_2 B^d + \frac{C_4}{t} B^{d-s} k \right). \end{aligned}$$

We take  $t = k^{-\gamma(s-d)/d}$  for  $\gamma \in (0, 1)$ , and choose  $B = k^{(\gamma/d)+(1/s)(1-\gamma)}$  to obtain

$$\mathbb{P}[\deg_{n-1}(v_n) = k, \zeta_{n-1}(X_n) > k^{-\gamma(s-d)/d} \mid \tilde{\mathcal{F}}_{n-1}] \leq C_5(q_k^{(n-1)} - q_{k+1}^{(n-1)})k^{\gamma+(d/s)(1-\gamma)}.$$

Now incorporating the case where  $\zeta_{n-1}(X_n)$  is small, using (5.13), gives, a.s.,

$$\mathbb{P}[\deg_{n-1}(v_n) = k \mid \tilde{\mathcal{F}}_{n-1}] \leq C_6e^{-u_1k^\gamma} + C_5(q_k^{(n-1)} - q_{k+1}^{(n-1)})k^\beta, \tag{5.16}$$

for all sufficiently large  $n$ , where we have set  $\beta = \gamma + (d/s)(1 - \gamma)$ . For any  $k$ , between times  $n - 1$  and  $n$ , the number of vertices of degree at least  $k$  either stays the same, or increases by exactly 1 if and only if  $\deg_{n-1}(v_n) = k - 1$ , so that  $\deg_n(v_n) = k$ . Thus,

$$\mathbb{E}[q_{k+1}^{(n)} \mid \tilde{\mathcal{F}}_{n-1}] - q_{k+1}^{(n-1)} = \frac{1}{n+1}(nq_{k+1}^{(n-1)} + \mathbb{P}[\deg_{n-1}(v_n) = k \mid \tilde{\mathcal{F}}_{n-1}] - q_{k+1}^{(n-1)}),$$

and we may express (5.16) as

$$\begin{aligned} \mathbb{E}[q_{k+1}^{(n)} \mid \tilde{\mathcal{F}}_{n-1}] - q_{k+1}^{(n-1)} &\leq \frac{1}{n+1}(C_6e^{-u_1k^\gamma} + C_5(q_k^{(n-1)} - q_{k+1}^{(n-1)})k^\beta - q_{k+1}^{(n-1)}) \\ &= \frac{1}{n+1}(C_6e^{-u_1k^\gamma} + q_k^{(n-1)}C_5k^\beta - q_{k+1}^{(n-1)}(1 + C_5k^\beta)). \end{aligned} \tag{5.17}$$

If we suppose that  $q_k^{(n)} \leq \tau_k$  for some  $\tau_k$  and all sufficiently large  $n$  (which we can, of course, always do for  $\tau_k = 1$ ) then (5.17) gives, for large enough  $n$ ,

$$\mathbb{E}[q_{k+1}^{(n)} \mid \tilde{\mathcal{F}}_{n-1}] - q_{k+1}^{(n-1)} \leq \frac{1}{n+1}(C_6e^{-u_1k^\gamma} + \tau_k C_5k^\beta - q_{k+1}^{(n-1)}(1 + C_5k^\beta)). \tag{5.18}$$

The final step in the proof of Theorem 2.4 is an analysis of (5.18) that will enable us iteratively to improve the bound  $\tau_k$ ; this uses the following stochastic approximation result, which is related to Lemma 2.6 of [15] and of some independent interest.

**Lemma 5.4.** *Let  $(\mathcal{G}_n; n \in \mathbb{Z}_+)$  be a filtration. Let  $g$  be a bounded function on  $\mathbb{R}_+$ . For  $n \in \mathbb{Z}_+$ , let  $Y_n, r_n, \xi_n$  be  $\mathcal{G}_n$ -measurable random variables, with  $Y_n \in \mathbb{R}_+$ , and*

$$Y_{n+1} - Y_n \leq \gamma_n(g(Y_n) + \xi_{n+1} + r_n) \tag{5.19}$$

for constants  $\gamma_n > 0$ . Suppose also that

- (i)  $\mathbb{E}[\xi_{n+1} \mid \mathcal{G}_n] = 0$  and  $\mathbb{E}[\xi_{n+1}^2 \mid \mathcal{G}_n] \leq C$  for some constant  $C < \infty$ ;
- (ii)  $\sum_n \gamma_n = \infty, \sum_n \gamma_n^2 < \infty$ , and  $\sum_n \gamma_n |r_n| < \infty$  a.s.;
- (iii)  $g(y) < -\delta$  for  $y > y_0$  for constants  $\delta > 0$  and  $y_0 \in \mathbb{R}_+$ .

Then  $\limsup_{n \rightarrow \infty} Y_n \leq y_0$ , a.s.

*Proof.* Summing (5.19) we obtain  $Y_n - Y_0 \leq M_n + A_n$  for any  $n \in \mathbb{Z}_+$ , where

$$M_n = \sum_{k=0}^{n-1} \gamma_k \xi_{k+1} \quad \text{and} \quad A_n = \sum_{k=0}^{n-1} \gamma_k (g(Y_k) + r_k);$$

the  $\mathcal{G}_n$ -martingale  $M_n$  and  $\mathcal{G}_{n-1}$ -measurable  $A_n$  constitute the Doob decomposition of the process whose increments are the right-hand side of (5.19). By (i),

$$\mathbb{E}[M_{n+1}^2 - M_n^2 \mid \mathcal{G}_n] = \mathbb{E}[(M_{n+1} - M_n)^2 \mid \mathcal{G}_n] \leq C\gamma_n^2, \text{ a.s.,}$$

which is summable, by (ii), so the increasing process associated with  $M_n$  is a.s. bounded. Hence,  $M_n \rightarrow M_\infty$  a.s., for some finite limit  $M_\infty$ . Also, writing  $R_n = \sum_{k=0}^{n-1} \gamma_k r_k$ , we have  $R_n \rightarrow R_\infty$  a.s. for some finite limit  $R_\infty$ , by (ii). In particular, for any  $\varepsilon > 0$ , there exists an a.s. finite  $N$  such that,

$$\sup_{n \geq N} \sup_{m \geq 0} |M_{n+m} - M_n| \leq \frac{\varepsilon}{4} \quad \text{and} \quad \sup_{n \geq N} \sup_{m \geq 0} |R_{n+m} - R_n| \leq \frac{\varepsilon}{4}.$$

Consider some  $n \geq N$  for which  $Y_n > y_0$ . Let  $\kappa_n$  be the first time after  $n$  for which  $Y. \leq y_0$ . Then, by summing (5.19) again, for  $m \geq 0$ ,

$$\begin{aligned} Y_{(n+m) \wedge \kappa_n} - Y_n &\leq M_{(n+m) \wedge \kappa_n} - M_n + R_{(n+m) \wedge \kappa_n} - R_n + \sum_{k=n}^{(n+m) \wedge \kappa_n - 1} \gamma_k g(Y_k) \\ &\leq \frac{\varepsilon}{2} - \delta \sum_{k=n}^{(n+m) \wedge \kappa_n - 1} \gamma_k. \end{aligned}$$

In particular, on  $\{\kappa_n = \infty\}$ , letting  $m \rightarrow \infty$  the left-hand side of the last display remains bounded below by  $-Y_n$  while the right-hand side tends to  $-\infty$ , by (ii); hence  $\kappa_n < \infty$  a.s., and the process returns to the interval  $[0, y_0]$  without exceeding  $Y_n + \varepsilon$ . Moreover,  $Y_{n+1} - Y_n \leq \varepsilon/2 + \gamma_n g(Y_n) < \varepsilon$ , for all large enough  $n \geq N$ , since  $g$  is bounded and  $\gamma_n \rightarrow 0$ . Hence,  $Y_n \leq y_0$  infinitely often, and, for all but finitely many such  $n$ , any exit from  $[0, y_0]$  cannot exceed  $y_0 + \varepsilon$ ; but starting from  $[y_0, y_0 + \varepsilon]$  the process returns to  $[0, y_0]$  before reaching  $y_0 + 2\varepsilon$ . Hence,  $\limsup_{n \rightarrow \infty} Y_n \leq y_0 + 2\varepsilon$ , a.s. Since  $\varepsilon > 0$  was arbitrary, the result follows.

*Proof of Theorem 2.4.* Setting

$$\begin{aligned} \mathcal{G}_n &= \tilde{\mathcal{F}}_n, \quad Y_n = q_{k+1}^{(n)}, \quad \gamma_n = \frac{1}{n+2}, \quad r_n = 0, \\ g(y) &= C_6 e^{-u_1 k^\gamma} + \tau_k C_5 k^\beta - y(1 + C_5 k^\beta), \end{aligned}$$

and

$$\xi_{n+1} = (n+2)(q_{k+1}^{(n+1)} - \mathbb{E}[q_{k+1}^{(n+1)} \mid \tilde{\mathcal{F}}_n]),$$

we apply Lemma 5.4 to (5.18). Note that, since  $N_n^{\text{GPA}}(k)$  is  $\tilde{\mathcal{F}}_n$ -measurable,

$$\begin{aligned} \xi_{n+1} &= N_{n+1}^{\text{GPA}}(k+1) - \mathbb{E}[N_{n+1}^{\text{GPA}}(k+1) \mid \tilde{\mathcal{F}}_n] \\ &= N_{n+1}^{\text{GPA}}(k+1) - N_n^{\text{GPA}}(k+1) - \mathbb{E}[N_{n+1}^{\text{GPA}}(k+1) - N_n^{\text{GPA}}(k+1) \mid \tilde{\mathcal{F}}_n], \end{aligned}$$

which is uniformly bounded, since  $0 \leq N_{n+1}^{\text{GPA}}(k) - N_n^{\text{GPA}}(k) \leq 1$ , a.s. Hence, the conditions of Lemma 5.4 are satisfied for any

$$y_0 > \frac{C_6 e^{-u_1 k^\gamma} + \tau_k C_5 k^\beta}{1 + C_5 k^\beta},$$

and we deduce that

$$\limsup_{n \rightarrow \infty} q_{k+1}^{(n)} \leq \frac{C_6 e^{-u_1 k^\gamma} + \tau_k C_5 k^\beta}{1 + C_5 k^\beta}. \tag{5.20}$$

In particular, if  $q_k^{(n)} \leq \tau_k$  for all but finitely many  $n$ , a.s., then (5.20) implies that  $q_{k+1}^{(n)} \leq \tau_{k+1}$  for all but finitely many  $n$ , a.s., where

$$\tau_{k+1} = \frac{2C_6 e^{-u_1 k^\gamma} + \tau_k C_5 k^\beta}{1 + C_5 k^\beta}. \tag{5.21}$$

Defining  $\sigma_k > 0$  via  $\tau_k = 2C_6 \sigma_k e^{-u_1 k^\gamma}$ , we obtain from (5.21), after some algebra,

$$\sigma_{k+1} - \sigma_k = \left( 1 - a_{k+1} + \frac{a_{k+1}}{1 + C_5 k^\beta} \right) (1 - \sigma_k) - (1 - a_{k+1}),$$

where  $a_{k+1} := \exp\{u_1((k+1)^\gamma - k^\gamma)\} = 1 + \gamma u_1 k^{\gamma-1} + O(k^{\gamma-2})$  as  $k \rightarrow \infty$ . Then, assuming that  $\beta < 1 - \gamma$ , it is straightforward to check that

$$1 - a_{k+1} + \frac{a_{k+1}}{1 + C_5 k^\beta} \sim \frac{1}{C_5 k^\beta} \quad \text{as } k \rightarrow \infty.$$

Hence, we may apply Lemma 1 of [10] to see that  $\lim_{k \rightarrow \infty} \sigma_k = 1$ , provided  $\beta < 1 - \gamma$ , i.e.  $\gamma < (s - d)/(2s - d)$ . For any such  $\gamma$ , we thus obtain  $\limsup_{n \rightarrow \infty} q_k^{(n)} \leq \tau_k \leq 3C_6 e^{-u_1 k^\gamma}$ , a.s., for all sufficiently large  $k$ , giving the almost sure statement in the theorem. Then the reverse Fatou lemma yields the statement on expectations.

### 6. Proofs for the on-line nearest-neighbour graph

In this section we prove Theorem 2.1. Our argument extends the 2-dimensional argument of [1, Section 3.1], which considered the uniform distribution on the square.

Recall the definition of the Voronoi cell  $\mathcal{V}_n(i)$  from (5.1). Then

$$\mathcal{V}_{n+1}(i) = \mathcal{V}_n(i) \cap \{x \in S : \rho(x, X_i) < \rho(x, X_{n+1})\} \subseteq \mathcal{V}_n(i). \tag{6.1}$$

A key fact is provided by the following lemma, which will be used to show that the volume of a Voronoi cell associated with a vertex in the ONG shrinks, on average, by a positive fraction whenever a new vertex lands in the cell.

**Lemma 6.1.** *Let  $R \subseteq S$  be convex, and let  $X$  be a random point in  $S$  distributed according to the probability density  $f$  satisfying (2.1). For  $x_0 \in R$ , let  $R' = \{x \in R : \rho(x, x_0) < \rho(x, X)\}$ . Then there exists  $\delta > 0$  not depending on  $R$  or  $x_0$  such that*

$$\mathbb{E}[|R'| \mid X \in R] \leq (1 - \delta)\mathbb{E}[|R|].$$

*Proof.* Without loss of generality, take  $x_0 = 0 \in R$ . Partition  $R$  according to the Cartesian orthants as  $R_1, \dots, R_{2d}$ . For each  $j$ , any two points  $x, y \in R_j$  have the same signs in corresponding coordinates, so  $\|x - y\| \leq \|x + y\|$ , and, hence,  $(x + y)/2$  is closer to  $x$  (and to  $y$ ) than to  $0$ . Thus, given  $X \in R_j$ , any point  $x$  of  $R'_j := \{(X + y)/2 : y \in R_j\}$  has  $\|x - X\| \leq \|x - 0\|$ , and, by convexity,  $R'_j \subseteq R_j$ . Hence, given  $X \in R_j$ ,  $R' \subseteq R \setminus R'_j$ . By

construction,  $R''_j$  is a translate of  $R_j$  scaled by a factor of  $\frac{1}{2}$ , so

$$\begin{aligned} \mathbb{E}[|R'| \mid X \in R] &\leq |R| - \sum_{j=1}^{2^d} 2^{-d} |R_j| \mathbb{P}[X \in R_j \mid X \in R] \\ &\leq |R| - 2^{-d} \left(\frac{\lambda_0}{\lambda_1}\right) |R|^{-1} \sum_{j=1}^{2^d} |R_j|^2, \end{aligned}$$

using (2.1). Now, by Jensen’s inequality,  $\sum_{j=1}^{2^d} |R_j|^2 \geq 2^{-d} (\sum_{j=1}^{2^d} |R_j|)^2 = 2^{-d} |R|^2$ , and the claimed result follows with  $\delta = 2^{-2d} \lambda_0 / \lambda_1$ .

Next, we provide bounds on expectations for  $N_n^{\text{ONG}}(k)$ .

**Lemma 6.2.** *Let  $d \in \mathbb{N}$ . Suppose that (2.1) holds. Then there exist finite positive constants  $A, A', C$ , and  $C'$  such that, for all  $k \in \mathbb{N}$ ,*

$$A' e^{-C'k} \leq \liminf_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \leq \limsup_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \leq A e^{-Ck}. \tag{6.2}$$

Moreover, more precisely,

$$-\liminf_{k \rightarrow \infty} \left( k^{-1} \log \left( \liminf_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \right) \right) \leq 1, \tag{6.3}$$

and, in the case where  $f$  is the uniform density on  $S$ ,

$$-\limsup_{k \rightarrow \infty} \left( k^{-1} \log \left( \limsup_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \right) \right) \geq \frac{1}{2} \log(1 + (2^{2d} - 1)^{-1}). \tag{6.4}$$

*Proof.* First we prove the upper bound in (6.2), using an argument based in part on [1, Section 3.1]. Fix  $i \in \mathbb{Z}_+$ . Let  $t_0 = i$  and for  $j \in \mathbb{N}$  define recursively  $t_j = \min\{t > t_{j-1} : X_t \in \mathcal{V}_{t-1}(i)\}$ , so that  $t_1, t_2, \dots$  are the times at which edges to  $X_i$  are created. Following [1, p. 311], let  $\mathcal{W}_j = \mathcal{V}_{t_j}(i)$ .

Observe that if  $i$  has degree greater than  $k$  in the ONG on  $(X_0, \dots, X_n)$ ,  $n \geq i$ , then necessarily  $t_k \leq n$ , and so also  $|\mathcal{V}_n(i)| \leq |\mathcal{V}_{t_k}(i)| = |\mathcal{W}_k|$ , by (6.1). Hence,

$$\mathbb{P}[\text{deg}_n(i) > k] \leq \mathbb{P}[|\mathcal{W}_k| \geq z] + \mathbb{P}[|\mathcal{V}_n(i)| \leq z] \quad \text{for any } z > 0. \tag{6.5}$$

We bound each of the probabilities on the right-hand side of (6.5) in turn, and then optimize the choice of  $z$ .

By definition,  $X_{t_j}$  is distributed according to the density  $f$ , conditioned to fall in the convex set  $\mathcal{V}_{t_{j-1}}(i) \subseteq \mathcal{V}_{t_{j-1}}(i) \subseteq S$ . Hence, from Lemma 6.1, it follows that  $\mathbb{E}[|\mathcal{W}_j|] \leq (1 - \delta) \mathbb{E}[|\mathcal{W}_{j-1}|]$ , where  $\delta \in (0, 1)$  depends only on  $d$  and  $\lambda_0 / \lambda_1$ , and

$$\mathbb{E}[|\mathcal{W}_j|] \leq (1 - \delta)^j \mathbb{E}[|\mathcal{V}_i(i)|] = \frac{1}{i + 1} (1 - \delta)^j,$$

since the vector  $(|\mathcal{V}_i(0)|, \dots, |\mathcal{V}_i(i)|)$  is exchangeable and its components sum to 1, so  $\mathbb{E}[|\mathcal{V}_i(j)|] = 1 / (i + 1)$ . Markov’s inequality implies that, for any  $z > 0$ ,

$$\mathbb{P}[|\mathcal{W}_j| \geq z] \leq \frac{1}{z} \frac{1}{i + 1} (1 - \delta)^j. \tag{6.6}$$

The final term in (6.5) is bounded above by (5.2). Combining (6.5) with (6.6) and (5.2), we obtain, for any  $z > 0$ ,  $\mathbb{P}[\text{deg}_n(i) > k] \leq (1/z(i + 1))(1 - \delta)^k + Cnz$ , where  $C < \infty$  depends only on  $d, S$ , and  $\lambda_1$ . The optimal bound is obtained on taking  $z = (1 - \delta)^{k/2}/\sqrt{Cn(i + 1)}$ , and we conclude

$$\mathbb{P}[\text{deg}_n(i) > k] \leq 2(1 - \delta)^{k/2} \sqrt{\frac{Cn}{i + 1}}. \tag{6.7}$$

The upper bound in (6.2) follows from (6.7), since

$$\mathbb{E}[N_n^{\text{ONG}}(k)] = \sum_{i=0}^n \mathbb{P}[\text{deg}_n(i) \geq k] \leq C'n(1 - \delta)^{k/2},$$

for some  $C' < \infty$  not depending on  $k$  or  $n$ . The statement (6.4) also follows, since when  $\lambda_0 = \lambda_1$ , we have from the proof of Lemma 6.1 that we may take  $\delta = 2^{-2d}$ .

To prove the lower bound in (6.2) as well as (6.3), we use a similar idea to that briefly outlined for the analogous argument in [1, p. 311], but filling in the details takes some work, and we must be more careful with our estimates to obtain the quantitative bound (6.3). First note that, for  $j > i$ , the (unconditional) probability that  $X_j$  is joined to  $X_i$  is  $\mathbb{P}[\eta_1(j) = i] = \mathbb{P}[X_j \in \mathcal{V}_{j-1}(i)] = 1/j$ . Write  $d_n(i) := \mathbb{E}[\text{deg}_n(i)]$ . Then, for  $i \in \mathbb{N}$ ,

$$d_n(i) = 1 + \sum_{j=i+1}^n \mathbb{P}[\eta_1(j) = i] \geq \sum_{j=i}^n \frac{1}{j} \geq \int_i^n \frac{1}{y} dy = \log\left(\frac{n}{i}\right).$$

Let  $\theta > 1$ . For  $k \in \mathbb{Z}_+$ , let  $H_{n,k}^\theta := \mathbb{N} \cap [1, e^{-\theta k}n]$ . Then for any  $i \in H_{n,k}^\theta$ ,  $d_n(i) \geq \log(n/i) \geq \theta k$ . It follows that

$$\mathbb{E}[N_n^{\text{ONG}}(k)] \geq \sum_{i \in H_{n,k}^\theta} \mathbb{P}[\text{deg}_n(i) \geq k] \geq \sum_{i \in H_{n,k}^\theta} \mathbb{P}[\text{deg}_n(i) \geq \theta^{-1}d_n(i)]. \tag{6.8}$$

Let  $w \in (1, \infty)$ , to be specified later. Then  $w > 1 > 1/\theta$ , and

$$\begin{aligned} wd_n(i)\mathbb{P}[\text{deg}_n(i) \geq \theta^{-1}d_n(i)] &\geq \mathbb{E}[\text{deg}_n(i) \mathbf{1}\{\text{deg}_n(i) \geq \theta^{-1}d_n(i)\}] \\ &\quad - \mathbb{E}[\text{deg}_n(i) \mathbf{1}\{\text{deg}_n(i) > wd_n(i)\}] \\ &\geq (1 - \theta^{-1})d_n(i) - \mathbb{E}[\text{deg}_n(i) \mathbf{1}\{\text{deg}_n(i) > wd_n(i)\}], \end{aligned} \tag{6.9}$$

using the fact that  $\mathbb{E}[X \mathbf{1}\{X \geq x\}] \geq \mathbb{E}[X] - x$  for any  $x \geq 0$  and any nonnegative random variable  $X$ . By the Cauchy–Schwarz inequality, the final term in (6.9) satisfies

$$\mathbb{E}[\text{deg}_n(i) \mathbf{1}\{\text{deg}_n(i) > wd_n(i)\}] \leq (\mathbb{E}[\text{deg}_n(i)^2] \mathbb{P}[\text{deg}_n(i) > wd_n(i)])^{1/2}. \tag{6.10}$$

We claim that, given  $\theta > 1$ , there exists  $w = w(\theta) \in (1, \infty)$  such that

$$\sup_{i \in H_{n,k}^\theta} (\mathbb{E}[\text{deg}_n(i)^2] \mathbb{P}[\text{deg}_n(i) > wd_n(i)])^{1/2} \leq e^{-\theta k} \quad \text{for all } n, k \in \mathbb{N}. \tag{6.11}$$

Given (6.11), which we verify at the end of this proof, we obtain from (6.9), (6.10), and (6.11) that, for any  $n \in \mathbb{N}$  and any  $k \in \mathbb{N}$ ,

$$\begin{aligned} w \inf_{i \in H_{n,k}^\theta} \mathbb{P}[\text{deg}_n(i) \geq \theta^{-1}d_n(i)] &\geq (1 - \theta^{-1}) - e^{-\theta k} \sup_{i \in H_{n,k}^\theta} \frac{1}{d_n(i)} \\ &\geq (1 - \theta^{-1}) - \frac{e^{-\theta k}}{\theta k}, \end{aligned} \tag{6.12}$$

using the fact that  $d_n(i) \geq \theta k$  for  $i \in H_{n,k}^\theta$ . To prove the lower bound in (6.2), it is enough to fix  $\theta = 2$ . Then (6.12) becomes, for any  $n \in \mathbb{N}$  and any  $k \in \mathbb{N}$ ,

$$w \inf_{i \in H_{n,k}^2} \mathbb{P}[\deg_n(i) \geq \frac{1}{2}d_n(i)] \geq \frac{1}{2}(1 - e^{-2}) \geq \frac{3}{8},$$

say, where  $w = w(2)$  is constant. Hence, from (6.8) we obtain, for all  $n, k \in \mathbb{N}$ ,

$$\mathbb{E}[N_n^{\text{ONG}}(k)] \geq w^{-1} \sum_{i \in H_{n,k}^2} \frac{3}{8} \geq \frac{3}{8w}(e^{-2k}n - 1),$$

which gives  $\liminf_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)] \geq (3/8w)e^{-2k}$ .

To prove (6.3), we adapt the preceding argument. For any  $\theta > 1$ , there exists  $k_0 \in \mathbb{N}$  such that, for all  $k \geq k_0$ , the final expression on the right-hand side of (6.12) exceeds  $(1 - \theta^{-1})/2 > 0$ , say. Then, similarly to before, we obtain, for all  $k \geq k_0$  and  $n \in \mathbb{N}$ ,

$$\mathbb{E}[N_n^{\text{ONG}}(k)] \geq w^{-1} \sum_{i \in H_{n,k}^\theta} \left(\frac{1 - \theta^{-1}}{2}\right) \geq w^{-1} \left(\frac{1 - \theta^{-1}}{2}\right)(e^{-\theta k}n - 1).$$

First letting  $n \rightarrow \infty$  and then  $k \rightarrow \infty$ , it follows that

$$\limsup_{k \rightarrow \infty} \left(-k^{-1} \log(\liminf_{n \rightarrow \infty} n^{-1} \mathbb{E}[N_n^{\text{ONG}}(k)])\right) \leq \theta.$$

Since  $\theta > 1$  was arbitrary, (6.3) follows.

It remains to establish the claim (6.11). To this end, an application of (6.7) shows that, for constants  $C_1, C_2 < \infty$  and  $c > 0$ , for all  $n \in \mathbb{N}$  and  $1 \leq i \leq n$ ,

$$\mathbb{E}[\deg_n(i)^2] = \sum_{k=0}^\infty \mathbb{P}[\deg_n(i) > \sqrt{k}] \leq C_1 \sqrt{\frac{n}{i}} \sum_{k=0}^\infty e^{-c\sqrt{k}} \leq C_2 \sqrt{\frac{n}{i}}.$$

Another application of (6.7) shows that, for some constant  $C_3 < \infty$ , for any  $w > 0$ ,

$$\mathbb{P}[\deg_n(i) > wd_n(i)] \leq C_3 \sqrt{\frac{n}{i}} e^{-cw \log(n/i)} = C_3 \left(\frac{n}{i}\right)^{(1/2)-cw}.$$

Hence, we obtain, for all  $1 \leq i \leq n$ ,

$$(\mathbb{E}[\deg_n(i)^2] \mathbb{P}[\deg_n(i) > wd_n(i)])^{1/2} \leq C_4 \left(\frac{i}{n}\right)^{(cw-1)/2},$$

where  $C_4 < \infty$  is constant. Taking  $w > 3/c$ , we have, for any  $i \in H_{n,k}^\theta$ ,

$$C_4 \left(\frac{i}{n}\right)^{(cw-1)/2} \leq C_4 e^{-\theta k} e^{-(cw-3)\theta k/2},$$

since  $i/n \leq e^{-\theta k}$  for  $i \in H_{n,k}^\theta$ . In particular, for all  $k \in \mathbb{N}$ , we can choose  $w$  (depending on  $c, C_4$ , and  $\theta$ ) such that  $C_4 e^{-(cw-3)\theta k/2} \leq C_4 e^{-(cw-3)\theta/2} \leq 1$ . This verifies (6.11).

Next we have a concentration result for  $N_n^{\text{ONG}}(k)$ .

**Lemma 6.3.** *Let  $d \in \mathbb{N}$ . Suppose that (2.1) holds. Then*

$$\limsup_{n \rightarrow \infty} n^{-1} \sup_{k \in \mathbb{N}} |N_n^{\text{ONG}}(k) - \mathbb{E}[N_n^{\text{ONG}}(k)]| = 0, \text{ a.s.} \tag{6.13}$$

*Proof.* We use the resampling idea described before Lemma 5.3 and a modification of the Azuma–Hoeffding inequality. Set  $D_{n,i} = \mathbb{E}[N_n^{\text{ONG}}(k) \mid \mathcal{G}_i] - \mathbb{E}[N_n^{\text{ONG}}(k) \mid \mathcal{G}_{i-1}]$ , where  $\mathcal{G}_i = \sigma(X_0, X_1, \dots, X_i)$ . In words,  $-D_{n,i}$  is equal to the expected (conditional on  $\mathcal{G}_i$ ) change in  $N_n^{\text{ONG}}(k)$  on independently resampling  $X_i$ , and  $D_{n,i}, 1 \leq i \leq n$  is a martingale difference sequence with  $\sum_{i=1}^n D_{n,i} = N_n^{\text{ONG}}(k) - \mathbb{E}[N_n^{\text{ONG}}(k)]$ .

We bound  $|D_{n,i}|$  in terms of  $\text{deg}_n(i)$  and  $\text{deg}_n^i(i)$ , the degree of vertex  $i$  in the ONG on  $\mathcal{X}_n$  and  $\mathcal{X}_n^i$ , respectively. On replacement of  $X_i$  by  $X'_i$ , the degree of vertex  $i$  may change, leading to a change of  $\pm 1$  in  $N_{n,i}^{\text{ONG}}(k)$  compared to  $N_n^{\text{ONG}}(k)$ . The degrees of at most  $\text{deg}_n(i) - 1$  other vertices increase (namely, those vertices that gain incoming edges that were previously connected to  $X_i$ ), while the degrees of at most  $\text{deg}_n^i(i) - 1$  vertices decrease (namely, those vertices that lose incoming edges reassigned to  $X'_i$ ).

Hence,  $|D_{n,i}| \leq \text{deg}_n(i) + \text{deg}_n^i(i)$ . Now, for any  $r > 0$ ,

$$\mathbb{P}[|D_{n,i}| > r] \leq \mathbb{P}\left[\text{deg}_n(i) > \frac{r}{2}\right] + \mathbb{P}\left[\text{deg}_n^i(i) > \frac{r}{2}\right] = 2\mathbb{P}\left[\text{deg}_n(i) > \frac{r}{2}\right],$$

since  $\text{deg}_n^i(i)$  and  $\text{deg}_n(i)$  are identically distributed. Hence, by (6.7),  $\mathbb{P}[|D_{n,i}| > D \log n] = O(n^{-5})$ , uniformly in  $i$ , choosing sufficiently large  $D \in (0, \infty)$ ; note that this bound is also uniform in  $k$ . By a modification of the Azuma–Hoeffding inequality due to Chalker *et al.* [4, Lemma 1], it follows that

$$\mathbb{P}[|N_n^{\text{ONG}}(k) - \mathbb{E}[N_n^{\text{ONG}}(k)]| > r] \leq \left(1 + \frac{4n}{r}\right)n^{-4} + 2 \exp\left\{-\frac{r^2}{32D^2n(\log n)^2}\right\}$$

for any  $r > 0$ . Taking  $r = n^{3/4}$ , say, shows that  $\mathbb{P}[|N_n^{\text{ONG}}(k) - \mathbb{E}[N_n^{\text{ONG}}(k)]| > n^{3/4}] = O(n^{-3})$ , uniformly in  $k \in \{1, \dots, n\}$ , while for  $k > n, N_n^{\text{ONG}}(k) = 0$  a.s. Hence,

$$\sum_{n=1}^{\infty} \sum_{k \in \mathbb{N}} \mathbb{P}[|N_n^{\text{ONG}}(k) - \mathbb{E}[N_n^{\text{ONG}}(k)]| > n^{3/4}] \leq C \sum_{n=1}^{\infty} n^{-2} < \infty.$$

The Borel–Cantelli lemma now yields (6.13).

*Proof of Theorem 2.1.* Penrose [16, Section 3.4] showed that functionals such as counts of vertices of a given degree in the ONG satisfy *stabilization* (a form of local dependence). This guarantees a law of large numbers of the form  $n^{-1}N_n^{\text{ONG}}(k) \rightarrow \mathbb{P}[\xi(0, U; \mathcal{H}) \geq k]$  *in probability*: concretely, one may apply results of [19] or [17]. The bounded convergence theorem gives  $n^{-1}\mathbb{E}[N_n^{\text{ONG}}(k)] \rightarrow \mathbb{P}[\xi(0, U; \mathcal{H}) \geq k]$ , and Lemma 6.3 then shows that convergence in probability can be replaced by almost sure convergence. The  $L^1$  convergence follows from the bounded convergence theorem again. Thus, (2.3) holds.

Then, applying Lemma 6.2 with (2.3), (2.4) follows from (6.2). Given (2.3), the upper bound in (2.5) follows from (6.3). Similarly, the lower bound in (2.5) follows from (6.4), noting that the limit  $\rho_k$  is independent of the choice of  $f$ .

It is easy to see that  $\rho_k$  is nonincreasing with  $\rho_1 = 1$ . Since  $\sum_{k \in \mathbb{N}} N_n^{\text{ONG}}(k) = 2n$ , twice the number of edges in the ONG, dividing both sides of this last equality by  $n$  and letting

$n \rightarrow \infty$  we must have  $\sum_{k \in \mathbb{N}} \rho_k = 2$ ; hence, also  $\lim_{k \rightarrow \infty} \rho_k = 0$ . For the final statement of the theorem, we have from (6.7) that for any  $k > 0$ ,

$$\mathbb{P}\left[\max_{0 \leq i \leq n} \deg_n(i) > k\right] \leq (n+1) \max_{0 \leq i \leq n} \mathbb{P}[\deg_n(i) > k] \leq Cn^{3/2}e^{-ck}$$

for some absolute constants  $c, C \in (0, \infty)$ . Taking  $k = D \log n$ , we can choose  $D \in (0, \infty)$  for which this last bound is  $O(n^{-2})$ ; the Borel–Cantelli lemma then gives (2.6).

### Acknowledgement

The authors are grateful to the anonymous referee for a careful reading of the first version of this paper.

### References

- [1] BERGER, N. *et al.* (2007). Degree distribution of the FKP network model. *Theoret. Comput. Sci.* **379**, 306–316.
- [2] BOLLOBÁS, B., RIORDAN, O., SPENCER, J. AND TUSNÁDY, G. (2001). The degree sequence of a scale-free random graph process. *Random Structures Algorithms* **18**, 279–290.
- [3] BROCKWELL P. J. AND BROWN, B. M. (1981). High-efficiency estimation for the positive stable laws. *J. Amer. Statist. Assoc.* **76**, 626–631.
- [4] CHALKER, T. K. *et al.* (1999). On the size of a random sphere of influence graph. *Adv. Appl. Prob.* **31**, 596–609.
- [5] FABRIKANT, A., KOUTSOPIAS, E. AND PAPADIMITRIOU, C. H. (2002). Heuristically optimized trade-offs: a new paradigm for power laws in the internet. In *Automata, Languages and Programming* (Lecture Notes Comput. Sci. **2380**), Springer, Berlin, pp. 110–122.
- [6] FERRETTI, L. AND CORTELEZZI, M. (2011). Preferential attachment in growing spatial networks. *Phys. Rev. E* **84**, 016103.
- [7] FLAXMAN, A. D., FRIEZE, A. M. AND VERA, J. (2006). A geometric preferential attachment model of networks. *Internet Math.* **3**, 187–205.
- [8] FLAXMAN, A. D., FRIEZE, A. M. AND VERA, J. (2007). A geometric preferential attachment model of networks. II. *Internet Math.* **4**, 87–111.
- [9] JACOB, E. AND MÖRTERS, P. (2015). Spatial preferential attachment networks: power laws and clustering coefficients. *Ann. Appl. Prob.* **25**, 632–662.
- [10] JORDAN, J. (2006). The degree sequences and spectra of scale-free random graphs. *Random Structures Algorithms* **29**, 226–242.
- [11] JORDAN, J. (2010). Degree sequences of geometric preferential attachment graphs. *Adv. Appl. Prob.* **42**, 319–330.
- [12] MANNA, S. S. AND SEN, P. (2002). Modulated scale-free network in Euclidean space. *Phys. Rev. E* **66**, 066114.
- [13] MCDIARMID, C. (1998). Concentration. In *Probabilistic Methods for Algorithmic Discrete Mathematics*, Springer, Berlin, pp. 195–248.
- [14] MOLLOY, M. AND REED, B. (2002). *Graph Colouring and the Probabilistic Method*. Springer, Berlin.
- [15] PEMANTLE, R. (2007). A survey of random processes with reinforcement. *Prob. Surv.* **4**, 1–79.
- [16] PENROSE, M. D. (2005). Multivariate spatial central limit theorems with applications to percolation and spatial graphs. *Ann. Prob.* **33**, 1945–1991.
- [17] PENROSE, M. D. (2007). Laws of large numbers in stochastic geometry with statistical applications. *Bernoulli* **13**, 1124–1150.
- [18] PENROSE, M. D. AND WADE, A. R. (2010). Random directed and on-line networks. In *New Perspectives in Stochastic Geometry*, Oxford University Press, pp. 248–274.
- [19] PENROSE, M. D. AND YUKICH, J. E. (2003). Weak laws of large numbers in geometric probability. *Ann. Appl. Prob.* **13**, 277–303.