

<i>Schistosoma bovis</i> ♀ crossed with <i>S. haematobium</i> ♂ (Iran) ...	298
crossed with <i>S. haematobium</i> ♂ (Nigeria)	294
crossed with <i>S. mansoni</i> ♂	281
<i>mansoni</i> crossed with <i>S. rodhaini</i>	301
testes number in diagnosis of	69
<i>mattheei</i> crossed with <i>S. bovis</i>	282
♀ crossed with <i>S. haematobium</i> ♂ (Nigeria)	288
crossed with <i>S. mansoni</i>	272
Schistosomes, African, characteristics of	258
hybridisation experiments with	272
<i>Spinitectus</i> sp. from fish in Ghana	64
<i>Spironoura</i> , key to species of	68
<i>petrei</i> n.sp. from <i>Distichodus rostratus</i>	64
Southern Sinai, helminths of rodents in	246
Sporocysts, physiology of starvation in	75
<i>Tautogolabrus adspersus</i> , acanthocephala from	185
cestodes from	179
metacercariae from	189
nematodes from	183
trematodes from	173
Technique for recovering nematode larvae from herbage ...	219
Tobacco roots infected with <i>Aphelenchoides</i> n.sp.	323
<i>Paraphelenchus</i> n.sp.	326
<i>Trichinella spiralis</i> , activity of cambendazole against ...	211
rate of larval production of	357
removal of, with methyridine	358
<i>Trichostrongylus axei</i> , acquired immunity in lambs	202

DATES OF PUBLICATION OF VOLUME XLIII

Parts 1/2 ...	11th May, 1969
Parts 3/4 ...	29th December, 1969

CORRIGENDUM IN VOLUME XLIII

Page 284, Table I, Line 10, 3rd column, for 2·5 read ·125.