

EPP1308

A mixed-methods analysis of patient safety incidents involving opioid substitution treatment with methadone or buprenorphine in community-based care in England and Wales

R. Gibson^{1*}, N. Macleod², L. Donaldson³, H. Williams², A. Sheikh⁴ and A. Carson-Stevens²

¹Community Mental Health Team, Livewell Southwest, Plymouth, United Kingdom; ²Division Of Population Medicine, Cardiff University, Cardiff, United Kingdom; ³Department Of Non-communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, United Kingdom and ⁴School Of Medicine, University of Edinburgh, Edinburgh, United Kingdom

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1488

Introduction: There is a paucity of knowledge and understanding of medical error in opioid substitution treatment programmes.

Objectives: To characterise patient safety incidents involving opioid-substitution treatment with methadone or buprenorphine in community-based care to identify the sources and nature of harm, describe and interpret themes and use this qualitative analysis to identify priorities to focus future improvement work.

Methods: We undertook a mixed-methods study examining incidents involving opioid substitution treatment with methadone or buprenorphine in community-based care submitted between 2005 and 2015 from the National Reporting and Learning System, a repository of incident reports from England and Wales. We analysed each report using four frameworks to identify incident type, contributory factors, incident outcome and severity of harm. Analysis involved detailed data coding and iterative generation of data summaries using descriptive statistical and thematic analysis.

Results: 2,284 reports were identified. We found that most risks of harm came from failure in one of four processes of care delivery: prescribing opiate-substitution (n=151); supervised dispensing errors (n=248); non-supervised dispensing errors (n=318); and monitoring and communication activities (n=1544). Most incidents resulting in harm involved supervised or non-supervised dispensing (n=91/127, 72%). Staff- (e.g. mistakes, not following protocols) and organisation-related (e.g. poor working conditions or poor continuity of care between services) contributory factors were present for over half of incidents.

Conclusions: We have identified four processes of care delivery and associated contributory factors, which represent potential target areas for healthcare systems worldwide to develop interventions to improve the safe delivery of opioid substitution treatment.

Keywords: Opioid; Patient safety; Mixed methods

EPP1307

Family functioning and therapeutic success in detoxification in dual diagnosis patients

S. Pombo Chorto

Medicine And Surgery, Rovira i Virgili University, Reus, Spain
doi: 10.1192/j.eurpsy.2021.1489

Introduction: Voluntary dropouts before completion of inpatient psychiatric treatment are common in patients with dual disorders.

However, the socioeconomic profile and family structure of patients who undergo a therapeutic program may correlate with their motivation and participation in the treatment received, as well as with their subsequent recovery.

Objectives: To evaluate the family typology of patients with dual diagnosis according to the degree of intrafamily cohesion and adaptability correlated with the prediction of therapeutic success and the risk of voluntary abandonment of hospital detoxification treatment.

Methods: A total of 211 patients admitted to an inpatient psychiatric unit with substance use disorders were studied. Data were obtained from two sources: (1) interview of participants, (2) review of participants' medical records using the Maudsley Addiction Profile (MAP) and the Family Cohesion and Adaptability Assessment Scale (FACES III).

Results: The 127 subjects who completed the hospital detoxification program had significantly lower MAP and FACES III scores at baseline than the 84 subjects who did not complete the study. Those who did not complete the admission reported greater addictive severity and poorer family functioning. Family cohesion and adaptability measured with FACES III and addictive severity assessed with MAP positively correlated with successful compliance with the hospital treatment program for dual diagnosis patients.

Conclusions: These findings reveal the association of psychosocial and family determinants and addictive severity with treatment completion and subsequent prognostic evolution. Recognizing these predictive characteristics may allow early identification of patients at higher risk of early dropout and prevent it by increasing the intensity of treatment.

Keywords: dual diagnosis; family functioning; patient dropout; drug detoxification

EPP1308

Latent by-product of substance use: Burden of care

A. Channa^{1*} and H. Mohsin²

¹Psychiatry, Heath Lane Hospital, Birmingham, United Kingdom and ²Liaison Psychiatry, St. Vincent's hospital, Dublin, Ireland

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1490

Introduction: Substance use affects it's user and also risks the health of the caregivers.



Objectives: Identify persons at risk of developing substance use disorder, assess the burden borne by the caregivers and development of psychiatric illness.

Methods: Clinical assessment based on DSM-V criteria was performed for SUD diagnosis. Data was recorded using Substance use risk profile scale (SURPs) on the patient and the caregivers were evaluated using M.I.N.I. International Neuropsychiatric Interview (M.I.N.I) and caregiver's strain index (CSI).

Results: 81 participants-96% were male, mean age 32.4 years, 53.1% married, 72.8% employed and 52% lived in joint family system). The substance use ascertained were alcohol 24.7%, benzodiazepines 21%, cannabis 34.6%, opioid 30.9% and others 4.8%. 50% had substance use lasting 2-9 years. 50.6% reported starting as a recreation and the perpetuating factor for 49.4%. was emotional distress. 44% quit due to family pressure. On SURP, 85.2% demonstrated anxiety sensitivity, 96.3% were hopeful, 66% sensation seeking and 77% were impulsive. Caregiver mean age was 37.8 years, with two-third being parents and spouses. The burden reported was sleep

disturbance 59.3%, inconvenience (61.7%) physical strain 46.9%, confining 50.6%, family adjustment 76.5%, plan changes 65.4%, emotional adjustment 88.9%, behavioral adjustment 74.1%, financial strain 80.2%, work adjustment 46.9%, 71.6% felt overwhelmed and 67.9% were upset about the changes from former self. Major depressive disorder was identified in 51.9% of the caregivers.

Conclusions:



SURP identified personality features linked with risk of developing substance use disorder. The study also provided evidence for significant burden on caregivers and an increased likelihood to develop a psychiatric disorder.

Keywords: Caregiver burden; SURP; Caregiver mental health; Substance use

EPP1309

Pregabalin addiction characteristics at oran: A cohort study.

C. Yasmine* and D. Aicha

Addictology Service, specialized hospital establishment in psychiatry sid chami, oran, Algeria

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1491

Introduction: Pregabalin is an analogue of gamma-aminobutyric acid (GABA). Recent reports suggest illicit pregabalin use may be increasing among youth, however its addictive potential has not been well established (1).

Objectives: Drug seeking behavior and chronic drug use are associated with deficits in glutamate clearance and activation of post-synaptic glutamatergic receptors (2). Based upon multiple studies, we compare here the addiction and misuse risks of pregabalin with those of traditional psychoactive substances (3).

Methods: Users of pregabalin were identified from 1st January 2019 to 31 December 2019 in Oran Addictology service, at west Algeria. The aim of the study was to establish the addictive potential of pregabalin and to compare the addiction risks of pregabalin with traditional psychoactive substances in west Algerian population. Clinical diagnosis was established according DSM-5 diagnosis criteria.

Results: A total of 92 cases of pregabalin abuse or dependence were identified. The principal population at risk consists of patients with other current or past substance use disorders, for the most part opioid and multi-drug users, the age group were between 17-38 years old, mostly single men. The mean daily dose of pregabalin was 1200 mg. Almost all patients experienced withdrawal symptoms when pregabalin was discontinued.

Conclusions: The misuse of pregabalin often leads to abuse and dependence, mostly in the context of multiple drug addiction, especially in youth population.

Keywords: addiction; pregabalin; misuse; abuse

EPP1310

Generalized problematic internet use, use of social networks, and appearance schemas in late adolescence

B. Rodrigues Maia^{1*}, H. Moreira², A. Macedo³ and A.T. Pereira⁴

¹Faculty Of Philosophy And Social Sciences, Universidade Católica Portuguesa, Braga, Portugal; ²Center For Research In Neuropsychology And Cognitive Behavioral Intervention, University of Coimbra, Coimbra, Portugal; ³Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal and ⁴Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1492

Introduction: Research on the association between internet use and concerns about body image is still scarce.

Objectives: To explore the associations between generalized problematic internet use, number of social networks, and appearance schemas.

Methods: 216 Portuguese late adolescents (89.3% females), with a mean age of 18.62 years old (SD = .488, range: 18-19) filled in an internet patterns of use questionnaire, the Generalized Problematic Internet Use Scale 2 (GPIUS2) and The Appearance Schemas Inventory-Revised (ASI-R).

Results: A total of 99.6% of the students use social networks. Subjects were divided into three groups (group1: 1-2 social networks; group 2: 3 social networks, and group 3: >3 social networks). There was a statistically difference in Motivational Saliency scores (ASI-R) for the three groups [F (2, 503) = 6.0, p = .003]. Post-hoc comparisons indicated that the mean score for group 3 (M = 28.29, SD = 4.95) was significantly different from group 1 (M = 26.69, SD