

Metallicity of M dwarfs: The link to exoplanets

V. Neves^{1,2}, X. Bonfils² and N. C. Santos^{1,3}

¹Centro de Astrofísica, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal
email: vasco.neves@astro.ua.pt

²Laboratoire d'Astrophysique, Observatoire de Grenoble, BP 53, F-38041 Grenoble Cédex 9,
France

³Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto,
Portugal

Abstract. The determination of the stellar parameters of M dwarfs is of prime importance in the fields of galactic, stellar and planetary astronomy. M stars are the least studied galactic component regarding their fundamental parameters. Yet, they are the most numerous stars in the galaxy and contribute to most of its total (baryonic) mass. In particular, we are interested in their metallicity in order to study the star-planet connection and to refine the planetary parameters. As a preliminary result we present a test of the metallicity calibrations of Bonfils *et al.* (2005), Johnson & Apps (2009), and Schlaufman & Laughlin (2010) using a new sample of 17 binaries with precise V band photometry.

Keywords. stars: fundamental parameters, planetary systems, stars: late-type, stars: abundances, stars: atmospheres

Preliminary Results

We tested the metallicity calibration of Bonfils *et al.* (2005) (hereafter B05), as well as the calibrations of Johnson & Apps (2009) (hereafter JA09) and Schlaufman & Laughlin (2010) (hereafter SL10), with a sample of 17 M dwarf secondaries with a wide (> 5 arcsec separation) physical FGK companion.

Following B05, three papers with different calibrations were published: JA09, SL10, and Rojas-Ayala *et al.* (2010) (hereafter RA10). Each work claims a calibration with a better precision than the previous ones, and in general, poor V photometry is identified as a serious limitation. In order to address the photometric limitation, only M stars with precise V photometry ($\sigma < 0.04$ mag) were selected. Most stars have V magnitude uncertainties of 0.01 or 0.02 mag. Note that the RA10 calibration was not tested because it requires IR indices that we do not have. This test will be done in the near future.

We found that the metallicity values of our stars (obtained from the FGK primary component) are in reasonable agreement with the [Fe/H] values obtained with all calibrations, as can be seen in Fig. 1. However, our calibrators are found to be more metal poor (on average) than both JA09 and SL10 calibrations.

A better photometry did not improve the dispersion measured around the different calibrations. This means that precision on V photometry may not be the main limitation in the derivation of the [Fe/H] calibration.

Table 1. shows a quantitative comparison between the calibrations. We note that the rms, RMS_P and the R_{ap}^2 values were offset-corrected. In general, the calibrations have similar offsets, rms, RMS_P , and correlation coefficients.

Table 1. Comparison of the residuals offset, *rms*, residual mean square (RMS_P), and adjusted square of the multiple correlation coefficient (R_{ap}^2) of the calibrations of Bonfils *et al.* (2005), Johnson & Apps (2009), and Schlafman & Laughlin (2010) applied to our data. RMS_P and R_{ap}^2 definitions were taken from Schlafman & Laughlin (2010).

Calibration Source + equation	offset (dex)	rms (dex)	RMS_P (dex)	R_{ap}^2
B05 (1) : $[Fe/H] = 0.196 - 1.527M_K + 0.091M_K^2 + 1.886(V - K) - 0.142(V - K)^2$	0.05	0.21	0.06	0.18
B05 (2) : $[Fe/H] = -0.149 - 6.508\Delta M$, $\Delta M = Mass_V - Mass_K$	0.07	0.23	0.06	0.17
JA09 : $[Fe/H] = 0.56\Delta M_K - 0.05$, $\Delta M_K = M_S - M_K$	0.19	0.22	0.05	0.27
SL10 : $[Fe/H] = 0.79\Delta(V - K) - 0.17$, $\Delta(V - K) = (V - K)_{obs} - (V - K)_{fit}$	0.06	0.22	0.05	0.28




Figure 1. *Left panel:* Plot of metallicity versus the difference between masses calculated from the V- and the K-band Mass-Luminosity equations of Delfosse *et al.* (2000). The black line represents the calibration of B05 (2). *Middle panel:* $[Fe/H]$ versus the difference between the mean value of M_K of M dwarfs and the value of M_K of each star, as defined by Johnson & Apps (2009). The black line represents the calibration of JA09. *Right panel:* $[Fe/H]$ versus the observed difference between V and K magnitudes ($V-K$) and the fit of the ($V-K$) corresponding to the horizontal distance (in the $V-K$, M_K plane) between the mean value of M_K of M dwarfs and the value of M_K of each star, as defined by Schlafman & Laughlin (2010). The black line represents the calibration of SL10.

Interestingly, the calibration of B05 (1) has the lowest offset and rms. However, the correlation coefficient is a bit lower than the values of JA09 and SL10. The results are inconclusive and require further study.

Acknowledgements: We acknowledge the support by the European Research Council/European Community under the FP7 through Starting Grant agreement number 239953. NCS also acknowledges the support from Fundação para a Ciência e a Tecnologia (FCT) through program Ciência2007 funded by FCT/MCTES (Portugal) and POPH/FSE (EC), and in the form of grant reference PTDC/CTE-AST/098528/2008. VN would also like to acknowledge the support from FCT in the form of the fellowship SFRH/BD/60688/2009.

References

- Delfosse, X., Forveille, T., Ségransan, D., Beuzit, J.-L., Udry, S., Perrier, C., & Mayor, M. 2000 *A&A*, 364, 217
- Bonfils, X., Delfosse, X., Udry, S., Santos, N. C., Forveille, T., & Ségransan, D. 2005 *A&A*, 442, 635
- Johnson, J. A. & Apps, K. 2009 *ApJ*, 699, 933
- Schlaufman, K. C. & Laughlin, G. 2010, *A&A*, 519, A105
- Rojas-Ayala, B., Covey, K. R., Muirhead, P. S., & Lloyd, J. P. 2010, *ApJ*, 720, L113