

on, decision-making across a mental health organization. With further empirical testing and revision, such a framework can be used to inform the improvement of approaches to making decisions.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Do Medical Students Adopt Problem-Focused or Emotion-Focused Coping Strategies?

Mrs Aisha Hawsawi*, Dr Elena Nixon and Dr Neil Nixon

University of Nottingham, Nottingham, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.160

Aims. The pursuit of a career in medicine, while potentially rewarding, is undeniably accompanied by demanding challenges. These challenges encompass not only rigorous academic demands and long work hours but also contend with a competitive academic environment, conflicts in maintaining a study-life balance, and a multitude of other stressors unique to the medical profession. Amidst this backdrop, concerns are growing worldwide about the mental health challenges that medical students face as they start their careers in medicine. Coping can play a pivotal role in overcoming these challenges. This study explores how coping is associated with wellbeing aspects, i.e., anxiety and depression, as well as personality, and looks into the coping strategies adopted by medical students, specifically focusing on whether they predominantly employ problem-focused or emotion-focused coping. Additionally, it aims to explore contextual factors influencing students' coping strategies, which is crucial for informing wellbeing interventions and support services.

Methods. This study used a mixed-methods approach, employing quantitative data on coping, personality, stress, anxiety and depression and qualitative data from semi-structured interviews with preclinical and clinical year medical students at the University of Nottingham.

Results. Regression findings revealed that medical students primarily used emotion-focused over problem-focused coping. Interestingly, thematic analysis showed that medical students employ problem-focused coping strategies in rigorous, academically challenging and controllable situations such as upcoming exams; they prioritise structured study schedules, seek additional academic resources, and actively engage with faculty to enhance their understanding of complex topics; conversely, emotion-focused coping emerged prominently in the face of personal or interpersonal stressors, particularly in situations perceived as uncontrollable. In such instances, like unexpected setbacks or health concerns, students may acknowledge and express their emotions and engage in activities for emotional relief, including seeking wellbeing support.

Conclusion. The study reveals a dynamic interplay between problem-focused and emotion-focused coping strategies in medical students. Recognising that medical students tend to adopt different coping strategies in different situations, medical education systems should aim to develop or tailor existing resources to provide appropriate academic and wellbeing support.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Prevalence of Low Mood, Thoughts of Self-Harm and Suicidal Ideation in Women Affected by the Perimenopause and Menopause

Ms Olivia Hendriks^{1*}, Ms Aini Kamal², Dr Dan Reisel², Dr Louise Newson² and Dr Pooja Saini¹

¹Liverpool John Moores University, Liverpool, United Kingdom and

²Newson Health, Stratford-upon-Avon, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.161

Aims. Low mood is a common clinical symptom during the perimenopause and menopause. However, the extent to which low mood in menopausal women intersects with thoughts of self-harm and suicidal ideation is largely unknown. In this study we aimed to explore these questions by using two different validated symptom scores.

Methods. We administered a modified version of the Greene Climacteric Symptom Questionnaire (Greene 1976) to all new patients attending the Newson Health Menopause and Wellbeing Clinic, Stratford-upon-Avon, between 1 September 2023 and 31 December 2023. Patients were also asked to complete the PHQ-9 symptom questionnaire, an instrument for diagnosing and measuring the severity of depression. Data were collected from electronic health records and analysed using descriptive statistics.

Results. 1,212 patients were included in the study and completed the Greene Climacteric and PHQ-9 questionnaires at baseline and after 3 months. Mood and mental health symptoms including self-reported anxiety and depression affected 98% of patients. 16% of respondents indicated that they had thoughts of self-harm or suicidal ideation on at least some days in the 2 weeks prior to their initial appointment (Question 9 of the PHQ-9).

Conclusion. The findings of our study demonstrate that negative mood symptoms are common in perimenopausal and menopausal women. 1 in 6 women reported thoughts of self-harm prior to initiation of HRT. Our observational data suggest that mood symptoms are highly prevalent and some women have severe symptoms and may experience suicidal ideation. Our findings should inform better mental health support and access to treatment for women experiencing negative mood symptoms in the menopause transition.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Healthcare Needs of an Older Adult Population Referred for Psychiatric Assessment in the Emergency Department of a University Hospital

Dr Fiona Hoare^{1*}, Dr Ann O'Donoghue², Dr Colm Sweeney³, Ms Darlene Tan⁴ and Mr Meshari Alabulrahman⁴

¹St Vincent's University Hospital, Dublin 4, Ireland; ²St Patrick's University Hospital, Dublin, Ireland; ³National Forensic Mental Health Service, Dublin, Ireland and ⁴UCD, Dublin, Ireland

*Presenting author.

doi: 10.1192/bjo.2024.162

Aims. It is estimated that approximately 15% of adults over 60 years old suffer from a mental disorder. Self-harm and suicidal ideation are associated with a range of mental disorders, and