

# ON A PROBLEM OF G. GOLOMB.

P. ERDÖS

(received 3 August 1960)

In his paper on sets of primes with intermediate density Golomb<sup>1</sup> proved the following theorem:

Let  $2 < P_1 < P_2 < \dots$  be any sequence of primes for which

$$(1) \quad P_j \not\equiv 1 \pmod{P_i}$$

for every  $i$  and  $j$ . Denote by  $A(x)$  the number of  $P$ 's not exceeding  $x$ . Then

$$(2) \quad \liminf_{x \rightarrow \infty} A(x)/x = 0.$$

It is not difficult to see that in some sense (2) is best possible since it is easy to construct a sequence of primes satisfying (1) for which

$$\limsup_{x \rightarrow \infty} A(x)/x > 0,$$

and in fact the  $\limsup$  can be as close to 1 as we wish. Golomb pointed out that in some ways the most natural sequence satisfying (1) can be obtained as follows:  $q_1 = 3, q_2 = 5, q_3 = 17, \dots, q_k$  is the smallest prime greater than  $q_{k-1}$  for which

$$q_k \not\equiv 1 \pmod{q_i}, \quad 1 \leq i < k.$$

Henceforth we will only consider this special sequence satisfying (1). We shall prove the following (as before  $A(x)$  denotes the number of  $q_i \leq x$ ).

THEOREM.

$$A(x) = (1 + o(1)) \frac{x}{\log x \log \log x}.$$

$\log_k x$  will denote the  $k$  times iterated logarithm,  $c_1, c_2, \dots$  will denote positive absolute constants.

Our method will be similar to the one used in our recent joint paper with Jabotinsky<sup>2</sup>, but we will also need Brun's method and the results on primes in short arithmetic progressions.

<sup>1</sup> S. Golomb, *Math. Scand.* 3 (1955), 264—74.

<sup>2</sup> P. Erdős and E. Jabotinsky, *Indig. Math.* 20 (1958), 115—128.

LEMMA 1. Denote by  $\pi(x, k, l)$  the number of primes  $p \leq x, p \equiv l \pmod k, (l, k) = 1$ . Then  $(\exp z = e^z)$

$$(3) \quad \pi(x, k, l) = \frac{x}{\varphi(k) \log x} \left( 1 + O\left(\frac{1}{\log x}\right) \right)$$

uniformly for all  $k < \exp(c_1 \log x / \log \log x)$ , except possibly for the multiples of a certain  $k^* = k^*(x)$  where  $k^* > (\log x)^A$  ( $A$  is an arbitrary constant, but the constant in  $O(1/\log x)$  depends on  $A$ ).

Lemma 1 is well known<sup>3</sup>.

LEMMA 2. Let  $2 = p_1 < p_2 < \dots$  be the sequence of consecutive primes, and let  $r$  be a fixed integer,  $0 \leq r_i < r$ . Denote by  $N_k(x)$  the number of integers  $1 \leq z \leq x$  for which  $z \equiv l \pmod k, (l, k) = 1$  and

$$z \not\equiv a_i^{(j)} \pmod{p_i}, \quad 1 \leq j \leq r_i$$

where the  $a_i^{(j)}$  are arbitrary residues and  $p_i \leq x$ . Then

$$N_k(x) < c_2 \frac{x}{k} \prod_{p_i \leq x/k} (1 - r_i/p_i).$$

The proof follows immediately from Brun's method<sup>4</sup>.

LEMMA 3. There exists a constant  $c_3$  so that

$$(4) \quad \log_3 x - c_3 < \sum_{q_i \leq x} 1/q_i < \log_3 x + c_3.$$

First we prove the upper bound. If the upper estimation in (4) would not hold then for every  $c$  there would be arbitrarily large values of  $x$  so that for every  $z < x$

$$(5) \quad \sum_{q_i \leq x} \frac{1}{q_i} - \log_3 x > \sum_{q_i \leq z} \frac{1}{q_i} - \log_3 z$$

and

$$(6) \quad \sum_{q_i \leq x} \frac{1}{q_i} > \log_3 x + c.$$

Let  $x^{1/2} < q_i \leq x$ . Clearly by the definition of the  $q$ 's  $q_i \not\equiv 0 \pmod p$  for all  $p < x^{1/2}$  and  $q_i \not\equiv 1 \pmod{q_j}$  for  $q_j < x^{1/2}$ . Thus by lemma 2 ( $k = 1$ )

$$(7) \quad A(x) < x^{1/2} + c_2 x \prod_{p_i \leq x^{1/2}} (1 - r_i/p_i)$$

where  $r_i = 2$  if  $p_i$  is a  $q$  and is 1 otherwise. From (7), (6) and from  $\prod_{p < x^{1/2}} (1 - 1/p) < c_4/\log x$

$$(8) \quad A(x) < c_5 \frac{x}{\log x} \prod_{q_i < x^{1/2}} \left( 1 - \frac{1}{q_i} \right) < c_6 x \exp(-c)/\log x \log_2 x.$$

<sup>3</sup> This is Theorem 2.3 p. 230 of Prachar's book *Primzahlverteilung* (Springer 1957) where the literature of this question can be found.

<sup>4</sup> See e.g. P. Erdős, *Proc. Cambridge Phil. Soc.* 34 (1957), 8.

The last inequality in (8) follows from  $\prod_{q_i < x} (1 - 1/q_i) < c_7 \exp(-\sum_{q_i < x} 1/q_i)$  and from (using (6))

$$\sum_{q_i < x^{1/2}} \frac{1}{q_i} > \sum_{q_i \leq x} \frac{1}{q_i} - \sum_{x^{1/2} \leq p \leq x} \frac{1}{p} > \log_3 x + c - c_8.$$

From (8) we have

$$(9) \quad \sum_{x/2 < q_i \leq x} \frac{1}{q_i} < \frac{2A(x)}{x} < 2c_6 \exp(-c) / \log x \log \log x.$$

But from (5) we have for  $z = x/2$

$$\sum_{x/2 < q_i \leq x} \frac{1}{q_i} > \log_3 x - \log_3 \frac{x}{2} > c_9 / \log x \log_2 x,$$

which contradicts (9) for sufficiently large  $c$ . Thus the upper bound in (4) is proved.

The proof of the lower bound will be more complicated. Put  $y = \exp(\log x / (\log \log x)^{10})$  and denote by  $A_y(x)$  the number of primes  $p \leq x$  satisfying

$$(10) \quad p \not\equiv 1 \pmod{q_i}, \quad 3 \leq q_i \leq y.$$

We evidently have

$$(11) \quad A_y(x) - \sum_{y < q_j < x} B(x, q_j) < A(x) < A_y(x) + y$$

where  $B(x, q_j)$  denotes the number of primes  $p \leq x$  satisfying

$$p \equiv 1 \pmod{q_j}, \quad p \not\equiv 1 \pmod{q_i}, \quad 3 \leq q_i \leq y.$$

Now we estimate  $A_y(x)$  by Brun's method.

LEMMA 4.

$$A_y(x) = (1 + o(1)) \frac{x}{\log x} \prod_{q_i \leq y} \left(1 - \frac{1}{q_i - 1}\right).$$

By the sieve of Eratosthenes we have

$$A_y(x) = \pi(x) - \sum \pi(x, q_i, 1) + \sum \pi(x, q_i, q_{i_1}, 1) - \dots$$

where  $3 \leq q_i \leq y$  and  $i$ 's are distinct. By the well known idea of Brun<sup>5</sup> we have  $(\sum_r = \sum \pi(x, q_{i_1} \cdot q_{i_2} \cdot \dots \cdot q_{i_r}, 1))$ .

$$(12) \quad \pi(x) - \Sigma_1 + \Sigma_2 - \Sigma_3 + \dots - \Sigma_{2k-1} < A_y(x) < \pi(x) - \Sigma_1 + \Sigma_2 - \dots + \Sigma_{2k}.$$

We now choose  $k = [10 \log_2 x]$ . We distinguish two cases. In the first case none of the numbers  $q_{i_1} \cdot \dots \cdot q_{i_r}, 1 \leq r \leq 2k$  are exceptional from the point

<sup>5</sup> See e.g. E. Landau, *Zahlentheorie* Vol. 1.

of view of Lemma 1. In this case we can estimate  $\Sigma_r$  by Lemma 1 and following say Landau's treatment of Brun's method<sup>5</sup> we obtain from (12) by a simple computation

$$(13) \quad A_\nu(x) = \frac{x}{\log x} \prod_{3 \leq q_i \leq \nu} \left(1 - \frac{1}{q_i - 1}\right) + O\left(\frac{x}{(\log x)^2}\right) \prod_{3 \leq q_i \leq \nu} \left(1 + \frac{1}{q_i - 1}\right) + o\left(\frac{x}{(\log x)^2}\right).$$

By the upper bound of (4) we have

$$\prod_{q_i \leq \nu} \left(1 + \frac{1}{q_i - 1}\right) < c_{10} \log_2 x \quad \text{and} \quad \prod_{q_i \leq \nu} \left(1 - \frac{1}{q_i - 1}\right) > c_{10}/\log_2 x,$$

thus from (13) we obtain Lemma 4 in the first case.

In the second case let  $d = q_{i_1} \cdot q_{i_2} \cdots q_{i_r}$  be the smallest exceptional number (i.e. for which Lemma 1 does not hold). By Lemma 1 we can assume that  $d > (\log x)^4$ . We estimate  $\pi(x, td, 1)$  from below by 0 and from above by  $x/td$ . Since

$$\sum_{t < x} \frac{x}{td} = O\left(\frac{x \log x}{d}\right) = o\left(\frac{x}{(\log x)^2}\right)$$

we can neglect this exceptional  $d$  and the proof of Lemma 4 is complete.

Now we complete the proof of Lemma 3. Assume that the lower bound in (4) is false. Then for every  $c_3$  there are infinitely many integers  $x$  satisfying for every  $z \leq x$

$$(14) \quad \sum_{q_i \leq x} \frac{1}{q_i} - \log_3 x < \sum_{q_i \leq z} \frac{1}{q_i} - \log_3 z$$

and

$$(15) \quad \sum_{q_i \leq x} \frac{1}{q_i} = \log_3 x - c_\infty, \quad c_\infty > c_3.$$

From (14) we have

$$(16) \quad \sum_{z < q_i < x} \frac{1}{q_i} < \log_3 x - \log_3 z.$$

By Lemma 4 and (16) (since  $\log_3 x - \log_3 y = o(1)$ )

$$(17) \quad A_\nu(x) - A_\nu\left(\frac{x}{2}\right) = (1 + o(1)) \frac{x}{2 \log x} \prod_{q_i \leq \nu} \left(1 - \frac{1}{q_i - 1}\right) > c_{11} \frac{x \exp c_\infty}{\log x \log_2 x}.$$

Thus from (11) and (17)

$$(18) \quad A(x) - A\left(\frac{x}{2}\right) > c_{11} \frac{x \exp c_\infty}{\log x \log_2 x} - y - \sum_{\nu < q_i \leq x} B(x, q_i).$$

Now we estimate  $\sum_{\nu < q_i \leq x} B(x, q_i)$ . Write

<sup>5</sup> See e.g. E. Landau, *Zahlentheorie* Vol. 1.

$$(19) \quad \sum_{\nu < q_j \leq x} B(x, q_j) = \Sigma_1 + \Sigma_2 + \Sigma_3$$

where in  $\Sigma_1 y < q_j \leq x \exp(-\log x / (\log_2 x)^{1/2})$  in  $\Sigma_2 x \exp(-\log x / (\log_2 x)^{1/2}) < q_j \leq x \exp(-\log x / (\log_2 x)^{5/4})$  and in  $\Sigma_3 x \exp(-\log x / (\log_2 x)^{5/4}) < q_j \leq x$ . From Lemma 2 we have for the  $q_j$  in  $\Sigma_1$  and  $\Sigma_2$ .

$$(20) \quad B(x, q_j) < c_2 \frac{x}{q_j \log \frac{x}{q_j}} \prod' \left(1 - \frac{1}{q_i}\right) < c_2 \frac{x}{q_j \log \frac{x}{q_j}} \prod_{q_i < \nu} \left(1 - \frac{1}{q_i}\right)$$

where in  $\prod' q_i < \min(q_j, x/q_j)$ . (20) holds since for the  $q_j$  in  $\Sigma_1$  and  $\Sigma_2 \min(q_j, x/q_j) > y$ . Now from (16)  $\sum_{\nu < q_i \leq x} 1/q_i < \log_3 x - \log_3 y = o(1)$ . Thus from (15)

$$(21) \quad \sum_{q_i < \nu} \frac{1}{q_i} = \log_3 x - c_x - o(1).$$

From (20) and (21) we have for the  $q_j$  in  $\Sigma_1$

$$(22) \quad B(x, q_j) < c_{12} \frac{x \exp c_x}{q_j \log \frac{x}{q_j} \log_2 x} < c_{12} \frac{x \exp c_x}{q_j \log x (\log_2 x)^{1/2}}.$$

But from (16)

$$(23) \quad \Sigma_1 \frac{1}{q_j} \leq \sum_{\nu < q_j \leq x} \frac{1}{q_j} < \log_3 x - \log_3 y < c_{13} \log_3 x / \log_2 x$$

Thus from (22) and (23)

$$(24) \quad \Sigma_1 < c_{12} \frac{x \exp c_x}{\log x (\log_2 x)^{1/2}} \Sigma_1 \frac{1}{q_j} < c_{12} c_{13} \frac{x \log_3 x \exp c_x}{\log x (\log_2 x)^{3/2}} = o\left(\frac{x \exp c_x}{\log x \log_2 x}\right).$$

Again from (20), (21) and (16) we obtain as in the estimation

$$(25) \quad \Sigma_2 < c_{14} \frac{x (\log_2 x)^{1/4} \exp c_x}{\log x} \Sigma_2 \frac{1}{q_j} < c_{14} c_{15} \frac{x \exp c_x}{\log x (\log_2 x)^{5/4}} = o\left(\frac{x \exp c_x}{\log x \log_2 x}\right).$$

To estimate  $\Sigma_3$  denote by  $N(a, x)$  the number of primes  $p < x/a, a < x^{1/2}$ , for which  $a \cdot p + 1$  is also a prime. A well known consequence of Brun's method implies that

$$(26) \quad N(a, x) < c_{16} \frac{x}{(\log x)^2} \prod_{p|a} \left(1 + \frac{1}{p}\right).$$

(26) easily follows from Lemma 2. From (26) we have by interchanging the order of summation ( $\sum'$  denotes that  $1 \leq a < \exp(\log x / (\log_2 x)^{5/4})$ )

$$(27) \quad \Sigma_3 \leq \sum' N(a, x) < c_{16} \frac{x}{(\log x)^2} \sum' \frac{\prod_{p/a} \left(1 + \frac{1}{p}\right)}{a} < c_{17} \frac{x}{\log x (\log_2 x)^{5/4}}.$$

The last inequality of (27) holds since it is well known that

$$(28) \quad \sum_{a=1}^z \frac{\prod_{p/a} \left(1 + \frac{1}{p}\right)}{a} < c_{18} \log z.$$

((28) follows easily from the well known result  $\sum_{a=1}^z \prod_{p/a} (1 + 1/p) < \sum_{a=1}^z \sigma(a)/a = (1 + o(1))\pi^2/6 \log z$  by partial summation). From (24), (25) and (27) we obtain

$$(29) \quad \sum_{y \leq q_i \leq x} B(x, q_i) = o\left(\frac{x \exp c_x}{\log x \log_2 x}\right).$$

From (18) and (29) we have

$$(30) \quad A(x) - A\left(\frac{x}{2}\right) > c_{19} \frac{x \exp c_x}{\log x \log_2 x}.$$

(30) implies that

$$(31) \quad \sum_{(x/2) < q_i < x} \frac{1}{q_i} > c_{19} \exp c_x / \log x \log_2 x.$$

On the other hand (16) implies that

$$\sum_{(x/2) < q_i < x} \frac{1}{q_i} < \log_3 x - \log_3 \frac{x}{2} < c_{20} / \log x \log_2 x$$

an evident contradiction for sufficiently large  $c_3$  ( $c_x > c_3$ ). Thus the upper bound of (4) is proved and the proof of Lemma 3 is complete.

From the upper bound in (4), (19), (24), (25) and (27) we immediately obtain (we now know that  $c_x < c_3$ )

$$(32) \quad \sum_{y \leq q_i \leq x} B(x, q_i) = o\left(\frac{x}{\log x \log_2 x}\right).$$

From (11), (32) and Lemmas 3 and 4 we obtain

$$(33) \quad \begin{aligned} A(x) &= (1 + o(1)) \frac{x}{\log x} \prod_{q_i \leq y} \left(1 - \frac{1}{q_i - 1}\right) + o\left(\frac{x}{\log x \log_2 x}\right) \\ &= (1 + o(1)) \frac{x}{\log x} \prod_{q_i \leq y} \left(1 - \frac{1}{q_i - 1}\right). \end{aligned}$$

The last inequality of (33) follows, since by the lower bound in (4)  $\prod_{q_i \leq y} (1 - 1/(q_i - 1)) > c_{21} / \log_2 x$ . From (33) and the lower bound in (4)

$$(34) \quad A(x) < c_{22} x / \log x \log_2 x \quad (\text{since } \prod_{q_i < y} \left(1 - \frac{1}{q_i - 1}\right) < c_{23} / \log_2 x).$$

Thus by a simple computation

$$(35) \quad \sum_{y \leq q_i \leq x} \frac{1}{q_i} = o(1).$$

From (33) and (35) we finally obtain

$$(36) \quad A(x) = (1 + o(1)) \frac{x}{\log x} \prod_{q_i \leq x} \left(1 - \frac{1}{q_i - 1}\right).$$

To complete the proof of our Theorem we only have to show that

$$(37) \quad \prod_{q_i \leq x} \left(1 - \frac{1}{q_i - 1}\right) = \frac{1 + o(1)}{\log_2 x}.$$

Assume that (37) does not hold. Assume first that

$$(38) \quad \limsup \log_2 x \prod_{q_i \leq x} \left(1 - \frac{1}{q_i - 1}\right) = c > 1.$$

The limit of the expression in (38) cannot exist. For if it would exist it would equal  $c > 1$ . But then by (36)

$$\lim \frac{A(x) \log x \log_2 x}{x} = c, \quad \text{or} \quad \lim \frac{q_n}{n \log n \log_2 n} = \frac{1}{c} < 1$$

which contradicts (38).

Since the limit in (38) does not exist it follows by a simple argument that there exists a constant  $c'$ ,  $1 < c' < c$  and two infinite sequences  $x_k < z_k$  so that

$$(39) \quad \lim_{k \rightarrow \infty} \log_2 x_k \prod_{q_i \leq x_k} \left(1 - \frac{1}{q_i - 1}\right) = c'$$

$$(40) \quad \lim_{k \rightarrow \infty} \log_2 z_k \prod_{q_i \leq z_k} \left(1 - \frac{1}{q_i - 1}\right) = c$$

and for every  $x_k < w < z_k$

$$(41) \quad \log_2 x_k \prod_{q_i \leq x_k} \left(1 - \frac{1}{q_i - 1}\right) < \log_2 w \prod_{q_i \leq w} \left(1 - \frac{1}{q_i - 1}\right)$$

From (34) we have for every  $\alpha > 1$

$$(42) \quad \prod_{x < q_i < \alpha x} \left(1 - \frac{1}{q_i - 1}\right) = 1 + o(1).$$

Thus from (39), (40) and (42)  $z_k/x_k \rightarrow \infty$ . Choose now  $w = (1 + \eta)x_k < z_k$  where  $\eta > 0$  is a sufficiently small constant. Put

$$U_k = A[(1 + \eta)x_k] - A(x_k).$$

From (41) we have

$$(43) \quad \frac{\log_2 x_k}{\log_2 [x_k(1 + \eta)]} < \prod_{x_k < q_i < (1+\eta)x_k} \left(1 - \frac{1}{q_i - 1}\right) < \left(1 - \frac{1}{(1 + \eta)x_k}\right)^{U_k}.$$

From (36), (39) and (42) we have

$$(44) \quad U_k = (1 + o(1)) \frac{c'(1 + \eta)x_k}{\log x_k \cdot \log_2 x_k} - (1 + o(1)) \frac{c'x_k}{\log x_k \log_2 x_k} = \frac{(1 + o(1))c'\eta x_k}{\log x_k \log_2 x_k}.$$

Now by a simple computation

$$(45) \quad \frac{\log_2 x_k}{\log_2 [x_k(1 + \eta)]} = 1 - \frac{\log(1 + \eta)}{\log x_k \log_2 x_k} + o\left(\frac{1}{\log x_k \log_2 x_k}\right).$$

From (43), (44) and (45) we have

$$(46) \quad \begin{aligned} 1 - \frac{\log(1 + \eta)}{\log x_k \log_2 x_k} + o\left(\frac{1}{\log x_k \log_2 x_k}\right) &< \left(1 - \frac{1}{(1 + \eta)x_k}\right)^{U_k} \\ &= 1 - \frac{c'\eta}{(1 + \eta) \log x_k \log_2 x_k} + o\left(\frac{1}{\log x_k \log_2 x_k}\right). \end{aligned}$$

But (46) is false for sufficiently small  $\eta$  (since  $c' > 1$ ). This contradiction shows that the  $\overline{\lim}$  in (38) equals 1. In the same way we can show that the  $\underline{\lim}$  of the expression in (38) is 1. Thus (37) is proved, and (36) implies our Theorem.

I do not know whether for infinitely many  $i$ 's  $q_{i+1}$  is the least prime greater than  $q_i$ .

By similar arguments we can prove the following more general result:

Let  $r \geq 1$ ,  $Q_1 > r + 1$ ,  $Q_1$  prime.  $Q_{i+1}$  is the smallest prime greater than  $Q_i$  so that  $Q_i \not\equiv t \pmod{Q_j}$ ,  $1 \leq j \leq i$ ,  $1 \leq t \leq r$ .

Denote by  $B_{Q_i, r}(x)$  the number of  $Q$ 's not exceeding  $x$ , then

$$(47) \quad B_{Q_i, r}(x) = (1 + o(1)) \frac{x}{\log x \log_2 x \cdots \log_{r+1} x}.$$

For  $Q_1 = 3$ ,  $r = 1$ ,  $A(x) = B_{Q_1, r}(x)$ , (47) is thus a generalisation of our Theorem.

Technion,  
Haifa.