



RESEARCH ARTICLE

Hyperpolygonal arrangements

Lorenzo Giordani¹, Paul Mücksch², Gerhard Röhrle¹  and Johannes Schmitt¹

¹Fakultät für Mathematik, Ruhr-Universität Bochum, D-44780, Bochum, Germany

²Technische Universität Berlin, Institut für Mathematik, D-10587, Berlin, Germany

Corresponding author: Gerhard Röhrle; Email: gerhard.roehrle@rub.de

Received: 28 February 2025; **Revised:** 2 June 2025; **Accepted:** 30 May 2025

Keywords: free arrangements; factored arrangements; formal arrangements; simplicial arrangements; $K(\pi, 1)$ arrangements

2020 Mathematics Subject Classification: Primary - 52C35, 14N20, 32S25, 32S22

Abstract

In [5], a particular family of real hyperplane arrangements stemming from hyperpolygonal spaces associated with certain quiver varieties was introduced which we thus call *hyperpolygonal arrangements* \mathcal{H}_n . In this note, we study these arrangements and investigate their properties systematically. Remarkably, the arrangements \mathcal{H}_n discriminate between essentially all local properties of arrangements. In addition, we show that hyperpolygonal arrangements are projectively unique and combinatorially formal.

We note that the arrangement \mathcal{H}_5 is the famous counterexample of Edelman and Reiner [17] of Orlik’s conjecture that the restriction of a free arrangement is again free.

1. Introduction and main results

In [5], Bellamy et al introduced a particular family of real hyperplane arrangements in \mathbb{R}^n stemming from hyperpolygon spaces realized as certain quiver varieties which we call *hyperpolygonal arrangements* \mathcal{H}_n , see Definition 1.1. In this note, we study these hyperpolygonal arrangements and investigate their properties in a systematic manner. It turns out that the arrangements \mathcal{H}_n differentiate essentially between all local properties of arrangements, see Theorem 1.2. In addition, we show that hyperpolygonal arrangements are projectively unique, see Theorem 1.5, and combinatorially formal, see Theorem 1.6.

We briefly indicate how the arrangements \mathcal{H}_n arise in [5] and show their connection to birational geometry, see [5] for details and references. The arrangement \mathcal{H}_n characterizes stability conditions on the parameter θ of the hyperpolygon space $X_n(\theta)$. Namely, $X_n(\theta)$ is smooth if and only if θ does not lie on any hyperplane in \mathcal{H}_n . As explained in [5], the varieties $X_n(0)$ are *conical symplectic varieties* and the map $X_n(\theta) \rightarrow X_n(0)$ is a crepant projective resolution (hence a symplectic projective resolution), if θ does not lie in \mathcal{H}_n . Furthermore, two such resolutions $X_n(\theta_1) \rightarrow X_n(0)$ and $X_n(\theta_2) \rightarrow X_n(0)$ are isomorphic if θ_1 and θ_2 lie in the same region of the complement of \mathcal{H}_n in \mathbb{R}^n . This construction works for all conical symplectic varieties by Namikawa [33]: For a conical symplectic variety Y , there is a certain “parameter space” containing a hyperplane arrangement that characterizes the isomorphism classes of crepant projective resolutions (or, in general, the \mathbb{Q} -factorial terminalizations) of Y . Very little is known about the hyperplane arrangements arising in this way, although they are essential in understanding the birational geometry of Y , see also [4], [7]. Special cases of conical symplectic varieties are *symplectic quotient singularities* V/G , where V is a symplectic vector space over \mathbb{C} and $G \leq \mathrm{Sp}(V)$ is a finite group maintaining the symplectic form. By [5], the hyperpolygon space $X_n(0)$ is not a quotient singularity for $n > 5$. In contrast, $X_4(0)$ is the Kleinian singularity of type D_4 , that is, the quotient of \mathbb{C}^2 by the quaternion group Q_8 . Further, $X_5(0)$ is the quotient of \mathbb{C}^4 by a symplectic reflection group of order 32. This quotient was extensively studied in [6] and [15].

In this note, we study the hyperpolygonal arrangements \mathcal{H}_n in a systematic manner. We note that \mathcal{H}_5 is the famous counterexample of Edelman and Reiner [17] of Orlik's conjecture that the restriction of a free arrangement is again free.

We first recall the definition from [5]. Fix $n \in \mathbb{N}$. Let $V = \mathbb{R}^n$. Let x_1, \dots, x_n be the dual basis in V^* of the standard \mathbb{R} -basis of V . For $I \subseteq [n] = \{1, \dots, n\}$, define the hyperplane

$$H_I := \ker \left(\sum_{i \in I} x_i - \sum_{j \in [n] \setminus I} x_j \right)$$

in V .

Definition 1.1. *With the notation as above, the hyperpolygonal arrangement \mathcal{H}_n in V is defined as*

$$\mathcal{H}_n := \{\ker x_i \mid i \in [n]\} \cup \{H_I \mid \emptyset \neq I \subseteq [n]\}.$$

For the various notions used in our main theorem, we refer the reader to Section 2.

Theorem 1.2. *Fix $n \in \mathbb{N}$. Then we have*

- (i) \mathcal{H}_n is supersolvable if and only if $n \leq 2$;
- (ii) \mathcal{H}_n is inductively factored if and only if $n \leq 3$;
- (iii) \mathcal{H}_n is inductively free if and only if $n \leq 4$;
- (iv) \mathcal{H}_n is free if and only if $n \leq 5$;
- (v) \mathcal{H}_n is simplicial if and only if $n \leq 4$;
- (vi) \mathcal{H}_n is not $K(\pi, 1)$ if $n \geq 6$.

It follows from Theorem 1.2(v) and Remark 2.19(i) that \mathcal{H}_n is $K(\pi, 1)$ for $n \leq 4$ and \mathcal{H}_n fails to be $K(\pi, 1)$ for $n \geq 6$ by part (vi). It is not known whether \mathcal{H}_5 is $K(\pi, 1)$.

In general, $K(\pi, 1)$ arrangements need not be free, for example see [34, Figure 5.4]. However, for hyperpolygonal arrangements, this does seem to be the case. While \mathcal{H}_5 is free, it is not known whether \mathcal{H}_5 is $K(\pi, 1)$. We can thus formulate

Corollary 1.3. *With the possible exception when $n = 5$, \mathcal{H}_n is $K(\pi, 1)$ if and only if \mathcal{H}_n is free.*

Thus with the possible exception of \mathcal{H}_5 , the hyperpolygonal arrangements \mathcal{H}_n satisfy Saito's Conjecture that for a complexified arrangement freeness implies $K(\pi, 1)$. Of course, the latter is known to be false in general [18].

Theorem 1.2(ii) and (v) imply the following which is in support of a conjecture due to Falk and Randell namely that every (complex) factored arrangement is $K(\pi, 1)$, [22, Probl. 3.12].

Corollary 1.4. *If \mathcal{H}_n is factored, then \mathcal{H}_n is $K(\pi, 1)$.*

A property for arrangements is said to be *combinatorial* if it only depends on the intersection lattice of the underlying arrangement. In this context, our next theorem shows that the class of hyperpolygonal arrangements is very special in the sense that essentially every property we may formulate for members of this class is combinatorial. This is formally captured by the notion of *projective uniqueness* due to Ziegler [49], see Definition 4.1.

Theorem 1.5. *For any $n \in \mathbb{N}$, \mathcal{H}_n is projectively unique.*

Theorem 1.5 implies that for the class of all real arrangements whose underlying matroid admits a realization over \mathbb{R} as a hyperpolygonal arrangement freeness is combinatorial. In particular, Terao's conjecture over \mathbb{R} is valid within this class, cf. [49, Prop. 2.3]. Likewise, asphericity is combinatorial within this class. Whether both these properties in general are longstanding and wide open problems, see [34, Conj. 4.138] and [22, Prob. 3.8].

A hyperplane arrangement is called *formal* provided all linear dependencies among the defining forms of the hyperplanes are generated by ones corresponding to intersections of codimension two. The significance of this notion stems from the fact that complex arrangements with aspherical complements are formal, [21, Thm. 4.2]. In addition, free arrangements are known to be formal, [47, Cor. 2.5], and factored arrangements are formal, [30, Thm. 1.1]. Thus all the properties studied in Theorem 1.2 entail formality. In our next result, we show that indeed all hyperpolygonal arrangements are *combinatorially formal*, see Definition 5.1.

Theorem 1.6. *For any $n \in \mathbb{N}$, \mathcal{H}_n is combinatorially formal.*

Theorem 1.6 is proved in §5, based on results from [30].

We end in §6 with a brief discussion of the rank-generating functions of the poset of regions of the free hyperpolygonal arrangements \mathcal{H}_n .

For general information about arrangements, we refer the reader to [34].

2. Preliminaries

2.1. Hyperplane arrangements

Let \mathbb{K} be a field and let $V = \mathbb{K}^\ell$ be an ℓ -dimensional \mathbb{K} -vector space. A *hyperplane arrangement* is a pair (\mathcal{A}, V) , where \mathcal{A} is a finite collection of hyperplanes in V . Usually, we simply write \mathcal{A} in place of (\mathcal{A}, V) . By Φ_ℓ we denote the empty arrangement in V .

The *lattice* $L(\mathcal{A})$ of \mathcal{A} is the set of subspaces of V of the form $H_1 \cap \dots \cap H_i$ where $\{H_1, \dots, H_i\}$ is a subset of \mathcal{A} . For $X \in L(\mathcal{A})$, we have two associated arrangements, firstly $\mathcal{A}_X := \{H \in \mathcal{A} \mid X \subseteq H\} \subseteq \mathcal{A}$, the *localization of \mathcal{A} at X* , and secondly, the *restriction of \mathcal{A} to X* , (\mathcal{A}^X, X) , where $\mathcal{A}^X := \{X \cap H \mid H \in \mathcal{A} \setminus \mathcal{A}_X\}$. The lattice $L(\mathcal{A})$ is a partially ordered set by reverse inclusion: $X \leq Y$ provided $Y \subseteq X$ for $X, Y \in L(\mathcal{A})$.

A property for arrangements is said to be *local* if it is passed to every localization of the ambient arrangement. For instance, freeness, asphericity, and factoredness are all local (cf. [34, Thm. 4.37], [35, Lem. 1.1], and the proof of [46, Cor. 2.11], respectively). In contrast, this fails for formal arrangements of rank at least 4 (cf. the example following [47, Def. 2.3]).

Throughout, we only consider arrangements \mathcal{A} such that $0 \in H$ for each H in \mathcal{A} . These are called *central*. In that case the *center* $T(\mathcal{A}) := \bigcap_{H \in \mathcal{A}} H$ of \mathcal{A} is the unique maximal element in $L(\mathcal{A})$ with respect to the partial order. A *rank function* on $L(\mathcal{A})$ is given by $r(X) := \text{codim}_V(X)$. The *rank* of \mathcal{A} is defined as $r(\mathcal{A}) := r(T(\mathcal{A}))$.

The *Poincaré polynomial* $\pi(\mathcal{A}, t) \in \mathbb{Z}[t]$ of \mathcal{A} is defined by

$$\pi(\mathcal{A}, t) := \sum_{X \in L(\mathcal{A})} \mu(X)(-t)^{r(X)},$$

and the *characteristic polynomial* $\chi(\mathcal{A}, t) \in \mathbb{Z}[t]$ of \mathcal{A} is defined by

$$\chi(\mathcal{A}, t) := t^\ell \pi(\mathcal{A}, -t^{-1}) = \sum_{X \in L(\mathcal{A})} \mu(X)t^{\dim X},$$

where μ is the Möbius function of $L(\mathcal{A})$, see [34, Def. 2.48, Def. 2.52].

We recall the concept of a generic arrangement from [34, Def. 5.22].

Definition 2.1. *An ℓ -arrangement \mathcal{A} with $r(\mathcal{A}) = r$ is called *generic* if every subarrangement \mathcal{B} of \mathcal{A} of cardinality ℓ is linearly independent and $|\mathcal{A}| > r$, [34, Def. 5.22].*

2.2. Supersolvable arrangements

Let \mathcal{A} be an arrangement. Following [34, §2], we say that $X \in L(\mathcal{A})$ is *modular* provided $X + Y \in L(\mathcal{A})$ for every $Y \in L(\mathcal{A})$, cf. [34, Def. 2.32, Cor. 2.26]. The following notion is due to Stanley [42].

Definition 2.2. Let \mathcal{A} be a central (and essential) ℓ -arrangement. We say that \mathcal{A} is *supersolvable* provided there is a maximal chain

$$V = X_0 < X_1 < \dots < X_{\ell-1} < X_\ell = \{0\}$$

of modular elements X_i in $L(\mathcal{A})$.

Remark 2.3.

- (i) By [34, Ex. 2.28], V , $\{0\}$ and the members in \mathcal{A} are always modular in $L(\mathcal{A})$. It follows that all 0-, 1-, and 2-arrangements are supersolvable.
- (ii) Supersolvability is a local property [42, Prop. 3.2].

2.3. Free arrangements

Free arrangements play a crucial role in the theory of arrangements; see [34, §4] for the definition and basic properties. If \mathcal{A} is free, then we can associate with \mathcal{A} the multiset of its *exponents*, denoted $\exp \mathcal{A}$.

Remark 2.4.

- (i) Generic arrangements are not free, for example see [39, §4.4].
- (ii) Freeness is a local property [34, Thm. 4.37].

Terao's *Factorization Theorem* [44] shows that the Poincaré polynomial of a free arrangement \mathcal{A} factors into linear terms given by the exponents of \mathcal{A} (cf. [34, Thm. 4.137]):

Theorem 2.5. Suppose that \mathcal{A} is free with $\exp \mathcal{A} = \{b_1, \dots, b_\ell\}$. Then

$$\pi(\mathcal{A}, t) = \prod_{i=1}^{\ell} (1 + b_i t).$$

Terao's celebrated *Addition-Deletion Theorem* [43] plays a fundamental role in the study of free arrangements, [34, Thm. 4.51].

Theorem 2.6. Suppose that $\mathcal{A} \neq \Phi_\ell$. Let $H_0 \in \mathcal{A}$. Let $(\mathcal{A}, \mathcal{A}' = \mathcal{A} \setminus \{H_0\}, \mathcal{A}'' = \mathcal{A}^{H_0})$ be a triple of arrangements. Then any two of the following statements imply the third:

- (i) \mathcal{A} is free with $\exp \mathcal{A} = \{b_1, \dots, b_{\ell-1}, b_\ell\}$;
- (ii) \mathcal{A}' is free with $\exp \mathcal{A}' = \{b_1, \dots, b_{\ell-1}, b_\ell - 1\}$;
- (iii) \mathcal{A}'' is free with $\exp \mathcal{A}'' = \{b_1, \dots, b_{\ell-1}\}$.

Theorem 2.6 motivates the notion of an *inductively free* arrangement, [34, Def. 4.53].

Definition 2.7. The class \mathcal{IF} of inductively free arrangements is the smallest class of arrangements subject to

- (i) $\Phi_\ell \in \mathcal{IF}$ for each $\ell \geq 0$;
- (ii) if there exists a hyperplane $H_0 \in \mathcal{A}$ such that both \mathcal{A}' and \mathcal{A}'' belong to \mathcal{IF} , and $\exp \mathcal{A}'' \subseteq \exp \mathcal{A}'$, then \mathcal{A} also belongs to \mathcal{IF} .

Remark 2.8.

- (i) *Inductively free arrangements are free. However, the latter class properly contains the former, cf. [34, Ex. 4.59].*
- (ii) *Inductive freeness is also a local property, thanks to [26, Thm 1.1].*
- (iii) *Supersolvable arrangements are inductively free, [34, Thm. 4.58].*

2.4. Nice arrangements

The notion of a *nice* or *factored* arrangement goes back to Terao [46]. It generalizes the concept of a supersolvable arrangement. We recall the relevant notions and results from [46] (cf. [34, §2.3]).

Definition 2.9. Let $\pi = (\pi_1, \dots, \pi_s)$ be a partition of \mathcal{A} . Then π is called *independent*, provided for any choice $H_i \in \pi_i$ for $1 \leq i \leq s$, the resulting s hyperplanes are linearly independent, that is we have $r(H_1 \cap \dots \cap H_s) = s$.

Definition 2.10. Let $\pi = (\pi_1, \dots, \pi_s)$ be a partition of \mathcal{A} and let $X \in L(\mathcal{A})$. The induced partition π_X of \mathcal{A}_X is given by the non-empty blocks of the form $\pi_i \cap \mathcal{A}_X$.

Definition 2.11. The partition π of \mathcal{A} is *nice* for \mathcal{A} or a *factorization* of \mathcal{A} provided

- (i) π is independent, and
- (ii) for each $X \in L(\mathcal{A}) \setminus \{V\}$, the induced partition π_X admits a block which is a singleton.

If \mathcal{A} admits a factorization, then we also say that \mathcal{A} is *factored* or *nice*.

Remark 2.12. The class of nice arrangements is closed under taking localizations; cf. the proof of [46, Cor. 2.11].

In [46, Thm. 2.8], Terao proved that a partition π of \mathcal{A} gives rise to a tensor factorization of the Orlik-Solomon algebra of \mathcal{A} if and only if π is nice for \mathcal{A} , see [34, Thm. 3.87]. We record a consequence of this fact for our purposes.

Corollary 2.13. Let $\pi = (\pi_1, \dots, \pi_s)$ be a factorization of \mathcal{A} . Then the following hold:

- (i) $s = r = r(\mathcal{A})$ and

$$\pi(\mathcal{A}, t) = \prod_{i=1}^r (1 + |\pi_i|t);$$

- (ii) the multiset $\{|\pi_1|, \dots, |\pi_r|\}$ only depends on \mathcal{A} ;
- (iii) for any $X \in L(\mathcal{A})$, we have

$$r(X) = |\{i \mid \pi_i \cap \mathcal{A}_X \neq \emptyset\}|.$$

Remark 2.14. It follows from Corollary 2.13 that the question whether \mathcal{A} is factored is a purely combinatorial property and only depends on the lattice $L(\mathcal{A})$.

Moreover, the following is immediate from Corollary 2.13 and Theorem 2.5.

Lemma 2.15. Let (\mathcal{A}, π) be a factored arrangement which is also free. Then $\exp \mathcal{A} = \{|\pi_1|, \dots, |\pi_\ell|\}$.

2.5. Inductively factored arrangements

Following Jambu and Paris [27] and Hoge and Röhrle [24], we introduce further notation. Suppose that \mathcal{A} is non-empty and let $\pi = (\pi_1, \dots, \pi_s)$ be a partition of \mathcal{A} . Let $H_0 \in \pi_1$ and let $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ be the triple associated with H_0 . We have the *induced partition* π' of \mathcal{A}' consisting of the non-empty parts $\pi'_i := \pi_i \cap \mathcal{A}'$. Further, we have the *restriction map* $\varrho = \varrho_{\pi, H_0}: \mathcal{A} \setminus \pi_1 \rightarrow \mathcal{A}''$ given by $H \mapsto H \cap H_0$, depending on π and H_0 . Let $\pi''_i := \varrho(\pi_i)$ for $i = 2, \dots, s$. Clearly, imposing that $\pi'' = (\pi''_2, \dots, \pi''_s)$ is again a partition of \mathcal{A}'' entails that ϱ is onto.

Here is the analogue for nice arrangements of Terao's Addition-Deletion Theorem (cf. Theorem 2.6) for free arrangements from [24].

Theorem 2.16. *Suppose $\pi = (\pi_1, \dots, \pi_s)$ is a partition of $\mathcal{A} \neq \Phi_\ell$. Let $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ be the triple associated with $H_0 \in \pi_1$. Then any two of the following statements imply the third:*

- (i) π is nice for \mathcal{A} ;
- (ii) π' is nice for \mathcal{A}' ;
- (iii) $\varrho: \mathcal{A} \setminus \pi_1 \rightarrow \mathcal{A}''$ is bijective and π'' is nice for \mathcal{A}'' .

The Addition-Deletion Theorem 2.16 for nice arrangements motivates the following stronger notion of factorization, cf. [27].

Definition 2.17 [24, Def. 3.8]. *The class \mathcal{IFAC} of inductively factored arrangements is the smallest class of pairs (\mathcal{A}, π) of arrangements \mathcal{A} along with a partition π subject to*

- (i) $(\Phi_\ell, (\emptyset)) \in \mathcal{IFAC}$ for each $\ell \geq 0$;
- (ii) *if there exists a partition π of \mathcal{A} and a hyperplane $H_0 \in \pi_1$ such that for the triple $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ associated with H_0 the restriction map $\varrho = \varrho_{\pi, H_0}: \mathcal{A} \setminus \pi_1 \rightarrow \mathcal{A}''$ is bijective and for the induced partitions π' of \mathcal{A}' and π'' of \mathcal{A}'' both (\mathcal{A}', π') and (\mathcal{A}'', π'') belong to \mathcal{IFAC} , then (\mathcal{A}, π) also belongs to \mathcal{IFAC} .*

If (\mathcal{A}, π) is in \mathcal{IFAC} , then we say that \mathcal{A} is *inductively factored with respect to π* , or else that π is an *inductive factorization* of \mathcal{A} . Usually, we say \mathcal{A} is *inductively factored without reference to a specific inductive factorization* of \mathcal{A} .

Remark 2.18.

- (i) If \mathcal{A} is inductively factored, then \mathcal{A} is inductively free, by [24, Prop. 3.14].
- (ii) Thanks to [31, Thm. 1.1], inductive factoredness is preserved under localizations.
- (iii) If \mathcal{A} is supersolvable, then it is inductively factored, see [27] or [24, Prop. 3.11].

2.6. Simplicial arrangements

A real arrangement \mathcal{A} is *simplicial* provided each chamber of the complement of \mathcal{A} is an open simplicial cone in the ambient space. Simpliciality is a combinatorial property. For, thanks to [10, Cor. 2.4], a central essential real ℓ -arrangement \mathcal{A} is simplicial if and only if

$$\ell \cdot \chi(\mathcal{A}, -1) + 2 \sum_{H \in \mathcal{A}} \chi(\mathcal{A}^H, -1) = 0. \quad (2.1)$$

Remark 2.19. *Simpliciality is preserved under localizations, see [13, Lem. 2.17(1)].*

2.7. $K(\pi, 1)$ -arrangements

A complex ℓ -arrangement \mathcal{A} is called *aspherical*, or a $K(\pi, 1)$ -arrangement (or that \mathcal{A} is $K(\pi, 1)$ for short), provided the complement $M(\mathcal{A})$ of the union of the hyperplanes in \mathcal{A} in \mathbb{C}^ℓ is aspherical, that is, is a $K(\pi, 1)$ -space. That is, the universal covering space of $M(\mathcal{A})$ is contractible and the fundamental group $\pi_1(M(\mathcal{A}))$ of $M(\mathcal{A})$ is isomorphic to the group π . This is an important topological property, for the cohomology ring $H^*(X, \mathbb{Z})$ of a $K(\pi, 1)$ -space X coincides with the group cohomology $H^*(\pi, \mathbb{Z})$ of π . The crucial point here is that the intersections of codimension 2 determine the fundamental group $\pi_1(M(\mathcal{A}))$ of $M(\mathcal{A})$.

Remark 2.20.

- (i) By Deligne's seminal work [14], complexified simplicial arrangements are $K(\pi, 1)$. Likewise for complex supersolvable arrangements, cf. [20] and [45] (cf. [34, Prop. 5.12, Thm. 5.113]).
- (ii) Thanks to an observation by Oka, asphericity is preserved under localizations, for example see [35, Lem. 1.1].
- (iii) By work of Hattori, generic arrangements are not $K(\pi, 1)$, [34, Cor. 5.23].

3. Proof of Theorem 1.2

We begin by identifying the small rank hyperpolygonal arrangements \mathcal{H}_n with known ones. Clearly, \mathcal{H}_2 is just the reflection arrangement of the Weyl group of type B_2 .

It follows from the next lemma that \mathcal{H}_3 is linearly isomorphic to the connected subgraph arrangement \mathcal{A}_G , where $G = C_3$ is the cycle graph on three vertices. See [12] for the class of *connected subgraph arrangements*.

Lemma 3.1. *The arrangements \mathcal{H}_3 and \mathcal{A}_{C_3} are linearly isomorphic.*

Proof. One checks that (up to scalar multiples) the map on linear forms

$$\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}^3 \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mapsto \begin{bmatrix} x_1 + x_2 \\ x_1 + x_3 \\ x_2 + x_3 \end{bmatrix}$$

gives a linear isomorphism between \mathcal{H}_3 and \mathcal{A}_{C_3} . □

We also note that \mathcal{H}_3 is linearly isomorphic to the cone over the Shi arrangement of type A_2 , by means of Lemma 3.1 and [12, Prop. 3.1].

Our next lemma shows that \mathcal{H}_4 is linearly isomorphic to the reflection arrangement $\mathcal{A}(D_4)$ of the Weyl group of type D_4 .

Lemma 3.2. *The arrangements \mathcal{H}_4 and $\mathcal{A}(D_4)$ are linearly isomorphic.*

Proof. One checks that (up to scalar multiples) the map on linear forms

$$\varphi : \mathbb{R}^4 \rightarrow \mathbb{R}^4 \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \mapsto \begin{bmatrix} x_1 + x_2 + x_3 \\ x_2 \\ x_2 + x_3 + x_4 \\ -x_1 - x_2 - x_4 \end{bmatrix}$$

gives a linear isomorphism between \mathcal{H}_4 and $\mathcal{A}(D_4)$. □

Our next lemma gives that the reverse implications in the statements in Theorem 1.2 hold for $n \geq 6$.

Lemma 3.3. *Let $n \geq 6$. Then there exists a generic rank 3 localization of \mathcal{H}_n . As a consequence, for $n \geq 6$, \mathcal{H}_n is not free (and so is not supersolvable, not inductively free, and not inductively factored), it is also not $K(\pi, 1)$ (and so is also not simplicial).*

Proof. For $n = 6$, let $I_1 := \{1\}$, $I_2 := \{1, 2, 3\}$, $I_3 := \{1, 4, 5\}$, $I_4 := \{1, 2, 3, 4, 5\}$, and define $X = \bigcap_{i=1}^4 H_{I_i} \in L(\mathcal{H}_6)$. Then, as the localization $(\mathcal{H}_6)_X$ consists of precisely the four hyperplanes H_{I_1}, \dots, H_{I_4} of \mathcal{H}_6 , we infer that $(\mathcal{H}_6)_X$ is generic of rank 3, see [39, Ex. 4.5.6].

Generalizing this example for $n > 6$, for $X = \bigcap_{i=1}^4 H_{I_i} \in L(\mathcal{H}_n)$, the very same argument shows that also the localization $(\mathcal{H}_n)_X$ in \mathcal{H}_n is still generic of rank 3.

It follows from Remark 2.4(i) that $(\mathcal{H}_n)_X$ is not free and from Remark 2.20(iii) that $(\mathcal{H}_n)_X$ is not $K(\pi, 1)$. Consequently, $(\mathcal{H}_n)_X$ is not supersolvable, not inductively free, not inductively factored, and not simplicial. And as all of the latter are local properties, thanks to Remarks 2.3(ii), 2.4(ii), 2.8(ii), 2.18(ii), 2.19, and 2.20(ii), the lemma follows. \square

Thus, it remains to show the forward implications in the statements in Theorem 1.2 and the remaining reverse implications for $n \leq 5$.

Part (i). Thanks to Remark 2.3 (i), \mathcal{H}_2 is supersolvable. It follows from Lemma 3.1 and [12, Cor. 8.11] that \mathcal{H}_3 is not supersolvable. By [25, Lem. 3.2], $\mathcal{A}(D_4)$ is not factored (thus not inductively factored), thus, thanks to Lemma 3.2, neither is \mathcal{H}_4 , so the latter is not supersolvable, by Remark 2.18 (iii). Thanks to [17], \mathcal{H}_5 is not inductively free, thus by Remark 2.8 (iii) it is not supersolvable. Thus Theorem 1.2 (i) now follows from Lemma 3.3.

Part (ii). It follows from [23, Thm. 1.7] that \mathcal{A}_{C_3} is inductively factored, thus so is \mathcal{H}_3 , by Lemma 3.1. We have already observed above that \mathcal{H}_4 is not inductively factored. Since \mathcal{H}_5 is not inductively free, it is also not inductively factored, by Remark 2.18 (i). Consequently, Theorem 1.2 (ii) follows again from Lemma 3.3.

Part (iii). For $n \leq 4$, the arrangements \mathcal{H}_n are inductively free: For $n \leq 3$, this follows from Theorem 1.2 (ii) and Remark 2.18 (i). For \mathcal{H}_4 this follows from Lemma 3.2 and [28, Ex. 2.6]. Edelman and Reiner [17] have observed that \mathcal{H}_5 is free but not inductively free. Consequently, Theorem 1.2 (iii) follows from Lemma 3.3.

Part (iv). By Theorem 1.2 (iii) and [17], \mathcal{H}_n is free for $n \leq 5$. Whence Theorem 1.2 (iv) is once again a consequence of Lemma 3.3.

Part (v). With the aid of the formula (2.1), it is straightforward to check that \mathcal{H}_n is simplicial for $n \leq 4$ and that \mathcal{H}_5 is no longer simplicial. So Theorem 1.2 (v) follows from Lemma 3.3.

Finally, Theorem 1.2 (vi) is immediate from Lemma 3.3.

In closing this section, we show that \mathcal{H}_5 is still *recursively free*, a concept originally due to Ziegler [48]; see [34, Def. 4.60].

Remark 3.4. *While \mathcal{H}_5 is not inductively free, one can show that it is still recursively free. We sketch the argument. First one checks that $\mathcal{A} := \mathcal{H}_5 \cup \{\ker(x_2 - x_4)\}$ is inductively free with $\exp(\mathcal{A}) = \{1, 5, 5, 5, 6\}$. Next one checks that $\mathcal{B} := \mathcal{A}^{\ker(x_2 - x_4)}$ is itself again recursively free with $\exp(\mathcal{B}) = \{1, 5, 5, 5\}$. Owing to [34, Def. 4.60], \mathcal{H}_5 is recursively free. To see in turn that \mathcal{B} is recursively free, one first checks that $\mathcal{C} := \mathcal{B} \cup \{\ker(x_1 - x_3)\}$ is inductively free with $\exp(\mathcal{C}) = \{1, 5, 5, 6\}$. Then one checks that $\mathcal{D} := \mathcal{C}^{\ker(x_1 - x_3)}$ is free with $\exp \mathcal{D} = \{1, 5, 5\}$. Thanks to [2, Thm. 1.1], \mathcal{D} is recursively free. Thus, again by [34, Def. 4.60], \mathcal{B} is recursively free.*

It thus follows from the paragraph above about \mathcal{H}_5 and Theorem 1.2 (iv) that \mathcal{H}_n is free if and only if \mathcal{H}_n is recursively free. In general, these two notions differ, see [11].

4. Projective uniqueness: proof of Theorem 1.5

Definition 4.1. Let \mathcal{A}, \mathcal{B} be two arrangements in a finite-dimensional \mathbb{R} -vector space V .

- (i) \mathcal{A} and \mathcal{B} are linearly isomorphic if there is a $\varphi \in \text{GL}(V)$ such that $\mathcal{B} = \{\varphi(H) \mid H \in \mathcal{A}\}$; denoted by $\mathcal{A} \cong \mathcal{B}$.
- (ii) \mathcal{A} and \mathcal{B} are L -equivalent if $L(\mathcal{A})$ and $L(\mathcal{B})$ are isomorphic as posets; denoted by $\mathcal{A} \cong_L \mathcal{B}$.
- (iii) \mathcal{A} is projectively unique if for any arrangement \mathcal{C} in V we have: $\mathcal{C} \cong_L \mathcal{A}$ implies $\mathcal{C} \cong \mathcal{A}$.

We recall some results from [23]. Let \mathcal{A} be an \mathbb{R} -arrangement. We specify what we mean by a subarrangement of \mathcal{A} being generated by a subarrangement of \mathcal{A} .

Definition 4.2. Let $\emptyset \neq S \subseteq \mathcal{A}$. Set $\text{Gen}_0(\mathcal{A}, S) := S$ and inductively

$$\text{Gen}_{i+1}(\mathcal{A}, S) := \left\{ H \in \mathcal{A} \mid \exists J \subseteq L(\text{Gen}_i(\mathcal{A}, S)) : H = \sum_{X \in J} X \right\}$$

for $i \geq 0$. Then we say that

$$\langle S \rangle_{\mathcal{A}} := \bigcup_{i \geq 0} \text{Gen}_i(\mathcal{A}, S) \subseteq \mathcal{A}$$

is the subarrangement of \mathcal{A} generated by S . If $\langle S \rangle_{\mathcal{A}} = \mathcal{A}$, then we say that S generates \mathcal{A} .

Lemma 4.3 [23, Lem. 3.5]. Let \mathcal{A} and \mathcal{B} be two arrangements in V . Suppose $\emptyset \neq S \subseteq \mathcal{A}$, $\emptyset \neq T \subseteq \mathcal{B}$ such that $\langle S \rangle_{\mathcal{A}} = \mathcal{A}$ and $\langle T \rangle_{\mathcal{B}} = \mathcal{B}$, that is S generates \mathcal{A} and T generates \mathcal{B} . If \mathcal{A} and \mathcal{B} are L -equivalent via a poset isomorphism $\psi: L(\mathcal{A}) \rightarrow L(\mathcal{B})$ and $\varphi \in \text{GL}(V)$ such that $\psi(S) = \varphi(S) = T$ and $\psi(H) = \varphi(H)$ for all $H \in S$, then φ extends to a linear isomorphism between the whole arrangements, that is $\varphi(\mathcal{A}) = \mathcal{B}$.

The following is a consequence of Lemma 4.3.

Proposition 4.4 [23, Prop. 3.6]. Let \mathcal{A} be an essential and irreducible arrangement in $V \cong \mathbb{R}^\ell$. Suppose there is a subset S of \mathcal{A} such that $\langle S \rangle_{\mathcal{A}} = \mathcal{A}$ and $|S| = \ell + 1$. Then \mathcal{A} is projectively unique.

Finally, the following result gives Theorem 1.5.

Proposition 4.5. Let $n \geq 1$ and let

$$S := \{\ker x_1, \dots, \ker x_{n-1}, \ker(x_1 + \dots + x_n), \ker(x_1 + \dots + x_{n-1} - x_n)\} \subseteq \mathcal{H}_n.$$

Then $\langle S \rangle_{\mathcal{H}_n} = \mathcal{H}_n$. In particular, \mathcal{H}_n is projectively unique over \mathbb{R} .

Proof. Let $\emptyset \neq I \subseteq [n]$. Without loss, we may assume that $n \notin I$. For $i \in [n-1]$, set $H_i := \ker x_i$, and for $\alpha := x_1 + \dots + x_n$ and $\beta := x_1 + \dots + x_{n-1} - x_n$, set $H_\alpha := \ker \alpha$ and $H_\beta := \ker \beta$. Then $H_I = \ker(\sum_{i \in I} x_i - \sum_{j \in [n] \setminus I} x_j) = \ker \gamma$, where

$$\gamma = -\alpha + 2 \sum_{i \in I} x_i = \beta - 2 \sum_{j \in [n-1] \setminus I} x_j. \quad (4.6)$$

It follows from (4.6) that

$$X := H_\alpha \cap \bigcap_{i \in I} H_i \subseteq H_I \text{ and}$$

$$Y := H_\beta \cap \bigcap_{j \in [n-1] \setminus I} H_j \subseteq H_I.$$

Since $\dim X = n - |I| - 1$, $\dim Y = |I|$, and $\dim(X \cap Y) = 0$, we have $H_I = X + Y$. Since both X and Y belong to the lattice of intersections of the subarrangement S of \mathcal{H}_n , we infer $H_I = X + Y \in \langle S \rangle_{\mathcal{H}_n}$. Consequently, the result follows thanks to Proposition 4.4. \square

5. Combinatorial Formality: proof of Theorem 1.6

A property for arrangements is said to be *combinatorial* if it only depends on the intersection lattice of the underlying arrangement. Yuzvinsky [47, Ex. 2.2] demonstrated that formality is not combinatorial, answering a question raised by Falk and Randell [21] in the negative. Yuzvinsky's insight motivates the following notion from [30].

Definition 5.1. Suppose \mathcal{A} is a formal arrangement. We say \mathcal{A} is *combinatorially formal* if every arrangement with an intersection lattice isomorphic to the one of \mathcal{A} is also formal.

The following definitions, which are originally due to Falk for matroids [19], were adapted for arrangements in [30, §2.4]. Let $\mathcal{B} \subset \mathcal{A}$ be a subset of hyperplanes. We say \mathcal{B} is *closed* if $\mathcal{B} = \mathcal{A}_Y$ for $Y = \bigcap_{H \in \mathcal{B}} H$. We call \mathcal{B} *line-closed* if for every pair $H, H' \in \mathcal{B}$ of hyperplanes, we have $\mathcal{A}_{H \cap H'} \subset \mathcal{B}$. The *line-closure* $\text{lc}(\mathcal{B})$ of \mathcal{B} is defined as the intersection of all line-closed subsets of \mathcal{A} containing \mathcal{B} . The arrangement \mathcal{A} is called *line-closed* if every line-closed subset of \mathcal{A} is closed. With these notions, we have the following criterion for combinatorial formality, see [19, Cor. 3.8], [30, Prop. 3.2]:

Proposition 5.2. Let \mathcal{A} be an arrangement of rank r . Suppose $\mathcal{B} \subseteq \mathcal{A}$ consists of r hyperplanes such that $r(\mathcal{B}) = r$ and $\text{lc}(\mathcal{B}) = \mathcal{A}$. Then \mathcal{A} is combinatorially formal.

A subset $\mathcal{B} \subseteq \mathcal{A}$ as in Proposition 5.2 is called an *lc-basis* of \mathcal{A} .

Proof of Theorem 1.6. Let $\mathcal{B} = \{\ker x_1, \dots, \ker x_{n-1}, \ker(x_1 + \dots + x_n)\} \subseteq \mathcal{H}_n$. Then it is easy to see that successively all H_I for $\emptyset \neq I \subseteq [n]$ belong to the line-closure $\text{lc}(\mathcal{B})$ of \mathcal{B} , as follows. For $i \in [n]$, set $H_i := \ker x_i$, and for $\alpha := x_1 + \dots + x_n$ set $H_\alpha := \ker \alpha$. Then for $I = [n] \setminus \{i\}$, we have $H_I = \ker(\alpha - 2x_i) \supset H_\alpha \cap H_i$. Thus for each I of cardinality $n - 1$, H_I belongs to $\text{lc}(\mathcal{B})$. Now iterate the argument above for I of successively smaller cardinality. Consequently, $\text{lc}(\mathcal{B}) = \mathcal{H}_n$. Since $\text{rk}(\mathcal{H}_n) = n = |\mathcal{B}| = \text{rk}(\mathcal{B})$, it follows from Proposition 5.2 that \mathcal{H}_n is combinatorially formal. \square

There is a stronger notion of formality for an arrangement \mathcal{A} , that of *k-formality* for $1 \leq k \leq \text{rk}(\mathcal{A})$ due to Brandt and Terao [9]. In view of Theorem 1.6 and in view of the fact that all free arrangements are not just formal but are *k-formal* for all k , by [9, Thm. 4.15], one might ask for this stronger notion of *k-formality* among hyperpolygonal arrangements. Computational evidence for further non-free hyperpolygonal arrangements suggests the following.

Conjecture 5.3. For any $n \in \mathbb{N}$, \mathcal{H}_n is *k-formal* for any k .

6. Rank-generating functions of the poset of regions

Let \mathcal{A} be a hyperplane arrangement in the real vector space $V = \mathbb{R}^\ell$. A *region* of \mathcal{A} is a connected component of the complement $M(\mathcal{A}) := V \setminus \bigcup_{H \in \mathcal{A}} H$ of \mathcal{A} . Let $\mathcal{R} := \mathcal{R}(\mathcal{A})$ be the set of regions of \mathcal{A} . For $R, R' \in \mathcal{R}$, we let $S(R, R')$ denote the set of hyperplanes in \mathcal{A} separating R and R' . Then with respect to a choice of a fixed base region B in \mathcal{R} , we can partially order \mathcal{R} as follows:

$$R \leq R' \quad \text{if} \quad S(B, R) \subseteq S(B, R').$$

Endowed with this partial order, we call \mathcal{R} the *poset of regions of \mathcal{A} (with respect to B)* and denote it by $P(\mathcal{A}, B)$. This is a ranked poset of finite rank, where $\text{rk}(R) := |\mathcal{S}(B, R)|$, for R a region of \mathcal{A} , [16, Prop. 1.1]. The *rank-generating function* of $P(\mathcal{A}, B)$ is defined to be the following polynomial in $\mathbb{Z}_{\geq 0}[t]$

$$\zeta(P(\mathcal{A}, B); t) := \sum_{R \in \mathcal{R}} t^{\text{rk}(R)}.$$

This poset along with its rank-generating function was introduced by Edelman [16].

Thanks to work of Björner, Edelman, and Ziegler [8, Thm. 4.4] (see also Paris [37]), respectively Jambu and Paris [27, Prop. 3.4, Thm. 6.1], in case of a real arrangement \mathcal{A} which is supersolvable, respectively inductively factored, there always exists a suitable base region B so that $\zeta(P(\mathcal{A}, B); t)$ admits a multiplicative decomposition which is determined by the exponents of \mathcal{A} , that is

$$\zeta(P(\mathcal{A}, B); t) = \prod_{i=1}^{\ell} (1 + t + \dots + t^{e_i}), \quad (6.1)$$

where $\{e_1, \dots, e_{\ell}\} = \exp \mathcal{A}$ is the set of exponents of \mathcal{A} .

Quite remarkably, many classical real arrangements do satisfy the factorization identity (6.1), the most prominent ones being Coxeter arrangements.

Let $W = (W, S)$ be a Coxeter group with associated reflection arrangement $\mathcal{A} = \mathcal{A}(W)$ which consists of the reflecting hyperplanes of the reflections in W in the real space $V = \mathbb{R}^n$, where $|S| = n$. The *Poincaré polynomial* $W(t)$ of the Coxeter group W is the polynomial in $\mathbb{Z}[t]$ defined by

$$W(t) := \sum_{w \in W} t^{\ell(w)}, \quad (6.2)$$

where ℓ is the length function of W with respect to S . Then $W(t)$ coincides with the rank-generating function of the poset of regions $\zeta(P(\mathcal{A}, B); t)$ of the underlying reflection arrangement $\mathcal{A} = \mathcal{A}(W)$ with respect to B being the dominant Weyl chamber of W in V ; see [8] or [27].

The following factorization of $W(t)$ is due to Solomon [40]:

$$W(t) = \prod_{i=1}^n (1 + t + \dots + t^{e_i}), \quad (6.3)$$

where $\{e_1, \dots, e_n\}$ is the set of exponents of W , that is, the set of exponents of $\mathcal{A}(W)$; see also [29]. So by the comments above, (6.3) coincides with the factorization in (6.1).

Let W be a Coxeter group again with reflection arrangement $\mathcal{A} = \mathcal{A}(W)$, let X be a member of the intersection lattice $L(\mathcal{A})$, and consider the restricted reflection arrangement \mathcal{A}^X . In general, \mathcal{A}^X is no longer a reflection arrangement. It was shown in [32, Thm. 1.3] that there always exists a suitable base region B of \mathcal{A}^X in X so that also $\zeta(P(\mathcal{A}^X, B); t)$ satisfies (6.1) provided W is not of type E_8 . In case W is of type E_8 , then $\zeta(P(\mathcal{A}^X, B); t)$ satisfies (6.1) provided X has rank at most 3 with only two exceptions or $\mathcal{A}^X \cong (E_8, D_4)$.

Moreover, also the rank-generating function of the poset of regions $\zeta(P(\mathcal{A}_{\mathcal{I}}, B); t)$ for a so-called *ideal arrangement* $\mathcal{A}_{\mathcal{I}}$ also obeys the factorization identity (6.1). Ideal arrangements $\mathcal{A}_{\mathcal{I}}$ stem from ideals \mathcal{I} in the poset of positive roots associated with a Weyl group; see [41], [1], [38], and [3].

We close with a comment on the rank-generating function of the poset of regions of the free hyperpolygonal arrangements \mathcal{H}_n .

Remark 6.1. It follows from Theorem 1.2(ii) and [27, Prop. 3.4, Thm. 6.1] that \mathcal{H}_n satisfies (6.1) for $n \leq 3$, and for \mathcal{H}_4 this follows from Lemma 3.2 and (6.3). One can check that in contrast (6.1) fails for \mathcal{H}_5 .

Acknowledgments. We are grateful to T. Hoge and S. Wiesner for helpful discussions on material in this paper. The research of this work was supported in part by the DFG (Grant #RO 1072/25-1 (project number: 539865068) to G. Röhrle).

References

- [1] T. Abe, M. Barakat, M. Cuntz, T. Hoge and H. Terao, The freeness of ideal subarrangements of Weyl arrangements, *JEMS* **18**(6) (2016), 1339–1348.
- [2] T. Abe, M. Cuntz, H. Kawanoue and T. Nozawa, Non-recursive freeness and non-rigidity, *Discrete Math.* **339**(5) (2016), 1430–1449.
- [3] T. Abe, T. Horiguchi, M. Masuda, S. Murai and T. Sato, Hessenberg varieties and hyperplane arrangements, *J. Reine Angew. Math.* **764** (2020), 241–286.
- [4] G. Bellamy, Counting resolutions of symplectic quotient singularities, *Compos. Math.* **152**(1) (2016), 99–114.
- [5] G. Bellamy, A. Craw, S. Rayan, T. Schedler and H. Weiss, All 81 crepant resolutions of a finite quotient singularity are hyperpolygon spaces, *J. Algebraic Geom.* **33**(4) (2024), 757–793.
- [6] G. Bellamy and T. Schedler, A new linear quotient of \mathbb{C}^4 admitting a symplectic resolution, *Math. Z.* **273**(3–4) (2013), 753–769.
- [7] G. Bellamy, T. Schedler and U. Thiel, Hyperplane arrangements associated to symplectic quotient singularities, *Phenomenological approach to algebraic geometry*, vol. 116 Banach Center Publ., Polish Acad. Sci. Inst. Math. (2018), 25–45.
- [8] A. Björner, P. Edelman and G. Ziegler, Hyperplane arrangements with a lattice of regions, *Discrete Comput. Geom.* **5**(3) (1990), 263–288.
- [9] K. A. Brandt and H. Terao, Free arrangements and relation spaces, *Discrete Comput. Geom.* **12**(1) (1994), 49–63.
- [10] M. Cuntz and D. Geis, Combinatorial simpliciality of arrangements of hyperplanes, *Beitr. Algebra Geom.* **56**(2) (2015), 439–458.
- [11] M. Cuntz and T. Hoge, Free but not recursively free arrangements, *Proc. Amer. Math. Soc.* **143**(1) (2015), 35–40.
- [12] M. Cuntz and L. Kühne, On arrangements of hyperplanes from connected subgraphs, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **5**, 22. <https://journals.sns.it/index.php/annaliscienze/article/view/6134/1418>
- [13] M. Cuntz and P. Mücksch, Supersolvable simplicial arrangements, *Adv. in Appl. Math.* **107** (2019), 32–73.
- [14] P. Deligne, Les immeubles des groupes de tresses généralisés, *Invent. Math.* **17**(4) (1972), 273–302.
- [15] M. Donten-Bury and J. A. Wiśniewski, On 81 symplectic resolutions of a 4-dimensional quotient by a group of order 32, *Kyoto J. Math.* **57**(2) (2017), 395–434.
- [16] P. H. Edelman, A partial order on the regions of \mathbb{R}^n dissected by hyperplanes, *Trans. Amer. Math. Soc.* **283**(2) (1984), 617–631.
- [17] P. H. Edelman and V. Reiner, A counterexample to Orlik’s conjecture, *Proc. Amer. Math. Soc.* **118**(3) (1993), 927–929.
- [18] P. H. Edelman and V. Reiner, Not all free arrangements are $K(\pi, 1)$, *Bull. Amer. Math. Soc. (N.S.)* **32**(1) (1995), 61–65.
- [19] M. Falk, Line-closed matroids, quadratic algebras, and formal arrangements, *Adv. in Appl. Math.* **28** (2002), 250–271.
- [20] M. Falk and R. Randell, The lower central series of a fiber-type arrangement, *Invent. Math.* **82**(1) (1985), 77–88.
- [21] M. Falk and R. Randell, On the homotopy theory of arrangements, Complex analytic singularities, 101–124, Adv. Stud. Pure Math., 8, North-Holland, Amsterdam, 1987.
- [22] M. Falk and R. Randell, On the homotopy theory of arrangements. II. Arrangements—Tokyo 1998, 93–125, Adv. Stud. Pure Math., 27, Kinokuniya, Tokyo, 2000.
- [23] L. Giordani, T. Möller, P. Mücksch, G. Röhrle, On connected subgraph arrangements, preprint 2025. <http://arxiv.org/abs/2502.18144>.
- [24] T. Hoge and G. Röhrle, Addition–Deletion theorems for factorizations of Orlik–Solomon algebras and nice arrangements, *Eur. J. Comb.* **55** (2016), 20–40.
- [25] T. Hoge and G. Röhrle, Nice Reflection Arrangements, *Electron. J. Combin.* **23**(2) (2016), #P2.9.
- [26] T. Hoge, G. Röhrle and A. Schauenburg, Inductive and recursive freeness of localizations of multiarrangements, in *Algorithmic and experimental methods in Algebra, Geometry, and Number Theory*. (Springer Verlag, 2017).
- [27] M. Jambu and L. Paris, Combinatorics of inductively factored arrangements, *Eur. J. Comb.* **16** (1995), 267–292.
- [28] M. Jambu and H. Terao, Free arrangements of hyperplanes and supersolvable lattices, *Adv. in Math.* **52**(3) (1984), 248–258.
- [29] I. G. Macdonald, The Poincaré series of a Coxeter group, *Math. Ann.* **199** (1972), 161–174.
- [30] T. Möller, P. Mücksch and G. Röhrle, On Formality and Combinatorial Formality for Hyperplane Arrangements, *Discrete Comput. Geom.* **72**(1) (2024), 73–90.
- [31] T. Möller and G. Röhrle, Localizations of inductively factored arrangements, in *Algorithmic and experimental methods in Algebra, Geometry, and Number Theory*, (Springer Verlag, 2017), 495–501.
- [32] T. Möller and G. Röhrle, Counting chambers in restricted Coxeter arrangements, *Archiv Math.* **112**(4) (2019), 347–359.
- [33] Y. Namikawa, Poisson deformations and birational geometry, *J. Math. Sci. Univ. Tokyo* **22**(1) (2015), 339–359.
- [34] P. Orlik and H. Terao, *Arrangements of hyperplanes*, (Springer-Verlag, Berlin, 1992).
- [35] L. Paris, The Deligne complex of a real arrangement of hyperplanes, *Nagoya Math. J.* **131** (1993), 39–65.
- [36] L. Paris, Topology of factored arrangements of lines, *Proc. Amer. Math. Soc.* **123**(1) (1995), 7–261.
- [37] L. Paris, The counting polynomial of a supersolvable arrangement, *Adv. Math.* **116**(2) (1995), 356–364.
- [38] G. Röhrle, Arrangements of ideal type, *J. Algebra* **484** (2017), 126–167.
- [39] L. L. Rose and H. Terao, A free resolution of the module of logarithmic forms of a generic arrangement, *J. Algebra* **136**(2) (1991), 376–400.

- [40] L. Solomon, The orders of the finite Chevalley groups, *J. Algebra* **3**(3) (1966), 376–393.
- [41] E. Sommers and J. Tymoczko, Exponents for B -stable ideals, *Trans. Amer. Math. Soc.* **358**(8) (2006), 3493–3509.
- [42] R. P. Stanley, Supersolvable lattices, *Algebra Univ.* **2**(1) (1972), 197–217.
- [43] H. Terao, Arrangements of hyperplanes and their freeness I, II, *J. Fac. Sci. Univ. Tokyo* **27** (1980), 293–320.
- [44] H. Terao, Generalized exponents of a free arrangement of hyperplanes and Shepherd-Todd-Brieskorn formula, *Invent. Math.* **63**(1) (1981), 159–179.
- [45] H. Terao, Modular elements of lattices and topological fibrations, *Adv. in Math.* **62**(2) (1986), 135–134.
- [46] H. Terao, Factorizations of the Orlik-Solomon algebras, *Adv. in Math.* **92**(1) (1992), 45–53.
- [47] S. Yuzvinsky, The first two obstructions to the freeness of arrangements, *Trans. Amer. Math. Soc.* **335**(1) (1993), 231–244.
- [48] G. M. Ziegler, Algebraic combinatorics of hyperplane arrangements, Ph.D. Thesis. (MIT, Boston USA, 1987).
- [49] G. M. Ziegler, Matroid representations and free arrangements, *Trans. Amer. Math. Soc.* **320**(2) (1990), 525–541.