

Varieties of topological groups

Sidney A. Morris

We define a variety \underline{V} of topological groups to be a family of topological groups closed under cartesian products, quotient groups and subgroups. The class of groups, which with some topology appear in \underline{V} , is a variety of groups. A topological group $F(X, \underline{V})$ with generating space X is a free topological group of \underline{V} on X if any continuous mapping of X into any G in \underline{V} can be extended to a continuous homomorphism of $F(X, \underline{V})$ into G . Algebraically $F(X, \underline{V})$ is the free group on the set X of the underlying algebraic variety.

If \underline{V} is the variety of all (all abelian) topological groups then $F(X, \underline{V})$ is the free (free abelian) topological group defined by Markov. We generalize many of Markov's results; in particular, a necessary and sufficient condition for $F(X, \underline{V})$ to exist is found and uniqueness is proved. Further, if \underline{V} is non-indiscrete, $F(X, \underline{V})$ is disconnected. Any topological group Y in \underline{V} is a quotient group of $F(Y, \underline{V})$.

A full variety \underline{V} is one which contains every topological group algebraically isomorphic to a group in \underline{V} . If X is any Tychonoff space then $F(X, \underline{V})$ exists, is Hausdorff and has X as a closed subset. Algebraically fully invariant subgroups of $F(X, \underline{V})$ are closed. A topological group $F_M(X)$ is moderately free if its topology is maximal (with respect to X) and it is algebraically relatively free on X . We see that $F(X, \underline{V})$ is moderately free on X . If $F_M(X)$ is in any variety \underline{W} , then it is a quotient group of $F(X, \underline{W})$. Further, $F_M(X)$ is a free topological group of the variety it generates. We prove that finitely generated subgroups of $F_M(X)$ have the discrete topology. If X

Received 27 July 1970. Thesis submitted to The Flinders University of South Australia, February 1970. Degree approved July 1970. Supervisor: Professor I. Kluvanek. The author thanks Dr I.D. Macdonald for drawing his attention to many of the problems considered.

is compact Hausdorff, then we show $F_M(X)$ is a normal space. In fact we explicitly determine the topology of $F_M(X)$ in this case.

A β -variety \underline{V} is one for which $F(X, \underline{V})$ exists and is Hausdorff for each compact Hausdorff space X . For any β -variety \underline{V} and Tychonoff space X , $F(X, \underline{V})$ exists, is Hausdorff and has X as a closed subset. We find a necessary and sufficient condition for a variety to be a β -variety. Whilst every full variety is a β -variety we show that the converse is false.

We prove that the family of all topological varieties with a given underlying algebraic variety is not a set. In fact the family of all β -varieties with a given underlying algebraic variety is not a set. We also prove the striking result that a variety generated by a family of topological groups of bounded cardinal is not full.

Finally, the varieties $\underline{V}(R)$ and $\underline{V}(T)$ generated by the additive group of reals and the circle group respectively, each with its usual topology, are examined. Each is shown to be a β -variety but not a full variety. Further, it is shown that a locally compact Hausdorff abelian group is in $\underline{V}(T)$ if and only if it is compact. Thus R is not in $\underline{V}(T)$ and consequently $\underline{V}(R)$ properly contains $\underline{V}(T)$.

References

- [1] A.A. Markov, "On free topological groups", *C.R. (Doklady) Acad. Sci. URSS, (N.S.)* 31 (1941), 299-301. *Bull. Acad. Sci. URSS Sér. Math. [Izvestiya Akad. Nauk SSSR]* 9 (1945), 3-64 (Russian. English summary). English transl., *Amer. Math. Soc. Translation no. 30* (1950), 11-88; reprint *Amer. Math. Soc. Transl. (1)* 8 (1962), 195-273.
- [2] Sidney A. Morris, "Varieties of topological groups", *Bull. Austral. Math. Soc.* 1 (1969), 145-160.
- [3] Sidney A. Morris, "Varieties of topological groups II", *Bull. Austral. Math. Soc.* 2 (1970), 1-13.

- [4] Sidney A. Morris, "Varieties of topological groups III", *Bull. Austral. Math. Soc.* 2 (1970), 165-178.