

Effect of Palmer amaranth (*Amaranthus palmeri*) time of emergence on furrow-irrigated rice yields and weed seed production

Research Article

Cite this article: King TA, Norsworthy JK, Butts TR, Fernandes SB, Drescher GL, Avent TH (2025). Effect of Palmer amaranth (*Amaranthus palmeri*) time of emergence on furrow-irrigated rice yields and weed seed production. *Weed Sci.* **73**(e21), 1–8. doi: [10.1017/wsc.2024.102](https://doi.org/10.1017/wsc.2024.102)

Received: 16 July 2024
Revised: 29 October 2024
Accepted: 14 December 2024

Associate Editor:

Muthukumar V. Bagavathiannan, Texas A&M University

Keywords:

Competition; influence; interference; seedbank

Corresponding author:

Tanner A. King; Email: tak196@msstate.edu

Tanner A. King¹ , Jason K. Norsworthy² , Thomas R. Butts³ , Samuel B. Fernandes⁴ , Gerson L. Drescher⁵  and Tristen H. Avent¹ 

¹Graduate Research Assistant, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA; ²Distinguished Professor and Elms Farming Chair of Weed Science, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA; ³Clinical Assistant Professor and Extension Specialist, Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN, USA; ⁴Assistant Professor of Agricultural Statistics and Quantitative Genetics, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA and ⁵Assistant Professor of Soil Fertility, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA

Abstract

Furrow-irrigated rice (*Oryza sativa* L.) hectares are increasing in the Midsouth. The lack of sustained flooding creates a favorable environment for weed emergence and persistence, which makes Palmer amaranth (*Amaranthus palmeri* S. Watson) difficult to control throughout the growing season. The negative yield impacts associated with season-long *A. palmeri* interference in corn (*Zea mays* L.), cotton (*Gossypium hirsutum* L.), and soybean [*Glycine max* (L.) Merr.] have been evaluated. However, there is limited knowledge of the weed's ability to influence rice grain yield. Research was initiated in 2022 and 2023 to determine the effect of *A. palmeri* time of emergence relative to rice on weed seed production and grain yield. Cotyledon-stage *A. palmeri* plants were marked every 7 d, beginning 1 wk before rice emergence through 4 wk after rice emergence. *Amaranthus palmeri* seed production decreased exponentially as emergence timing was delayed relative to rice, and seed production increased by 447 seed plant⁻¹ for every 1-g increase in weed biomass. Without rice competition and from the earliest emergence timing, *A. palmeri* produced 540,000 seeds plant⁻¹. *Amaranthus palmeri* that emerged 1 wk before the crop had the greatest spatial influence on rice, with grain yield loss of 5% and 50% at a distance of 1.4 m and 0.40 m from the weed, respectively. As *A. palmeri* emergence was delayed, the area of influence decreased. However, *A. palmeri* plants emerging 3.5 wk after rice emergence still negatively affected grain yield and produced sufficient seed to replenish the soil seedbank, potentially impacting long-term crop management decisions. These results show that the time of *A. palmeri* emergence is a crucial factor influencing rice grain yield and weed seed production, which can be used to determine the consequences of escapes in rice.

Introduction

For decades, Palmer amaranth (*Amaranthus palmeri* S. Watson) has ranked among the most troublesome and common weed species in most southern row-crop production systems (Norsworthy et al. 2014; Van Wychen 2022; Webster and Nichols 2012). The pernicious effects of *A. palmeri* are a result of its high competitiveness with the crop for available resources, rapid growth rate, and prolific seed production (Chandi et al. 2012; Horak and Loughin 2000). With a prolonged emergence period that aligns with most row-crop production systems and noteworthy growth rate when in competition for available resources, *A. palmeri* can accumulate sufficient aboveground biomass to interfere with crop development substantially (Bell et al. 2015; Jha and Norsworthy 2009; Klingaman and Oliver 1994; Mahoney et al. 2021). As a result, negative yield impacts associated with the season-long survival of *A. palmeri* have been well documented in corn (*Zea mays* L.) (Massinga et al. 2001), cotton (*Gossypium hirsutum* L.) (MacRae et al. 2013; Norsworthy et al. 2016b), and soybean [*Glycine max* (L.) Merr.] (Klingaman and Oliver 1994). However, the competitive ability of *A. palmeri* is not restricted to the growing season in which it emerges, because its high fecundity can affect management strategies in future years due to sufficient replenishment of the soil seedbank (Schwartz et al. 2016).

To understand the impact of *A. palmeri* on crop yields, it is important to recognize the interactions between the crop and weed at developmental stages, which will also ensure successful crop management (Myers et al. 2004). Crop yield loss is primarily influenced by weed density and interspecific interference between two species (Shekhawat et al. 2020;

© University of Arkansas, 2024. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Smith et al. 1997). Additionally, the timing of weed emergence relative to the crop is a critical factor affecting yield loss and is the most vital component in crop–weed interactions (Knezevic et al. 1994; Kropff and Spitters 1991; Swanton et al. 2015). Thus, successful and timely weed management strategies can be applied with awareness of weed biology and establishment time relative to the crop (Korres et al. 2017; Wyse 1992).

Traditionally, *A. palmeri* has not been a problematic weed in conventional flood-irrigated rice production systems due to the continual flood acting as a weed-suppression mechanism for most terrestrial weeds (Bagavathiannan et al. 2011). However, a non-flooded, furrow-irrigated rice production system has increased prominence by more than 18-fold since 2015 (Hardke 2022). As a result, new weed control challenges emerge due to furrow-irrigated rice production practices being comparable to those associated with cotton, soybean, and corn production (Norsworthy et al. 2008, 2011). In a furrow-irrigated system, *A. palmeri* is problematic throughout the entirety of the growing season due to the aerobic conditions providing a conducive environment for weed emergence, which makes weed management increasingly problematic (Beesinger et al. 2022; Norsworthy et al. 2008, 2011).

A 2020 survey of Arkansas rice producers and consultants showed that *A. palmeri* was the fifth and second most problematic weed in flood- and furrow-irrigated rice, respectively (Butts et al. 2022). The problematic nature of *A. palmeri* in a furrow-irrigated rice systems can cause herbicide expenditures to be elevated if left uncontrolled throughout the growing season (Bagavathiannan et al. 2011; Barber et al. 2021). Additionally, *A. palmeri* has evolved resistance to many previously effective preemergence and post-emergence rice herbicides (Norsworthy et al. 2016a). Therefore, growers must utilize a well-rounded approach that includes chemical, cultural, biological, and mechanical control methods to reduce weed infestations and focus on controlling *A. palmeri* early in the year (DeVore et al. 2013; Harker and O'Donovan 2013).

The primary goal when creating management strategies for controlling *A. palmeri* is to prevent weed seed production and the potential for herbicide resistance spread (Shekhawat et al. 2020). With *A. palmeri* having the innate ability to emerge over an extended period, effective weed management is crucial during the summer months (Jha and Norsworthy 2009). Although an economic threshold approach has been considered for some weeds (Jones and Medd 2000), researchers have recommended a “zero-tolerance” approach for *A. palmeri* because of its potential to spread over the landscape quickly (Barber et al. 2015; Norsworthy et al. 2014). Therefore, it is crucial to not rely on chemical applications alone for controlling *A. palmeri* due to herbicide resistance and lack of control under optimal climatic conditions (Bagavathiannan and Norsworthy 2012; Butts et al. 2022).

In Arkansas, soybean is commonly rotated with rice due to the convenient implementation of herbicide programs that target troublesome monocot weeds in a dicot crop (Burgos et al. 2008, 2021; Nalley et al. 2022). With *A. palmeri* being among the most problematic weed species in Arkansas soybean, flood-irrigated rice tended to reduce the soil seedbank because of the weed's inability to survive anaerobic conditions (Beesinger et al. 2022; Riar et al. 2013). With the increased popularity of furrow-irrigated rice and widespread herbicide-resistant weed species across both cropping systems, *A. palmeri* infestations are likely to occur each year, which may lead to increased soil seedbank inputs (Butts et al. 2022; Norsworthy et al. 2013). However, rice production systems have not quantified *A. palmeri* interference and seed production.

Therefore, this research aimed to (1) evaluate the impact of *A. palmeri* interference on rice growth and yield and (2) assess *A. palmeri* biology in response to competition with rice.

Materials and Methods

Amaranthus palmeri Area of Influence in Furrow-irrigated Rice

A field experiment was conducted in 2022 and 2023 at the Milo J. Shult Agriculture Research and Extension Center in Fayetteville, AR (coordinates: 36.09352 N, 94.16890 W), to determine the impact of *A. palmeri* emergence time relative to rice on *A. palmeri* seed production and rice grain yield in a furrow-irrigated system. The experiment was set up as a completely randomized design, and the experimental area was ~0.10 ha. *Amaranthus palmeri* emergence was random and could not be blocked, but management thereafter was controlled. In both site-years, the soil was a Leaf silt loam (fine, mixed, active, thermic Typic Albaquults) composed of 18% sand, 69% silt, 13% clay, and 1.6% organic matter with a pH of 6.6. Before rice planting, the fields were disked, tilled, and hipped into 91-cm-wide beds. The soil was amended for fertility before planting based on the University of Arkansas System Division of Agriculture Marianna Soil Test Lab fertility recommendations (Roberts et al. 2018). Additionally, the entire experimental area in both years was over-sprayed with clomazone (Command® 3ME, FMC, Philadelphia, PA 19104) at 336 g ai ha⁻¹ before planting to minimize the occurrence of annual grasses. All herbicide applications were made with a CO₂-pressurized backpack sprayer calibrated to deliver 140 L ha⁻¹ at 276 kPa using four AIXR 110015 (TeeJet® Technologies, Glendale Heights, IL 60139) nozzles at 4.8 km h⁻¹.

On April 17, 2022, a hybrid long-grain cultivar ‘RT 7301’ (RiceTec, Alvin, TX 77512) was planted at 36 seeds m⁻¹ of row at a 1-cm depth in a 19-cm row spacing; and on April 22, 2023, a hybrid long-grain cultivar ‘Full Page RT 7321FP’ (RiceTec) was sown using the same planting methods. Rice emerged on May 9 and May 8 in 2022 and 2023, respectively. Within each trial, a natural population of *A. palmeri* was allowed to germinate and emerge throughout the growing season. The experiment consisted of six *A. palmeri* emergence timings, beginning 1 wk before rice emergence to 4 wk after. At each emergence timing, 10 cotyledon-stage *Amaranthus palmeri* plants were marked randomly and considered as replications within the experiment. There were 129, 123, 110, 64, and 54 replications for week -1, week 0, week 1, week 2, and week 3 to 4 evaluation timings, respectively. To reduce competition from adjacent weeds, marked plants were separated by at least 5 m from one another. At each evaluation, the marked *A. palmeri* plants were covered with buckets, and the trial was over-sprayed with propanil (STAM, UPL, King of Prussia, PA 19406) at 4,486 g ai ha⁻¹ to remove unwanted weed species while still allowing new *A. palmeri* plants to emerge. Once the rice reached the V5 growth stage, the trials were irrigated using standard furrow-irrigated rice methods, and nitrogen, as urea (460 g N kg⁻¹), was applied at a total of 135 kg N ha⁻¹ in three separate applications at 2-wk intervals (Barber et al. 2021).

At rice harvest, the height of each surviving *A. palmeri* plant was recorded. Additionally, each surviving *A. palmeri* plant was cut at the soil surface, bagged, and dried at 66 C for 2 wk to constant mass, and dry biomass was recorded. Each female plant was then threshed, and the residual material was separated from the seeds




Figure 1. Rough rice yield collection as a function of distance from *Amaranthus palmeri* at each emergence timing relative to rice. The numbers inside the ladder represent each quadrant from which rice grain was collected. Quadrat 1 was not duplicated because the ladder was only turned in a different direction to obtain yield from a separate location.

using a 20-mesh sieve and a vertical air column seed cleaner (Miranda et al. 2021). After being cleaned, 200 seeds from three plants at each emergence timing were counted and weighed, and the average weight of each subsample was divided by the total weight of seed for each plant in order to quantify the total seed produced. Rice grain yield was collected using rice sickles. A square ladder was made from a polyvinyl chloride (PVC) pipe measuring 2.4-m long, and quadrats within the ladder were 0.30-m wide by 0.30-m long and were used for portioning rice grain into 8 sections radiating in two directions from the center of the *A. palmeri* (Figure 1). After the *A. palmeri* plants were removed from the field, the center of the first quadrat was placed directly on top of the origin where the weed emerged. Rice panicles were harvested by hand in each quadrat in opposite directions and later averaged to determine rice yield loss as a function of distance from the weed. The rice panicles were threshed using an Almaco small bundle thresher (Almaco, Nevada, IA 50201), and weighed to calculate yield in each quadrat for each marked *A. palmeri* plant. After the weight of rice in each quadrat was obtained, three random quadrat samples were combined and inserted into a DICKEY-john® mini GAC™ 2500 portable grain moisture analyzer (DICKEY-john, Auburn, IL 62615) to obtain an average moisture content. Rice yields from each quadrat were then adjusted to 12% moisture.

Statistical Analysis

All data were analyzed in JMP Pro v. 17.0 (SAS Institute, Cary, NC 27513). Regression analysis was used to quantify *A. palmeri* interference in furrow-irrigated rice as a function of time of emergence relative to the crop. Because two different rice hybrids were planted in separate years, a three-parameter logistic model (logistic 3P) curve was fit by year to determine the maximum yield potential for each rice cultivar, which helped account for year-to-year variation. The initial exploratory analysis included probit, Gompertz, Weibull, and logistic curves, with the logistic 3P

(Equation 1) having the lowest Akaike information criterion (AIC) of 8598.25,

$$\text{maximum yield potential} = \frac{\text{Asymptote}}{\{1 + \exp[-\text{growth rate} \times (\text{distance} - \text{inflection point})]\}} \quad [1]$$

where maximum yield potential (kg ha^{-1}) is the dependent variable; asymptote, growth rate, and inflection points are the parameters; and distance (m) from the *A. palmeri* plant is the independent variable. Parameters to fit the logistic 3P curve can be found in Table 1. The model included male and female *A. palmeri* plants after determining that gender did not influence rice yield loss based on all model parameters not being statistically different at $\alpha = 0.05$ (data not shown). The maximum yield in each year was based on the asymptote of the model with hybrids RT 7301 and RT 7321 FP reaching a maximum yield potential of 8,700 and 10,600 kg ha^{-1} , respectively.

After maximum yield potential was determined, the rice grain yield of each quadrat was made relative to this potential within each year. To determine rice yield loss (%) as a function of distance from the *A. palmeri* plants, two- and three-parameter curves were fit within the Fit Curve platform in JMP with *A. palmeri* time of emergence relative to rice as a grouping variable. The three-parameter exponential decay (exponential 3P) model (Equation 2) achieved the lowest AIC and an $R^2 = 0.63$.

$$\text{Yield loss} = \text{Asymptote} + \text{scale} \times \exp(\text{growth rate} \times \text{distance}) \quad [2]$$

where yield loss is the dependent variable; asymptote, scale, and growth rate are parameters; and the distance in meters is the independent variable (Figure 2). Parameter estimates for the exponential 3P decay model can be found in Table 2. Inverse predictions were made for the distance in meters required from *A. palmeri* plants to observe 5% and 50% yield loss for each time of emergence relative to rice (Table 3). Prediction estimates were compared using mean 95% confidence intervals. Relative grain yield and *A. palmeri* data were combined over site-years to increase the number of observations and provide a stronger prediction estimate. Additionally, relative yield and weed data were pooled for *A. palmeri* plants emerging 3 and 4 wk after the crop to make 3.5 wk due to mortality of some plants and few observations at 3 and 4 wk than at earlier emergence times. The increased number of observations also allowed for more accurate predictions.

A linear regression model was utilized to fit *A. palmeri* seed production per female plant by *A. palmeri* biomass per female plant (Norsworthy et al. 2016b) using the Fit Curve platform in JMP Pro v. 17 (Equation 3) (Figure 3).

$$Y = a + bX \quad [3]$$

where Y is the dependent variable (seed production per female *A. palmeri* plant), b is the slope of the line, X is the independent variable (*A. palmeri* biomass per female plant), and a is the intercept when X is equal to zero. There was also a nonlinear relationship between *A. palmeri* seed production per female plant and *A. palmeri* time of emergence relative to the crop; therefore, a two-curve exponential decay (exponential 2P) model (Equation 4) was used to fit the relationship, considering it produced the lowest AIC and an $R^2 = 0.55$ (Figure 4).

Table 1. List of model parameters for the logistic 3P curve of yield (kg ha⁻¹) predicted by distance (m) from *Amaranthus palmeri* for each site-year, with R² value presented to display the percentage of variability explained by the model.

Year ^b	— Model parameters ^a —				n	R ²
	Asymptote ^c	Inflection point	Growth rate			
2022	8,678	0.3062	5.688		241	0.63
95% CI	(8,377, 8980)	(0.2560, 0.3563)	(4.110, 7.267)			
2023	10,592	0.3973	5.040		239	
95% CI	(10267, 10917)	(0.3551, 0.4395)	(4.000, 6.081)			

^aAll model parameters were significant χ^2 ($P < 0.0001$). Sigmoidal model and parameters determined using JMP Pro v. 17 with the Fit Curve Platform.

^bAbbreviation: CI, confidence interval.

^cAsymptote was used as the maximum yield potential for each site-year.




Figure 2. Three-parameter exponential decay model [$y = a + b * \exp(c * \text{distance})$], where a = asymptote, b = scale, and c = growth rate, to determine yield loss data as a function of distance from *Amaranthus palmeri* in 2022 and 2023. Inverse predictions were made from the fitted lines, giving an accurate representation of the required distance from *A. palmeri* to observe 5% and 50% rice yield loss (Table 3). (A-E) The individual predicted line for each weed emergence timing and corresponding 95% confidence interval, highlighted by the solid and dotted lines, respectively. (F) The predicted lines of the entire model for the five emergence timings of *A. palmeri* relative to rice.

Table 2. List of model parameters for the exponential 3P decay model of percent yield loss by distance from *Amaranthus palmeri* at each emergence timing, with R² value presented to display the percentage of variability explained by the model.

Emergence timing	Model parameters ^a			n	R ²
	Asymptote	Scale	Growth rate		
Week -1	-0.1714	122.4752	-2.2879	129	0.63
P-value	0.9558	<0.0001	<0.0001		
Week 0 ^b	0.1710	134.3482	-3.3137	123	
P-value	0.9458	<0.0001	<0.0001		
Week 1	-4.9869	111.3632	-2.1217	110	
P-value	0.1718	<0.0001	<0.0001		
Week 2	-3.0286	102.2000	-3.1168	64	
P-value	0.4033	<0.0001	0.0006		
Week 3-4	-5.0742	104.1803	-3.7269	54	
P-value	0.1713	<0.0001	0.0045		

^aExponential 3P decay model and parameters determined using JMP Pro v. 17 with the Fit Curve Platform.

^bThe week *A. palmeri* emerged simultaneously with rice.

Table 3. The predicted distance from *Amaranthus palmeri* to observe 5% and 50% yield loss at each *A. palmeri* emergence timing relative to rice.

Yield loss	Emergence	Predicted distance ^a	CI of mean ^b
5		m	lower, upper
	Week -1	1.38*	(1.06, 1.70)
	Week 0 ^c	1.00	(0.76, 1.25)
	Week 1	1.13	(0.93, 1.34)
	Week 2	0.81	(0.54, 1.09)
	Week 3-4	0.63*	(0.37, 0.88)
50	Week -1	0.39*	(0.33, 0.45)
	Week 0	0.30	(0.25, 0.34)
	Week 1	0.33	(0.28, 0.39)
	Week 2	0.21*	(0.15, 0.27)
	Week 3-4	0.17*	(0.11, 0.23)

^aThe distance in meters from *A. palmeri* to reach the predicted percent yield loss. Asterisks (*) show the confidence limits of an emergence timing not overlapping with the week *A. palmeri* emerged with the crop at the same percent yield loss.

^bDisplays the 95% confidence limits of the distance required from *A. palmeri* to reach the predicted percent yield loss. CI, confidence interval.

^cThe week *A. palmeri* emerged simultaneously with the rice crop.

$$Y = \text{Scale} \times \exp(\text{growth rate} \times \text{emergence}) \quad [4]$$

where *Y* is the dependent variable (seed production plant⁻¹), scale and growth rate are parameters, and *A. palmeri* emergence timing relative to rice is the explanatory variable.

Results and Discussion

Impact of *Amaranthus palmeri* on Furrow-irrigated Rice Yields

Maximum rice yield potential was statistically higher in 2023 than in 2022 based on the 95% confidence intervals not overlapping for the asymptote and inflection point model parameters; however, the growth rate parameter estimate was similar for both site-years, indicating that yield loss responses were consistent for both years of the experiment (Table 1). Maximum yield potential differences are attributed to the genetics of the two rice hybrids and environmental differences between years (JKN, personal observations). When the week of emergence was considered within the model, the relationship between rice yield loss and distance from the *A. palmeri* plant accounted for 62.7% of the variation in rice

yield loss. Additionally, differences occurred between the week of *A. palmeri* emergence relative to rice and rice yield loss as a function of distance from the weed (Figure 2F).

Regardless of when *A. palmeri* emerged relative to the crop, rice yield loss was >50% within 0.15 m from the weed (Figure 2). These results are the opposite of those observed by Bensch et al. (2003), who reported that soybean yield loss was not affected by *A. palmeri* emerging within 15 cm of the crop 19 to 38 d after planting. However, these results likely differ due to variations in crop row spacing between planting methods and subsequent proximity to *A. palmeri* plants. Nevertheless, there was a reduction in rice yield loss as distance increased from the *A. palmeri* plant and when weed emergence was delayed relative to the crop (Figure 2).

Rice yield loss was most severe when *A. palmeri* emerged 1 wk before the crop, which captures a worst-case scenario for a producer in the event a burndown application fails to control all weeds before planting. At this time, inverse predictions estimate rice yield loss to be 50% and 5% at distances of 0.40 m (±0.03 m) and 1.4 m (±0.16 m) from *A. palmeri*, respectively (Table 3). Based on the model predicting 5% or greater rice yield loss, the crop was negatively affected in the 6.2 m² surrounding each *A. palmeri* plant (π × 1.40²). Weeds that emerged 3.5 wk after rice were predicted to cause 50% and 5% yield loss at distances of 0.17 m (±0.03 m) and 0.63 m (±0.13 m), an ~5-fold reduction in area where rice yield was negatively affected compared with *A. palmeri* emerging before rice. A single *A. palmeri* emerging within 3 to 4 wk of rice emergence has the potential to impact at least 1.2 m² of rice, assuming a 95% yield preservation. These results show that *A. palmeri* emerging 3 to 4 wk after rice must be removed from the crop to prevent rough rice yield loss. It should be noted that these results are from a scenario in which no additional control measures are taken once *A. palmeri* emerges; hence, future research should investigate the effect of different removal timings and the subsequent influence on rice yield.

Amaranthus palmeri generally had a greater influence on rice yield when emergence coincided with rice emergence (Figure 2A). These results display the effect of early-emerging *A. palmeri* on rice yields and the extent to which the weed can effectively compete with the rice crop for available resources. Additionally, *A. palmeri* that emerged 3.5 wk after rice still caused significant yield losses, which was supported by Massinga et al. (2001), who observed a 7% reduction in corn yield from 0.5 *A. palmeri* plants m⁻¹ of row




Figure 3. Relationship between *Amaranthus palmeri* dry biomass and seed production per female plant in field studies conducted in 2022 and 2023. The solid line represents the fit of a linear regression model, and the dotted lines represent the 95% confidence interval of the fitted line. R^2 value displays the percentage of variability explained by the fit of the line.




Figure 4. Two-parameter exponential decay model [$y = a * \exp(b * \text{emergence})$], where a = scale and b = growth rate, to estimate *Amaranthus palmeri* seed production per female plant as a function of *A. palmeri* time of emergence relative to rice. The solid line represents the fit of the two-parameter exponential decay model, and the dotted lines represent the 95% confidence interval of the fitted line. Week 0 is the week *A. palmeri* emerged with the crop.

emerging at the 7-leaf growth stage of the crop. One factor that could influence the ability of *A. palmeri* to negatively impact rice yields is its C_4 photosynthetic pathway and rapid root augmentation (Black et al. 1969; Massinga et al. 2003; Wiese 1968); however, additional research is needed to confirm the effect of specific yield-limiting characteristics of *A. palmeri* in a furrow-irrigated rice system.

Amaranthus palmeri Seed Production as Influenced by Weed Biomass

Results of the linear regression model indicate that *A. palmeri* seeds per plant had a positive relationship with the dry weight of each female plant across all emergence timings and both site-years

(Figure 3). Previous research in other crops has identified a strong relationship between *A. palmeri* dry biomass per plant and seed production, indicating that weed seed set increases as *A. palmeri* weight increases (Mahoney et al. 2021; Schwartz et al. 2016; Spaunhorst et al. 2018). Seed production information may be useful in predicting the quantity of weed seed replenishment in the soil seedbank.

A 1-g increase in *A. palmeri* biomass plant^{-1} increased seed production by 447 seeds plant^{-1} up to a maximum of 2,500 g of dry biomass and 1.2 million seeds plant^{-1} (Figure 4). Similarly, Webster and Grey (2015) documented that *A. palmeri*, competing with cotton, produced 330 seeds for every gram increase in weed biomass. In general, weeds that emerge with or before the crop often produce more biomass relative to late-emerging weeds

(Korres et al. 2019), suggesting that seed production could be moderated if weed emergence was delayed until after the crop. The results of the previously mentioned studies and this research indicate a high fecundity potential for *A. palmeri* in rice; hence, there is an emphasis on preventing escapes and increases in soil seedbank (Norsworthy et al. 2012).

Amaranthus palmeri Seed Production as a Function of Time of Emergence

Across site-years, *A. palmeri* seed production decreased exponentially as *A. palmeri* emergence relative to the crop was delayed, as described by the exponential 2P model (Figure 4). On average, *A. palmeri* emerging 1 wk before rice produced 540,000 seeds plant⁻¹. These findings support other researchers who reported *A. palmeri* produced 446,000 to 613,000 seeds plant⁻¹ without crop competition (Keeley et al. 1987; Webster and Grey 2015). *Amaranthus palmeri* seed production per plant was significantly reduced when the weed emerged simultaneously with rice, producing an average of 115,000 seeds plant⁻¹. Hence, as much as a 4-fold decrease in weed seed production occurred when there was interference between *A. palmeri* and rice (Figure 4).

Amaranthus palmeri fecundity did not differ among emergence dates 1, 2, and 3.5 wk after rice emergence (Figure 4). *A. palmeri* emerging as late as 3.5 wk after rice could still produce 500 seeds plant⁻¹, indicating that the weed can disperse sufficient seed to contribute to increases in the soil seedbank (Crow et al. 2015; Norsworthy et al. 2020). *Amaranthus palmeri* emerging 10 wk after cotton has been shown to produce up to 880 viable seeds plant⁻¹ (Norsworthy et al. 2016b). In general, *A. palmeri* seed production decreased as weed emergence was delayed relative to rice, indicating the competitive ability of the crop for limited resources. Although late-emerging *A. palmeri* can become less prolific, the offspring produced later in the season can still significantly contribute to the soil seedbank and potentially impact management strategies in subsequent years; hence, producers should not allow any *A. palmeri* to escape control, which supports the recommended “zero-tolerance” threshold and aids in diminishing the seedbank (Norris 2007; Norsworthy et al. 2014).

Overall, no peer-reviewed publications quantify furrow-irrigated rice yield loss from season-long *A. palmeri* interference. Findings from this research support that the timing of *A. palmeri* emergence relative to the crop is important for determining yield loss (Chickoye et al. 1995; Knezevic et al. 1994; Massinga et al. 2001), but that rice yields are also greatly affected outside the immediate area of the weed. As herbicide options for control of *A. palmeri* become less available and furrow-irrigated rice hectares increase, developing an understanding of the weed’s biological effects on rice yields will prove vital in incorporating successful weed management programs into a production system (Beesinger et al. 2022; Korres and Norsworthy 2017).

Based on the *A. palmeri* seed production data presented here, female *A. palmeri* plants that emerge 3 to 4 wk after rice can still successfully contribute offspring to the soil seedbank (Figure 4). As a result, producers must focus on minimizing returns to the soil seedbank, which will subsequently help mitigate the additional emergence of *A. palmeri* seedlings in future growing seasons (Norris 2007; Webster and Grey 2015). Moreover, preventing *A. palmeri* from reaching reproductive maturity will require a combination of control measures throughout the growing season, including the use of residual preemergence herbicides, considering effective weed control options decline once weeds become

established in a field (Jha and Norsworthy 2009; Webster and Grey 2015). As a result of the consequences of allowing *A. palmeri* to compete with rice throughout the growing season and the need to prevent weed seed dispersal at harvest, producers should place extreme emphasis on the “zero-tolerance” approach regardless of when the weed emerges (Norsworthy et al. 2016b). Although not assessed in this study, the monetary losses associated with *A. palmeri* interference in furrow-irrigated rice are exacerbated when harvest efficiency and quality are negatively impacted and the economic impact of *A. palmeri* escapes in rice will extend well beyond the year in which plants are allowed to compete and produce seed.

Funding statement. This research received no specific grant from any funding agency or the commercial or not-for-profit sectors.

Competing interests. The authors declare no conflicts of interest.

References

- Bagavathiannan MV, Norsworthy JK (2012) Late-season seed production in arable weed communities: management implications. *Weed Sci* 60:325–334
- Bagavathiannan MV, Norsworthy JK, Scott RC (2011) Comparison of weed management program for furrow-irrigated and flooded hybrid rice production in Arkansas. *Weed Technol* 25:556–562
- Barber LT, Bateman N, Butts T, Hamilton M, Henry C, Lorenz G, Mazzanti R, Norsworthy J, Roberts T, Wamishe Y, Watkins B (2021) Pages 33–36 in *Arkansas Furrow-irrigated Rice Handbook*. University of Arkansas System Division of Agriculture Research and Extension
- Barber LT, Smith KL, Scott RC, Norsworthy JK, Vangilder AM (2015) Zero Tolerance: A Community-based Program for Glyphosate-Resistant Palmer Amaranth Management. University of Arkansas Cooperative Extension. 6 p
- Beesinger JW, Norsworthy JK, Butts TR, Roberts TL (2022) Palmer amaranth control in furrow-irrigated rice with florypyrauxifen-benzyl. *Weed Technol* 36:490–496
- Bell HD, Norsworthy JK, Scott RC (2015) Effect of drill-seeded soybean density and residual herbicide on Palmer amaranth (*Amaranthus palmeri*) emergence. *Weed Technol* 29:697–706
- Bensch CN, Horak MJ, Peterson D (2003) Interference of redroot pigweed (*Amaranthus retroflexus*), Palmer amaranth (*A. palmeri*), and common waterhemp (*A. rudis*) in soybean. *Weed Sci* 51:37–43
- Black CC, Chen TM, Brown RH (1969) Biochemical basis for plant competition. *Weed Sci* 17:338–344
- Burgos NR, Butts TR, Werle IS, Bottoms S, Mauromoustakos A (2021) Weedy rice update in Arkansas, USA, and adjacent locales. *Weed Sci* 69:514–525
- Burgos NR, Norsworthy JK, Scott RC, Smith KL (2008) Red rice (*Oryza sativa*) status after 5 years of imidazolinone-resistant rice technology in Arkansas. *Weed Technol* 22:200–208
- Butts TR, Kouame KB-J, Norsworthy JK, Barber LT (2022) Arkansas rice: herbicide resistance concerns, production practices, and weed management costs. *Front Agron* 4:881667
- Chandi A, Jordan DL, York AC, Milla-Lewis SR, Burton JD, Culpepper AS, Whitaker JR (2012) Interference of selected Palmer amaranth (*Amaranthus palmeri*) biotypes in soybean (*Glycine max*). *Int J Agron* 2012:1–7
- Chickoye DS, Weise SF, Swanton CJ (1995) Influence of common ragweed (*Ambrosia artemisiifolia*) time of emergence and density on white bean (*Phaseolus vulgaris*). *Weed Sci* 43:375–380
- Crow WD, Steckel LE, Hayes RM, Mueller TC (2015) Evaluation of post-harvest herbicide applications for seed prevention of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*). *Weed Technol* 29:405–411
- DeVore JD, Norsworthy JK, Bryce KR (2013) Influence of deep tillage, a rye cover crop, and various soybean production systems on Palmer amaranth emergence in soybean. *Weed Technol* 27:263–270
- Hardke JT (2022) Trends in Arkansas rice production. Pages 11–18 in *Arkansas Rice Research Studies 2022*. Little Rock: University of Arkansas System Division of Agriculture, Cooperative Extension Service. <https://scholarwo>

- rks.uark.edu/cgi/viewcontent.cgi?article=1217&context=aaesser. Accessed: January 7, 2024
- Harker KN, O'Donovan JT (2013) Recent weed control, weed management, and integrated weed management. *Weed Technol* 27:1–11
- Horak MJ, Loughin TM (2000) Growth analysis of four *Amaranthus* species. *Weed Sci* 48:347–355
- Jha P, Norsworthy JK (2009) Soybean canopy and tillage effects on emergence of Palmer amaranth (*Amaranthus palmeri*) from a natural seed bank. *Weed Sci* 57:644–651
- Jones RE, Medd RW (2000) Economic thresholds and the case for longer term approaches to population management of weeds. *Weed Technol* 14:337–350
- Keeley PE, Carter CH, Thullen RJ (1987) Influence of planting date on growth of Palmer amaranth (*Amaranthus palmeri*). *Weed Sci* 35:199–204
- Klingaman TE, Oliver LR (1994) Palmer amaranth (*Amaranthus palmeri*) interference in soybeans (*Glycine max*). *Weed Sci* 42:523–527
- Knezevic SZ, Weise SF, Swanton CJ (1994) Interference of redroot pigweed (*Amaranthus retroflexus* L.) in corn (*Zea mays* L.). *Weed Sci* 42:568–573
- Korres NE, Norsworthy JK (2017) Palmer amaranth (*Amaranthus palmeri*) demographic and biological characteristics in wide-row soybean. *Weed Sci* 65:491–503
- Korres NE, Norsworthy JK, Brye KR, Skinner V, Mauromoustakos A (2017) Relationships between soil properties and the occurrence of the most agronomically important weed species in the field margins of eastern Arkansas—implications for weed management infield margins. *Weed Res* 57:159–171
- Korres NE, Norsworthy JK, Mauromoustakos A (2019) Effects of Palmer amaranth (*Amaranthus palmeri*) establishment time and distance from the crop row on biological and phenological characteristics of the weed: implications on soybean yield. *Weed Sci* 67:126–135
- Kropff MJ, Spitters CJT (1991) A simple model of crop loss by weed competition from early observations on relative leaf area of the weed. *Weed Res* 31:97–105
- MacRae AW, Webster TM, Sosnoskie LM, Culpepper AS, Kichler JM (2013) Cotton yield loss potential in response to length of Palmer amaranth (*Amaranthus palmeri*) interference. *J Cotton Sci* 17:227–232
- Mahoney DJ, Jordan DL, Hare AT, Leon RG, Roma-Burgos N, Vann MC, Jennings KM, Everman WJ, Cahoon CW (2021) Palmer amaranth (*Amaranthus palmeri*) growth and seed production when in competition with peanut and other crops in North Carolina. *Agronomy* 11:article 1734
- Massinga RA, Currie RS, Horak MJ, Boyer J (2001) Interference of Palmer amaranth in corn. *Weed Sci* 49:202–208
- Massinga RA, Currie RS, Trooien TP (2003) Water use and light interception under Palmer amaranth (*Amaranthus palmeri*) and corn competition. *Weed Sci* 51:523–531
- Miranda JWA, Jhala AJ, Bradshaw J, Lawrence NC (2021) Palmer amaranth (*Amaranthus palmeri*) interference and seed production in dry edible bean. *Weed Technol* 35:995–100
- Myers MM, Curran WS, VanGessel MJ, Calvin DD, Mortensen DA, Majek BA, Karsten HD, Roth GW (2004) Predicting weed emergence for eight annual species in northeastern United States. *Weed Sci* 52:913–919
- Nalley LL, Massey J, Durand-Morat A, Shew A, Parajuli R, Tsioboe F (2022) Comparative economic and environmental assessments of furrow- and flood-irrigated rice production systems. *Agric Water Manag* 274:107964
- Norris RF (2007) Weed fecundity: current status and future needs. *Crop Prot* 26:182–188
- Norsworthy JK, Bond J, Scott RC (2013) Weed management practices and needs in Arkansas and Mississippi rice. *Weed Technol* 27:623–630
- Norsworthy JK, Green JK, Barber T, Roberts TL, Walsh MJ (2020) Seed destruction of weeds in southern US crops using heat and narrow-windrow burning. *Weed Technol* 34:589–596
- Norsworthy JK, Griffith GM, Griffin T, Bagavathiannan M, Gbur EE (2014) In-field movement of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) and its impact on cotton lint yield: evidence supporting a zero-threshold strategy. *Weed Sci* 62:237–249
- Norsworthy JK, Griffith GM, Scott RC (2008) Imazethapyr use with and without clomazone for weed control in furrow-irrigated, imidazolinone-tolerant rice. *Weed Technol* 22:217–221
- Norsworthy JK, Korres NE, Walsh MJ, Powles SB (2016a) Integrating herbicide program with harvest weed seed control and other fall management practices for the control of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*). *Weed Sci* 64:540–550
- Norsworthy JK, Schrage BW, Barber TL, Schwartz LM (2016b) Emergence date influences growth and fecundity of Palmer amaranth in cotton. *J Cotton Sci* 20:263–270
- Norsworthy JK, Scott RC, Bangarwa SK, Griffith GM, Wilson MJ, McClelland M (2011) Weed management in a furrow-irrigated imidazolinone-resistant hybrid rice production system. *Weed Technol* 25:25–29
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60(SP1):31–62
- Riar DS, Norsworthy JK, Steckel LE, Stephenson DO, Eubank TW, Scott RC (2013) Assessment of weed management practices and problem weeds in the mid-south United States—soybean: a consultant perspective. *Weed Technol* 27:612–622
- Roberts T, Slaton N, Wilson C, Norman R (2018) Soil fertility. *In* Arkansas Rice Production Handbook. Little Rock: University of Arkansas Division of Agriculture, Cooperative Extension Service. Pp 69–102
- Schwartz LM, Norsworthy JK, Young BG, Bradley KW, Kruger GR, Davis VM, Steckel LE, Walsh MJ (2016) Tall waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*) seed production and retention at soybean maturity. *Weed Technol* 30:284–290
- Shekhawat K, Rathore SS, Chauhan BS (2020) Weed management in dry direct-seeded rice: a review on challenges and opportunities for sustainable rice production. *Agron J* 10:1264
- Smith RJ, Flinchum WT, Seaman DE (1977) Weed Control in U.S. Rice Production. U.S. Department of Agriculture Handbook 497. Washington, DS: Government Printing Office. P 82
- Spaunhorst DJ, Devkota P, Johnson WG, Smeda RJ, Meyer CJ, Norsworthy JK (2018) Phenology of five Palmer amaranth (*Amaranthus palmeri*) populations grown in northern Indiana and Arkansas. *Weed Sci* 66:457–469
- Swanton CJ, Nkoa R, Blackshaw RE (2015) Experimental methods for crop-weed competition studies. *Weed Sci* 63(SP):2–11
- Van Wychen L (2022) 2022 Survey of the Most Common and Troublesome Weeds in Broadleaf Crops, Fruits, & Vegetables in the United States and Canada. Weed Science Society of America National Weed Survey Dataset. <http://wssa.net/wp-content/uploads/2022-Weed-Survey-Broadleaf-crops.xlsx>. Accessed: September 11, 2023
- Webster TM, Grey TL (2015) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) morphology, growth, and seed production in Georgia. *Weed Sci* 63:264–272
- Webster TM, Nichols RL (2012) Changes in the prevalence of weed species in the major agronomic crops of the southern United States: 1994/1995 to 2008/2009. *Weed Sci* 60:145–157
- Wiese AF (1968) Rate of weed root elongation. *Weed Sci* 16:11–13
- Wyse DL (1992) Future of weed science research. *Weed Technol* 6:162–165