

CONVOLUTION OPERATORS AND HOMOMORPHISMS OF LOCALLY COMPACT GROUPS

CÉDRIC DELMONICO

(Received 1 September 2005; accepted 25 April 2007)

Communicated by G. Willis

Abstract

Let $1 < p < \infty$, let G and H be locally compact groups and let ω be a continuous homomorphism of G into H . We prove, if G is amenable, the existence of a linear contraction of the Banach algebra $CV_p(G)$ of the p -convolution operators on G into $CV_p(H)$ which extends the usual definition of the image of a bounded measure by ω . We also discuss the uniqueness of this linear contraction onto important subalgebras of $CV_p(G)$. Even if G and H are abelian, we obtain new results. Let G_d denote the group G provided with a discrete topology. As a corollary, we obtain, for every discrete measure, $\|\mu\|_{CV_p(G)} \leq \|\mu\|_{CV_p(G_d)}$, for G_d amenable. For arbitrary G , we also obtain $\|\mu\|_{CV_p(G_d)} \leq \|\mu\|_{CV_p(G)}$. These inequalities were already known for $p = 2$. The proofs presented in this paper avoid the use of the Hilbertian techniques which are not applicable to $p \neq 2$. Finally, for G_d amenable, we construct a natural map of $CV_p(G)$ into $CV_p(G_d)$.

2000 *Mathematics subject classification*: primary 43A22; secondary 43A15, 43A25, 43A45, 47B38.

Keywords and phrases: convolution operators, pseudomeasures, p -multipliers, transform theorem, Herz algebra.

1. Introduction

In 1965, de Leeuw [5] studied the transfer of p -multipliers from the circle \mathbb{T} to \mathbb{R} and from \mathbb{R}_d to \mathbb{R} . These results were extended in part to locally compact abelian groups by Saeki [22], Lohoué [16–18] and Lust-Piquard [19]. The present paper investigates this problem for nonabelian locally compact groups.

Let $1 < p < \infty$, $\omega : G \rightarrow H$ be a continuous homomorphism of locally compact groups and $CV_p(G)$ be the set of all continuous operators on $L^p(G)$ commuting with left translation; they are called the p -convolution operators on G . Provided with the operator norm, denoted $\|\cdot\|_p$, $CV_p(G)$ is a Banach algebra. If G is abelian, $CV_p(G)$ is isomorphic to the Banach algebra of all p -multipliers of \widehat{G} .

The first part of this paper is devoted to the transfer of convolution operators. We show in Theorem 3.1, if G is amenable, that there is a linear contraction of $CV_p(G)$

into $CV_p(H)$ which generalizes the transfer of bounded measures. This map is unique for convolution operators with compact support (see Theorem 4.6). We give a global and a new point of view of the problem; our approach completely avoids the use of the structure theory of locally compact abelian groups and methods of Hilbert spaces. Moreover, we obtain new results even in the abelian case; we give a generalization of Reiter’s theorem of relativization of the Beurling spectrum [21] in Scholium 5.5.

Theorem 3.1 gives us, if G_d is amenable, a map of $CV_p(G_d)$ into $CV_p(G)$. In Theorem 6.6, in analogy to the Bohr compactification for G abelian, we are able to construct a natural new map of $CV_p(G)$ into $CV_p(G_d)$, even if G is nonabelian. On the way, we compare the operator norm of the discrete measures on G and on G_d . Theorem 6.1 shows that $\|\mu\|_{CV_p(G_d)} \leq \|\mu\|_{CV_p(G)}$, for every discrete measure μ and that equality holds if G_d is amenable. This result is already known for $p = 2$ (see [3, 4]), but the proof cannot be adapted to $p \neq 2$.

The ultraweak closure of the bounded measures in $CV_p(G)$ is denoted $PM_p(G)$ and called the Banach algebra of the p -pseudomeasures of G . If $p = 2$, $PM_2(G)$ is the von Neumann algebra $VN(G)$ of G . We recall that $PM_2(G) = VN(G) = CV_2(G)$. In this case, the study of the convolution operators is related to the theory of von Neumann algebras and Hilbert spaces. For example, the map $a \mapsto \lambda_G^2(\delta_a)$, where δ_a is the Dirac measure, is the left regular representation of G on $L^2(G)$. These techniques are not applicable to $p \neq 2$.

2. Preliminaries

Let $1 < p < \infty$, $p' = p/(p - 1)$, and let G, H be two locally compact groups.

For any function f on G , we define ${}_a f(x) = f(ax)$, $f_a(x) = f(xa)$, $\check{f}(x) = f(x^{-1})$ and $\tilde{f}(x) = \overline{f(x^{-1})}$. For any measure μ on G , we define $\check{\mu}(f) = \mu(\check{f})$, $\bar{\mu}(f) = \mu(\tilde{f})$ and $\tilde{\mu}(f) = \mu(\tilde{\check{f}})$. We define an isometric involution of $L^p(G)$ via $\tau_p \varphi = \Delta_G^{1/p'} \check{\varphi}$, where Δ_G denotes the modular function of G .

Let $M^1(G)$ denote the Banach algebra of the bounded measures of G . The map λ_G^p , defined via $\lambda_G^p(\mu)(\varphi) = \varphi \star \Delta_G^{1/p'} \check{\mu}$, where $\mu \in M^1(G)$ and $\varphi \in C_{oo}(G)$, is an injection of $M^1(G)$ into $CV_p(G)$.

We recall that $A_p(G)$ is the set of the bounded functions on G ,

$$u = \sum_{n=1}^{\infty} \tilde{f}_n \star \check{g}_n \quad \text{where } f_n \in L^p(G), \quad g_n \in L^{p'}(G) \text{ and } \sum_{n=1}^{\infty} \|f_n\|_p \|g_n\|_{p'} < \infty,$$

$PM_p(G)$ is the dual of $A_p(G)$ and $CV_p(G) = PM_p(G)$, if G is amenable or $p = 2$.

Let $\langle \cdot, \cdot \rangle_{L^p(G), L^{p'}(G)}$ denote the duality of $L^p(G)$ and $L^{p'}(G)$. We recall that the duality of $A_p(G)$ and $PM_p(G)$ is given by

$$\langle u, T \rangle_{A_p(G), PM_p(G)} = \sum_{n=1}^{\infty} \overline{\langle T(\tau_p f_n), \tau_{p'} g_n \rangle_{L^p(G), L^{p'}(G)}},$$

where $u = \sum_{n=1}^{\infty} \tilde{f}_n \star \check{g}_n$.

DEFINITION 2.1. For each $T \in PM_p(G)$, the support of T is the set, denoted $\text{supp}(T)$, of all $x \in G$ such that, for every neighborhood V of x , there is $v \in A_p(G)$ such that $\text{supp}(v) \subset V$ and $\langle v, T \rangle_{A_p(G), PM_p(G)} \neq 0$.

3. A transfer theorem for convolution operators

Our first main result is the following theorem.

THEOREM 3.1. Let $1 < p < \infty$, let G, H be two locally compact groups with G amenable and let ω be a continuous homomorphism of G into H . Then there is a linear contraction

$$\omega : CV_p(G) \rightarrow CV_p(H)$$

which satisfies

$$\omega(\lambda_G^p(\mu)) = \lambda_H^p(\omega(\mu)) \quad \text{for each bounded measure } \mu \text{ of } G.$$

To prove this theorem, we need the following preliminaries.

Let $\omega : G \rightarrow H$ be a continuous homomorphism. For each $T \in CV_p(G)$, $f, g \in C_{oo}(G)$, $\varphi \in L^p(H)$ and $\psi \in L^{p'}(H)$, we consider the function of H

$$h \mapsto \langle T(\tau_p(f((\tau_p\varphi)_h \circ \omega))), \tau_{p'}(g((\tau_{p'}\psi)_h \circ \omega)) \rangle_{L^p(G), L^{p'}(G)}.$$

This function is integrable and continuous on H with its L^1 -norm bounded by $\|T\|_p \|f\|_p \|g\|_{p'} \|\varphi\|_p \|\psi\|_{p'}$. Then for each $T \in CV_p(G)$ and $f, g \in C_{oo}(G)$, there is a unique p -convolution operator on H , denoted $\omega_{f,g}(T)$, such that, for all $(\varphi, \psi) \in L^p(H) \times L^{p'}(H)$,

$$\begin{aligned} & \langle \omega_{f,g}(T)\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} \\ &= \int_H \langle T(\tau_p(f((\tau_p\varphi)_h \circ \omega))), \tau_{p'}(g((\tau_{p'}\psi)_h \circ \omega)) \rangle_{L^p(G), L^{p'}(G)} dh. \end{aligned}$$

PROPOSITION 3.2. Let G and H be two locally compact groups (not necessary amenable) and $\omega : G \rightarrow H$ be a continuous homomorphism. Let $f, g \in C_{oo}(G)$. Then $\omega_{f,g}$ is a linear map of $CV_p(G)$ into $CV_p(H)$ and $\|\omega_{f,g}\| \leq \|f\|_p \|g\|_{p'}$. Moreover, for each $\mu \in M^1(G)$ and $f, g \in C_{oo}(G)$,

$$\langle \omega_{f,g}(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} = \overline{\tilde{\mu}(\bar{f} \star \check{g}(\overline{\tau_p\varphi} \star (\tau_{p'}\check{\psi})) \circ \omega)}.$$

We can immediately compare this result with

$$\langle \lambda_H^p(\omega(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} = \overline{\tilde{\mu}(\overline{(\tau_p\varphi} \star (\tau_{p'}\check{\psi})) \circ \omega)},$$

and see that, if $\bar{f} \star \check{g}$ is close to 1, $\lambda_H^p(\omega(\mu))$ is close to $\omega_{f,g}(\lambda_G^p(\mu))$.

REMARK 3.3. The special cases where ω is the inclusion of a closed subgroup or the projection on a quotient were already treated in [1, 2, 6–9]. Combining these two cases, it is possible to treat open continuous homomorphisms. The study of a general continuous homomorphism requires new ideas.

PROOF OF THEOREM 3.1.

Let $f, g \in C_{oo}(G)$. For all $T \in CV_p(G)$ and $(\varphi, \psi) \in L^p(H) \times L^{p'}(H)$, we define

$$\Omega_{f,g}(T, \varphi, \psi) = \langle \omega_{f,g}(T)\varphi, \psi \rangle_{L^p(H), L^{p'}(H)}.$$

In fact, $\Omega_{f,g}$ is a continuous form on $CV_p(G) \times L^p(H) \times L^{p'}(H)$ which is bilinear on the first two factors and conjugate linear on the last. Let \mathcal{B} denote the set of these forms provided with the weak topology of duality with $CV_p(G) \times L^p(H) \times L^{p'}(H)$.

For every compact $K \subset G$ and $\varepsilon > 0$, we define

$$\begin{aligned} \mathcal{U}_{K,\varepsilon} &= \{U \subset G : U \text{ compact, } m(U) > 0, m(xU\Delta U) < \varepsilon m(U) \forall x \in K\} \\ \mathcal{A}_{K,\varepsilon} &= \{\Omega_{f,g} : f = m(U)^{-1/p} 1_U, g = m(U)^{-1/p'} 1_U, U \in \mathcal{U}_{K,\varepsilon}\}. \end{aligned}$$

By the Banach–Alaoglu theorem,

$$\begin{aligned} \mathcal{S} &= \{F \in \mathcal{B} : |F(T, \varphi, \psi)| \leq \|T\|_p \|\varphi\|_p \|\psi\|_{p'}, \\ &\text{for all } (T, \varphi, \psi) \in CV_p(G) \times L^p(H) \times L^{p'}(H)\} \end{aligned}$$

is a compact subset of \mathcal{B} . Since G is amenable, it satisfies the property (F) of Følner [20, Theorem 7.3], so the $\mathcal{U}_{K,\varepsilon}$ are all nonempty. Then, the family $\{\overline{\mathcal{A}_{K,\varepsilon}}\}$ (where $\overline{\mathcal{A}_{K,\varepsilon}}$ denotes the weak closure of $\mathcal{A}_{K,\varepsilon}$) have the property of finite intersection. However, each $\mathcal{A}_{K,\varepsilon} \subset \mathcal{S}$ and \mathcal{S} is a compact set, so

$$\bigcap_{\substack{K \subset G \text{ compact,} \\ \varepsilon > 0}} \overline{\mathcal{A}_{K,\varepsilon}} \neq \emptyset.$$

Let

$$\Omega \in \bigcap_{\substack{K \subset G \text{ compact,} \\ \varepsilon > 0}} \overline{\mathcal{A}_{K,\varepsilon}}.$$

There is a unique continuous linear operator $\omega(T)$ on $L^p(G)$ such that, for all $(\varphi, \psi) \in L^p(H) \times L^{p'}(H)$,

$$\langle \omega(T)\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} = \Omega(T, \varphi, \psi).$$

By construction, $\omega(T) \in CV_p(H)$ and ω is a contraction.

Let $\mu \in M^1(G)$. We prove that

$$\omega(\lambda_G^p(\mu)) = \lambda_H^p(\omega(\mu)).$$

Let $(\varphi, \psi) \in L^p(H) \times L^{p'}(H)$ and $\varepsilon > 0$. We consider

$$0 < \delta < \varepsilon[1 + \omega(|\mu|)(\Delta_H^{1/p} |\varphi \star \tilde{\psi}|) + 2\|(\Delta_H^{1/p} \varphi \star \tilde{\psi}) \circ \omega\|_\infty]^{-1}.$$

There is a compact subset $K_\delta \subset G$ such that $|\mu|(G \setminus K_\delta) < \delta$. By definition of Ω , there is a compact subset $U \in \mathcal{U}_{K_\delta, \delta}$ such that

$$|\Omega_{f,g}(\lambda_G^p(\mu), \varphi, \psi) - \Omega(\lambda_G^p(\mu), \varphi, \psi)| < \frac{\varepsilon}{2},$$

where $f = m(U)^{-1/p} 1_U$ and $g = m(U)^{-1/p'} 1_U$. In fact, for all $x \in K_\delta^{-1}$,

$$0 \leq 1 - \frac{m(x^{-1}U \cap U)}{m(U)} < \frac{\delta}{2}.$$

Let $(\varphi, \psi) \in L^p(H) \times L^{p'}(H)$. On the one hand,

$$\begin{aligned} & \left| \int_{K_\delta} \left(1 - \frac{m(x^{-1}U \cap U)}{m(U)}\right) \Delta_H^{1/p}(\omega(x)) \varphi \star \tilde{\psi}(\omega(x)) \, d\mu(x) \right| \\ & \leq \frac{\delta}{2} \omega(|\mu|)(\Delta_H^{1/p} |\varphi \star \tilde{\psi}|). \end{aligned}$$

On the other hand,

$$\begin{aligned} & \left| \int_{G \setminus K_\delta} \left(1 - \frac{m(x^{-1}U \cap U)}{m(U)}\right) \Delta_H^{1/p}(\omega(x)) \varphi \star \tilde{\psi}(\omega(x)) \, d\mu(x) \right| \\ & \leq \|\Delta_H^{1/p} \circ \omega \varphi \star \tilde{\psi} \circ \omega\|_\infty |\mu|(G \setminus K_\delta) \leq \|(\Delta_H^{1/p} \varphi \star \tilde{\psi}) \circ \omega\|_\infty \delta. \end{aligned}$$

Finally,

$$\begin{aligned} & |\langle \omega(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} - \langle \lambda_H^p(\omega(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)}| \\ & \leq |\langle \omega(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} - \langle \omega_{f,g}(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)}| \\ & \quad + |\langle \omega_{f,g}(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)} - \langle \lambda_H^p(\omega(\mu))\varphi, \psi \rangle_{L^p(H), L^{p'}(H)}| \\ & \leq \frac{\varepsilon}{2} + \left| \int_G \left(1 - \frac{m(x^{-1}U \cap U)}{m(U)}\right) \Delta_H^{1/p}(\omega(x)) \varphi \star \tilde{\psi}(\omega(x)) \, d\mu(x) \right| \\ & < \frac{\varepsilon}{2} + \frac{\delta}{2} \omega(|\mu|)(\Delta_H^{1/p} |\varphi \star \tilde{\psi}|) + \|(\Delta_H^{1/p} \varphi \star \tilde{\psi}) \circ \omega\|_\infty \delta < \varepsilon. \quad \square \end{aligned}$$

REMARK 3.4. Instead of Følner’s property, we could use the property (P_p) of Reiter [20, Proposition 6.12]. It is sufficient to consider the set

$$\begin{aligned} \mathcal{R}_{K, \varepsilon} = \{ \Omega_{f,g} : f, g > 0, \|f\|_p = \|g\|_{p'} = 1, \\ \|af - f\|_p < \varepsilon \text{ and } \|ag - g\|_{p'} < \varepsilon \text{ for all } a \in K \}. \end{aligned}$$

With the same arguments, we obtain that

$$\bigcap_{\substack{K \subset G \text{ compact,} \\ \varepsilon > 0}} \overline{\mathcal{R}_{K,\varepsilon}} \neq \emptyset.$$

REMARK 3.5.

- (1) The definition of the convolution operator $\omega_{f,g}(T)$ does not require the amenability of G .
- (2) Using duality techniques of Herz [11, 12], one can give a shorter proof of Theorem 3.1. We have presented the above proof as it uses more basic ideas. We use duality arguments in the next section.

4. Image of a pseudomeasure and the A_p algebras

We show now that $\omega(T)$ is uniquely defined for T in the norm closure of the set of all compactly supported convolution operators. This Banach algebra is denoted $cv_p(G)$. We recall that $cv_2(G) = C_u^b(\widehat{G})$, since G is abelian.

For $u \in A_p(G)$ and $T \in PM_p(G)$, it is useful to define $\omega_u(T)$ by

$$\omega_u(T) = \sum_{n=1}^{\infty} \omega_{f_n, g_n}(T) \quad \text{where } u = \sum_{n=1}^{\infty} \bar{f}_n \star \check{g}_n.$$

The map ω_u is well defined because

$$\begin{aligned} & \sum_{n=1}^{\infty} \overline{\langle \omega_{f_n, g_n}(T)(\tau_p \varphi), \tau_{p'} \check{\psi} \rangle_{L^p(H), L^{p'}(H)}} \\ &= \left\langle (\bar{\varphi} \star \check{\psi}) \circ \omega \sum_{n=1}^{\infty} \bar{f}_n \star \check{g}_n, T \right\rangle_{A_p(G), PM_p(G)}. \end{aligned}$$

The following proposition is similar to Proposition 3.2.

PROPOSITION 4.1. *Let $1 < p < \infty$ and $u \in A_p(G)$. Then, ω_u is a linear map of $PM_p(G)$ into $PM_p(H)$ such that $\|\omega_u(T)\|_p \leq \|T\|_p \|u\|_{A_p}$.*

REMARK 4.2. Let us assume that G is amenable. Theorem 3.1 implies that, for every $T \in PM_p(G)$, $\varepsilon > 0$, $v \in A_p(H)$, there is $u \in A_p(G)$ such that

$$|\langle v, \omega(T) \rangle_{A_p(G), PM_p(G)} - \langle v, \omega_u(T) \rangle_{A_p(G), PM_p(G)}| < \varepsilon.$$

Let MA_p denote the set of the multipliers of A_p (that is, $v \in MA_p$, if $vu \in A_p$, for all $u \in A_p$). It is well known that $MA_p(G)$ multiplies $PM_p(G)$ in the sense of

$$\langle v, uT \rangle_{A_p(G), PM_p(G)} = \langle uv, T \rangle_{A_p(G), PM_p(G)},$$

for all $u \in MA_p(G)$, $v \in A_p(G)$ and $T \in PM_p(G)$. We recall that $\omega(u) \in MA_p(H)$, for all $u \in MA_p(G)$, see [13].

PROPOSITION 4.3. *Let $T \in PM_p(G)$ and $u \in MA_p(H)$. If G is amenable, then*

$$\omega((u \circ \omega)T) = u \omega(T).$$

PROOF. Let $\varepsilon > 0$ and $w \in A_p(H)$.

There is $v \in A_p(G)$ such that

$$|\langle w, \omega((u \circ \omega)T) \rangle_{A_p(H), PM_p(H)} - \langle w, \omega_v((u \circ \omega)T) \rangle_{A_p(H), PM_p(H)}| < \frac{\varepsilon}{2}$$

and

$$|\langle uw, \omega(T) \rangle_{A_p(H), PM_p(H)} - \langle uw, \omega_v(T) \rangle_{A_p(H), PM_p(H)}| < \frac{\varepsilon}{2}.$$

However, $\langle w, \omega_v((u \circ \omega)T) \rangle_{A_p(H), PM_p(H)} = \langle uw, \omega_v(T) \rangle_{A_p(H), PM_p(H)}$. □

LEMMA 4.4. *Let $T \in PM_p(G)$ and $u \in A_p(G)$. If $h \in \text{supp}(\omega_u(T))$, then for every neighborhood V of h , there is $v \in A_p(G)$ with $\text{supp}(v) \subset \omega^{-1}(V)$ such that*

$$\langle v, T \rangle_{A_p(G), PM_p(G)} \neq 0.$$

THEOREM 4.5. *Let $T \in PM_p(G)$. If G is amenable, then*

$$\text{supp}(\omega(T)) \subset \overline{\omega(\text{supp}(T))}.$$

PROOF. Let $u \in A_p(G)$. First, we prove that $\text{supp}(\omega_u(T)) \subset \overline{\omega(\text{supp}(T))}$.

Let $h \in \text{supp}(\omega_u(T))$ and suppose $h \notin \overline{\omega(\text{supp}(T))}$. Then there exists a closed neighborhood V of h in H such that

$$V \cap \omega(\text{supp}(T)) = \emptyset.$$

Let $v \in A_p(H)$ with $\text{supp}(v) \subset V$. For each $x \in G$ with $((v \circ \omega)u)(x) \neq 0$, we have $x \in \omega^{-1}(V)$, so $\text{supp}((v \circ \omega)u) \subset \omega^{-1}(V)$. However, $\omega^{-1}(V) \cap \text{supp}(T) = \emptyset$. Then, $((v \circ \omega)u)T = 0$, and by the amenability of G ,

$$\langle (v \circ \omega)u, T \rangle_{A_p(G), PM_p(G)} = 0,$$

which contradicts Lemma 4.4.

Finally, we prove that

$$\text{supp}(\omega(T)) \subset \overline{\bigcap_{u \in A_p(G)} \text{supp}(\omega_u(T))}.$$

Let $h_0 \in \text{supp}(\omega(T))$. Suppose

$$h_0 \notin \overline{\bigcap_{u \in A_p(G)} \text{supp}(\omega_u(T))}.$$

Then there exists a closed neighborhood V_0 of h_0 in H such that, for all $u \in A_p(G)$, $V_0 \cap \text{supp}(\omega_u(T)) = \emptyset$. Let $v \in A_p(H)$ with $\text{supp}(v) \subset V_0$. For each $u \in A_p(G)$, $\text{supp}(v) \cap \text{supp}(\omega_u(T)) = \emptyset$, then $v \omega_u(T) = 0$ and by the amenability of G ,

$$\langle v, \omega_u(T) \rangle_{A_p(H), PM_p(H)} = 0.$$

It follows that

$$\langle v, \omega(T) \rangle_{A_p(H), PM_p(H)} = 0,$$

which contradicts Lemma 4.4. □

We now want to prove that the transfer mapping is uniquely defined on a larger class of convolution operators, notably on $cv_p(G)$. We recall that, if G is amenable, then $cv_p(G) = A_p(G)PM_p(G)$, as a direct consequence of the Cohen–Hewitt theorem [14, Ch. VIII, Paragraph 32].

THEOREM 4.6. *Let $T \in PM_p(G)$ and $u \in A_p(G)$. If G is amenable, then*

$$\omega(uT) = \omega_u(T).$$

In fact, there is a unique linear contraction $\omega : cv_p(G) \rightarrow cv_p(H)$ which generalizes the transfer of bounded measures.

PROOF. Let $T \in PM_p(G)$, $u \in A_p(G)$, $v \in A_p(H)$ and $\varepsilon > 0$. There is $w \in A_p(G)$ such that

$$|\langle v, \omega(uT) \rangle_{A_p(H), PM_p(H)} - \langle v, \omega_w(uT) \rangle_{A_p(H), PM_p(H)}| < \frac{\varepsilon}{2}$$

and

$$|\langle v \circ \omega u, T \rangle_{A_p(G), PM_p(G)} - \langle v \circ \omega u w, T \rangle_{A_p(G), PM_p(G)}| < \frac{\varepsilon}{2}.$$

However,

$$\begin{aligned} \langle v, \omega_w(uT) \rangle_{A_p(H), PM_p(H)} &= \langle v \circ \omega w, uT \rangle_{A_p(G), PM_p(G)} \\ &= \langle v \circ \omega u w, T \rangle_{A_p(G), PM_p(G)}. \end{aligned}$$

Then,

$$\begin{aligned} \langle v, \omega(uT) \rangle_{A_p(H), PM_p(H)} &= \langle v \circ \omega u, T \rangle_{A_p(G), PM_p(G)} \\ &= \langle v, \omega_u(T) \rangle_{A_p(H), PM_p(H)}. \end{aligned}$$

Finally, we prove that $\omega(T) \in cv_p(G)$. There is a sequence $(T_n)_{n=1}^\infty$ of convolution operators with compact support such that $\|T_n - T\|_p \rightarrow 0$, and $(K_n)_{n=1}^\infty$ is a sequence of compact subsets of G with $\text{supp}(T_n) \subset K_n$. For each $n \in \mathbb{N}$, $\text{supp}(\omega(T_n)) \subset \omega(K_n)$. However, $\omega : CV_p(G) \rightarrow CV_p(H)$ is ultraweak continuous, then $\|\cdot\|_p$ -continuous. So

$$\lim_{n \rightarrow \infty} \omega(T_n) = \omega\left(\lim_{n \rightarrow \infty} T_n\right) = \omega(T). \quad \square$$

EXAMPLE 4.7. Let H be a closed amenable subgroup of G and let $\omega = i : H \rightarrow G$ be the canonical inclusion. For all $T \in PM_p(H)$ and $v \in A_p(G)$,

$$\langle v, i(T) \rangle_{A_p(G), PM_p(G)} = \langle \text{Res}_H v, T \rangle_{A_p(H), PM_p(H)}.$$

Derighetti obtained this result without supposing the amenability of the subgroup H (see [8, Theorem 2, p. 76]). However, he used techniques which cannot be applied to arbitrary continuous homomorphisms.

EXAMPLE 4.8. Let G be an amenable locally compact group and $\omega : G \rightarrow \{e\}$ be the trivial homomorphism. Then there is a linear contraction

$$\omega : CV_p(G) \rightarrow \mathbb{C}$$

with the following properties:

- (1) $\omega(\lambda_G^p(\mu)) = \mu(G)$ for each bounded measure μ of G ;
- (2) $\omega(uT) = \langle u, T \rangle_{A_p(G), PM_p(G)}$ for each $u \in A_p(G)$.

In fact, this defines a kind of integral on $CV_p(G)$!

EXAMPLE 4.9. Let G be an arbitrary Lie group. Then, for each X in its Lie algebra, there is a continuous homomorphism of \mathbb{R} into G defined by $t \mapsto \exp(tX)$. Hence, we are able to transfer every $T \in CV_p(\mathbb{R})$ into $CV_p(G)$.

5. The abelian case

The aim of this section is to compute the Fourier transform of $\omega(T)$.

Let G and H be two locally compact *abelian* groups and $\omega : G \rightarrow H$ be a continuous homomorphism. Here $A(G)$ denotes the Fourier algebra of G (we recall that $A(G) = A_2(G)$) and \widehat{G} be the character group of G . We denote by $\varepsilon_G : G \rightarrow \widehat{\widehat{G}}$ the canonical isomorphism defined by $\varepsilon_G(x)(\chi) = \chi(x)$, for all $x \in G$ and $\chi \in \widehat{G}$. We define an isometric isomorphism $\Phi_{\widehat{G}} : L^1(\widehat{G}) \rightarrow A_2(G)$ by

$$\Phi_{\widehat{G}}(f)(x) = \int_{\widehat{G}} f(\chi) \varepsilon_G(x)(\chi) d\chi,$$

for all $x \in G$ and the Fourier transform $\widehat{\cdot} : L^1(\widehat{G}) \rightarrow A_2(\widehat{\widehat{G}})$ by

$$\widehat{f}(\xi) = \int_{\widehat{G}} f(\chi) \overline{\xi(\chi)} d\chi,$$

for all $\xi \in \widehat{\widehat{G}}$. Let $\mathcal{F} : L^2(\widehat{G}) \rightarrow L^2(\widehat{\widehat{G}})$ denote the extension of $\widehat{\cdot}$ on $L^2(\widehat{G})$.

Let $\widehat{\omega} : \widehat{H} \rightarrow \widehat{G}$ denote the dual homomorphism defined by $\widehat{\omega}(\chi') = \chi' \circ \omega$, for all $\chi' \in \widehat{H}$. For each $T \in CV_2(G)$, \widehat{T} denotes the Fourier transform of T , that is the unique function of $L^\infty(\widehat{G})$ such that, for all $\varphi, \psi \in L^2(\widehat{G})$,

$$\langle T\varphi, \psi \rangle_{L^2(G), L^2(G)} = \langle \widehat{T} \mathcal{F}(\varphi), \mathcal{F}(\psi) \rangle_{L^2(\widehat{\widehat{G}}), L^2(\widehat{\widehat{G}})}.$$

Let $1 < p < \infty$. We define a contractive monomorphism $\alpha_p : CV_p(G) \rightarrow CV_2(G)$ such that, for all $\varphi \in L^2(G) \cap L^p(G)$, $\alpha_p(T)(\varphi) = T(\varphi)$. For $T \in CV_p(G)$, the Fourier transform of T is defined by

$$\widehat{T} = \widehat{\alpha_p(T)}. \tag{5.1}$$

From these definitions we have the following lemma.

LEMMA 5.1. *Let $1 < p < \infty$, $u \in A(G)$ and $T \in CV_p(G)$. Then*

$$\omega_u(\alpha_p(T)) = \omega_u(T)$$

and

$$\widehat{\omega_u(T)} = (\widehat{T} \star \widetilde{\Phi^{-1}(u)}) \circ \widehat{\omega}.$$

THEOREM 5.2. *Let $T \in PM_p(G)$ with \widehat{T} continuous on \widehat{G} . Then,*

$$\widehat{\omega(T)} = \widehat{T} \circ \widehat{\omega}.$$

PROOF. First, we consider $S = \alpha_p(T) \in CV_2(G)$. Let $\varepsilon > 0$ and $f \in L^1(\widehat{H})$. By hypothesis, \widehat{S} is a continuous function on \widehat{G} . So, for all $\chi \in \widehat{G}$, there is a compact neighborhood C of $e \in \widehat{G}$ such that

$$|\widehat{S}(\chi\chi') - \widehat{S}(\chi)| < \frac{\varepsilon}{4(1 + \|f\|_1)},$$

for all $\chi' \in C$.

There is $\delta > 0$ and a compact $K \subset G$ such that, for all $U \in \mathcal{U}_{K^{-1},\delta}$,

$$\int_{\widehat{G} \setminus C} \Phi^{-1}(v)(\chi) d\chi < \frac{\varepsilon}{8(1 + \|\widehat{S}\|_\infty)(1 + \|f\|_1)}$$

and

$$|\langle u, \omega(S) \rangle_{A_p(H), PM_p(H)} - \langle u, \omega_v(S) \rangle_{A_p(H), PM_p(H)}| < \delta,$$

where $v = m(U)^{-1} 1_U \star \check{1}_U \in A(G)$. Then,

$$\|\widehat{S} \star \widetilde{\Phi^{-1}(v)} - \widehat{S}\|_\infty < \frac{\varepsilon}{2(1 + \|f\|_1)}.$$

On the one hand,

$$\begin{aligned} & | \langle f, \widehat{\omega(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{\omega_v(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} | \\ & = | \langle \Phi(f), \omega(S) \rangle_{A(H), PM_2(G)} - \langle \Phi(f), \omega_v(S) \rangle_{A(H), PM_2(G)} | < \frac{\varepsilon}{2}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & |\langle f, \widehat{\omega_v(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{S} \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| \\ &= |\langle f, (\widehat{S} \star \widehat{\Phi^{-1}(v)}) \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{S} \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| \\ &= |\langle f, (\widehat{S} \star \widehat{\Phi^{-1}(v)} - \widehat{S}) \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| < \|f\|_1 \frac{\varepsilon}{2(1 + \|f\|_1)} < \frac{\varepsilon}{2}. \end{aligned}$$

Finally,

$$\begin{aligned} & |\langle f, \widehat{\omega(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{S} \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| \\ &\leq |\langle f, \widehat{\omega(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{\omega_v(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| \\ &\quad + |\langle f, \widehat{\omega_v(S)} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})} - \langle f, \widehat{S} \circ \widehat{\omega} \rangle_{L^1(\widehat{H}), L^\infty(\widehat{H})}| < \varepsilon \end{aligned}$$

and

$$\widehat{\omega(S)} = \widehat{S} \circ \widehat{\omega}.$$

We conclude by applying (5.1). □

REMARK 5.3. Theorem 5.2 was previously proved by Lohoué [17, Theorem I.1] and [15]. Nonabelian methods allow us to give a new proof.

REMARK 5.4. Let G be a locally compact abelian group and consider the homomorphism of Example 4.8,

$$\omega : G \rightarrow \{e\}.$$

Then, for each $T \in CV_p(G)$ with \widehat{T} continuous,

$$\omega(\widehat{T}) = \widehat{T}(1).$$

Let $\varphi \in L^\infty(G)$. We recall that the spectrum of φ is the set

$$\text{sp}(\varphi) = \{\chi \in \widehat{G} : \widehat{f}(\chi) = 0 \text{ for all } f \in L^1(G) \text{ with } f \star \varphi = 0\},$$

and that

$$\varepsilon_G(\text{supp}(T)) = (\text{sp}(\widehat{T}))^{-1}.$$

SCHOLIUM 5.5. Let $T \in CV_p(G)$ with \widehat{T} continuous. Then,

$$\text{sp}(\widehat{T} \circ \widehat{\omega}) \subset \overline{\widehat{\omega}(\text{sp}(\widehat{T}))}.$$

PROOF. By Theorem 4.5, we have $\text{supp}(\omega(T)) \subset \overline{\omega(\text{supp}(T))}$ and then

$$\varepsilon_G(\text{supp}(\omega(T))) \subset \overline{\widehat{\omega}(\varepsilon_G(\text{supp}(T)))}.$$

By Theorem 5.2,

$$\text{sp}(\widehat{T} \circ \widehat{\omega}) \subset \overline{(\widehat{\omega}((\text{sp}(\widehat{T}))^{-1}))^{-1}} \subset \overline{\widehat{\omega}((\text{sp}(\widehat{T})))}. \quad \square$$

REMARK 5.6. In [21, Theorem 7.2.2, p. 200], Reiter proves a result called ‘relativisation of the spectrum’. It is, in fact, a particular case of the Scholium 5.5 where \widehat{H} is a closed subgroup of \widehat{G} and $\widehat{\omega}$ is the inclusion.

6. Relations between $CV_p(G)$ and $CV_p(G_d)$

We know that deep relations exist between the harmonic analysis of G and G_d . In [10, 12], Eymard and Herz investigated the relationship between $B(G)$ and $B(G_d) \cap C(G)$. In this section, we study the relationship between $CV_p(G)$ and $CV_p(G_d)$. More precisely, we construct a new map of $CV_p(G)$ into $CV_p(G_d)$, for G_d amenable.

First, we give results about the operator norm of discrete measures. For each sequence $(c_n)_{n=1}^\infty \in \ell^1$ and $(a_n)_{n=1}^\infty$ on G , we consider the measure $\mu = \sum_{n=1}^\infty c_n \delta_{a_n}$, where δ_a is the Dirac measure on a . Here μ is a bounded measure on both G and on G_d with $\omega(\mu) = \mu$. All of these measures are called discrete measures of G .

THEOREM 6.1. *Let $1 < p < \infty$, G be a locally compact group and let μ be a discrete measure of G . Then,*

$$\|\lambda_{G_d}^p(\mu)\|_p \leq \|\lambda_G^p(\mu)\|_p.$$

Moreover, if G_d is amenable,

$$\|\lambda_{G_d}^p(\mu)\|_p = \|\lambda_G^p(\mu)\|_p.$$

The proof of the first inequality is based on the following construction and the second is a corollary of Theorem 3.1.

DEFINITION 6.2. Let W be a relatively compact neighborhood of e in G . For each $k \in C_{oo}(G_d)$, we define

$$T_W^p(k) = m(W)^{-1/p} \sum_{x \in G} k(x)_{x^{-1}}(1_W).$$

It is straightforward to prove the following properties:

- (1) $T_W^p : C_{oo}(G_d) \rightarrow L^p(G)$;
- (2) $\|T_W^p(k)\|_p \leq \|k\|_1$;
- (3) $T_W^p({}_a k) = {}_a(T_W^p(k))$ for all $a \in G$.

The second property can be improved on, as follows.

LEMMA 6.3. *Let $k \in C_{oo}(G_d)$ with $\text{supp}(k) = \{x_1, \dots, x_n\}$ and let W be a relatively compact neighborhood of e in G such that $x_i W \cap x_j W = \emptyset$, for each $1 \leq i, j \leq n$ with $i \neq j$. Then*

$$\|T_W^p(k)\|_p = \|k\|_p.$$

PROOF. For each $y \in G$, there is $j_y \in \{1, \dots, n\}$ such that $x_{j_y}^{-1}y \in W$. Then

$$\left| \sum_{j=1}^n k(x_j) 1_W(x_j^{-1}y) \right|^p = \sum_{j=1}^n |k(x_j)|^p 1_W(x_j^{-1}y) \quad \text{and}$$

$$\|T_W^p(k)\|_p^p = m(W)^{-1} \sum_{j=1}^n |k(x_j)|^p \int_G 1_W(y) dy = \|k\|_p. \quad \square$$

LEMMA 6.4. *Let $k, l \in C_{oo}(G_d)$ and let μ be a bounded measure on G with finite support (that is, $\mu = \sum_{i=1}^n c_i \delta_{a_i}$, where $c_i \in \mathbb{C}$ and $a_i \in G$). Then there exists a neighborhood W of e in G such that*

$$\langle \bar{k} \star \check{l}, \lambda_{G_d}^p(\mu) \rangle_{A_p(G_d), PM_p(G_d)} = \langle (T_W^p(\bar{k})) \star (T_W^{p'}(l)), \lambda_G^p(\mu) \rangle_{A_p(G), PM_p(G)}.$$

PROOF. Suppose that $\mu = \delta_a$, for any $a \in G$. Let $\text{supp}(k) = \{x_1, \dots, x_n\}$ and $\text{supp}(l) = \{y_1, \dots, y_m\}$.

We define $E = \{(i, j) \in \mathbb{N}_n \times \mathbb{N}_m : ax_i = y_j\}$ and consider W a neighborhood of e such that $(x_i^{-1}a^{-1}y_j)W \cap W = \emptyset$. Then,

$$\begin{aligned} & \langle (T_W^p(\bar{k})) \star (T_W^{p'}(l)), \lambda_G^p(\mu) \rangle_{A_p(G), PM_p(G)} \\ &= m(W)^{-1} \sum_{i=1}^n \sum_{j=1}^m \bar{k}(x_i) l(y_j) \int_G 1_W(x_i^{-1}a^{-1}y_j y) 1_W(y) dy \\ &= \sum_{(i,j) \in E} \bar{k}(a^{-1}y_j) l(y_j) = \sum_{j=1}^m \bar{k}(a^{-1}y_j) l(y_j) \\ &= \langle \bar{k} \star \check{l}, \lambda_{G_d}^p(\mu) \rangle_{A_p(G_d), PM_p(G_d)}. \end{aligned}$$

The result now follows by linearity. □

PROOF OF THEOREM 6.1. We prove that $\|\lambda_{G_d}^p(\mu)\|_p \leq \|\lambda_G^p(\mu)\|_p$.

Let $r \in C_{oo}(G_d)$ with $\|r\|_1 \leq 1$. We define $f = r^{1/p}$ and $g = r^{1/p'}$. Let ν be a bounded measure with finite support. There is a neighborhood W of e in G such that

$$\begin{aligned} & \langle \overline{\tau_p f} \star (\tau_p g), \lambda_{G_d}^p(\nu) \rangle_{A_p(G_d), PM_p(G_d)} \\ &= \langle T_W^p(\overline{\tau_p f}) \star T_W^{p'}((\tau_p g)), \lambda_G^p(\nu) \rangle_{A_p(G), PM_p(G)}. \end{aligned}$$

Then

$$\begin{aligned} & |\langle \lambda_{G_d}^p(\nu) f, g \rangle_{\ell^p(G), \ell^{p'}(G)}| \\ &= |\langle \lambda_G^p(\nu) (\tau_p(T_W^p(\overline{\tau_p f}))), \tau_{p'}(T_W^{p'}(\tau_p g)) \rangle_{L^p(G), L^{p'}(G)}| \\ &\leq \|\lambda_{G_d}^p(\nu)\|_p \|T_W^p(\overline{\tau_p f})\|_p \|T_W^{p'}(\tau_p g)\|_{p'} \leq \|\lambda_G^p(\nu)\|_p \|f\|_p \|g\|_{p'}. \end{aligned}$$

Finally, from $\|f\|_p \leq 1$ and $\|g\|_{p'} \leq 1$, we obtain $\|\lambda_{G_d}^p(v)\|_p \leq \|\lambda_G^p(v)\|_p$.

There is a $(v_n)_{n=1}^\infty$ sequence of bounded measures of G with finite support such that $\lim \|v_n - \mu\| = 0$.

$$\begin{aligned} \|\lambda_{G_d}^p(\mu)\|_p &\leq \|\lambda_{G_d}^p(\mu) - \lambda_{G_d}^p(v_n)\|_p + \|\lambda_{G_d}^p(v_n)\|_p \\ &\leq \|\lambda_{G_d}^p(\mu - v_n)\|_p + \|\lambda_{G_d}^p(v_n)\|_p \\ &\leq \|\mu - v_n\| + \|\lambda_G^p(v_n) - \lambda_G^p(\mu)\|_p + \|\lambda_G^p(\mu)\|_p \\ &\leq 2\|\mu - v_n\| + \|\lambda_G^p(\mu)\|_p. \end{aligned}$$

Assume that G_d is amenable. The inequality $\|\lambda_{G_d}^p(\mu)\|_p \geq \|\lambda_G^p(\mu)\|_p$ is then a direct consequence of Theorem 3.1. □

REMARK 6.5. The map $a \mapsto \lambda_G^2(\delta_a)$ is the left regular representation of G on $L^2(G)$. Theorem 6.1 is a version when $p \neq 2$ of the result of [3, Lemma 2, p. 606] and [4, Theorem 2, p. 3152]. The Hilbert space methods used to prove the version when $p = 2$ are not applicable when $p \neq 2$. Our proof requires another approach.

THEOREM 6.6. *Let $1 < p < \infty$ and G be a locally compact group. Assume that G_d is amenable. Then there is a linear contraction*

$$\sigma : CV_p(G) \rightarrow CV_p(G_d)$$

such that, for all discrete measures μ on G ,

$$\sigma(\lambda_G^p(\mu)) = \lambda_{G_d}^p(\mu).$$

The proof of this theorem is based on Definition 6.2 and the following construction.

DEFINITION 6.7. Let $1 < p < \infty$, let $T \in CV_p(G)$, let W be a relatively compact neighborhood of e in G and let $k, l \in C_{oo}(G_d)$. We define $\sigma_{W,k,l}(T)$ by

$$\begin{aligned} &\langle \sigma_{W,k,l}(T)\varphi, \psi \rangle_{L^p(G_d), L^{p'}(G_d)} \\ &= \sum_{x \in G} \langle T(\tau_p(T_W^p(k(x\check{\varphi}))), \tau_{p'}(T_W^{p'}(l(x\check{\psi}))) \rangle_{L^p(G), L^{p'}(G)} \end{aligned}$$

for all $\varphi \in L^p(G_d)$ and $\psi \in L^{p'}(G_d)$.

Let \mathcal{W} denote the set of pairs (W, r) where W is a relatively compact neighborhood of e in G , $r \in C_{oo}(G_d)$ such that $x_i W \cap x_j W = \emptyset$, for all $i \neq j$, $1 \leq i, j \leq n$, where $\{x_1, \dots, x_n\} = \text{supp}(r)$.

LEMMA 6.8. *Let $(W, r) \in \mathcal{W}$, $k = r^{1/p}$ and $l = r^{1/p'}$. Then $\sigma_{W,k,l}$ is a linear map of $CV_p(G)$ into $CV_p(G_d)$ with $\|\sigma_{W,k,l}(T)\|_p \leq \|T\|_p \|k\|_p \|l\|_{p'}$.*

LEMMA 6.9. *Let μ be a bounded measure of G with finite support. Then there is $(W, r) \in \mathcal{W}$ such that, for all $\varphi, \psi \in C_{oo}(G_d)$,*

$$\langle \sigma_{W,k,l}(\lambda_G^p(\mu))\varphi, \psi \rangle_{L^p(G_d), L^{p'}(G_d)} = \overline{\mu^*((\bar{k} \star \check{l})(\tilde{\varphi} \star \psi))},$$

where $k = r^{1/p}$ and $l = r^{1/p'}$.

PROOF OF THEOREM 6.6. For each $(W, r) \in \mathcal{W}$, $k = r^{1/p}$ and $l = r^{1/p'}$, we define

$$\Sigma_{W,k,l}(T, \varphi, \psi) = \langle \sigma_{W,k,l}(T)\varphi, \psi \rangle_{L^p(G_d), L^{p'}(G_d)},$$

where $T \in CV_p(G)$ and $\varphi, \psi \in C_{oo}(G_d)$. Here $\Sigma_{W,k,l}$ is a continuous form on $CV_p(G) \times L^p(G_d) \times L^{p'}(G_d)$, which is bilinear in the two first factors and conjugate linear on the third. Let \mathcal{B} denote the set of these forms with the weak topology of duality with $CV_p(G) \times L^p(G_d) \times L^{p'}(G_d)$. By the Banach–Alaoglu theorem, $\mathcal{S} = \{F \in \mathcal{B} : |F(T, \varphi, \psi)| \leq \|T\|_p \|\varphi\|_p \|\psi\|_{p'}\}$ is a compact subset of \mathcal{B} . For each K finite subset of G , $\varepsilon > 0$ and U neighborhood of e in G , we define

$$\begin{aligned} \mathcal{A}_{K,\varepsilon,U} = \{ \Sigma_{W,k,l} : (W, r) \in \mathcal{W}, k = r^{1/p}, l = r^{1/p'}, r \geq 0, \|r\|_1 = 1, \\ \|_{x^{-1}k - k}\|_p < \varepsilon \quad \forall x \in K, W \subset U \}. \end{aligned}$$

The $\mathcal{A}_{K,\varepsilon,U}$ are all nonempty, because G_d is amenable. It easy to show that for all $n \in \mathbb{N}$, $K_1, \dots, K_n \subset G$ finite, $\varepsilon_1, \dots, \varepsilon_n > 0$ and U_1, \dots, U_n neighborhood of e on G , $\bigcap_{i=1}^n \mathcal{A}_{K_i, \varepsilon_i, U_i} \neq \emptyset$. However, \mathcal{S} is compact, so there is

$$\Sigma \in \bigcap_{\substack{K \in G \text{ finite} \\ \varepsilon > 0 \\ U \text{ neighbor of } e}} \overline{\mathcal{A}_{K,\varepsilon,U}}.$$

For each $T \in CV_p(G)$, $\varphi \in L^p(G_d)$ and $\psi \in L^{p'}(G_d)$, we define

$$\Sigma(T, \varphi, \psi) = \langle \sigma(T)\varphi, \psi \rangle_{L^p(G_d), L^{p'}(G_d)}. \quad \square$$

This extends Lust-Piquard’s result [19, Theorem 4.1]. The techniques used for the proof are completely different and are not applicable to nonabelian groups. This problem was also treated by Lohoué in [17, 18] for special kinds of convolution operators, with strong use of structure theory of locally compact abelian groups.

REMARK 6.10. For G amenable, the map defined in Theorem 6.6 could be considered as a substitute for the map of the p -multipliers of \widehat{G} into the p -multipliers of the Bohr compactification of \widehat{G} .

Acknowledgements

This work was part of the author’s PhD Thesis defended in July 2003 at the University of Lausanne. I am very grateful to Professor A. Derighetti for his continuous support and for his good advice which helped me to produce my PhD Thesis and the present paper.

References

- [1] J.-P. Anker, 'Aspects de la p -induction en analyse harmonique', PhD Thesis, Université de Lausanne, 1982.
- [2] ———, 'Applications de la p -induction en analyse harmonique', *Comment. Math. Helv.* **58**(4) (1983), 622–645.
- [3] E. Bédos, 'On the C^* -algebra generated by the left regular representation of a locally compact group', *Proc. Amer. Math. Soc.* **120**(2) (1994), 603–608.
- [4] M. B. Bekka, E. Kaniuth, A. T. Lau and G. Schlichting, 'On C^* -algebras associated with locally compact groups', *Proc. Amer. Math. Soc.* **124**(10) (1996), 3151–3158.
- [5] K. de Leeuw, 'On L_p multipliers', *Ann. of Math. (2)* **81** (1965), 364–379.
- [6] J. Delaporte, 'Convoluteurs continus et topologie stricte', in: *Harmonic Analysis (Luxembourg, 1987)* (Springer, Berlin, 1988), pp. 135–141.
- [7] J. Delaporte and A. Derighetti, ' p -pseudomeasures and closed subgroups', *Monatsh. Math.* **119**(1–2) (1995), 37–47.
- [8] A. Derighetti, 'Relations entre les convoluteurs d'un groupe localement compact et ceux d'un sous-groupe fermé', *Bull. Sci. Math. (2)* **106**(1) (1982), 69–84.
- [9] ———, 'À propos des convoluteurs d'un groupe quotient', *Bull. Sci. Math. (2)* **107**(1) (1983), 3–23.
- [10] P. Eymard, 'L'algèbre de Fourier d'un groupe localement compact', *Bull. Soc. Math. France* **92** (1964), 181–236.
- [11] C. Herz, 'The theory of p -spaces with an application to convolution operators', *Trans. Amer. Math. Soc.* **154** (1971), 69–82.
- [12] ———, 'Harmonic synthesis for subgroups', *Ann. Inst. Fourier (Grenoble)* **23**(3) (1973), 91–123.
- [13] ———, 'Une généralisation de la notion de transformée de Fourier–Stieltjes', *Ann. Inst. Fourier (Grenoble)* **24**(3) (1974), 145–157.
- [14] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis, Vol. II: Structure and Analysis for Compact Groups. Analysis on Locally Compact Abelian Groups* (Springer, New York, 1970).
- [15] N. Lohoué, 'Sur certains ensembles de synthèse dans les algèbres $A_p(G)$ ', *C. R. Acad. Sci. Paris Sér. A–B* **270** (1970), A589–A591.
- [16] ———, 'Sur le critère de S. Bochner dans les algèbres $B_p(R^n)$ et l'approximation bornée des convoluteurs', *C. R. Acad. Sci. Paris Sér. A–B* **271** (1970), A247–A250.
- [17] ———, 'Algèbres $A_p(G)$ et convoluteurs de $L^p(G)$ '. Doctorat d'état, Université Paris-Sud, Centre d'Orsay, 1971.
- [18] ———, 'Approximation et transfert d'opérateurs de convolution', *Ann. Inst. Fourier (Grenoble)* **26**(4) (1976), 133–150.
- [19] F. Lust-Piquard, 'Means on $CV_p(G)$ -subspaces of $CV_p(G)$ with RNP and Schur property', *Ann. Inst. Fourier (Grenoble)* **39**(4) (1989), 969–1006.
- [20] J.-P. Pier, *Amenable Locally Compact Groups* (John Wiley & Sons, New York, 1984).
- [21] H. Reiter and J. D. Stegeman, *Classical Harmonic Analysis and Locally Compact Groups*, 2nd edn (The Clarendon Press/Oxford University Press, New York, 2000).
- [22] S. Saeki, 'Translation invariant operators on groups', *Tôhoku Math. J. (2)* **22** (1970), 409–419.

CÉDRIC DELMONICO, EPFL SB IACS, Station 8,

CH-1015 LAUSANNE, Switzerland

e-mail: cedric.delmonico@a3.epfl.ch