



Falling clouds

Élisabeth Guazzelli[†]

Université Paris Cité, CNRS, Matière et Systèmes Complexes (MSC) UMR 7057, Paris, France

(Received 29 June 2024; accepted 30 August 2024)

The featured article ‘Break-up of a falling drop containing dispersed particles’ (Nitsche and Batchelor, *J. Fluid Mech.*, 1997, vol. 340, pp. 161–175) is G. K. Batchelor’s last published paper with his former postdoctoral associate J. M. Nitsche. The objective of the study was to investigate the randomness of the velocities of interacting rigid particles falling under gravity through a viscous fluid at a small Reynolds number and its consequence for the breakup of a falling cloud of particles. The study focused on a quintessential problem of the collective dynamics of interacting particles and has been an inspiration for subsequent work.

Key words: suspensions, particle/fluid flow

1. Batchelor & Nitsche’s falling clouds

Nitsche & Batchelor (1997) examined the temporal evolution of an initially spherical cloud of particles falling in a quiescent fluid in the Stokes regime. Their questions of interest were as follows: Do particles leave the cloud, and if so, how? What is the lifetime of the cloud as a cohesive entity? Their analysis was substantiated by a numerical simulation of interacting particles (with a maximum particle number of $N = 320$), in which the particles were supposed to act as Stokeslets. This approximation assumed that spherical particles could be treated as point particles that interact with their leading fluid velocity disturbance in Stokes flows, which decays as one over the distance to their centre. The cloud was observed to maintain its initial shape, while particles were observed to leak from its rear in a vertical tail that eventually led to its disintegration.

Notwithstanding the small gravitational slip of the particle phase (Batchelor 1974; Ekiel-Jeżewska, Metzger & Guazzelli 2006), the flow around the suspension cloud is, in fact, related to that of the settling of a spherical drop of heavy fluid in an otherwise lighter fluid as described by Hadamard (1911) and Rybczyński (1911). Consequently, the cloud falls like a dense, effective-fluid drop, with no surface tension, at a settling speed $V_c \sim NV_sa/R$, where a is the particle radius, R the cloud radius and $V_S = 2(\rho_p - \rho_f)a^2g/9\mu$ is the Stokes velocity of an individual particle of density ρ_p settling in the quiescent fluid of viscosity μ and density ρ_f under the gravitational acceleration g . In the reference

[†] Email address for correspondence: elisabeth.guazzelli@u-paris.fr




Figure 1. A schematic representation of a falling particle cloud illustrating the toroidal circulation of particles within the cloud and the particle leakage in the reference frame of the moving cloud (from a drawing of Sylvie Pic in Guazzelli & Morris 2012).

frame of the moving cloud, the particles experience a circular motion along closed toroidal trajectories (the well-known Hadamard–Rybczyński toroidal flow) within the entire volume of the cloud. The hydrodynamic interactions between the particles cause random displacements superposed on this toroidal motion. These random displacements ultimately result in the particle crossing over into the region of the cloud where there is no particle presence, as depicted in figure 1. In this region, the streamlines sweep around the cloud surface and are no longer closed (as they are inside), but extend to infinity. Once particles are swept around to the rear, they fall behind the faster-moving cloud and never catch up again, and thus are lost in the aforementioned tail.

Having identified the mechanism of the cloud disintegration, Nitsche & Batchelor (1997) sought a law governing the rate of particle loss. They proposed that the leakage rate is given by $-dN/dt \propto V_c/d$, considering that the rate-determining factor is the fall velocity of the cloud, V_c , and that the relevant unit of length is the mean particle spacing, $d = (4\pi/3N)^{1/3}R$, as it describes the chaotic displacements of the particles which may result in their escape from the cloud internal circulation. This was confirmed in their simulations, which demonstrated a linear increase in the leakage rate over time.

2. Later fate of the clouds

The clouds described by Nitsche & Batchelor (1997) were composed of a relatively small number of particles ($20 \lesssim N \lesssim 320$) and exhibited a high degree of cohesion until they disintegrated due to the constant loss of particles. In contrast, clouds comprising a larger number of particles ($N \gtrsim 500$) were subsequently studied and were seen to become unstable (see e.g. Adachi, Kiriyama & Yoshioka 1978; Machu *et al.* 2001; Metzger, Nicolas & Guazzelli 2007). These clouds initially remained roughly spherical, with a leakage of particles in a vertical tail, and then slowly evolved into a torus which subsequently broke up into two droplets in a repeating cascade, as shown in figure 2.

Falling clouds




Figure 2. A schematic representation of the evolution of the cloud into a torus and subsequent breakup (the time, t^* , is normalised by the time for the spherical cloud to fall its radius) (from a drawing of Sylvie Pic in Guazzelli & Morris 2012).

The leakage rate found by Nitsche & Batchelor (1997) was at least confirmed as the cloud remains spherical while falling through the first 10 cloud diameters. At larger times, when the cloud has evolved towards a toroidal shape, the loss slowed down, and a $t^{-1/3}$ leakage rate was observed (Metzger *et al.* 2007). What is more intriguing is the later development that took place long after the early times of cloud evolution studied by Nitsche & Batchelor (1997). This process involved the robust evolution of the initial spherical cloud into a torus, followed by its subsequent breakup, which occurs even in the complete absence of inertia and without the need to perturb the initial cloud shape. The previously described leakage of particles from the outer streamlines of the toroidal circulation results in a deficit of particles near the vertical axis in the central region of the cloud, which in turn gives rise to the formation of a torus. The toroidal cloud is then observed to expand and to break into two droplets for a critical horizontal to vertical aspect ratio. While the precise mechanism by which the toroidal cloud expands remains unclear, the breakup of the torus is a consequence of the change in flow configuration created by the particles when the aspect ratio reaches a critical value (Metzger *et al.* 2007).

The point-particle approach, pioneered by Nitsche & Batchelor (1997), has proven to be highly effective in capturing the evolution of the cloud. Subsequent numerical simulations, which have employed a range of varying degrees of sophistication, have been conducted to replicate this evolution (see, for example, one of the latest modelling attempt by Zhan & Wenxiao 2024). These more evolved simulations capture the multi-body and lubrication effects that are missed in the Stokeslet approximation and are certainly more accurate at modelling more concentrated clouds. However, the primary objective of the simplified Stokeslet simulation is to demonstrate the minimal physics required to describe the long-range hydrodynamic interactions between the particles and to illustrate how the

coupling between hydrodynamics and the microscopic arrangement of the particles gives rise to a collective dynamics.

Up to this point, our attention has been directed towards clouds of spherical particles. A comparable evolution and breakup has been observed in clouds of fibres, although at a faster rate due to the self-motion of the anisotropic particles (Park *et al.* 2010).

3. Beyond Stokes flows

In §§ 1 and 2, inertia was ignored. However, in most phenomena involving the dispersion of particles, such as turbidity currents, volcanic clouds, particle sedimentation in river beds or dust particle transport in the atmosphere, the particulate flow is dominated by inertial forces.

As the inertia of the falling cloud is increased, a transition occurs to a regime dominated by macro-scale inertia. This transition happens when the inertia at the scale of the cloud becomes large, which is indicated by the cloud Reynolds number, $Re_c = \rho_f V_c R / \mu$, reaching a value of approximately one, i.e. $Re_c \sim 1$ (Bosse *et al.* 2005). The subsequent transition is toward a micro-scale inertial regime when the inertia at the particle scale becomes important, i.e. when the individual particle wakes interact within the cloud boundaries. In its most basic form, particles interact through their steady Oseen velocity fields within the cloud and the micro-scale inertial regime occurs when the inertial length is of the order of the cloud radius, i.e. $a/Re_a \sim R$, where $Re_a = \rho_f V_s a / \mu$ is the particle Reynolds number (Subramanian & Koch 2008). In both inertial regimes, the cloud deforms into a flat torus that eventually destabilises and breaks up into a number of secondary droplets. However, particle leakage is much weaker if not null. While this evolution resembles that observed in the Stokes regime, the physical mechanisms involved are qualitatively different (Pignatel, Nicolas & Guazzelli 2011). In the inertial regimes, the evolution towards a torus shape is due to fluid inflow at the rear of the cloud, which leads to a decrease in particle leakage. The breakup process also differs, occurring at a larger aspect ratio of the torus within the inertial regime.

The aforementioned studies consider finite-but-moderate-Reynolds-number clouds. However, there are additional complexities in even larger-Reynolds-number clouds, including turbulent clouds of particles whose behaviour may deviate from a turbulent thermal model due to inertial particulate effects (see e.g. Kriaa *et al.* 2022). Furthermore, the interactions between particles and flow structures, as well as the collective effects between particles, are also important when particle clouds are settling in a complex flowing fluid rather than a quiescent fluid (see e.g. Marchetti, Bergougnoux & Guazzelli 2011).

Although these cloud behaviours are considerably more complex than those studied by Nitsche & Batchelor (1997), their work evidencing collective effects remains a great source of inspiration.

Declaration of interests. The author reports no conflict of interest.

Author ORCIDs.

 Élisabeth Guazzelli <https://orcid.org/0000-0003-3019-462X>.

REFERENCES

- ADACHI, K., KIRIYAMA, S. & YOSHIOKA, N. 1978 The behavior of a swarm of particles moving in a viscous fluid. *Chem. Engng Sci.* **33**, 115–121.
BATCHELOR, G.K. 1974 Low-Reynolds-number bubbles in fluidised beds. *Arch. Mech.* **26**, 339.
BOSSE, T., KLEISER, L., HÄRTEL, C. & MEIBURG, E. 2005 Numerical simulation of finite Reynolds number suspension drops settling under gravity. *Phys. Fluids* **17**, 037101.

Falling clouds

- EKIEL-JEŻEWSKA, M.L., METZGER, B. & GUAZZELLI, É. 2006 Spherical cloud of point particles falling in a viscous fluid. *Phys. Fluids* **18**, 038104.
- GUAZZELLI, É. & MORRIS, J.F. 2012 *A Physical Introduction to Suspension Dynamics*. Cambridge University Press.
- HADAMARD, J.S. 1911 Mouvement permanent lent d'une sphère liquide et visqueuse dans un liquide visqueux. *C. R. Acad. Sci. (Paris)* **152**, 1735–1738.
- KRIAA, Q., SUBRA, E., FAVIER, B. & LE BARS, M. 2022 Effects of particle size and background rotation on the settling of particle clouds. *Phys. Rev. Fluids* **7**, 124302.
- MACHU, G., MEILE, W., NITSCHE, L.C. & SCHAFLINGER, U. 2001 Coalescence, torus formation and break-up of sedimenting clouds: experiments and computer simulations. *J. Fluid Mech.* **447**, 299–336.
- MARCHETTI, B., BERGOUGNOUX, L. & GUAZZELLI, É. 2011 Falling clouds of particles in vortical flows. *J. Fluid Mech.* **908**, A30.
- METZGER, B., NICOLAS, M. & GUAZZELLI, É. 2007 Falling clouds of particles in viscous fluids. *J. Fluid Mech.* **580**, 283–301.
- NITSCHE, J.M. & BATCHELOR, G.K. 1997 Break-up of a falling drop containing dispersed particles. *J. Fluid Mech.* **340**, 161–175.
- PARK, J., METZGER, B. & GUAZZELLI, É. & BUTLER, J.E. 2010 A cloud of rigid fibres sedimenting in a viscous fluid. *J. Fluid Mech.* **648**, 351–362.
- PIGNATEL, F., NICOLAS, M. & GUAZZELLI, É. 2011 A falling cloud of particles at a small but finite Reynolds number. *J. Fluid Mech.* **671**, 34–51.
- RYBCZYŃSKI, W. 1911 Über die fortschreitende Bewegung einer flüssigen Kugel in einem zähen Medium. *Bull. Acad. Sci. Cracovie A*, 40–46.
- SUBRAMANIAN, G. & KOCH, D.L. 2008 Evolution of clusters of sedimenting low-Reynolds-number particles with Oseen interactions. *J. Fluid Mech.* **603**, 63–100.
- ZHAN, M. & WENXIAO, P. 2024 Shape deformation, disintegration, and coalescence of suspension drops: efficient simulation enabled by graph neural networks. *Intl J. Multiphase Flow* **176**, 104845.