

**Objectives:** The current study objective is to investigate the sleep quality of polypharmacy older adults and its association with depression and anxiety in this population.

**Methods:** A cross-sectional study was conducted in Iraq from June to August 2024. A total of 245 participants were selected from internal medicine wards in hospitals and private clinics. All participants were aged 50 years or older and had been taking five or more medications for at least 90 days. The study questionnaire comprised three main sections: demographic data, the Patient Health Questionnaire-4 (PHQ-4), and the Pittsburgh Sleep Quality Index (PSQI). A PSQI score greater than 5 was used to identify poor sleepers. The association between poor sleep and other factors was assessed by using chi-squared tests and binary logistic regression.

**Results:** The sample consisted of 245 polypharmacy patients, with a mean age of  $61.5 \pm 12.4$  years. Of the participants, 150 (61%) were female and 95 (39%) were male. Anxiety was observed in 95 (38.8%) participants, depression in 96 (39.2%), and poor sleep quality in 189 (77.1%). The study found significant associations between both anxiety (OR = 3.4 [95% CI: 1.55-7.57],  $p = 0.002$ ) and depressive symptoms (OR = 2.43 [95% CI: 1.15-5.15],  $p = 0.020$ ) with poor sleep quality.

**Conclusions:** Our study suggests that Iraqi polypharmacy older adults suffer from poor sleep quality, with depression and anxiety potentially exacerbating this issue. The findings suggest that mental health support is necessary for older adults with polypharmacy.

**Disclosure of Interest:** None Declared

## EPV1910

### Sleep disorders and stress in children and adolescents with ASD

A. Alvarez<sup>1\*</sup>, N. Santamaria<sup>1</sup>, V. Bote<sup>1</sup>, R. Medina<sup>1</sup>, J. A. Monreal<sup>1</sup>, B. Sánchez<sup>1</sup> and A. Hervás<sup>1</sup>

<sup>1</sup>Mental Health, University Hospital Mutua Terrassa, Terrassa, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2324

**Introduction:** Sleep disorders are very common in children with neurodevelopmental disorders such as autism. Poor sleep can have detrimental effects on cognitive processes, attention, memory, language, and regulation of mood and behavior. Many people with autism have difficulty correctly processing sensory information that reaches them from both the environment and their own body. Depending on how the person with autism perceives the sensations, these can impact on sleep, causing nighttime awakenings or poor sleep conciliation. It is vitally important to evaluate nighttime rest and have knowledge of appropriate tools to improve sleep in patients with ASD.

**Objectives:** In this work we aim to evaluate, in a child-youth population diagnosed with ASD who have been admitted to our ASD Day Hospital during the year 2024, the prevalence of sleep disorders and their characteristics. As well as generating strategies to improve sleep conciliation and factors to take into account or avoid that may be affecting our patients.

**Methods:** A review is carried out of the clinical history of patients admitted to the TEA Day Hospital during the year 2024. Sleep problems are analyzed, as well as their relationship with stress. Results and measures to improve the quality of night rest in these patients are presented.

**Results:** From January to September 2024, a total of 26 patients with autism have been treated at the ASD Day Hospital of the University Hospital of Mutua Terrassa, in the following programs: intensive, eating problems and low functionality. Of these, 32% were women, with an average age of 11.64 and an average stay of 40 days.

More than half of the patients had sleep problems, where the majority had more than one. Among the most frequent: difficulties falling asleep (12), maintenance insomnia (8), frequent awakenings (4), night terrors (6), and the need for the presence of parents and/or sensory interventions to be able to fall asleep (12).

The main interventions carried out have consisted of: establishing adequate sleeping habits, avoiding naps, avoiding highly stimulating activities, feeling tired beforehand, adapting environmental conditions to the needs of the child and giving proprioceptive information before sleeping.

Among the most commonly used drugs we found atypical antipsychotics: olanzapine, risperidone. Some antidepressants: mirtazapine and occasionally benzodiazepines. The vast majority of children took melatonin.

**Conclusions:** Sleep disorders are highly prevalent in patients with ASD, but correct intervention can improve comorbid symptoms such as anxiety, stress and discomfort, which are largely related to poor sleep.

Sensory-perceptive interventions are essential to improve nighttime rest in children with ASD

**Disclosure of Interest:** None Declared

## EPV1911

### Daytime Sleepiness Among Healthcare Workers Following SARS-CoV-2 Infection

E. Bechrifa<sup>1\*</sup>, D. Brahim<sup>1</sup>, N. Mechergui<sup>2</sup>, I. Youssef<sup>3</sup>, M. Mersni<sup>1</sup>, G. Bahri<sup>1</sup>, H. Ben Said<sup>4</sup>, M. Bani<sup>1</sup> and N. Laadhari<sup>1</sup>

<sup>1</sup>Occupational Medicine Department, Charles Nicolle Hospital;

<sup>2</sup>Occupational Medicine Department, Habib Thameur Hospital;

<sup>3</sup>Occupational Medicine Department, Mongi Slim Hospital and

<sup>4</sup>Occupational Medicine Department, Maternity and Neonatology Center, Tunis, Tunisia

\*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2325

**Introduction:** COVID-19 infections had a variety of symptoms, including a range of sleep disorders such as daytime sleepiness.

**Objectives:** The objective of the study was to determine the frequency and persistence of daytime Sleepiness among HCWs post-COVID-19 and analysis of related risk factors.

**Methods:** A prospective descriptive study was conducted over six months (January-July 2022) among HCWs at Charles-Nicollé Hospital in Tunis who contracted COVID-19. Daytime sleepiness was evaluated at three intervals: during the infection (T0), at three months, and at six months post-infection, using the Epworth Sleepiness Scale (ESS) in its French version. Anxiety and depressive symptoms were assessed with the Hospital Anxiety and Depression Scale (HAD) to examine their association with sleep disturbances.

**Results:** The study included 155 HCWs, with an average age of  $40.2 \pm 10.3$  years and an average work seniority of  $14.1 \pm 10$  years. The assessment of anxiety-depressive disorders in the study population using the HAD scale showed that anxiety symptoms were certain in 27.3% of the personnel, while depressive symptoms were certain in 21.4% of cases. Additionally, 48.1% of participants showed no