

Bristol-Myers-Squibb, CIBERSAM, FIS- ISCiii, Janssen Cilag, Lundbeck, Otsuka, Servier and Medtronic., Consultant of: Has been a consultant to AB-Biotics, AstraZeneca, Bristol-Myers-Squibb, CIBERSAM, FIS- ISCiii, Janssen Cilag, Lundbeck, Otsuka, Servier and Medtronic., M. T. Campillo: None Declared, G. Martínez-Alés Grant / Research support from: American Foundation for Suicide Prevention grant ECR-1-101-23, Brain and Behavior Research Foundation grant NARSAD 31312 and National Institutes of Mental Health grant 1R25MH129256-01A1, L. Cano: None Declared, D. Guinart Grant / Research support from: CM21/00033, Consultant of: Has been a consultant and/or advisor or has received honoraria from: Angelini, Otsuka, Lundbeck and Teva., P. Mortier Grant / Research support from: This work was supported by grant CP21/00078 from the ISCIII-FSE Miguel Servet

## EPV1966

### Depression symptom profile in a nationally-representative cohort of older adults: Relationships with suicide-related outcomes

I. Pérez-Diez<sup>1,2,\*</sup>, A. de la Torre-Luque<sup>2,3</sup>, J. L. Ayuso-Mateos<sup>1,2,4</sup>, B. Dolz del Castellar<sup>1,2,4</sup>, M. Miret<sup>1,2</sup>, J. Domènech-Abella<sup>2,5</sup>, B. Olaya<sup>2,5</sup> and J. Maria Haro<sup>2,5</sup>

<sup>1</sup>Department of Psychiatry, Universidad Autónoma de Madrid;

<sup>2</sup>Instituto de Salud Carlos III, Centro de Investigación Biomédica en Red de Salud Mental. CIBERSAM; <sup>3</sup>Department of Legal Medicine, Psychiatry and Pathology, Universidad Complutense de Madrid; <sup>4</sup>Instituto de Investigación Sanitaria Princesa (IIS-Princesa), Madrid and <sup>5</sup>Sant Joan de Déu Research Institute, Barcelona, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2366

**Introduction:** Depression among older adults is a significant public health issue that often goes underrecognized, despite its profound impact on their quality of life and mental well-being. With advancing age, many individuals encounter challenges such as loneliness, chronic illness, and bereavement, which heighten their vulnerability to depressive symptoms. These symptoms are not only detrimental to overall health but are also strongly associated with an increased risk of suicidal behaviours. Although these suicidal behaviours are preventable, much research should be done to disentangle development pathways in older adults.


**Objectives:** The aim of this research is to identify the symptomatic profiles of depression in the elderly population and to relate this profile to suicidal outcomes.

**Methods:** We utilized data from the “*Edad con Salud*” cohort study, a longitudinal household survey conducted on a nationally representative sample in Spain (aged 18+ years). Participants who could not be personally interviewed and those under 50 years old were excluded, resulting in a final sample of 3,533 participants (54.31% female, age = 66.42(10.27)). Suicidal behaviours and depressive symptoms were assessed using the Composite International Diagnostic Interview (CIDI 3.0). A Latent Class Analysis (LCA) was performed at base level to identify participant profiles based on 8 depressive symptoms. We followed this with a survival analysis to assess differences between profiles in the progression of suicidal ideation and attempts in the different waves. Finally, Cox regression was used to examine the impact of sociodemographic and clinical covariates on suicidal behaviours.

**Results:** The three-cluster model provided the best fit. The first class (72.52%) included most participants without depressive

symptoms; the second class (13.67%) included participants with three symptoms (depressive mood, diminished interest or pleasure and fatigue); and the third class (13.81%) included participants with all eight symptoms (see Image 1). Survival analysis revealed significant differences between the groups over time for both suicidal ideation ( $\chi^2=1332$ ,  $p < .01$ ) and suicide attempts ( $\chi^2=215$ ,  $p < .01$ ), with the third group having the highest number of cases. Cox regression indicated that suicidal ideation was associated with the second ( $HR = 5.38$ ) and third classes ( $HR = 59.8$ ), age ( $HR = 0.97$ ), and loneliness ( $HR = 0.91$ ), among other variables. Suicide attempts were associated with the second ( $HR = 6.08$ ) and third classes ( $HR = 25.3$ ), age ( $HR = 0.92$ ), and loneliness ( $HR = 1.02$ ).

**Image 1:**



**Conclusions:** Participants with depressive symptoms, even at a subclinical level, have an increased risk of suicidal behaviours, especially when other risk factors such as loneliness are present.

**Disclosure of Interest:** None Declared

## EPV1967

### Shifting Perspectives: The Importance of Assessing Suicide Risk in Medically Ill Patients Without Psychiatric History

V. H. Santos<sup>1,2,\*</sup>, F. M. Tehrani<sup>3</sup>, M. Pires<sup>4</sup>, R. P. Andrade<sup>5</sup>, B. Sousa<sup>1</sup>, Z. C. e Sá<sup>1</sup>, T. Carvalhão<sup>1</sup> and S. Fontes<sup>1</sup>

<sup>1</sup>Department of Psychiatry and Mental Health, Cova da Beira Local Health Unit; <sup>2</sup>Faculty of Health Sciences, University of Beira Interior, Covilhã, Portugal; <sup>3</sup>Child and Adolescent Mental Health Center, Mental Health Services in the Capital Region of Denmark, Copenhagen, Denmark; <sup>4</sup>Department of Psychiatry and Mental Health, Guarda Local Health Unit, Guarda and <sup>5</sup>Department of Psychiatry and Mental Health, Viseu Dão-Lafões Local Health Unit, Viseu, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2367

**Introduction:** Chronic medical conditions are increasingly recognized as significant contributors to suicide risk, especially in patients without prior psychiatric diagnoses (Østergaard *et al.* JAMA Psychiatry 2024). This case examines the psychiatric impact in a 65-year-old male admitted to general surgery for abdominal pain, who subsequently underwent an ileocelectomy for suspected gastrointestinal malignancy.

**Objectives:** To explore the psychiatric impact of medical morbidity on suicide risk, emphasizing recent findings suggesting heightened attention for patients without a psychiatric history.

**Methods:** The patient, with no prior psychiatric history, was observed by liaison psychiatry after verbalizing suicide ideation