

Research Paper

Cite this article: Lee S-K, Lee EJ (2025). Effects of Korean water deer (*Hydropotes inermis argyropus*) feces on seed germination and early seedling growth: insights into their contribution to seed dispersal. *Seed Science Research* **35**, 27–32. <https://doi.org/10.1017/S096025852400028X>

Received: 11 March 2024

Revised: 7 June 2024

Accepted: 16 July 2024

First published online: 24 January 2025

Keywords:

endozoochory; long-distance dispersal; seed dispersal; seed germination; seedling growth; post-dispersal events

Corresponding author:

Eun Ju Lee;

Email: ejlee@snu.ac.kr

Effects of Korean water deer (*Hydropotes inermis argyropus*) feces on seed germination and early seedling growth: insights into their contribution to seed dispersal

Seung-Kyung Lee^{1,2}  and Eun Ju Lee¹ 

¹School of Biological Sciences, Seoul National University, Seoul, Republic of Korea and ²Institute of Health and Environment, Seoul National University, Seoul, Republic of Korea

Abstract

Endozoochory, the dispersal of seeds through the animal gut passage, plays a significant role in vegetation dynamics. The success of endozoochorous seed dispersal depends on each stage of the process: ingestion by animals, gut passage, and post-dispersal events after defecation. After the deposition of seeds through feces, the effects of feces on the initial stages of seedling establishment, including seed germination and seedling growth, can significantly impact overall survival. The pattern of fecal effects on plant species depends on the animal species. In this study, we investigated the effects of feces presence on seed germination and early seedling growth using feces of the Korean water deer (*Hydropotes inermis argyropus*). We conducted a germination experiment on 12 plant species belonging to 10 plant families, which are known to germinate in the feces of Korean water deer. The study compared the seed germination rate and seedling length after germination between seeds sown with and without feces of the Korean water deer. In general, we found that the presence of deer feces *per se* had no significant effects on seed germination and early growth stages. However, additional research on post-dispersal events such as long-term growth, fecal type, and germination conditions is needed to fully understand the costs and benefits of endozoochory.

Introduction

Endozoochory, the dispersal of seeds through the gut passage of animals, facilitates the dispersal of seeds from numerous plant species, facilitating the colonization in new habitats, and the expansion of plant populations (McConkey et al., 2012; Baltzinger et al., 2019). The success of the seed dispersal mechanism is influenced by various factors at each stage of the dispersal process (Cain et al., 2000; Baltzinger et al., 2019). In each stage of seed dispersal (i.e., from emigration, transfer, and to immigration), seed dispersal vectors contribute to plant dispersal in terms of both quantity and quality components (Schupp, 1993; van Leeuwen et al., 2022). Seed dispersal effectiveness depends on the quantity of seeds dispersed by animals and whether these seeds can germinate and establish in new habitats (Schupp et al., 2010). For endozoochory by ungulates, seed dispersal effectiveness has been widely examined in each stage, from ingestion of seeds, transportation of seeds through the animal gut, and to the dissemination stage (Baltzinger et al., 2019).

During the dissemination stage, seeds deposited through endozoochory encounter diverse environmental conditions (i.e., deposit quality) that significantly impact their survival and establishment (Baltzinger et al., 2019). These conditions include moisture level changes following the fecal desiccation, fecal types, feeding regime of animals, nutrient availability and seedling competition within fecal deposits (e.g., Milotić and Hoffmann, 2016b, 2017; Baltzinger et al., 2019; Karimi et al., 2020). Animal feces can serve as potential nutrient source containing elements such as nitrogen and potassium (Sakadevan et al., 1993; Williams and Haynes, 1995; Dai, 2000). Simultaneously, animal feces may contain harmful compounds that prevent seed germination (Marambe et al., 1993; Meyer and Witmer, 1998). The effects of fecal matter on seed germination and plant growth vary depending on the specific plant and animal species involved due to their fecal morphological types and digestion (Milotić and Hoffmann, 2016b; Guevara-Torres and Facelli, 2023; Ramos et al., 2024). These vector animal species-specific results emphasize the need of testing the response to each factor, which affects the successful results of endozoochorous seed dispersal. Understanding these effects is crucial for comprehending the dynamics of endozoochorous seed dispersal and early seedling establishment.

Hydropotes inermis argyropus, commonly known as the Korean water deer, is the most prevalent ungulate species on the Korean peninsula. This deer play a significant role in the dispersal of various plant species, especially forbs and those originating from open habitats

(Lee et al., 2021, 2022). A study by Park and Lee (2014) examined the influence of Korean water deer feces on the soil condition and growth of *Zea mays* seedlings. The authors found that the amount of feces added significantly affected the growth of *Zea mays* seedlings in nutrient-poor soil. This finding suggests that the fecal components of the deer can provide nutrients to the soil for plants to grow. However, at the same time, animal feces may also contain phytotoxic elements that could impede seed germination (e.g., frugivorous birds, Meyer and Witmer (1998); cattle and horse, (Milotić and Hoffmann, 2016b, 2017); cattle, chicken and pig, Marambe et al. (1993)), and still little is known about how the presence of the Korean water deer feces affects the germination and growth of plant species dispersed through the deer endozoochory. Given the relevance of endozoochory, it is imperative to systematically investigate and analyze the specific effects of feces on seed germination within this ecological context.

The objective of this study was to assess the effects of feces itself on seed germination and the early growth of seeds dispersed by the Korean water deer endozoochory through germination tests. Specifically, we investigated whether the presence of feces have an effect on (i) seed germination of species known to be dispersed through endozoochory and (ii) on the early growth of seedlings of 12 plant species previously reported to germinate from the feces of Korean water deer (Lee and Lee, 2020; Lee et al., 2022).

Materials and methods

Seed preparation

Based on previous studies (Lee and Lee, 2020; Lee et al., 2022), we selected 12 plant species that have been observed to germinate from the feces of the Korean water deer (Table 1). These species had previously been observed to germinate from the feces of the Korean water deer. The seeds of selected plant species were obtained for research purposes from the National Institute of Biological Resources. For species nomenclature, we followed a database managed by the National Institute of Biological

Resources (<https://species.nibr.go.kr/>) and to determine whether each target species was an alien plant in South Korea, we referred to Korea National Arboretum (2019).

Feces preparation

For the experiments, the feces of the Korean water deer were collected from Taehwa Research Forest and Civilian Control Zone, where previous sampling had been conducted (Lee and Lee, 2020; Lee, et al. 2022). The fecal pellet groups were individually stored in zipper bags, transported to the laboratory, and cleaned on their outer surface. Subsequently, the fecal pellets were allowed to air-dry at room temperature. To mitigate the potential effects of different sampling seasons and sites, all collected fecal samples were combined. The feces were ground to avoid possible contamination from external seeds and to ensure a consistent fecal amount could be used in the experiments.

Experimental setup

For the experimental setup, above mentioned 12 plant species were tested for germination under conditions of with (feces) and without the feces (control) of the Korean water deer. For each species, five seeds were sown in each pot, resulting in 10 replications for each treatment (with and without feces), and thus a total of 240 pots were tested. The pots (150 mm in diameter and 135 mm in height) were filled with commercial soil (Baroker, Seoul Bio Co., Ltd., Eumseong, Korea). In each pot with the treatment condition of adding the feces of Korean water deer, 5 g of air-dried and ground feces was added. All seeds were directly sown on the surface of feces or soil to facilitate light availability. The experiment was carried out at the temperature-controlled greenhouse facility at Seoul National University. The temperature of the greenhouse was set at 23°C during the experiments. The pots were watered on a regular basis and kept moist. During the experimental period, the pots were randomly allocated and redistributed every other day. The number of seedlings that germinated for each species (emergence rate) in each treatment was checked 60 days after the initial day of sowing, and the height of each seedling was measured.

Data analysis

To investigate the effects of fecal presence on the seedling emergence rate in each species, a Generalized Linear Model (GLM) of Poisson distribution was applied. To compare the effects of fecal presence on seedling height, the data were validated for homogeneity and homoscedasticity prior to analysis, and the *t*-test and Wilcoxon rank-sum tests were used. Visualization of the data was conducted using ggplot2 (Wickham, 2016) package. All analyses were conducted in R version 4.2.1 (R Core Team, 2022).

Results

Number of seedlings emerged

All 12 tested plant species germinated. Across all treatments (control and feces) and species, a median of one seed germinated per pot sown with five seeds. For all 12 species tested, the comparison of the seeds that were sown on the feces had no significantly different emergence rate from that of seeds that were sown in soil ($P > 0.05$; GLM model fitted with Poisson distribution; Fig. 1). The results showed that overall, the feces had no

Table 1. The list of plant species tested for germination, and their growth form

Species	Growth form	Note
Amaranthaceae <i>Amaranthus lividus</i>	Forb	
Asteraceae <i>Erigeron annuus</i>	Forb	Alien plant
Brassicaceae <i>Rorippa palustris</i>	Forb	
Caryophyllaceae <i>Stellaria aquatica</i>	Forb	
Chenopodiaceae <i>Chenopodium album</i> var. <i>centrorubrum</i>	Forb	Alien plant
Moraceae <i>Morus bombycis</i>	Woody plant	
Onagraceae <i>Ludwigia epilobioides</i>	Forb	
Plantaginaceae <i>Plantago asiatica</i>	Forb	
Poaceae <i>Panicum bisulcatum</i>	Graminoid	
	<i>Digitaria sanguinalis</i>	Graminoid
Urticaceae <i>Urtica angustifolia</i>	Forb	
	<i>Pilea mongolica</i>	Forb

The species that are categorized as alien plants are followed by Korea National Arboretum (2019).




Figure 1. The number of seedlings of each plant species germinated in a condition with feces (Feces) and without feces (Control) of the Korean water deer. n.s. denotes for not significant ($P > 0.05$) based on the GLM model with Poisson distribution.

significant impact on the germination rate of any species. Detailed statistics can be found in the Supplementary Table 1.

Seedling height

On the final day of the germination test, we measured the height of each seedling that germinated. We found that the height of all species that germinated showed no differences from that of control seedlings that were sown on the soil for 11 out of 12 plant species tested ($P > 0.05$; t -test; Fig. 2; Supplementary Table 2). In the case of *Urtica angustifolia* sown on the ground feces showed higher seedling height compared to that of the control condition (t -statistic = 2.657; d.f. = 12.138, $P = 0.021$; t -test).

Discussion

Understanding the effects of feces on seed germination and seedling growth enables us to comprehend the costs and benefits associated with the entire process of endozoochorous seed dispersal. This study aimed to investigate the influence of Korean water deer feces itself on seed germination and initial seedling growth of plant species dispersed through Korean water deer endozoochory. Among the 12 species studied, there were no significant differences in seed performance between those sown with feces and those sown without feces, and no differences in seedling growth for 11 of these species, except for *U. angustifolia*. Our findings indicate that negligible impact of feces *per se* on seed germination and initial seedling growth in the context of the Korean water deer endozoochory.

Previous studies suggest that the response of plant species depends on the plant species and physiology of animal species involved (e.g., ruminant and hindgut-fermenting). The results of this study could be associated with the gastrointestinal

physiology of Korean water deer, a ruminant species. While the effects of feces on seed germination differed among plant species, certain species exhibited lower germination rates when exposed to feces (Milotić and Hoffmann, 2016b). These results are also affected by the fermented gut characteristics of cows and horses (Milotić and Hoffmann, 2016b). Additionally, in a study examining *Neltuma flexuosa* seed germination and seedling growth with feces from cow, horse and mara, the results varied depending on the animal species (Ramos et al., 2024). Horse and mara feces showed reduced germination rates, while cow feces had no discernible effects. Regarding seedling growth, the native and wild mara species showed negligible effects, whereas domestic cow and horse species exhibited increased seedling growth compared to normal conditions. Moreover, the effects of native herbivore kangaroo and domestic sheep on different plant species also varied, promoting the growth of aboveground biomass in wallaby grass and reducing wild oat aboveground biomass, depending on the animal species involved (Guevara-Torres and Facelli, 2023). Therefore, it is essential to analyze and compare the results specific to both animal and plant species, to understand the ecological correlation between seed dispersal through endozoochory and associated animals and plants. Furthermore, it is crucial to consider not only the seed germination rate but also the timing of seed germination. Previous research has shown that feces can delay seed germination (Meyer and Florence, 1996; Ramos Font et al., 2015; Milotić and Hoffmann, 2016b). Additionally, certain species exhibited low germination rates, suggesting that their germination requirements may not have been met, including mortality shortly after germination. For example, *Morus bombycis* showed a particularly low germination rate.

Overall seedling height showed no significant differences with the addition of feces. However, for *U. angustifolia*, the addition of




Figure 2. Seedling height of each plant species germinated in a condition with feces (Feces) and without feces (Control) of the Korean water deer. n.s. denotes for not significant and * denotes for significant ($P < 0.05$) following the t -test.

feces showed higher growth (Fig. 2). It is important to note that the low germination rate for some species could have obscured the results, which could reduce the statistical power. In our study, we only assessed the early growth of the seedlings, which did not allow for the evaluation of the longer-term effects of feces on seeds of species dispersed through endozoochory. However, a study by Milotić and Hoffmann (2016a) demonstrated that over an extended growth period, plants with feces showed higher biomass production compared to those without feces. Related to this, Park and Lee (2014) tested the effects of the Korean water deer feces on the growth of *Zea mays*, demonstrating that the addition of feces resulted in greater plant material production than in plants without feces in nutrient-poor soil. Although we found negligible effects on early seedling stages, seed dispersal through the Korean water deer endozoochory may have positive consequences by promoting the growth of plant species in the longer term, as the fecal components can release the potential beneficial nutrients such as nitrogen, potassium, phosphorus (Rowarth et al., 1985; McDowell, 2006; Park and Lee, 2014; Park et al., 2015). Moreover, this pattern could also be related to the time it takes for nutrients to be released from fecal materials through physical breakdown and mineralization processes (Rowarth et al., 1985; Eichberg et al., 2007).

According to our previous feeding experiments of the Korean water deer, testing the effects of gut passage on seed germination, the seeds consumed by water deer exhibited a recovery rate of less than 30% across all tested species (Lee et al., 2021). It is evident that a considerable proportion of seeds undergo consumption during gut passage, which can be related to the seed traits including seed morphology (Mouissie et al., 2005; Lee et al., 2021). The low recovery rate of ingested plant seeds, combined with potential seedling competition, could significantly impact the overall survival and dispersal success of seeds through endozoochory. Based on the current results and our former research on gut passage effects (Lee et al., 2021), the deer is not an effective endozoochorous seed disperser. According to a meta-analysis study (Torres et al., 2020), the deer family (Cervidae) is not an effective seed dispersal vector for endozoochory, showing that gut passage by deer has no significant effects on seed germination. Furthermore, despite our efforts to assess the impact of Korean water deer feces on seed germination and seedling growth, a thorough experimental design using gut-passed seeds is essential to evaluate the effectiveness of endozoochorous seed dispersal, as the gut passage may alter the chemical and mechanical aspects of seeds (Samuels and Levey, 2005).

Endozoochorous seed dispersal is a comprehensive process, involving ingestion, defecation, and post-dispersal events. As the overall success of emergence and fruiting in real situations has low probabilities (Eichberg et al., 2007), final seedling emergence in real deposition events and fruiting stage also needs to be tested. After deposition, the feces are likely to undergo changes in their nutritional components and moisture (Holter, 1991, 2016). Moreover, we placed the seeds on the top parts of ground feces; however, in real deposition events, the water deer defecate the seeds in the structure of small fecal pellet groups, which may impose additional physical and biological barriers to seed germination. The feces type of deer is usually fecal pellet groups defecated with several small fecal pellets and is prone to desiccate after deposition (Welch, 1985; Eichberg et al. 2007; Milotić and Hoffmann, 2016b), affecting seed germination due to moisture loss.

After deposition through endozoochory, seeds encounter not only environmental factors such as habitat deposition site,

weather, and moisture (Dickinson and Craig, 1990), but also, biological components including secondary seed removal by dung beetles (Milotić et al., 2019; Urrea-Galeano et al., 2019) and seedling competitions (Milotić and Hoffmann, 2017), and each stage determining the success of endozoochorous seed dispersal. Furthermore, in outdoor environments, the seed germination rate would be decreased due to the fast desiccation in the outdoor environments, and the germination conditions in the outdoor environments compared to greenhouse results in a lower germination rate as well (Milotić and Hoffmann, 2016b; Karimi et al., 2020). Furthermore, we sowed five seeds to avoid possible intra-species competition; however, in our former Korean water deer results, the highest density showed 161 seedlings per fecal pellet group (Lee and Lee, 2020), suggesting possible higher competition among seedlings. Intra-species competition may have different effects on seedling dispersal patterns as well (Milotić and Hoffmann, 2017). Therefore, the comprehensive understanding of endozoochorous seed dispersal processes by the Korean water deer needs to be thoroughly investigated from ingestion to defecation to comprehensively understand the costs, benefits, and overall effectiveness.

Acknowledgements. We thank the members of the Plant Ecology Laboratory of Seoul National University for their help with the experiments. We appreciate anonymous reviewers for their constructive comments on the manuscript.

Funding statement. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No.2017R1A2B4006761).

Competing interests. The authors have no relevant financial or non-financial interests to disclose.

References

- Baltzinger C, Karimi S and Shukla U (2019) Plants on the move: hitch-hiking with ungulates distributes diaspores across landscapes. *Frontiers in Ecology and Evolution* 7, 38. doi:10.3389/fevo.2019.00038.
- Cain ML, Milligan BG and Strand AE (2000) Long-distance seed dispersal in plant populations. *American Journal of Botany* 87, 1217–1227.
- Dai X (2000) Impact of cattle dung deposition on the distribution pattern of plant species in an alvar limestone grassland. *Journal of Vegetation Science* 11, 715–724. doi:10.2307/3236578.
- Dickinson C and Craig G (1990) Effects of water on the decomposition and release of nutrients from cow pats. *New Phytologist* 115, 139–147.
- Eichberg C, Storm C and Schwabe A (2007) Endozoochorous dispersal, seedling emergence and fruiting success in disturbed and undisturbed successional stages of sheep-grazed inland sand ecosystems. *Flora-Morphology, Distribution, Functional Ecology of Plants* 202, 3–26.
- Guevara-Torres DR and Facelli JM (2023) Choose local: dung addition from native herbivores can produce substantial positive effects on the growth of native grasses compared to livestock dung. *Journal of Soil Science and Plant Nutrition* 23, 4647–4655. doi:10.1007/s42729-023-01380-7.
- Holter P (1991) Concentration of oxygen, carbon-dioxide and methane in the air within dung pats. *Pedobiologia* 35, 381–386.
- Holter P (2016) Herbivore dung as food for dung beetles: elementary coprology for entomologists. *Ecological Entomology* 41, 367–377.
- Karimi S, Hemami M-R, Esfahani MT and Baltzinger C (2020) Endozoochorous dispersal by herbivores and omnivores is mediated by germination conditions. *BMC Ecology* 20, 1–14.
- Korea National Arboretum (2019) *Checklist of Alien Plants in Korea*. Seoul: Dooroo Happy Co. Ltd.
- Lee S-K and Lee EJ (2020) Internationally vulnerable Korean water deer (*Hydropotes inermis argyropus*) can act as an ecological filter by endozoochory. *Global Ecology and Conservation* 24, e01368. doi:10.1016/j.gecco.2020.e01368.

- Lee S-K, Shin W-J, Ahn S, Kim Y, Kim J-T and Lee EJ (2021) Seed recovery and germination rate after gut passage by Korean water deer (*Hydropotes inermis argyropus*). *Seed Science Research* **31**, 311–318.
- Lee S-K, Ryu Y and Lee EJ (2022) Endozoochorous seed dispersal by Korean water deer (*Hydropotes inermis argyropus*) in Taehwa Research Forest, South Korea. *Global Ecology and Conservation* **40**, e02325. doi:10.1016/j.gecco.2022.e02325.
- Marambe B, Nagaoka T and Ando T (1993) Identification and biological activity of germination-inhibiting long-chain fatty acids in animal-waste composts. *Plant and cell physiology* **34**, 605–612.
- McConkey KR, Prasad S, Corlett RT, Campos-Arceiz A, Brodie JF, Rogers H and Santamaria L (2012) Seed dispersal in changing landscapes. *Biological Conservation* **146**, 1–13. doi:10.1016/j.biocon.2011.09.018.
- McDowell R (2006) Contaminant losses in overland flow from cattle, deer and sheep dung. *Water, Air, and Soil Pollution* **174**, 211–222.
- Meyer JY and Florence J (1996) Tahiti's native flora endangered by the invasion of *Miconia calvescens* DC (Melastomataceae). *Journal of Biogeography* **23**, 775–781. doi:10.1111/j.1365-2699.1996.tb00038.x.
- Meyer GA and Witmer MC (1998) influence of seed processing by Frugivorous Birds on germination success of three North American Shrubs. *The American Midland Naturalist* **140**, 129–139, 111.
- Milotić T and Hoffmann M (2016a) Cost or benefit for growth and flowering of seedlings and juvenile grassland plants in a dung environment. *Plant Ecology* **217**, 1025–1042. doi:10.1007/s11258-016-0629-2.
- Milotić T and Hoffmann M (2016b) Reduced germination success of temperate grassland seeds sown in dung: consequences for post-dispersal seed fate. *Plant Biology* **18**, 1038–1047.
- Milotić T and Hoffmann M (2017) The impact of dung on inter- and intra-specific competition of temperate grassland seeds. *Journal of Vegetation Science* **28**, 774–786. doi:10.1111/jvs.12535.
- Milotić T, Baltzinger C, Eichberg C, Eycott AE, Heurich M, Müller J, Noriega JA, Menendez R, Stadler J and Ádám R (2019) Functionally richer communities improve ecosystem functioning: dung removal and secondary seed dispersal by dung beetles in the Western Palaearctic. *Journal of Biogeography* **46**, 70–82.
- Mouissie AM, Van Der Veen CEJ, Veen GF and Van Diggelen R (2005) Ecological correlates of seed survival after ingestion by Fallow Deer. *Functional Ecology* **19**, 284–290. doi:10.1111/j.0269-8463.2005.00955.x.
- Park H and Lee S (2014) Factor of plant growth in relation to Feces of Korean water deer and land use patterns. *Journal of Wetlands Research* **16**, 443–452.
- Park H, Chun S and Lee S (2015) Study on effect on CO₂ flux of wetland soil by feces of Korean water deer (*Hydropotes inermis*). *Journal of Wetlands Research* **17**, 283–292.
- Ramos LC, Campos CM, Cona MI and Giordano CV (2024) The quality of endozoochorous depositions: effect of dung on seed germination and seedling growth of *Neltuma flexuosa* (DC.) CE Hughes & GP Lewis. *Journal of Arid Environments* **220**, 105114.
- Ramos Font M, González Rebollar J and Robles Cruz A (2015) Endozoochorous dispersal of wild legumes: from seed recovery to field establishment. *Ecosistemas* **24**, 14–21.
- R Core Team (2022) *R: A Language and Environment for Statistical Computing*. Vienna, Austria, R Foundation for Statistical Computing.
- Rowarth J, Gillingham A, Tillman R and Syers J (1985) Release of phosphorus from sheep faeces on grazed, hill country pastures. *New Zealand journal of agricultural research* **28**, 497–504.
- Sakadevan K, Mackay A and Hedley M (1993) Influence of sheep excreta on pasture uptake and leaching losses of sulfur, nitrogen and potassium from grazed pastures. *Soil Research* **31**, 151–162.
- Samuels IA and Levey DJ (2005) Effects of gut passage on seed germination: do experiments answer the questions they ask? *Functional Ecology* **19**, 365–368. doi:10.1111/j.1365-2435.2005.00973.x.
- Schupp EW (1993) Quantity, quality and the effectiveness of seed dispersal by animals. *Vegetatio* **107**, 15–29.
- Schupp EW, Jordano P and Gómez JM (2010) Seed dispersal effectiveness revisited: a conceptual review. *New Phytologist* **188**, 333–353. doi:10.1111/j.1469-8137.2010.03402.x.
- Torres DA, Castano JH and Carranza-Quiceno JA (2020) Global patterns in seed germination after ingestion by mammals. *Mammal Review* **50**, 278–290. doi:10.1111/mam.12195.
- Urrea-Galeano LA, Andresen E, Coates R, Mora Ardila F, Díaz Rojas A and Ramos-Fernández G (2019) Horizontal seed dispersal by dung beetles reduced seed and seedling clumping, but did not increase short-term seedling establishment. *PLoS ONE* **14**, e0224366.
- van Leeuwen CH, Villar N, Mendoza Sagrera I, Green AJ, Bakker ES, Soons MB, Galetti M, Jansen PA, Nolet BA and Santamaría L (2022) A seed dispersal effectiveness framework across the mutualism–antagonism continuum. *Oikos* **2022**, e09254.
- Welch D (1985) Studies in the grazing of heather moorland in north-east Scotland. IV. Seed dispersal and plant establishment in dung. *Journal of Applied Ecology* **22**, 461–472.
- Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer.
- Williams P and Haynes R (1995) Effect of sheep, deer and cattle dung on herbage production and soil nutrient content. *Grass and Forage Science* **50**, 263–271.