

RATIONALITY OF THE GENERALIZED BINOMIAL COEFFICIENTS FOR A MULTIPLICITY FREE ACTION

CHAL BENSON and GAIL RATCLIFF

(Received 14 October 1998; revised 19 November 1999)

Communicated by T. Dooley

Abstract

Let V be a finite dimensional Hermitian vector space and K be a compact Lie subgroup of $U(V)$ for which the representation of K on $\mathbb{C}[V]$ is multiplicity free. One obtains a canonical basis $\{p_\alpha\}$ for the space $\mathbb{C}[V_{\mathbb{R}}]^K$ of K -invariant polynomials on $V_{\mathbb{R}}$ and also a basis $\{q_\alpha\}$ via orthogonalization of the p_α 's. The polynomial p_α yields the homogeneous component of highest degree in q_α . The coefficients that express the q_α 's in terms of the p_β 's are the generalized binomial coefficients of Yan. The main result in this paper shows that these numbers are rational.

2000 *Mathematics subject classification*: primary 05E35, 20G05.

1. Introduction

Throughout this paper, V denotes a finite dimensional complex vector space with Hermitian inner product $\langle \cdot, \cdot \rangle$ and norm $|\cdot|$. K denotes a compact Lie group which acts linearly and unitarily on V . We write kz for the action of $k \in K$ on a vector $z \in V$. The associated representation of K on the space $\mathbb{C}[V]$ of holomorphic polynomials on V is given by the formula $(k \cdot p)(z) = p(k^{-1}z)$. The action of K on V is said to be a (linear) *multiplicity free action* when this representation of K on $\mathbb{C}[V]$ is multiplicity free. That is, no irreducible representation of K occurs more than once in $\mathbb{C}[V]$. We assume the action of K on V is multiplicity free throughout this paper. As shown in [1, Proposition 2.2], it follows that the action of the identity component of K on V is also multiplicity free. Thus we can assume that K is connected. The multiplicity

This work was supported in part by the University of Missouri Research Board.

© 2000 Australian Mathematical Society 0263-6115/2000 \$A2.00 + 0.00

free actions have been completely classified (see [5, 12, 13]). We make use of this classification below in Section 4 of this paper.

We write the decomposition of $\mathbb{C}[V]$ into K -irreducible subspaces as

$$(1.1) \quad \mathbb{C}[V] = \sum_{\alpha \in \Lambda} P_\alpha.$$

Here Λ denotes a countably infinite index set. In Section 2, we specify Λ concretely as a set of highest weights. Decomposition (1.1) is canonical because the action of K is multiplicity free. The subspace $\mathcal{P}_m(V) \subset \mathbb{C}[V]$ of homogeneous polynomials of degree m is K -invariant. Thus each irreducible P_α is contained in some $\mathcal{P}_m(V)$. We write $|\alpha|$ for the degree of homogeneity of P_α , so that $P_\alpha \subset \mathcal{P}_{|\alpha|}(V)$.

There are no non-constant K -invariant polynomials in $\mathbb{C}[V]$. Indeed, the trivial representation of K occurs only once in $\mathbb{C}[V]$. There are, however, non-constant K -invariant polynomials on the underlying real space $V_{\mathbb{R}}$ for V . These are non-holomorphic. The algebra $\mathbb{C}[V_{\mathbb{R}}]^K$ of such invariant polynomials has a natural vector space basis. To describe this, we introduce the Fock (or Fischer) inner product given on both $\mathbb{C}[V]$ and $\mathbb{C}[V_{\mathbb{R}}]$ by

$$(1.2) \quad (f, g)_{\mathcal{F}} = \left(\frac{1}{2\pi}\right)^n \int_V f(z) \overline{g(z)} e^{-|z|^2/2} dz.$$

Here $n = \dim_{\mathbb{C}}(V)$ and ‘ dz ’ denotes Lebesgue measure on $V_{\mathbb{R}} \cong \mathbb{R}^{2n}$. For $\alpha \in \Lambda$ let $d_\alpha = \dim(P_\alpha)$, choose an orthonormal basis $\{v_1, v_2, \dots, v_{d_\alpha}\}$ for P_α , and set

$$(1.3) \quad p_\alpha(z) = \frac{1}{d_\alpha} \sum_{j=1}^{d_\alpha} v_j(z) \overline{v_j(z)}.$$

It is not difficult to see that this definition of the p_α ’s does not depend on the choice of orthonormal basis for P_α and that $\{p_\alpha \mid \alpha \in \Lambda\}$ is a basis for the vector space $\mathbb{C}[V_{\mathbb{R}}]^K$ (see [2, Proposition 3.9]).

One obtains a second basis for $\mathbb{C}[V_{\mathbb{R}}]^K$ by orthogonalization of the p_α ’s. More precisely, we

- (1) choose any ordering on the set of indices Λ that ensures α precedes β if $|\alpha| < |\beta|$,
- (2) use the Fock inner product and this ordering to perform Gram-Schmidt orthogonalization on the sequence $\{p_\alpha \mid \alpha \in \Lambda\}$, and
- (3) normalize the resulting polynomials q_α so that $q_\alpha(0) = 1$.

In [2, Proposition 4.2] we show that the resulting basis $\{q_\alpha \mid \alpha \in \Lambda\}$ does not depend on the ordering chosen for the indices $\{\alpha \in \Lambda \mid |\alpha| = m\}$ that arise from the decomposition of $\mathcal{P}_m(V)$, $m \in \mathbb{Z}_+$. Our proof of this fact in [2] involves the representation theory of the Heisenberg group. We can suppress the Heisenberg

group and outline the key ideas as follows. For $z \in V$ and $f \in \mathbb{C}[V]$ let $\pi(z)f$ be the function on V given as

$$\pi(z)f(w) = e^{-(w,z)/2 - |z|^2/4} f(w + z)$$

and define functions q_α via

$$(1.4) \quad q_\alpha(z) = \frac{1}{d_\alpha} \sum_{j=1}^{d_\alpha} \langle \pi(z)v_j, v_j \rangle_{\mathcal{F}}.$$

It is shown in [2] that

- (1) q_α is a well defined K -invariant polynomial on $V_{\mathbb{R}}$ with $q_\alpha(0) = 1$,
- (2) the q_α 's are pair-wise orthogonal with respect to the Fock inner product, and
- (3) the homogeneous component of highest degree in q_α is $(-1)^{|\alpha|} p_\alpha$.

It follows from these facts that if one chooses any ordering for Λ as above then orthogonalization yields the polynomials q_α given by (1.4).

We now have two canonical bases $\{p_\alpha \mid \alpha \in \Lambda\}$ and $\{q_\alpha \mid \alpha \in \Lambda\}$ for $\mathbb{C}[V_{\mathbb{R}}]^K$. The p_α 's are homogeneous (of degree $2|\alpha|$) and the q_α 's are orthogonal. We write the q_α 's as linear combinations of the p_α 's:

$$(1.5) \quad q_\alpha = \sum_{|\beta| \leq |\alpha|} (-1)^{|\beta|} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} p_\beta.$$

Since the p_α 's and q_α 's are real valued functions, the coefficients $\begin{bmatrix} \alpha \\ \beta \end{bmatrix}$ are real. We call these the *generalized binomial coefficients* for the multiplicity free action of K on V . They are defined via (1.5) for $|\beta| \leq |\alpha|$. We extend the definition to all of $\Lambda \times \Lambda$ by setting $\begin{bmatrix} \alpha \\ \beta \end{bmatrix} = 0$ when $|\beta| > |\alpha|$.

The inclusion of the sign ' $(-1)^{|\beta|}$ ' in (1.5) is motivated by the fact that $(-1)^{|\alpha|} p_\alpha$ is the homogeneous component of highest degree in q_α . Thus $\begin{bmatrix} \alpha \\ \alpha \end{bmatrix} = 1$ and $\begin{bmatrix} \alpha \\ \beta \end{bmatrix} = 0$ for $\alpha \neq \beta$ with $|\alpha| = |\beta|$. Moreover, Lemma 3.9 in [6] shows that, with this convention, the generalized binomial coefficients are non-negative. The main result in this paper asserts that these are, moreover, rational numbers.

THEOREM 1.1. $\begin{bmatrix} \alpha \\ \beta \end{bmatrix}$ is a non-negative rational number for all $\alpha, \beta \in \Lambda$.

The generalized binomial coefficients were introduced in the setting of multiplicity free actions by Yan in his unpublished work [17]. The simplest example of a multiplicity free action is given by the standard action of $K = U(n)$ on $V = \mathbb{C}^n$. In this case $\mathbb{C}[V]$ decomposes as $\mathbb{C}[V] = \sum_m \mathcal{P}_m(V)$ and one can show (see [17] or [6]) that $\begin{bmatrix} m \\ j \end{bmatrix} = \binom{m}{j}$, independent of n . This motivates the terminology. For multiplicity free actions that arise from Hermitian symmetric spaces, Yan has shown that the generalized binomial coefficients as defined above agree with those introduced by Herz, Dib

and Faraut-Koranyi (see [8, 9, 18]). Our paper [6] shows that the generalized binomial coefficients appear in the solutions to a variety of combinatorial and analytic problems that arise in connection with multiplicity free actions and related Gelfand pairs associated with the Heisenberg group. We indicate one such interconnection. From p_β , one obtains a Wick ordered polynomial coefficient differential operator ' $p_\beta(z, \partial/\partial z)$ '. This is K -invariant and hence scalar on each subspace P_α . The eigenvalue for this operator on P_α is $\begin{bmatrix} \alpha \\ \beta \end{bmatrix} / (2^{|\beta|} d_\beta)$. This relates the computation of generalized binomial coefficients to that of eigenvalues for K -invariant polynomial coefficient differential operators, a problem that has been studied recently in [14–16]. Theorem 1.1 shows that these eigenvalues are non-negative rational numbers.

The rest of this paper is structured as follows. In Section 2 we show that the basis $\{p_\alpha \mid \alpha \in \Lambda\}$ contains a canonical subset $\{\gamma_1, \dots, \gamma_r\}$ of 'fundamental invariants' that freely generate $\mathbb{C}[V_{\mathbb{R}}]^K$ as an algebra. We prove that one can replace the p_α 's by polynomials of the form $\gamma_1^{a_1} \cdots \gamma_r^{a_r}$ in the orthogonalization procedure used to determine the q_α 's. For this, we need to use a specific refinement of the partial ordering by degrees that is compatible with the weight ordering used to determine the fundamental invariants. This is the content of Theorem 2.1 below. In Section 3, we reduce the proof of Theorem 1.1 to the assertion that one can find some orthonormal basis for V so that each fundamental invariant γ_i is given by a polynomial with rational coefficients in the coordinates with respect to the basis. The proof uses Theorem 2.1 together with some results concerning generalized binomial coefficients from [6]. In Section 4 we complete the proof of Theorem 1.1 by showing the existence of such a basis. We do this via case-by-case analysis using a classification of multiplicity free actions and exhibiting rational fundamental highest weight vectors. This requires rather explicit knowledge of the decomposition for $\mathbb{C}[V]$ in each case. We believe that the details of this case-by-case analysis, which extends work of Howe and Umeda from [11], are of independent interest.

2. Fundamental invariants

Let T be a maximal torus in K and $G = K_{\mathbb{C}}$, $H = T_{\mathbb{C}}$ be the complexified groups with Lie algebras \mathfrak{g} and \mathfrak{h} respectively. Choose a positive system $\Delta^+ = \Delta^+(\mathfrak{g}, \mathfrak{h})$ of roots. We recall that these choices produce a simple ordering on the weights $\lambda \in \mathfrak{h}^* = \text{hom}(\mathfrak{h}, \mathbb{C})$ for any representation of K . We denote this weight ordering by $<$. Let $\Lambda \subset \mathfrak{h}^*$ denote the set of highest weights for the irreducible representations of K which occur in $\mathbb{C}[V]$. Thus Λ is the index set for decomposition (1.1), and the irreducible component P_α has highest weight $\alpha \in \Lambda$.

If $h_\alpha, h_\beta \in \mathbb{C}[V]$ are α - and β -highest weight vectors, then $h_\alpha h_\beta$ is an $(\alpha + \beta)$ -highest weight vector and thus $\Lambda \subset \mathfrak{h}^*$ is an additive semigroup. Following [11] we

call the primitive elements of Λ *fundamental highest weights*. These are the elements of Λ that cannot be expressed as sums $\alpha + \beta$ with $\alpha, \beta \in \Lambda$. We see that if h_α is a prime polynomial, then α is a fundamental highest weight. As explained in [11], the converse is also true. The fundamental highest weights are finite in number and freely generate Λ . Let

$$(2.1) \quad \{\alpha_1, \alpha_2, \dots, \alpha_r\}$$

be the fundamental highest weights listed in increasing order using $<$. We then have

$$\Lambda = \{a_1\alpha_1 + \dots + a_r\alpha_r \mid a_1, \dots, a_r \in \mathbb{Z}_+\},$$

where \mathbb{Z}_+ denotes the non-negative integers. The fundamental highest weights thus establish a semigroup isomorphism

$$(\mathbb{Z}_+)^r \cong \Lambda, \quad (a_1, \dots, a_r) \mapsto a_1\alpha_1 + \dots + a_r\alpha_r.$$

This correspondence is canonical having chosen the data T and Δ^+ . We sometimes write elements $\alpha \in \Lambda$ as ' $\alpha = (a_1, \dots, a_r)$ ' to mean $\alpha = a_1\alpha_1 + \dots + a_r\alpha_r$. In this notation we have $\alpha_1 = (1, 0, \dots, 0)$, $\alpha_2 = (0, 1, 0, \dots, 0)$ and so on.

DEFINITION 2.1. The *fundamental invariants* $\gamma_1, \dots, \gamma_r$ are defined as $\gamma_j = p_{\alpha_j}$ where as above $\alpha_1, \dots, \alpha_r \in \Lambda$ are the fundamental highest weights.

For $\alpha = (a_1, \dots, a_r) \in \Lambda$ let

$$\gamma_\alpha = \gamma_1^{a_1} \gamma_2^{a_2} \dots \gamma_r^{a_r}.$$

Note that γ_α is K -invariant and homogeneous of degree $2|\alpha| = \deg(p_\alpha)$. We know that the q_α 's can be obtained by orthogonalization of the p_α 's using any ordering compatible with degree. We show that the q_α 's can also be obtained by orthogonalization of the γ_α 's. For this, we need to use the weight ordering $<$ on Λ to refine the partial ordering by degrees.

THEOREM 2.1. *The polynomials $\{q_\alpha \mid \alpha \in \Lambda\}$ are obtained from the polynomials $\{\gamma_\alpha \mid \alpha \in \Lambda\}$ by using the Fock inner product to perform Gram-Schmidt orthogonalization and normalizing so that $q_\alpha(0) = 1$. Here we impose the ordering $<$ on Λ defined by*

$$\alpha < \beta \Leftrightarrow [(|\alpha| < |\beta|) \text{ or } (|\alpha| = |\beta| \text{ and } \alpha < \beta)].$$

We remark that, from a practical viewpoint, Theorem 2.1 substantially improves the orthogonalization result discussed in Section 1. In many examples, it is much easier to obtain explicit formulae for the fundamental invariants, which are finite in number, than it is to produce formulae for all of the p_α 's. In such cases, the orthogonalization procedure in Theorem 2.1 is concrete in the sense that the sequence of γ_α 's is explicit. We provide formulae for the fundamental invariants for many of the examples discussed below in Section 4. In [3], Theorem 2.1 was obtained in the context of one specific multiplicity free action. This is Example 4.1.2 below.

PROOF OF THEOREM 2.1. Given $\lambda \in \mathfrak{h}^*$, we write W_λ for the λ -weight space in $\mathbb{C}[V]$, $W_\lambda = \{f \in \mathbb{C}[V] \mid X \cdot f = \lambda(X)f \ \forall X \in \mathfrak{h}\}$ in $\mathbb{C}[V]$. Choose a highest weight vector $h_j \in P_{\alpha_j}$ for each of the fundamental highest weights $\alpha_1, \dots, \alpha_r$. The h_j 's are unique up to non-zero scalar multiples. We normalize to ensure $\langle h_j, \bar{h}_j \rangle_{\mathfrak{F}} = 1$. Since P_{α_j} is an orthogonal direct sum of its weight spaces, we can write the fundamental invariant $\gamma_j = p_{\alpha_j}$ in the form

$$(2.2) \quad \gamma_j = \frac{1}{d_{\alpha_j}} h_j \bar{h}_j + r_j,$$

where $r_j \in \sum_{\lambda < \alpha_j} W_\lambda \otimes \bar{W}_\lambda$. For $\alpha = (a_1, \dots, a_r) \in \Lambda$ we let $h_\alpha = h_1^{a_1} h_2^{a_2} \dots h_r^{a_r}$. This is a highest weight vector in P_α . From (2.2) we see that

$$(2.3) \quad \gamma_\alpha = \frac{1}{d_{\alpha_1}^{a_1} \dots d_{\alpha_r}^{a_r}} h_\alpha \bar{h}_\alpha + r_\alpha,$$

where $r_\alpha \in \sum_{\lambda < \alpha} W_\lambda \otimes \bar{W}_\lambda$. Similarly, we see that p_α can also be written in the form

$$(2.4) \quad p_\alpha = \frac{b_\alpha}{d_\alpha} h_\alpha \bar{h}_\alpha + r'_\alpha,$$

where $b_\alpha = 1/\|h_\alpha\|^2$ is a positive constant and $r'_\alpha \in \sum_{\lambda < \alpha} W_\lambda \otimes \bar{W}_\lambda$.

Comparing (2.3) and (2.4), we see that

$$(2.5) \quad p_\alpha = c_\alpha \gamma_\alpha + s_\alpha,$$

where c_α is a positive constant and $s_\alpha \in \sum_{\lambda < \alpha} W_\lambda \otimes \bar{W}_\lambda$. As both p_α and γ_α are K -invariant, so is s_α . Since $\{p_\alpha \mid \alpha \in \Lambda\}$ is a basis for $\mathbb{C}[V_{\mathbb{R}}]^K$, we have that $s_\alpha \in \text{Span}(p_\beta \mid \beta \in \Lambda, \beta < \alpha)$. As both p_α and γ_α are homogeneous of degree $2|\alpha|$, equation (2.5) also shows that s_α is homogeneous of degree $2|\alpha|$, provided $s_\alpha \neq 0$. Thus we must have

$$s_\alpha \in \text{Span}(p_\beta \mid |\beta| = |\alpha| \text{ and } \beta < \alpha) \subset \text{Span}(p_\beta \mid \beta < \alpha).$$

This fact together with (2.5) implies that application of Gram-Schmidt orthogonalization to the sequences $\{p_\alpha \mid \alpha \in \Lambda\}$ and $\{\gamma_\alpha \mid \alpha \in \Lambda\}$ ordered via $<$ yield the same result. Since we know that orthogonalization of the p_α 's yields the q_α 's, this completes the proof. \square

REMARK 2.1. We note that the proof of Theorem 2.1 shows that for fixed m and $\alpha = \min\{\beta \in \Lambda \mid |\beta| = m\}$ one has

$$p_\alpha = c_\alpha \gamma_\alpha,$$

where $c_\alpha = (d_{\alpha_1}^{a_1} \cdots d_{\alpha_r}^{a_r}) / (d_\alpha \|h_\alpha\|^2)$.

3. Proof of Theorem 1.1

Suppose that $\{e_1, \dots, e_n\}$ is an orthonormal basis for the Hermitian vector space V . We obtain isomorphisms $V \cong \mathbb{C}^n$ and

$$\mathbb{C}[V] \cong \mathbb{C}[z_1, \dots, z_n], \quad \mathbb{C}[V_{\mathbb{R}}] \cong \mathbb{C}[z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n],$$

where z_1, \dots, z_n are the coordinates with respect to the basis. We write $\mathbb{Q}[V]$ and $\mathbb{Q}[V_{\mathbb{R}}]$ for the \mathbb{Q} -subalgebras of $\mathbb{C}[V]$ and $\mathbb{C}[V_{\mathbb{R}}]$ that correspond to $\mathbb{Q}[z_1, \dots, z_n]$ and $\mathbb{Q}[z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n]$ under these identifications. Our notation conceals the fact that $\mathbb{Q}[V]$ and $\mathbb{Q}[V_{\mathbb{R}}]$ depend on the orthonormal basis used. In Section 4 we prove the following result.

LEMMA 3.1. *There is an orthonormal basis for V in which $\gamma_1, \dots, \gamma_r \in \mathbb{Q}[V_{\mathbb{R}}]$.*

This result is a key ingredient in our proof of Theorem 1.1. We assume Lemma 3.1 here and complete the proof of Theorem 1.1. Fix an orthonormal basis for V so that the fundamental invariants are rational polynomials as in Lemma 3.1.

Next we recall that the monomials $z^I := z_1^{i_1} \cdots z_n^{i_n}$ in $\mathbb{C}[V] \cong \mathbb{C}[z_1, \dots, z_n]$ are orthogonal with respect to $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ and satisfy $\langle z^I, z^I \rangle_{\mathcal{F}} = 2^I I! := 2^{i_1 + \cdots + i_n} i_1! \cdots i_n!$. (See for example [10, Theorem 1.63], although this reference uses a different normalization convention for the Fock inner product.) Thus we also have

$$\langle z^I \bar{z}^J, z^{I'} \bar{z}^{J'} \rangle_{\mathcal{F}} = \begin{cases} 2^{I+J'} (I+J')! & \text{if } I+J' = I'+J; \\ 0 & \text{otherwise.} \end{cases}$$

In particular, the inner product of any two monomials in $\mathbb{C}[V_{\mathbb{R}}]$ is integral. We conclude immediately that

LEMMA 3.2. *$\langle f, g \rangle_{\mathcal{F}} \in \mathbb{Q}$ for any $f, g \in \mathbb{Q}[V_{\mathbb{R}}]$.*

LEMMA 3.3. $q_\alpha \in \mathbb{Q}[V_{\mathbb{R}}]$ for all $\alpha \in \Lambda$.

PROOF. We use the ordering specified in Theorem 2.1 to perform Gram-Schmidt orthogonalization with the sequence of polynomials $\{\gamma_\alpha \mid \alpha \in \Lambda\}$. This yields an (unnormalized) sequence of orthogonal polynomials \tilde{q}_α with $\tilde{q}_0 = 1$ and

$$\tilde{q}_\alpha = \gamma_\alpha - \sum_{\beta < \alpha} \frac{\langle \gamma_\alpha, \tilde{q}_\beta \rangle_{\mathcal{F}}}{\langle \tilde{q}_\beta, \tilde{q}_\beta \rangle_{\mathcal{F}}} \tilde{q}_\beta$$

for $\alpha > 0$. Here $0 = (0, \dots, 0) \in \Lambda$ is the index for which $P_0 = \mathbb{C}$, the scalar polynomials. We have that $\tilde{q}_0 \in \mathbb{Q}[V_{\mathbb{R}}]$. Assume inductively that $\alpha > 0$ and that $\tilde{q}_\beta \in \mathbb{Q}[V_{\mathbb{R}}]$ for $\beta < \alpha$. Lemma 3.1 shows that $\gamma_\alpha \in \mathbb{Q}[V_{\mathbb{R}}]$ and Lemma 3.2 ensures that the coefficients in the expression for \tilde{q}_α are all rational. Thus $\tilde{q}_\alpha \in \mathbb{Q}[V_{\mathbb{R}}]$ for all $\alpha \in \Lambda$. A second induction on α shows that $\tilde{q}_\alpha(0) \in \mathbb{Q}$. Thus also $q_\alpha = \tilde{q}_\alpha / \tilde{q}_\alpha(0) \in \mathbb{Q}[V_{\mathbb{R}}]$. □

We can now complete the proof of Theorem 1.1 by using some combinatorial identities from [6]. Proposition 3.7 in that paper asserts that

$$\frac{|z|^2}{2} q_\beta = - \sum_{|\alpha|=|\beta|+1} \frac{d_\alpha}{d_\beta} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} q_\alpha + (2|\beta| + n)q_\beta - \sum_{|\alpha|=|\beta|-1} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} q_\alpha.$$

We know, moreover, that the q_α 's are orthogonal and the norms are given by $\langle q_\alpha, q_\alpha \rangle_{\mathcal{F}} = 1/d_\alpha$. (See [4, 17].) Thus, for $|\alpha| = |\beta| + 1$ we have

$$\left\langle \frac{|z|^2}{2} q_\beta, q_\alpha \right\rangle_{\mathcal{F}} = -\frac{1}{d_\beta} \begin{bmatrix} \alpha \\ \beta \end{bmatrix}.$$

Lemma 3.3 shows that q_α and $|z|^2 q_\beta / 2$ belong to $\mathbb{Q}[V_{\mathbb{R}}]$ and hence the left hand side of the last equation is rational by Lemma 3.2. It follows that $\begin{bmatrix} \alpha \\ \beta \end{bmatrix} \in \mathbb{Q}$ whenever $|\alpha| = |\beta| + 1$. Suppose more generally that $|\alpha| = |\beta| + k$. Equation (3.9) in [6] reads:

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \frac{1}{k!} \sum \begin{bmatrix} \varepsilon_1 \\ \beta \end{bmatrix} \begin{bmatrix} \varepsilon_2 \\ \varepsilon_1 \end{bmatrix} \cdots \begin{bmatrix} \varepsilon_{k-1} \\ \varepsilon_{k-2} \end{bmatrix} \begin{bmatrix} \alpha \\ \varepsilon_{k-1} \end{bmatrix},$$

where the sum is over all $(\varepsilon_1, \dots, \varepsilon_{k-1})$ with $|\varepsilon_j| = |\beta| + j$. Since all the generalized binomial coefficients appearing on the right hand side of this equation are rational, we conclude that $\begin{bmatrix} \alpha \\ \beta \end{bmatrix} \in \mathbb{Q}$.

4. Case-by-case analysis

In this section we prove Lemma 3.1. This involves case-by-case analysis working from the classification of multiplicity free actions contained in [5]. We begin with a

simple lemma that enables a substantial reduction in the work required for each case. First note that the concept of multiplicity free action depends only on the image of K in $U(V)$ under its representation on V . Thus we can regard K as a subgroup of $U(V)$ acting on V in the usual fashion. Recall that we use an orthonormal basis of V to write polynomials in coordinates and identify rational subalgebras $\mathbb{Q}[V]$ and $\mathbb{Q}[V_{\mathbb{R}}]$ of $\mathbb{C}[V]$ and $\mathbb{C}[V_{\mathbb{R}}]$. Such a basis can also be used to realize $U(V)$ and $GL(V)$ as the matrix groups $U(n)$ and $GL(n, \mathbb{C})$. The group K and its complexification $G = K_{\mathbb{C}}$ become subgroups of $GL(n, \mathbb{C})$ acting on \mathbb{C}^n in the standard fashion. The Lie algebra \mathfrak{g} of G becomes a subalgebra of the Lie algebra $gl(n, \mathbb{C})$ of $n \times n$ matrices. Let $\mathfrak{g}_{\mathbb{Q}} = \mathfrak{g} \cap gl(n, \mathbb{Q})$. This is a Lie algebra over \mathbb{Q} . One says that $\mathfrak{g}_{\mathbb{Q}}$ is a *rational form* for \mathfrak{g} if $\mathfrak{g} = \mathbb{C} \otimes \mathfrak{g}_{\mathbb{Q}}$. Equivalently, \mathfrak{g} has a basis (over \mathbb{C}) that is contained in the subset $\mathfrak{g}_{\mathbb{Q}}$.

LEMMA 4.1. *Suppose that $\mathfrak{g}_{\mathbb{Q}}$ is a rational form for \mathfrak{g} . Let $\alpha \in \Lambda$ and suppose that $P_{\alpha} \cap \mathbb{Q}[V] \neq \{0\}$. Then $p_{\alpha} \in \mathbb{Q}[V_{\mathbb{R}}]$.*

PROOF. Let $h \in P_{\alpha} \cap \mathbb{Q}[V]$ with $h \neq 0$. Since P_{α} is K -irreducible we have $P_{\alpha} = \mathcal{U}(\mathfrak{g})h$. Thus any basis $\{f_1, \dots, f_m\}$ for P_{α} ($m = d_{\alpha}$) can be written as

$$f_1 = D_1h, f_2 = D_2h, \dots, f_m = D_mh$$

for some $D_1, \dots, D_m \in \mathcal{U}(\mathfrak{g})$. Since $\mathcal{U}(\mathfrak{g}) = \mathcal{U}(\mathbb{C} \otimes \mathfrak{g}_{\mathbb{Q}}) = \mathbb{C} \otimes \mathcal{U}(\mathfrak{g}_{\mathbb{Q}})$, we can write

$$D_j = c_{j,1}D_{j,1} + \dots + c_{j,\ell_j}D_{j,\ell_j}$$

for some $c_{j,1}, \dots, c_{j,\ell_j} \in \mathbb{C}$ and $D_{j,1}, \dots, D_{j,\ell_j} \in \mathcal{U}(\mathfrak{g}_{\mathbb{Q}})$. Thus

$$\begin{aligned} P_{\alpha} &= \mathbb{C} - \text{Span}(f_1, \dots, f_m) \\ &= \mathbb{C} - \text{Span}(D_{1,1}h, \dots, D_{1,\ell_1}h, D_{2,1}h, \dots, D_{2,\ell_2}h, \dots, D_{m,1}h, \dots, D_{m,\ell_m}h) \end{aligned}$$

and we conclude that P_{α} has a basis consisting of vectors of the form $D_{i,j}h$. Let $\{g_1, \dots, g_m\}$ denote such a basis. Since $D_{i,j} \in \mathcal{U}(\mathfrak{g}_{\mathbb{Q}}) \subset \mathcal{U}(gl(n, \mathbb{Q}))$ and $gl(n, \mathbb{Q})$ preserves $\mathbb{Q}[V] = \mathbb{Q}[z_1, \dots, z_n]$, we have that $g_1, \dots, g_m \in \mathbb{Q}[V]$.

Next we orthogonalize the basis $\{g_1, \dots, g_m\}$ to obtain a basis $\{u_1, \dots, u_m\}$ defined as $u_1 = g_1$ and

$$u_j = g_j - \sum_{i=1}^{j-1} \frac{\langle g_j, u_i \rangle_{\mathcal{F}}}{\langle u_i, u_i \rangle_{\mathcal{F}}} u_i$$

for $i = 2, \dots, m$. Since the g_j 's belong to $\mathbb{Q}[V]$, we can use Lemma 3.2 and induction, as in the proof for Lemma 3.3, to conclude that $u_j \in \mathbb{Q}[V]$. Since $\{u_1, \dots, u_m\}$ is an

orthogonal basis for P_α , we can use $v_j = u_j / \|u_j\|$ in (1.3) to write

$$p_\alpha(z) = \frac{1}{m} \sum_{j=1}^m \frac{u_j(z)\overline{u_j(z)}}{\langle u_j, u_j \rangle_{\mathcal{F}}}$$

Here $\langle u_j, u_j \rangle_{\mathcal{F}} \in \mathbb{Q}$ since $u_j \in \mathbb{Q}[V]$. Thus we see that $p_\alpha \in \mathbb{Q}[V]$. □

To prove Lemma 3.1, it now suffices to show that for any multiplicity free action:

- (1) there is an orthonormal basis for V for which $\mathfrak{g}_{\mathbb{Q}} = \mathfrak{g} \cap \mathfrak{gl}(n, \mathbb{Q})$ is a rational form for \mathfrak{g} , and
- (2) one can find fundamental highest weight vectors $h_j \in P_\alpha$, ($j = 1, \dots, r$) that are rational polynomials in the coordinates with respect to this basis. That is, $h_j \in \mathbb{Q}[V]$ for $j = 1, \dots, r$.

This completes the proof. Indeed, Lemma 4.1 shows that using such a basis we have $\gamma_1, \dots, \gamma_r \in \mathbb{Q}[V_{\mathbb{R}}]$.

The group $G \subset GL(V)$ is connected reductive and complex algebraic. Decompose V as an orthogonal direct sum of G -irreducible subspaces:

$$(4.1) \quad V = V_1 \oplus \dots \oplus V_m.$$

For our purposes, we can replace G by $\tilde{G} = G' \times (\mathbb{C}^\times)^m$ where G' denotes the commutator subgroup and we have one copy of the scalars \mathbb{C}^\times acting on each subspace V_j . Indeed, the decompositions of $\mathbb{C}[V]$ under the actions of G and \tilde{G} coincide and hence the p_α 's, q_α 's and generalized binomial coefficients for the two actions are the same. We can assume, moreover, that the action of the semisimple group G' on V is *indecomposable*. This means that we can't write V as a direct sum $V = W_1 \oplus W_2$ and G' as a product $G' = G'_1 \times G'_2$ with G'_j acting independently on W_j . Indeed, suppose that given any such indecomposable action one can find an orthonormal basis with respect to which the fundamental invariants are rational polynomials. If the action of G' decomposes as a product of indecomposable actions of subgroups G'_j on subspaces W_j then one obtains an orthonormal basis for V that meets our requirements by concatenation of appropriately chosen bases for the W_j 's. In particular, these bases produce obvious inclusions $\mathbb{Q}[(W_j)_{\mathbb{R}}] \subset \mathbb{Q}[V_{\mathbb{R}}]$ and the fundamental invariants $\gamma_{j,i} \in \mathbb{Q}[(W_j)_{\mathbb{R}}]$ for the actions of each $G'_j \times (\mathbb{C}^\times)^{m_j}$ on W_j just combine to yield the (rational) fundamental invariants for the action of \tilde{G} .

We can now simplify the notation from the preceding paragraph and restrict our attention to actions of the following sort.

- $G \subset GL(V)$ is a connected semisimple complex algebraic group acting indecomposably on V .
- $V = V_1 \oplus \dots \oplus V_m$, where each V_j is G -irreducible.
- The joint action of $G \times (\mathbb{C}^\times)^m$ on V is multiplicity free.

TABLE 1.

Group	Degrees of fundamental highest weight vectors (number)	Example number
SL(n) ($n \geq 1$)	1 (1)	4.1.1
SO(n) ($n \geq 3$)	1, 2 (2)	4.1.2
Sp(2n) ($n \geq 2$)	1 (1)	4.1.1
S²(SL(n)) ($n \geq 2$)	1, 2, ..., n (n)	4.1.4
Λ^2(SL(n)) ($n \geq 4$)	1, 2, ..., $\lfloor n/2 \rfloor$ ($\lfloor n/2 \rfloor$)	4.1.5
SL(n) \otimes SL(m) ($n, m \geq 2$)	1, 2, ..., $\min(n, m)$ ($\min(n, m)$)	4.1.3
Sp(2n) \otimes SL(2) ($n \geq 2$)	1, 2, 2 (3)	4.1.6
Sp(2n) \otimes SL(3) ($n \geq 2$)	1, 2, 2, 3, 3, 4 (6)	4.1.7
Sp(4) \otimes SL(n) ($n \geq 4$)	1, 2, 2, 3, 4, 4 (6)	4.1.8
Spin(7)	1, 2 (2)	4.1.9
Spin(9)	1, 2, 2 (3)	4.1.10
Spin(10)	1, 2 (2)	4.1.11
G₂	1, 2 (2)	4.1.2
E₆	1, 2, 3 (3)	4.1.12

Such actions have been completely classified. In all cases, one has either $m = 1$, so that the action of G on V is irreducible, or $m = 2$. The irreducible multiplicity free actions were classified by Kac in [12]. The indecomposable non-irreducible actions were classified by the authors in [5] and independently by Leahy in [13]. Below, we examine each possibility in turn to complete the proof of Lemma 3.1.

4.1. Irreducible multiplicity free actions Table 1, taken from [12], lists all possibilities for semisimple groups $G \subset GL(V)$ acting irreducibly on V for which the action of $G \times (\mathbb{C}^\times)$ is multiplicity free. The notation, adopted from [12], indicates G as the image of a group under some irreducible representation on some vector space V . For example, $\Lambda^2(\mathbf{SL}(n))$ indicates the image of $SL(n, \mathbb{C})$ in $GL(\Lambda^2(\mathbb{C}^n))$ and $\mathbf{Sp}(2n) \otimes \mathbf{SL}(3)$ indicates the image of $Sp(2n, \mathbb{C}) \times SL(3, \mathbb{C})$ in $GL(\mathbb{C}^{2n} \otimes \mathbb{C}^3)$ under the obvious representations.

For all of the groups $G \subset GL(V)$ in Table 1, the vector space V has a standard basis that is orthonormal for a natural Hermitian inner product that determines the compact real form $K \subset U(V)$. For the classical groups, these are the natural bases for \mathbb{C}^n , $\Lambda^2(\mathbb{C}^n)$, $S^2(\mathbb{C}^n)$ and $\mathbb{C}^n \otimes \mathbb{C}^m$. Using these bases to realize G as a matrix group $G \subset GL(n, \mathbb{C})$, one sees easily that $\mathfrak{g}_{\mathbb{Q}} = \mathfrak{g} \cap gl(n, \mathbb{Q})$ is a rational form for \mathfrak{g} in each case. Thus, it remains to show that one can find fundamental highest weight vectors that are rational polynomials in the coordinates with respect to these natural bases. We do this below in each case. The fundamental highest weight vectors for

most of these actions were given explicitly by Howe and Umeda in [11]. In all cases, the number of fundamental highest weight vectors and their degrees (as homogeneous polynomials in $\mathbb{C}[V]$) are in [11]. We have incorporated this information into Table 1. In many cases, we are able to present explicit formulae for the fundamental invariants, at least up to scalar multiples. We follow the notational conventions in [11], to which we refer the reader for further details regarding the decomposition of $\mathbb{C}[V]$ for each of these examples.

4.1.1. $SL(n), Sp(2n)$ Here $G = SL(n, \mathbb{C})$ and $G = Sp(2n, \mathbb{C})$ act on $V = \mathbb{C}^n$ and $V = \mathbb{C}^{2n}$ by their defining representations. The G -irreducible subspaces of $\mathbb{C}[V]$ are the spaces $\mathcal{P}_m(V)$ of homogeneous polynomials of each fixed degree m . There is a single fundamental highest weight vector, $z_1 \in \mathcal{P}_1(V)$ and the associated fundamental invariant is $\gamma(z) = |z|^2/2$. As we have already noted, the generalized binomial coefficients for these examples are the usual binomial coefficients, motivating our terminology.

4.1.2. $SO(n), G_2$ Let $\varepsilon(z) = z_1^2 + \dots + z_n^2$ be the $SO(n, \mathbb{C})$ -invariant polynomial on $V = \mathbb{C}^n$ and let

$$D_\varepsilon = \left(\frac{\partial}{\partial z_1}\right)^2 + \dots + \left(\frac{\partial}{\partial z_n}\right)^2.$$

Then $\mathcal{H} = \{p \in \mathbb{C}[V] \mid D_\varepsilon p = 0\}$ is the space of ‘harmonics’, with $\mathcal{H} = \sum_{m=0}^\infty \mathcal{H}_m$, $\mathcal{H}_m = \mathcal{H} \cap \mathcal{P}_m(V)$. The decomposition of $\mathbb{C}[V]$ into $SO(n, \mathbb{C}) \times \mathbb{C}^\times$ -irreducibles reads:

$$\mathbb{C}[V] = \sum_{m,\ell} \mathcal{H}_m \varepsilon^\ell.$$

There are two fundamental highest weight vectors in $\mathbb{C}[V]$, z_1 and $\varepsilon(z)$. The associated fundamental invariants are $\gamma_1(z) = |z|^2/2$ and $\gamma_2(z) = |\varepsilon(z)|^2/4n$. For further details on this example, we refer the reader to [3].

The exceptional group G_2 acts on $V = \mathbb{C}^7$ as a subgroup of $SO(7, \mathbb{C})$. The subspaces $\mathcal{H}_m \varepsilon^\ell$ in $\mathbb{C}[V]$ are irreducible for the action of $G_2 \times \mathbb{C}^\times$. Thus the decomposition, fundamental highest weight vectors and invariants for this example are the same as those for $SO(7)$.

In the following examples, V is $\mathbb{C}^n \otimes \mathbb{C}^m$, $\Lambda^2(\mathbb{C}^n)$, $S^2(\mathbb{C}^n)$ or $\mathbb{C}^n \oplus \mathbb{C}^n$. We regard V as the space of $n \times m$ matrices, skew-symmetric $n \times n$ matrices, symmetric $n \times n$ matrices or $n \times 2$ matrices respectively. We write $z = (z_{ij})$ for the coordinates of $z \in V$ with respect to the standard basis for V and use the notation

$$\det_k(z) = \det \begin{pmatrix} z_{11} & \cdots & z_{1k} \\ \vdots & \ddots & \vdots \\ z_{k1} & \cdots & z_{kk} \end{pmatrix}.$$

4.1.3. $SL(n) \otimes SL(m)$ The decomposition of $\mathbb{C}[C^n \otimes C^m]$ under the action of $SL(n, \mathbb{C}) \times SL(m, \mathbb{C}) \times C^\times$ is given by

$$\mathbb{C}[C^n \otimes C^m] = \sum_D \rho_D^n \otimes \rho_D^m.$$

Here the sum is taken over all Young’s diagrams with at most $\min(n, m)$ rows and ρ_D^n, ρ_D^m are the representations of $SL(n, \mathbb{C})$ and $SL(m, \mathbb{C})$ corresponding to D . The fundamental highest weight vectors are

$$\det_k(z) \quad \text{for } k = 1, \dots, \min(n, m).$$

The fundamental invariants can be written up to normalization as

$$\gamma_k(z) = \sum_{|I|=|J|=k} |\det_{I,J}(z)|^2.$$

Here I, J denote subsets of $\{1, \dots, n\}$ and $\{1, \dots, m\}$ and $\det_{I,J}(z)$ is the subdeterminant of z obtained from rows I and columns J .

4.1.4. $S^2(SL(n))$ Here $\mathbb{C}[S^2(C^n)]$ decomposes under the action of $SL(n, \mathbb{C}) \times C^\times$ as $\mathbb{C}[S^2(C^n)] = \sum_D \rho_D^n$. The sum is over all Young’s diagrams D with at most n rows, all of even length. The fundamental highest weight vectors are $\det_k(z)$ for $k = 1, \dots, n$.

4.1.5. $\Lambda^2(SL(n))$ The decomposition is $\mathbb{C}[\Lambda^2(C^n)] = \sum_D \rho_D^n$, where the sum is over all Young’s diagrams D with at most n rows and each column is of even length. The fundamental highest weight vectors are $\zeta_k(z)$ for $k = 1, \dots, \lfloor n/2 \rfloor$, where $\zeta_k^2(z) = \det_{2k}(z)$. That is, $\zeta_k(z)$ is the Pfaffian of the skew symmetric matrix given by the first $2k$ rows and columns of z . This is written explicitly in [11] as

$$\zeta_k(z) = \sum_{\sigma \in S_{2k}/B_k} \text{sign}(\sigma) z_{\sigma(1)\sigma(2)} \cdots z_{\sigma(2k-1)\sigma(2k)},$$

where B_k is the subgroup of the symmetric group S_{2k} which preserves the pairs $\{\{1, 2\}, \dots, \{2k - 1, 2k\}\}$. In particular, we see that $\zeta_k(z)$ is a rational polynomial in the entries of z . Let $I = \{i_1, \dots, i_{2k}\} \subset \{1, \dots, n\}$ and let ζ_I be the Pfaffian computed with rows I and columns I . Up to normalization, the fundamental invariants are $\gamma_k(z) = \sum_{|I|=2k} \zeta_I(z)\zeta_I(\bar{z})$.

4.1.6. $Sp(2n) \otimes SL(2)$ Let $v_i \in C^{2n}$ be the columns of $z \in V = C^{2n} \otimes C^2, i = 1, 2$. The symplectic product $\varepsilon(z) = \omega(v_1, v_2)$ of the columns of z is an $Sp(2n, \mathbb{C})$ -invariant. Let $D_\varepsilon = \varepsilon(\partial/\partial z)$, and $\mathcal{H} = \{p \in \mathbb{C}[V] \mid D_\varepsilon p = 0\}$. We have

$$\mathbb{C}[V] = \sum_{\substack{\alpha_1 \geq \alpha_2 \geq 0 \\ \ell \geq 0}} \mathcal{H}_{\alpha_1, \alpha_2} \varepsilon^\ell,$$

where the irreducible subspace $\mathcal{H}_{\alpha_1, \alpha_2} \varepsilon^\ell$ has highest weight vector $z_{11}^{\alpha_1 - \alpha_2} \det_2(z)^{\alpha_2} \varepsilon^\ell$. Thus the fundamental highest weight vectors are z_{11} , $\det_2(z)$ and $\varepsilon(z)$ and we have fundamental invariants $\gamma_1(z) = |z|^2$ and $\gamma_2(z) = |\varepsilon(z)|^2$, modulo normalizations. In order to compute the third fundamental invariant, we look at the decomposition of $\mathcal{P}_2(V)$. Under the action of $GL(2n, \mathbb{C}) \times GL(2, \mathbb{C})$ we have

$$(4.2) \quad \mathcal{P}_2(V) = (\rho_{(2)}^{2n} \otimes \rho_{(2)}^2) \oplus (\rho_{(1,1)}^{2n} \otimes \rho_{(1,1)}^2).$$

When we restrict to $Sp(2n, \mathbb{C}) \times GL(2, \mathbb{C})$, the second component decomposes as $\mathcal{H}_{1,1} \oplus \mathbb{C}\varepsilon$. As we saw in Example 4.1.3, the (unnormalized) fundamental invariant corresponding to the second component in (4.2) is $\sum_{|I|=2} \det_I(z) \det_I(\bar{z})$, where I indicates the choice of rows. Thus, if γ'_2 is the invariant corresponding to $\mathcal{H}_{1,1}$, we can average over orthonormal bases for $\mathcal{H}_{1,1} \oplus \mathbb{C}\varepsilon$ in two different ways to obtain $\sum_{|I|=2} (1/8) \det_I(z) \det_I(\bar{z}) = \gamma'_2(z) + |\varepsilon(z)|^2 / (8n)$, so that

$$\gamma'_2(z) = \frac{1}{8} \left[\sum_{|I|=2} \det_I(z) \det_I(\bar{z}) - \frac{|\varepsilon(z)|^2}{n} \right].$$

4.1.7. $Sp(2n) \otimes SL(3)$ Again, we write the columns of $z \in \mathbb{C}^{2n} \otimes \mathbb{C}^3$ as $v_i \in \mathbb{C}^{2n}$, $i = 1, 2, 3$. We have the three $Sp(2n, \mathbb{C})$ -invariants $\varepsilon_{ij}(z) = \omega(v_i, v_j)$, $1 \leq i < j \leq 3$, and corresponding operators $D_{ij} = \varepsilon_{ij}(\partial/\partial z)$. The space of harmonic polynomials is $\mathcal{H} = \{p \in \mathbb{C}[V] \mid D_{ij}p = 0 \text{ for all } i, j\}$, which decomposes as $\mathcal{H} = \sum \mathcal{H}_{\alpha_1, \alpha_2, \alpha_3}$, where $\mathcal{H}_{\alpha_1, \alpha_2, \alpha_3}$ has highest weight vector $z_{11}^{\alpha_1 - \alpha_2} \det_2(z)^{\alpha_2 - \alpha_3} \det_3(z)^{\alpha_3}$. The full decomposition of $\mathbb{C}[V]$ is (from [11]):

$$\mathbb{C}[V] = \sum_{\substack{0 \leq b_1 \leq \alpha_1 - \alpha_2 \\ 0 \leq b_2 \leq \alpha_2 - \alpha_3 \\ b_3 \geq 0}} \sigma_{\alpha_1, \alpha_2, \alpha_3}^n \otimes \rho_{(\alpha_1 + b_2 + b_3, b_1 + \alpha_2 + b_3, b_1 + b_2 + \alpha_3)}^3,$$

where $\sigma_{\alpha_1, \alpha_2, \alpha_3}^n$ is the representation of $Sp(2n, \mathbb{C})$ on $\mathcal{H}_{\alpha_1, \alpha_2, \alpha_3}$ and ρ_D^3 is the representation of $SL(3, \mathbb{C})$ with Young's diagram D . By [11], we have six fundamental highest weight vectors:

$$\begin{aligned} \zeta_1 &= z_{11}, & \zeta'_2 &= \varepsilon_{12}(z), & \zeta'_3 &= z_{11} \varepsilon_{23}(z) - z_{12} \varepsilon_{13}(z) + z_{13} \varepsilon_{12}(z), \\ \zeta_2 &= \det_2(z), & \zeta_3 &= \det_3(z), & \zeta'_4 &= \det_{12,13}(z) \varepsilon_{12}(z) - \det_2(z) \varepsilon_{13}(z). \end{aligned}$$

We have the usual fundamental invariant $\gamma_1(z) = |z|^2$. The second degree fundamental invariants $\gamma_2(z)$ and $\gamma'_2(z)$ are found in the components $\sigma_{1,1}^n \otimes \rho_{(1,1)}^3$ and $\sigma_0^n \otimes \rho_{(1,1)}^3$ respectively. The sum of these two components is the $SL(2n, \mathbb{C}) \times SL(3, \mathbb{C})$ -irreducible $\rho_{(1,1)}^{2n} \otimes \rho_{(1,1)}^3$. As in the previous example, we take diagonal sums in two ways to obtain

$$\frac{1}{8} \sum_{|I|=|J|=2} \det_{I,J}(z) \det_{I,J}(\bar{z}) = \gamma_2(z) + \gamma'_2(z).$$

On the other hand, $\rho_0^{2n} \otimes \rho_{(1,1)}^3$ is spanned by the ε_{ij} 's, and hence:

$$\gamma_2'(z) = \frac{1}{8n} \sum \varepsilon_{ij}(z)\varepsilon_{ij}(\bar{z}), \quad \gamma_2(z) = \frac{1}{8} \sum_{|I|=|J|=2} \det_{I,J}(z)\det_{I,J}(\bar{z}) - \frac{1}{8n} \gamma_2'(z).$$

The highest weight vector ζ_3' lies in the the irreducible subspace $\sigma_{1,0,0}^n \otimes \rho_{(1,1,1)}^3$, which has dimension six. An orthogonal basis is given by $u_i' = z_{i1}\varepsilon_{23} - z_{i2}\varepsilon_{13} + z_{i3}\varepsilon_{12}$, for $i = 1, \dots, 6$. The first of these basis elements can be rewritten as

$$u_1'(z) = \det \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{n+2,1} & z_{n+2,2} & z_{n+2,3} \end{pmatrix} + \det \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{31} & z_{32} & z_{33} \\ z_{n+3,1} & z_{n+3,2} & z_{n+3,3} \end{pmatrix},$$

which is a sum of twelve distinct monomials, each of norm 8. Thus we obtain:

$$\gamma_3'(z) = \frac{1}{96} \sum_{i=1}^6 u_i'(z)u_i'(\bar{z}).$$

The sum of subspaces $(\sigma_{1,1,1}^n \otimes \rho_{(1,1,1)}^3) \oplus (\sigma_{1,0,0} \otimes \rho_{(1,1,1)}^3)$ is the $SL(2n, \mathbb{C}) \times SL(3, \mathbb{C})$ -irreducible $\rho_{(1,1,1)}^{2n} \otimes \rho_{(1,1,1)}^3$, and hence

$$\gamma_3(z) = \frac{1}{48} \sum_{|I|=3} \det_I(z)\det_I(\bar{z}) - \gamma_3'(z),$$

where the subscript I indicates a choice of rows. We have not derived a formula for the fundamental invariant γ_4' , which seems quite complicated.

4.1.8. $Sp(4) \otimes SL(n)$ The fundamental highest weight vectors are, from [11],

$$\begin{aligned} \zeta_1 &= z_{11}, & \zeta_2 &= \det_2(z), & \zeta_3 &= \det_3(z), \\ \zeta_4 &= \det_4(z), & \zeta_2' &= \varepsilon_{12}(z), & \zeta_4' &= \det_{12,13}(z)\varepsilon_{12}(z) - \det_2(z)\varepsilon_{13}(z). \end{aligned}$$

The highest weight vector ζ_2' determines the space spanned by $\varepsilon_{ij}(z) = \omega(v_i, v_j)$, $1 \leq i < j \leq m$, and thus

$$\gamma_2'(z) = \frac{1}{16} \sum |\varepsilon_{ij}(z)|^2.$$

As before, the spaces with highest weight vectors ζ_2 and ζ_2' sum to the $SL(4, \mathbb{C}) \times SL(n, \mathbb{C})$ -irreducible $\rho_{(1,1)}^4 \otimes \rho_{(1,1)}^m$, and hence

$$\gamma_2(z) = \frac{1}{48} \sum_{|I|=|J|=2} |\det_{I,J}(z)|^2 - \gamma_2'(z).$$

In degree three, the irreducible subspace containing ζ_3 is the $SL(4, \mathbb{C}) \times SL(n, \mathbb{C})$ -irreducible $\rho_{(1,1,1)}^4 \otimes \rho_{(1,1,1)}^n$, and hence

$$\gamma_3(z) = \frac{1}{8} \sum_{|I|=|J|=3} |\det_{I,J}(z)|^2.$$

Similarly,

$$\gamma_4(z) = \frac{1}{24 \cdot 16} \sum_{|J|=4} |\det_J(z)|^2,$$

where J indicates a choice of columns.

4.1.9. Spin(7) The action here is the spin representation of $Spin(7, \mathbb{C})$ on $V = \Lambda(\mathbb{C}^3)$. We use the usual basis $1, e_i, e_i \wedge e_j, e_1 \wedge e_2 \wedge e_3$ for V and let $f_0, f_i, f_{ij}, f_{123}$ respectively be the dual basis for V^* . The polynomial $\varepsilon = f_0 f_{123} - f_1 f_{23} + f_2 f_{13} - f_3 f_{12}$ defines a $Spin(7, \mathbb{C})$ -invariant inner product on V . So we have fundamental highest weight vectors f_0, ε and corresponding (unnormalized) fundamental invariants $\gamma_1(z) = |z|^2$ and $\gamma_2(z) = |\varepsilon(z)|^2$.

4.1.10. Spin(9) According to [11], there are three fundamental highest weight vectors, of degrees 1, 2 and 2. The space $\mathcal{P}_2(V)$ decomposes into three irreducible components, of dimensions 126, 9 and 1. The 9-dimensional irreducible is a copy of the standard representation of $SO(9, \mathbb{C})$, and we have a $Spin(9, \mathbb{C})$ -invariant. More explicitly, we take $V = \Lambda(\mathbb{C}^4)$ and use the natural basis for V with dual basis $f_0, f_i, f_{ij}, f_{ijk}, f_{1234}$ as in Example 4.1.9. Here f_0 is a highest weight vector for $\mathcal{P}_1(V) \cong V^*$, and f_0^2 is a highest weight vector for the 126-dimensional irreducible in $\mathcal{P}_2(V)$. A highest weight vector for the 9-dimensional irreducible is $f_0 f_{234} - f_2 f_{34} + f_3 f_{24} - f_4 f_{23}$, and the $Spin(9, \mathbb{C})$ -invariant inner product is given by the following pairing of coordinates:

$$f_0 f_{1234} + f_1 f_{234} - f_2 f_{134} + f_3 f_{124} - f_4 f_{123} - f_{12} f_{34} + f_{13} f_{24} - f_{14} f_{23}.$$

4.1.11. Spin(10) The spin representation for $Spin(10, \mathbb{C})$ can be realized in $V = \Lambda^{\text{even}}(\mathbb{C}^5) = \mathbb{C} \oplus \Lambda^2(\mathbb{C}^5) \oplus \Lambda^4(\mathbb{C}^5)$, and we use the natural bases for V and V^* as in the preceding two examples. We have fundamental highest weight vectors of degrees 1 and 2 given by f_0 and $f_{23} f_{45} - f_{24} f_{35} + f_{25} f_{34}$. The second of these generates a copy of the standard module for $SO(10, \mathbb{C})$ in $\mathcal{P}_2(V)$.

4.1.12. E₆ Following [7], we realize the standard representation of E_6 on a Jordan algebra V of dimension 27 with elements

$$X = \begin{bmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{bmatrix},$$

where $\xi_i \in \mathbb{C}$ and $x_i \in \mathcal{C}$, the 8-dimensional (complex) Cayley algebra. We have $so(8, \mathbb{C})$ as a subalgebra of the the Lie algebra of E_6 . Under the restriction to $so(8, \mathbb{C})$, V decomposes into three 8-dimensional irreducibles (the standard representation and the two inequivalent spin representations) and a 3-dimensional trivial subspace (corresponding to the diagonal entries in X).

Under an appropriate choice of positive roots, we find that $X \mapsto \xi_1$ is a highest weight vector for $\mathcal{P}_1(V)$. We are left with two remaining fundamental highest weight vectors, of degrees 2 and 3. The highest weight vector in degree two generates the 27-dimensional representation contragredient to $\mathcal{P}_1(V)$. Let V_3 be the subspace of V spanned by the matrices X with $x_3 \neq 0$, and all other entries zero. Then $so(8, \mathbb{C})$ acts on V_3 by one of the spin representations. There is an $so(8, \mathbb{C})$ -invariant inner product on V_3 given by pairing the appropriate coordinates, and this is the fundamental highest weight vector of degree two. In degree three, we have an E_6 -invariant, 'det', given explicitly by

$$X \mapsto \xi_1 \xi_2 \xi_3 + (x_1 x_2 x_3 + \overline{x_1 x_2 x_3}) - \xi_1 x_1 \bar{x}_1 - \xi_2 x_2 \bar{x}_2 - \xi_3 x_3 \bar{x}_3.$$

4.2. Indecomposable non-irreducible multiplicity free actions Table 2 lists the semisimple groups $G \subset GL(V)$ acting indecomposably on $V = V_1 \oplus V_2$ for which the action of $G \times (\mathbb{C}^\times)^2$ is multiplicity free. The subscripts on the direct sums indicate simple factors acting diagonally. Thus, for example, $SL(n) \oplus_{SL(n)} (SL(n) \otimes SL(m))$ denotes the image of $SL(n) \times SL(m)$ under the representation on $V = V_1 \oplus V_2 = (\mathbb{C}^n) \oplus (\mathbb{C}^n \otimes \mathbb{C}^m)$ where $SL(n, \mathbb{C})$ acts diagonally on V_1 and V_2 . In each case only one simple factor acts diagonally. For each such group G , the restrictions to V_1 and V_2 are irreducible multiplicity free actions and have thus been discussed above. In each case, we use the orthonormal basis for V obtained by adjoining the standard orthonormal bases for V_1 and V_2 , employed in our discussion of the irreducible multiplicity free actions. It is transparent that if we use such a basis to realize G as a matrix group $G \subset GL(n, \mathbb{C})$, then $\mathfrak{g}_\mathbb{Q} = \mathfrak{g} \cap gl(n, \mathbb{Q})$ is a rational form for \mathfrak{g} . In each case we present formulae for the fundamental highest weight vectors in $\mathbb{C}[V]$ that are rational in the coordinates with respect to this natural basis. Clearly, these include fundamental highest weight vectors for the actions of G on V_1 and V_2 . In each case, however, there are additional fundamental highest weights. We have listed the number of fundamental highest weight vectors and their degrees in Table 2. We follow the notational conventions in [5], to which we refer the reader for justification of the decompositions of $\mathbb{C}[V]$ described below.

4.2.1. $SL(n) \oplus_{SL(n)} SL(n)$ Here $G = SL(n, \mathbb{C})$ acts on $V = V_1 \oplus V_2 = \mathbb{C}^n \oplus \mathbb{C}^n$ via two copies of its defining representation. The decomposition is

$$\mathcal{P}_k(V_1) \otimes \mathcal{P}_\ell(V_2) = \sum_{j \leq \min(k, \ell)} \rho_{(k+\ell-j, j)}^n.$$

TABLE 2.

Group Example number	Degrees of fundamental highest weight vectors (number)
$\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} \mathbf{SL}(n)$ ($n \geq 2$) Example 4.2.1	1, 1, 2 (3)
$\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} \mathbf{SL}(n)^*$ ($n \geq 3$) Example 4.2.2	1, 1, 2 (3)
$\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} \Lambda^2(\mathbf{SL}(n))$ ($n \geq 4$) Example 4.2.3	1, 2, ..., $\lfloor n/2 \rfloor$, 1, 2, ..., $\lfloor (n + 1)/2 \rfloor$ (n)
$\mathbf{SL}(n)^* \oplus_{\mathbf{SL}(n)} \Lambda^2(\mathbf{SL}(n))$ ($n \geq 4$) Example 4.2.6	1, 2, ..., $\lfloor n/2 \rfloor$, 1, 2, ..., $\lfloor (n - 1)/2 \rfloor$ ($n - 1$)
$\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} (\mathbf{SL}(n) \otimes \mathbf{SL}(m))$ ($n, m \geq 2$) Example 4.2.4	1, 2, ..., $\min(n, m)$, 1, 2, ..., $\min(n, m + 1)$ ($\min(2n, 2m + 1)$)
$\mathbf{SL}(n)^* \oplus_{\mathbf{SL}(n)} (\mathbf{SL}(n) \otimes \mathbf{SL}(m))$ ($n \geq 3, m \geq 2$) Example 4.2.5	1, 2, ..., $\min(n, m)$, 1, 2, ..., $\min(n, m + 1)$ ($\min(2n, 2m + 1)$)
$\mathbf{SL}(2) \oplus_{\mathbf{SL}(2)} (\mathbf{SL}(2) \otimes \mathbf{Sp}(2n))$ ($n \geq 2$) Example 4.2.7	1, 1, 2, 2, 2 (5)
$(\mathbf{SL}(n) \otimes \mathbf{SL}(2)) \oplus_{\mathbf{SL}(2)} (\mathbf{SL}(2) \otimes \mathbf{SL}(m))$ ($n, m \geq 2$) Example 4.2.8	1, 1, 2, 2, 2 (5)
$(\mathbf{SL}(n) \otimes \mathbf{SL}(2)) \oplus_{\mathbf{SL}(2)} (\mathbf{SL}(2) \otimes \mathbf{Sp}(2m))$ ($n, m \geq 2$) Example 4.2.8	1, 1, 2, 2, 2, 2 (6)
$(\mathbf{Sp}(2n) \otimes \mathbf{SL}(2)) \oplus_{\mathbf{SL}(2)} (\mathbf{SL}(2) \otimes \mathbf{Sp}(2m))$ ($n, m \geq 2$) Example 4.2.8	1, 1, 2, 2, 2, 2, 2 (7)
$\mathbf{Sp}(2n) \oplus_{\mathbf{Sp}(2n)} \mathbf{Sp}(2n)$ ($n \geq 2$) Example 4.2.9	1, 1, 2, 2 (4)
$\mathbf{Spin}(8) \oplus_{\mathbf{Spin}(8)} \mathbf{SO}(8)$ Example 4.2.10	1, 1, 2, 2, 2 (5)

If we identify V with $\mathbb{C}^n \otimes \mathbb{C}^2$, we can describe the fundamental highest weight vectors as $\zeta_1 = z_{11}$, $\zeta'_1 = z_{12}$, and $\zeta_2 = \det_2(z)$. The irreducible component $\rho_{(k+\ell-j, j)}^n$ has highest weight vector $\zeta_1^{k-j} (\zeta'_1)^{\ell-j} \zeta_2^j$. The fundamental invariants are $|v_1|^2$, $|v_2|^2$, and $\sum_{|I|=2} |\det_I(z)|^2$, up to normalizations.

4.2.2. $\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} \mathbf{SL}(n)^*$ Here $\mathbf{SL}(n, \mathbb{C})$ acts on $V = V_1 \oplus V_2 = \mathbb{C}^n \oplus (\mathbb{C}^n)^*$ and we have

$$\mathcal{P}_k(V_1) \otimes \mathcal{P}_\ell(V_2) = \sum_{j \leq \min(k, \ell)} \rho_{(k+\ell-j, \ell, \dots, \ell, j)}^n.$$

Again, there are three fundamental highest weight vectors, of degrees 1, 1 and 2. The first two are coordinates in V and V^* respectively. The fundamental highest weight vector of degree two is given by the natural pairing $V_1 \times V_2 = \mathbb{C}^n \times (\mathbb{C}^n)^* \rightarrow \mathbb{C}$, which is $\mathbf{SL}(n)$ -invariant. The irreducible component $\rho_{(k+\ell-j, \ell, \dots, \ell, j)}^n$ has highest weight vector $\zeta_1^{k-j} (\zeta'_1)^{\ell-j} \zeta_2^j$ as in the preceding example.

4.2.3. $\mathbf{SL}(n) \oplus_{\mathbf{SL}(n)} \Lambda^2(\mathbf{SL}(n))$ We identify $V = V_1 \oplus V_2 = \mathbb{C}^n \oplus \Lambda^2(\mathbb{C}^n)$ with $\Lambda^2(\mathbb{C}^{n+1})$ by regarding the first row (or column) of an $(n+1) \times (n+1)$ skew symmetric matrix as an element of \mathbb{C}^n , and the remaining entries as an element of $\Lambda^2(\mathbb{C}^n)$. For $z \in \Lambda^2(\mathbb{C}^{n+1})$, we write z' for this element of $\Lambda^2(\mathbb{C}^n)$. That is, z' is obtained by removing the first row and column of z . Under this identification, the diagonal action of $\mathbf{SL}(n, \mathbb{C})$ on $V_1 \oplus V_2$ is realized on $\Lambda^2(\mathbb{C}^{n+1})$ by restricting the action of $\mathbf{SL}(n+1, \mathbb{C})$ to the subgroup $\mathbf{SL}(n, \mathbb{C}) \subset \mathbf{SL}(n+1, \mathbb{C})$ embedded as

$$\left\{ \left[\begin{array}{c|c} 1 & 0 \\ \hline 0 & A \end{array} \right] \mid A \in \mathbf{SL}(n, \mathbb{C}) \right\}.$$

The fundamental highest weight vectors arising from the separate actions of $\mathbf{SL}(n, \mathbb{C})$ on V_1 and V_2 are z_{12} and $\zeta'_k(z)$ for $k = 1, \dots, \lfloor n/2 \rfloor$, where $\zeta'_k(z)$ is the Pfaffian of the first $2k$ rows and columns of the $n \times n$ matrix z' . There are additional fundamental highest weight vectors, $\zeta_k(z)$ for $k = 1, \dots, \lfloor (n+1)/2 \rfloor$. These are the Pfaffians of the first $2k$ rows and columns of the $(n+1) \times (n+1)$ matrix z . Note that $\zeta_1(z) = z_{12}$, so our fundamental highest weight vectors are the ζ_k 's together with the ζ'_k 's. The discussion in Example 4.1.5 shows how these Pfaffians can be written explicitly as rational polynomials in the matrix entries. The decomposition of $\mathbb{C}[V]$ is $\mathbb{C}[V] = \sum_D \rho_D$, where D ranges over all Young's diagrams. For $D = (\lambda_1, \dots, \lambda_n)$, a highest weight vector for ρ_D in $\mathbb{C}[V]$ is

$$\zeta_1^{\lambda_1 - \lambda_2} (\zeta'_1)^{\lambda_2 - \lambda_3} \zeta_2^{\lambda_3 - \lambda_4} \dots \zeta_m^{\lambda_{n-1} - \lambda_n} (\zeta'_m)^{\lambda_n}$$

when $n = 2m$ is even, and

$$\zeta_1^{\lambda_1 - \lambda_2} (\zeta'_1)^{\lambda_2 - \lambda_3} \zeta_2^{\lambda_3 - \lambda_4} \dots \zeta_m^{\lambda_{n-2} - \lambda_{n-1}} (\zeta'_m)^{\lambda_{n-1} - \lambda_n} (\zeta_{m+1})^{\lambda_n}$$

when $n = 2m + 1$ is odd.

4.2.4. $SL(n) \oplus_{SL(n)} (SL(n) \otimes SL(m))$ We realize $V = V_1 \oplus V_2 = \mathbb{C}^n \oplus (\mathbb{C}^n \otimes \mathbb{C}^m)$ as $\mathbb{C}^n \otimes \mathbb{C}^{m+1}$ with $SL(n, \mathbb{C}) \times SL(m, \mathbb{C})$ acting via its embedding in $SL(n, \mathbb{C}) \times SL(m+1, \mathbb{C})$. For an $n \times (m+1)$ matrix z , write z' for the $n \times m$ matrix obtained by removing the first column. The fundamental highest weight vectors are $\zeta_k(z) = \det_k(z)$ for $k = 1, \dots, \min(n, m+1)$ and $\zeta'_k(z) = \det_k(z')$ for $k = 1, \dots, \min(n, m)$. Here $\zeta_1(z) = z_{11}$ and the $\zeta'_k(z)$'s are the fundamental highest weight vectors that arise from the separate actions of $SL(n, \mathbb{C}) \times SL(m)$ on V_1 and V_2 . The irreducible components of $\mathbb{C}[V]$ have the form $\rho_{(\lambda_1+\mu_1, \dots, \lambda_n+\mu_n)}^n \otimes \rho_\lambda^m$ where λ is a Young's diagram with at most $\min(n, m)$ rows, and $\mu = (\mu_1, \dots, \mu_n)$ satisfies $\mu_j \leq \lambda_{j-1} - \lambda_j$ for $j \geq 2$. The highest weight vector for this component is

$$\zeta_1^{\mu_1} \zeta_2^{\mu_2} \dots \zeta_n^{\mu_n} (\zeta'_1)^{\lambda_1-\lambda_2-\mu_2} (\zeta'_2)^{\lambda_2-\lambda_3-\mu_3} \dots (\zeta'_{n-1})^{\lambda_{n-1}-\lambda_n-\mu_n} (\zeta'_n)^{\lambda_n}$$

when $n \leq m$ and

$$\zeta_1^{\mu_1} \zeta_2^{\mu_2} \dots \zeta_{m+1}^{\mu_{m+1}} (\zeta'_1)^{\lambda_1-\lambda_2-\mu_2} (\zeta'_2)^{\lambda_2-\lambda_3-\mu_3} \dots (\zeta'_{m-1})^{\lambda_{m-1}-\lambda_m-\mu_m} (\zeta'_m)^{\lambda_m-\mu_{m+1}}$$

when $n > m$.

4.2.5. $SL(n)^* \oplus_{SL(n)} (SL(n) \otimes SL(m))$ To decompose $\mathbb{C}[V]$ under the action of $SL(n, \mathbb{C}) \times SL(m, \mathbb{C})$ on $V = V_1 \oplus V_2 = (\mathbb{C}^n)^* \oplus (\mathbb{C}^n \otimes \mathbb{C}^m)$, we must decompose $\rho_{k^{n-1}}^n \otimes (\rho_\lambda^n \otimes \rho_\lambda^m)$, where k^{n-1} denotes the Young's diagram with k boxes in each of $n-1$ rows and λ is a Young's diagram with at most $\min(n, m)$ rows. We obtain $\sum_\mu \rho_{(\mu_1+k, \dots, \mu_{n-1}+k, \mu_n)}^n \otimes \rho_\lambda^m$, where $\mu_1 + \dots + \mu_n = \lambda_1 + \dots + \lambda_n$, $\mu_j \leq \lambda_j$ for $j = 1, \dots, n-1$, $0 \leq \mu_n - \lambda_n \leq k$, and $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \lambda_n$. We write $z = (z_{ij})$ for elements of $V_2 = \mathbb{C}^n \otimes \mathbb{C}^m$ and ξ for elements in $V_1 = (\mathbb{C}^n)^*$. Let z' be the $(n+1) \times m$ matrix

$$z' = \left[\begin{array}{ccc} \xi(v_1) & \dots & \xi(v_m) \\ z & & \end{array} \right],$$

where v_i is the i 'th column of z . Then the action of $SL(n, \mathbb{C}) \times SL(m, \mathbb{C})$ on z' is given by embedding $SL(n, \mathbb{C}) \times SL(m, \mathbb{C})$ in $SL(n+1, \mathbb{C}) \times SL(m, \mathbb{C})$. The standard coordinate function ξ_n is a highest weight vector for the representation of $SL(n, \mathbb{C})$ on $\mathcal{P}_1(V_1) \cong V_1^* \cong \mathbb{C}^n$. Our fundamental highest weight vectors are thus $\xi_n, \det_k(z)$ $k = 1, \dots, \min(n, m)$ and $\det_k(z')$ for $k = 1, \dots, \min(n-1, m)$. Here $\det_k(z)$ is a polynomial of degree k on V and $\det_k(z')$ has degree $k+1$. A highest weight vector for the irreducible $\rho_{(\mu_1+k, \dots, \mu_{n-1}+k, \mu_n)}^n \otimes \rho_\lambda^m$ in $\mathbb{C}[V]$ can be expressed in terms of the fundamental highest weight vectors as:

$$\det_1(z')^{\lambda_1-\mu_1} \det_2(z')^{\lambda_2-\mu_2} \dots \det_{n-1}(z')^{\lambda_{n-1}-\mu_{n-1}} \times \det_1(z)^{\mu_1-\lambda_2} \det_2(z)^{\mu_2-\lambda_3} \dots \det_n(z)^{\lambda_n} \xi_n^{k-\mu_n+\lambda_n}.$$

We remark that the number and degrees of the fundamental highest weight vectors for this example coincide with those for the related Example 4.2.4.

4.2.6. $SL(n)^* \oplus_{SL(n)} \Lambda^2(SL(n))$ Here $V = V_1 \oplus V_2 = (\mathbb{C}^n)^* \oplus \Lambda^2(\mathbb{C}^n)$. We write $z = (z_{ij}) \in \Lambda^2(\mathbb{C}^n)$, $\xi = (\xi_i) \in (\mathbb{C}^n)^*$, and let

$$z' = \left[\begin{array}{c|ccc} \mathbf{0} & \xi(v_1) & \cdots & \xi(v_n) \\ \hline -\xi(v_1) & & & \\ \vdots & & & \\ -\xi(v_n) & & & z \end{array} \right] \in \Lambda^2(\mathbb{C}^{n+1}),$$

where v_i is the i 'th column of z . The group $SL(n, \mathbb{C})$ acts on z' via the embedding $SL(n, \mathbb{C}) \hookrightarrow SL(n + 1, \mathbb{C})$, as in Example 4.2.3. Our fundamental highest weight vectors are ξ_n, ζ_k for $k = 1, \dots, \lfloor (n - 1)/2 \rfloor$ and ζ'_k for $k = 1, \dots, \lfloor n/2 \rfloor$. Here ζ_k and ζ'_k are the Pfaffians of the first $2k$ rows and columns of z and z' respectively. Note that ζ_k is a homogeneous polynomial of degree k on V , whereas ζ'_k has degree $k + 1$.

To understand the decomposition of $\mathbb{C}[V]$, we consider the cases n even and n odd separately. If $n = 2m$, we need to consider the decomposition of $\rho_{k^{n-1}}^n \otimes \rho_\lambda^n$ where $\lambda = (\lambda_1, \lambda_1, \lambda_2, \lambda_2, \dots, \lambda_m, \lambda_m)$. We obtain irreducible components corresponding to Young's diagrams $(k + \lambda_1, k + \mu_1, \dots, k + \lambda_m, \mu_m + \lambda_m)$, where $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \mu_{m-1} \geq \lambda_m, \mu_m \leq k$ and $\mu_1 + \dots + \mu_m = \lambda_1 + \dots + \lambda_{m-1}$. The highest weight vector for this component is

$$(\zeta'_1)^{\lambda_1 - \mu_1} (\zeta'_2)^{\lambda_2 - \mu_2} \dots (\zeta'_{m-1})^{\lambda_{m-1} - \mu_{m-1}} \xi_n^{k - \mu_m} \zeta_1^{\mu_1 - \lambda_2} \zeta_2^{\mu_2 - \lambda_3} \dots \zeta_{m-1}^{\mu_{m-1} - \lambda_m} \zeta_m^{\lambda_m}.$$

For $n = 2m + 1$, we have $\rho_{k^{2m}}^n \otimes \rho_\lambda^n$ with $\lambda = (\lambda_1, \lambda_1, \lambda_2, \lambda_2, \dots, \lambda_m, \lambda_m, 0)$, whose irreducible components correspond to diagrams $(k + \lambda_1, k + \mu_1, \dots, k + \lambda_m, k + \mu_m, \mu_{m+1})$, where $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \lambda_m \geq \mu_m, \mu_{m+1} \leq k$, and $\mu_1 + \dots + \mu_{m+1} = \lambda_1 + \dots + \lambda_m$. This component has highest weight vector

$$(\zeta'_1)^{\lambda_1 - \mu_1} (\zeta'_2)^{\lambda_2 - \mu_2} \dots (\zeta'_m)^{\lambda_m - \mu_m} \xi_n^{k - \mu_{m+1}} \zeta_1^{\mu_1 - \lambda_2} \zeta_2^{\mu_2 - \lambda_3} \dots \zeta_{m-1}^{\mu_{m-1} - \lambda_m} \zeta_m^{\mu_m}.$$

It is interesting to note that here we have one fewer fundamental highest weight vector than in the related untwisted Example 4.2.3.

4.2.7. $SL(2) \oplus_{SL(2)} (SL(2) \otimes Sp(2n))$ We identify $V = V_1 \oplus V_2 = \mathbb{C}^2 \oplus (\mathbb{C}^{2n} \otimes \mathbb{C}^2)$ with $\mathbb{C}^{2n+1} \otimes \mathbb{C}^2$ as in Example 4.2.4, and embed $G = Sp(2n, \mathbb{C}) \times SL(2, \mathbb{C})$ in $SL(2n + 1, \mathbb{C}) \times SL(2, \mathbb{C})$. Write $z = (z_{ij})$ for a $(2n + 1) \times 2$ matrix in V and let z' denote the $2n \times 2$ matrix obtained by removing the first row of z . The fundamental highest weight vectors are $z_{11}, z_{21}, \varepsilon', \det_2(z)$ and $\det_2(z')$, where ε' is the symplectic product of the columns of z' . The irreducible components are of the form

$$\rho_{(\alpha_1 + \alpha_3 + \beta_1, \alpha_2 + \alpha_3 + \beta_2)}^n \otimes \sigma_{\alpha_1, \alpha_2} \subset \mathcal{P}_{\beta_1 + \beta_2}(\mathbb{C}^2) \otimes \mathcal{P}_{\alpha_1 + \alpha_2 + 2\alpha_3}(\mathbb{C}^{2n} \otimes \mathbb{C}^2)$$

with the restrictions $\alpha_1 \geq \alpha_2, \beta_2 \leq \alpha_1 - \alpha_2$. The highest weight vector for this component is

$$z_{11}^{\beta_1} z_{12}^{\alpha_1 - \alpha_2 - \beta_2} (\varepsilon')^{\alpha_3} \det_2(z)^{\beta_2} \det_2(z')^{\alpha_2}.$$

4.2.8. $(\mathbf{SL}(n) \otimes \mathbf{SL}(2)) \oplus_{\mathbf{SL}(2)} (\mathbf{SL}(2) \otimes \mathbf{SL}(m))$ Here we examine together the actions of

- (a) $G = \mathbf{SL}(n, \mathbb{C}) \times \mathbf{SL}(2, \mathbb{C}) \times \mathbf{SL}(m, \mathbb{C})$ on $V = V_1 \oplus V_2 = (\mathbb{C}^n \otimes \mathbb{C}^2) \oplus (\mathbb{C}^2 \otimes \mathbb{C}^m)$,
- (b) $G = \mathbf{SL}(n, \mathbb{C}) \times \mathbf{SL}(2, \mathbb{C}) \times \mathbf{Sp}(2m, \mathbb{C})$ on $V = V_1 \oplus V_2 = (\mathbb{C}^n \otimes \mathbb{C}^2) \oplus (\mathbb{C}^2 \otimes \mathbb{C}^{2m})$,
- (c) $G = \mathbf{Sp}(2n, \mathbb{C}) \times \mathbf{SL}(2, \mathbb{C}) \times \mathbf{Sp}(2m, \mathbb{C})$ on $V = V_1 \oplus V_2 = (\mathbb{C}^{2n} \otimes \mathbb{C}^2) \oplus (\mathbb{C}^2 \otimes \mathbb{C}^{2m})$.

In each case, we identify V with $\mathbb{C}^{N+M} \otimes \mathbb{C}^2$, where $N = n$ or $2n$, $M = m$ or $2m$. We write elements of V as

$$z = \begin{bmatrix} x \\ y \end{bmatrix},$$

where $x \in \mathbb{C}^N \otimes \mathbb{C}^2$ and $y \in \mathbb{C}^M \otimes \mathbb{C}^2$. In cases (b) and (c), we have the symplectic invariants $\varepsilon(x), \varepsilon(y)$ given by the symplectic inner products of the columns of x and y respectively.

In case (a), $\mathbb{C}[V]$ decomposes into irreducibles of the form $\rho_\lambda^n \otimes \rho_\nu^2 \otimes \rho_\mu^m$ where $|\mu| = |\lambda| + |\nu|$, $\mu_2 \leq \lambda_1 + \nu_2$, $\mu_1 - \lambda_1 \geq \nu_2$, and $\mu_2 \geq \lambda_2 + \nu_2$. The fundamental highest weight vectors are $x_{11}, y_{11}, \det_2(x), \det_2(y)$, and $x_{11}y_{12} - x_{12}y_{11}$. The irreducible component above has highest weight vector

$$x_{11}^{\lambda_1 + \nu_2 - \mu_2} y_{11}^{\mu_1 - \lambda_1 - \nu_2} \det_2(x)^{\lambda_2} \det_2(y)^{\nu_2} (x_{11}y_{12} - x_{12}y_{11})^{\mu_2 - \nu_2 - \lambda_2}.$$

In case (b), the irreducibles are of the form

$$\rho_\lambda^n \otimes \rho_\mu^2 \otimes \sigma_\nu^m \subset \mathcal{P}_{|\lambda|}(\mathbb{C}^n \otimes \mathbb{C}^2) \otimes \mathcal{P}_{|\nu|+2j}(\mathbb{C}^{2m} \otimes \mathbb{C}^2),$$

where $j \geq 0$, $|\mu| = |\lambda| + |\nu| + 2j$, $\mu_1 \geq \nu_1 + j$, $\lambda_2 + \nu_2 + j \leq \mu_2 \leq \nu_1 + \lambda_2 + j$. The fundamental highest weight vectors are $x_{11}, y_{11}, \det_2(x), \det_2(y), \varepsilon(y)$ and $x_{11}y_{12} - x_{12}y_{11}$. The typical irreducible component above has highest weight vector

$$x_{11}^{\lambda_1 + \nu_2 + j - \mu_2} y_{11}^{\nu_1 + \lambda_2 + j - \mu_2} \det_2(x)^{\lambda_2} \det_2(y)^{\nu_2} \varepsilon(y)^j (x_{11}y_{12} - x_{12}y_{11})^{\mu_2 - \nu_2 - \lambda_2 - j}.$$

In case (c), the fundamental highest weight vectors are $x_{11}, y_{11}, \det_2(x), \det_2(y), \varepsilon(x), \varepsilon(y)$ and $x_{11}y_{12} - x_{12}y_{11}$. The irreducibles are of the form

$$\sigma_\lambda^n \otimes \rho_\mu^2 \otimes \sigma_\nu^m \subset \mathcal{P}_{|\lambda|+2a}(\mathbb{C}^{2n} \otimes \mathbb{C}^2) \otimes \mathcal{P}_{|\nu|+2b}(\mathbb{C}^{2m} \otimes \mathbb{C}^2),$$

where $|\mu| = |\lambda| + |\nu| + 2a + 2b$, $\mu_2 \geq \lambda_2 + \nu_2 + a + b$, $\mu_2 \leq \lambda_1 + \nu_2 + a + b$ and $\mu_2 \leq \lambda_2 + \nu_1 + a + b$. This has highest weight vector

$$x_{11}^{\lambda_1 + \nu_2 + a + b - \mu_2} y_{11}^{\nu_1 + \lambda_2 + a + b - \mu_2} \det_2(x)^{\lambda_2} \det_2(y)^{\nu_2} \varepsilon(x)^a \varepsilon(y)^b (x_{11}y_{12} - x_{12}y_{11})^{\mu_2 - \nu_2 - \lambda_2 - a - b}.$$

4.2.9. $\mathbf{Sp}(2n) \oplus_{\mathbf{Sp}(2n)} \mathbf{Sp}(2n)$ We identify $V = V_1 \oplus V_2 = \mathbb{C}^{2n} \oplus \mathbb{C}^{2n}$ with $\mathbb{C}^{2n} \otimes \mathbb{C}^2$ and write points $z \in V$ as $z = (z_{ij})$. The group $G = \mathbf{Sp}(2n, \mathbb{C})$ acts on V from the left and the symplectic product $\varepsilon(z)$ of the columns is an invariant. The fundamental highest weight vectors are $z_{11}, z_{12}, \varepsilon(z)$, and $\det_2(z)$. The irreducible components of $\mathcal{P}_k(\mathbb{C}^{2n}) \otimes \mathcal{P}_\ell(\mathbb{C}^{2n})$ are of the form $\sigma_{\lambda_1, \lambda_2}^n$, with $\lambda_1 \geq \lambda_2$, and $\lambda_1 + \lambda_2 = k + \ell - 2m$ for $m \geq 0$. The highest weight vector for this irreducible is

$$z_{11}^{k-\lambda_2-m} z_{12}^{\ell-\lambda_2-m} \varepsilon(z)^m \det_2(z)^{\lambda_2}.$$

4.2.10. $\mathbf{SO}(8) \oplus_{\mathbf{Spin}(8)} \mathbf{Spin}(8)$ In this example, $\mathbf{Spin}(8, \mathbb{C})$ acts on $V = V_1 \oplus V_2 = \mathbb{C}^8 \oplus \Lambda^{\text{even}}(\mathbb{C}^4)$ via the standard representation of $\mathbf{SO}(8, \mathbb{C})$ on V_1 and the spin representation on $V_2 \cong \mathbb{C}^8$. These two representations can be written as ω_{1000} and $\omega_{\frac{1}{2}\frac{1}{2}\frac{1}{2}\frac{1}{2}}$, where the subscripts indicate the highest weights for the representations. We have the decompositions

$$\mathcal{P}_\ell(V_1) \otimes \mathcal{P}_{2k}(V_2) = \sum_{i,j,m} \omega_{k+\ell-i-2j-m, k-i, k-i, k-i-m}, \quad \text{and}$$

$$\mathcal{P}_\ell(V_1) \otimes \mathcal{P}_{2k+1}(V_2) = \sum_{i,j,m} \omega_{k+\ell-i-2j-m+\frac{1}{2}, k-i+\frac{1}{2}, k-i+\frac{1}{2}, k-i-m+\frac{1}{2}}.$$

We use the standard basis for V_2^* as in Example 4.1.11 and give V_1 coordinates z_1, \dots, z_8 , which are weight vectors for the maximal torus. We have fundamental highest weight vectors corresponding to the highest weight vectors for each V_i , the $\mathbf{Spin}(8, \mathbb{C})$ -invariants in each V_i and a fifth highest weight vector which generates the other spin representation in $\mathcal{P}_2(V)$. Explicitly, these are

$$\begin{aligned} \zeta_1 &= z_1, & \zeta'_1 &= f_0, & \zeta_2 &= z_1 z_5 + z_2 z_6 + z_3 z_7 + z_4 z_8, \\ \zeta'_2 &= f_0 f_{1234} - f_{12} f_{34} + f_{13} f_{24} - f_{14} f_{13}, & \zeta''_2 &= z_8 f_0 + z_1 f_{14} + z_2 f_{24} + z_3 f_{34}. \end{aligned}$$

The irreducible component $\omega_{k+\ell-i-2j-m, k-i, k-i, k-i-m}$ has highest weight vector

$$\zeta_1^{\ell-2j-m} \zeta_2^j (\zeta'_1)^{2k-2i-m} (\zeta'_2)^i (\zeta''_2)^m,$$

and $\omega_{k+\ell-i-2j-m+\frac{1}{2}, k-i+\frac{1}{2}, k-i+\frac{1}{2}, k-i-m+\frac{1}{2}}$ has highest weight vector

$$\zeta_1^{\ell-2j-m} \zeta_2^j (\zeta'_1)^{2k-2i-m+1} (\zeta'_2)^i (\zeta''_2)^m.$$

References

[1] C. Benson, J. Jenkins, R. Lipsman and G. Ratcliff, 'A geometric criterion for Gelfand pairs associated with the Heisenberg group', *Pacific J. Math.* **178** (1997), 1–36.

- [2] C. Benson, J. Jenkins and G. Ratcliff, 'Bounded K -spherical functions on Heisenberg groups', *J. Funct. Anal.* **105** (1992), 409–443.
- [3] ———, ' $O(n)$ -spherical functions on Heisenberg groups', *Contemp. Math.* **145** (1993), 181–197.
- [4] ———, 'The spherical transform of a Schwartz function on the Heisenberg group', *J. Funct. Anal.* **154** (1998), 379–423.
- [5] C. Benson and G. Ratcliff, 'A classification for multiplicity free actions', *J. Algebra* **181** (1996), 152–186.
- [6] ———, 'Combinatorics and spherical functions on the Heisenberg group', *Representation Theory* **2** (1998), 79–105.
- [7] C. Chevalley and R. Shafer, 'The exceptional Lie algebras F_4 and E_6 ', *Proc. Nat. Acad. Sci. Amer.* **36** (1950), 137–141.
- [8] H. Dib, 'Fonctions de Bessel sur une algèbre de Jordan', *J. Math. Pures Appl.* **69** (1990), 403–448.
- [9] J. Faraut and A. Koranyi, *Analysis on symmetric cones* (Oxford University Press, New York, 1994).
- [10] G. Folland, *Harmonic analysis in phase space* (Princeton University Press, New Jersey, 1989).
- [11] R. Howe and T. Umeda, 'The Capelli identity, the double commutant theorem and multiplicity-free actions', *Math. Ann.* **290** (1991), 565–619.
- [12] V. Kac, 'Some remarks on nilpotent orbits', *J. Algebra* **64** (1980), 190–213.
- [13] A. Leahy, 'A classification of multiplicity free representations', *J. Lie Theory* **8** (1998), 367–391.
- [14] A. Okounkov and G. Olshanski, 'Shifted Jack polynomials, binomial formula, and applications', *Math. Res. Letters* **4** (1997), 69–78.
- [15] ———, 'Shifted Schur functions II. The binomial formula for characters of classical groups and its applications', *Amer. Math. Soc. Transl. Ser. 2* **181** (1998), 245–271.
- [16] S. Sahi, 'The spectrum of certain invariant differential operators associated to Hermitian symmetric spaces', in: *Lie theory and geometry* (ed. J. L. Brylinski), Progress in Math. 123 (Birkhäuser, Boston, 1994) pp. 569–576.
- [17] Z. Yan, 'Special functions associated with multiplicity-free representations', preprint.
- [18] ———, 'Generalized hypergeometric functions and Laguerre polynomials in two variables', *Contemp. Math.* **138** (1992), 239–259.

Department of Mathematics and Computer Science

University of Missouri-St. Louis

St. Louis, MO 63121

USA

e-mail: benson@arch.umsl.edu, ratcliff@arch.umsl.edu