

Review Article

Cite this article: Harvie M and Haiba M (2025). The impact of intermittent energy restriction on women's health. *Proceedings of the Nutrition Society*, page 1 of 10. doi: [10.1017/S0029665125000059](https://doi.org/10.1017/S0029665125000059)

Received: 14 October 2024

Revised: 20 January 2025

Accepted: 20 January 2025

Keywords:

Intermittent energy restriction; Time-restricted eating; Women's health; Weight loss

Corresponding author:

Michelle Harvie;

Email: michelle.harvie@manchester.ac.uk

The impact of intermittent energy restriction on women's health

Michelle Harvie^{1,2,3}  and Mai Haiba^{1,2}

¹Division of Cancer Sciences, The University of Manchester, Manchester, UK; ²Department of Dietetics, Wythenshawe Hospital, Manchester University NHS Foundation Trust, Manchester M23 9LT, UK and ³Manchester Breast Centre, Oglesby Cancer Research Centre, The Christie NHS Foundation Trust, University of Manchester, Manchester, UK

Abstract

Intermittent energy-restricted diets are used amongst women with overweight and obesity and a healthy weight. For those with overweight and obesity weight control is typically achieved through daily energy restriction (DER) which has reduced adherence and attenuated metabolic benefits over time. Several intermittent energy restriction (IER) regimens have been developed aiming to promote maintained weight loss and additional weight independent metabolic benefits including the 5:2 diet, alternate day fasting (ADF) and time-restricted eating (TRE). This review summarises the potential benefits or harms of these regimens for managing women's health. 5:2 and ADF diets have equivalent long term (≥ 6 -month) adherence, weight loss and metabolic benefits to DER. Current limited evidence suggests IER is a safe weight loss intervention for women which does not affect reproductive or bone health, increase eating disorders or disturb sleep. Adherence and weight loss with both IER and DER are lower amongst younger women compared to older women and men. Weight loss with ADF and TRE has, respectively, improved symptoms of polycystic ovarian syndrome and premenstrual syndrome, but there is no evidence of weight-independent effects of IER on these conditions. There is little evidence of the benefits and/or harms of IER amongst healthy weight women in whom there is a greater potential for adverse effects on reproductive and bone health, fat free mass, eating disorders and sleep. Further research benefits of IER for weight control and metabolic health as well as harms are required.

Incidence and consequences of overweight and obesity amongst women

Overweight and obesity pose a major health problem for women. In 2023 an estimated 61 % of adult females in the UK were living with either overweight or obesity. Twenty-nine percent of these were living with obesity, which had increased from 26 % in 2013 and 23 % in 2003⁽¹⁾. Increasing rates of obesity are evident among younger females (11–15 years) where currently 20 % living with obesity compared to 17 % in the year 2013 and 6.6 % in year 2003⁽²⁾.

There is an ethnic disparity with higher rates of obesity amongst women from black Caribbean (44 %), Pakistani (33 %) and black African (37 %) backgrounds compared to women in the white British population (28 %) in 2019⁽³⁾. These obesity estimates are based on BMI > 30 kg/m² so are likely to underestimate the true impact of obesity related health conditions within ethnic groups since these groups experience obesity related comorbidities at a lower BMI than white women, that is, overweight is defined at 23–27 kg/m² and obesity at 27 kg/m² in these groups⁽⁴⁾. Women from socially deprived backgrounds experience higher rates of obesity. In 2021 40 % of women in the most deprived areas were living with obesity as compared to 19 % of women in the least deprived areas⁽⁵⁾.

Overweight and obesity amongst women are linked to higher risk of 11 cancers including breast and endometrial cancer⁽⁶⁾. Obesity poses a greater cancer burden for women amongst whom 55 % of cancers are related to obesity compared to only 24 % of cancers among men⁽⁷⁾. Obesity also contributes to other health issues such as CVD, type 2 diabetes, osteoarthritis and reproductive complications including infertility which significantly impacts their quality of life as well as increasing healthcare costs^(8,9).

Intermittent energy-restricted diets

Energy-restricted diets are the cornerstone of the management of overweight and obesity and weight related metabolic disease. Current evidence based guidelines recommend daily energy restriction (DER), typically a 600 kcal energy deficit for sustainable weight loss⁽⁴⁾. DER can be effective for weight loss but has reduced adherence over time. Also, the metabolic benefits of DER and weight loss are attenuated once weight is reduced and an individual is euenergetic, that is, in energy balance at a lower weight^(10,11). Several IER regimens have been developed which attempt to overcome these issues, that is, to promote maintained dietary adherence and weight

© The Author(s), 2025. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



loss as well as potential additional metabolic benefits beyond those seen with weight loss. Intermittent diets which include periods of limited energy intakes interspersed with normal ad lib eating have gained in popularity in the past decade. The most popular methods involve a 5:2 diet with two consecutive or non-consecutive days of low energy diet (500–850 kcal/d) and 5 d of normal eating each week, or alternate day fasting (ADF) which usually involves 3–4 d of low energy diet (500–650 kcal/d) interspersed with days of normal eating across the week. Also, time-restricted eating (TRE) where individuals fast for between 12 to 18 h and eat in a 6–12-hour window each day. TRE aims to restrict energy intake and align eating with the circadian rhythm. TRE usually involves ad lib eating during the eating window (adlib TRE), although some research has tested an energy-restricted TRE in which the daily energy-restricted diet needs to be eaten in a defined eating window rather than across the day (energy-restricted TRE). The most popular IER regimens are summarised in Figure 1.

IER is also popular amongst people who are a healthy weight who are motivated by the purported health benefits of IER for health and disease prevention. A recent US survey of 3000 Americans aged 18–80 years including 17% with overweight obesity reported IER as the third most popular diet practiced by 13% of the population⁽¹²⁾.

This paper will review evidence for potential benefits and/or harms for each of these three popular IER regimens amongst women who are living with overweight or obesity or who are a healthy weight. Also the potential utility of these diets for managing women's health conditions will be explored.

Weight loss and metabolic effects of intermittent energy restriction compared to daily energy restriction in subjects with overweight and obesity

A recent meta-analysis summarised studies which have randomised people with overweight/obesity with or without type-2 diabetes to an IER (either 5:2 or ADF) or an energy matched DER⁽¹³⁾. The review included 11 randomised controlled trials (RCTs) and 850 participants, 67% were female and 33% were male. In this review IER achieved slightly better weight loss (7%) compared with DER (5%) in short-term studies (< 6 months), but weight loss was comparable between IER and DER in longer-term studies (both ~5–6%)⁽¹³⁾. These results suggest superior short-term adherence and weight loss success with IER compared to DER. However, this is not sustained, suggesting IER is hampered by the same behavioural and physiological drivers which promote weight regain with DER, that is, satiety, hunger, reward, adipose immune cells, adaptive thermogenesis and reduced lipolysis and lipid oxidation lipid metabolism⁽¹⁴⁾.

TRE can either ad lib TRE which has an eating window but no prescribed energy restriction, or an energy-restricted TRE diet with a prescribed eating window. Ad lib TRE diets typically reduce energy intake by 200–300 kcal/d and are associated with a modest weight loss of 3% in short-term studies⁽¹⁵⁾ and ~1% in longer-term studies⁽¹⁶⁾. Studies of TRE diets *v.* matched energy-restricted diets consumed across the day have reported both superior^(17,18) or equivalent^(19,20) weight loss with TRE. These data do not confirm superiority of TRE alongside energy restriction, but show that an eating window can sometimes increase adherence and weight loss with DER.

A further question is whether IER confers additional weight independent effects on metabolic health evidenced by markers including blood pressure, lipids and insulin resistance. IER may

have short-term beneficial effects on these markers during the repeated spells of energy restriction each week. In addition, there could be more sustained beneficial effects across the week if for example there were preferential reductions in body fat and ectopic fat stores with IER *v.* DER. Current evidence does not however support this assertion. Shubel et al reported equivalent weight loss paralleled with proportional reductions in visceral and subcutaneous fat stores with IER *v.* DER⁽²¹⁾. Cioffi et al reviewed 11 RCTs of 5:2, ADF (not TRE) studies including 630 patients (range 8–24 weeks) and concluded there were small favourable differences in metabolic markers with IER *v.* DER⁽²²⁾. These effects were modest and of doubtful clinical significance, that is, IER had 0.05 mmol/l (3%) higher HDL and 0.9 mmol/l (15% greater reductions in insulin) than DER. These beneficial effects were mainly reported in short-term studies and may simply reflect slightly greater weight loss in these studies. Also, two of these studies involving 223 of the 630 (34%) of participants were testing an intermittent low carbohydrate diet which may have superior glycaemic effects⁽²²⁾.

Studies of TRE have reported modest improvements in some glycaemic parameters which are largely driven by reduced energy intakes and weight loss. Some potential weight independent metabolic benefits have been reported when TRE has an early eating window and avoids the adverse metabolic effects of late-night eating. Whilst promising, early TRE can be difficult to integrate into family/social lives⁽²³⁾.

Weight loss and dietary adherence with intermittent energy restriction in women *v.* men

Lower dietary adherence and weight loss across a range of behavioural weight loss interventions have been reported amongst women *v.* men, amongst younger *v.* older subjects⁽²⁴⁾ and amongst parents with children living at home⁽²⁵⁾. Barriers to adherence in younger women include competing demands on time, stress, multiple role expectations, fatigue, limited family support and often unhealthy home food environments which include high energy foods, snacks and drinks⁽²⁶⁾.

Consistent with this data adherence and weight loss success with 12 weeks of ADF has been reported to be lower amongst premenopausal women ($-4.6 \pm 3.2\%$) compared to postmenopausal women ($-6.5 \pm 3.2\%$) and men ($-6.2 \pm 4.4\%$)⁽²⁷⁾. Similarly, Barnowsky et al. reported lower weight loss with 6 months of ADF amongst premenopausal ($-6.0 \pm 1.1\%$) *v.* postmenopausal women $11.6 \pm 2.3\%$ ⁽²⁸⁾.

In contrast to this, Schroor undertook a systematic review and meta-analyses of 28 RCTs of 5:2, ADF and TRE diets⁽²⁹⁾. The review concluded the three different IER diets resulted in comparable weight loss and cardiometabolic risk markers change compared with DER diets. However, a sub-group analysis in 7 studies which involved women only (*n* 507) showed IER had greater effects on reductions in body weight, body fat and waist circumference than DER. Body weight (weighted mean differences WMD -1.01 kg; 95% CI: -1.52 , -0.50), body fat (WMD: -1.08 kg; 95% CI: -1.68 , -0.48) and waist circumference (WMD: -1.40 cm; 95% CI: -2.64 , -0.15). No significant differences between IER and DER were observed in studies with men only or mixed cohorts. Caution is required when interpreting cross study comparisons. The findings in women may reflect different features of the IER and DER regimens and study design in the women only studies, rather than being evidence of a gender-specific effect of IER.

5:2 diet

- Limit energy intake 2 days a week (days together or apart), eat healthily 5 days a week

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
12am	500-850 Kcal Food or total diet replacements	500-850 Kcal Food or total diet replacements	Healthy eating	Healthy eating	Healthy eating	Healthy eating	Healthy eating
4am							
8am							
12pm							
4pm							
8pm							
12am							

Ad Lib Alternate day fasting

- Limit energy intake 3-4 days a week, eat healthily 3-4 days a week

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
12am	Healthy eating	500-850 Kcal Food or total diet replacements	Healthy eating	500-850 Kcal Food and/or total diet replacements	Healthy eating	500-850 Kcal Food or total diet replacements	Healthy eating
4am							
8am							
12pm							
4pm							
8pm							
12am							

Time Restricted Eating (TRE)

16:8 (shown) or 18:6 or 12:12

Fast (no food or energy containing drinks) for 12-18 hours a day and eat normally 6-12 hours a day
Usually ad-lib / normal eating in the eating window (ad-lib TRE) but can be an energy restricted diet (Energy restricted TRE).

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
12am	Fast	Fast	Fast	Fast	Fast	Fast	Fast
4am							
8am							
12pm	Eat Normally	Eat Normally	Eat Normally	Eat Normally	Eat Normally	Eat Normally	Eat Normally
4pm							
8pm	Fast	Fast	Fast	Fast	Fast	Fast	Fast
12am							

Figure 1. Popular regimes for intermittent energy restriction regimens.

Cyclic changes in hormones in premenopausal women are likely to influence appetite and energy expenditure and dietary adherence across each month. Energy intake is often increased in the luteal phase due to cravings for high fat and/or carbohydrate foods⁽³⁰⁾, making this a potentially challenging time for adherence to a low energy diet for some women. Adherence to the different IER diets across the menstrual cycle is not known. However a menstrual cycle adapted DER weight loss programme which attempted to align with these cyclic variations has had limited success compared to a standardised DER across the month⁽³¹⁾. Premenopausal women are reported to have a greater lipolytic response and higher plasma free fatty acids with extended overnight fasting compared to men and postmenopausal women which has a negative impact on postprandial glycaemia, summarised in⁽³²⁾. However the clinical significance of this normal physiological response to fasting is unclear.

Potential harms of intermittent energy restriction in women

Effects of intermittent energy restriction on Fat-Free Mass

One concern is whether IER leads to greater losses of fat free mass (FFM) for a given weight loss than seen with DER. In people with overweight/obesity around 25 % of weight loss with daily energy-restricted diets is loss of FFM⁽³³⁾. IER could lead to greater reductions in FFM for a given weight loss as a result of spontaneous decreases in physical activity during energy-restricted periods⁽³⁴⁾, insufficient protein intakes⁽³⁵⁾, or sub-optimal regularity of protein intake to optimise muscle protein synthesis⁽³⁶⁾; the latter being particularly relevant with TRE. Greater losses of FFM with energy restriction are seen amongst subjects with lower fat mass, that is, lean compared to those with overweight/obesity, men compared to

Table 1. Loss of weight and fat free mass amongst women undertaking intermittent energy restriction v. daily energy restriction

Reference	Cohort Sample size	Body composition measurement	Duration/ weeks	IER/ DER regimens	DER		IER	
	Age BMI				Weight loss- kg	% weight loss as FFM	Weight loss- kg	% weight loss as FFM
Alternate day fasting								
Hutchison et al 2019 ⁽⁴²⁾	<i>n</i> 88 (All female) Age 50 ± 1 years, BMI 32.3 ± 0.5 kg/m ²	DEXA	3 week	Both 30 % energy restricted diets: ADF 70 % ER & energy requirements on non-fasting days DER 30 % ER	−3.9 (0.4)* (<i>n</i> 24)	19.0 (6 %)*	−5.4 (0.5)* (<i>n</i> 22)	19.0 (7.0) *
Beaulieu et al 2020 ⁽⁴³⁾	<i>n</i> 54 (All female) Age 35 ± 11 years, BMI 29 ± 2.5 kg/m ²	Air displacement plethysmography (Bod Pod)	12 week	ADF alternate days 75 % ER/ adlib DER 25 % ER	−4.0	32.5 %	−5.3	24.5 %
Coutinho et al 2017 ⁽⁴⁴⁾	<i>n</i> 35 (79 % female) Age 39.4 ± 11.0 years BMI 35.6 ± 3.2 kg/m ²	Air displacement plethysmography (Bod Pod)	12 week	Both 33 % energy restricted diets ADF alternate days 550–660 kcal formula diet & prescribed food based diet DER 33 % ER	−8.2	23.2 %	−13.9	20.1 %
Bowen et al 2018 ⁽⁴⁵⁾	<i>n</i> 164 (81 % female) Age 40.0 ± 8.3 years BMI 35.7 ± 5.9 kg/m ²	DEXA	16 week	ADF alternate days 1200 kcal and 600 kcal meal replacement and food diet & 1 ad lib eating day DER 1200 kcal meal replacement diet and prescribed food diet	−12.4	15.3 %	−12	14.0 %
5:2 diet								
Harvie et al 2011 ⁽⁴⁶⁾	<i>n</i> 107 Premenopausal women Age 40.1 ± 4.1 years BMI 30.7(5) kg/m ²	Single frequency bioelectrical Impedance (leg to leg)	26 weeks	5:2 :75 % ER 2 d (650 kcal) DER 25 % ER	−6.4 (−7.9, −4.8)**	21.0 %	−5.6 (−6.9, −4.4)**	21.0 %
Harvie 2013 et al. ⁽⁴⁷⁾	<i>n</i> 115(All female) Age 45.6 ± 8.3 years BMI 29.7 ± 4.1 kg/m ²	Multi- frequency bioelectrical impedance 5 kHz/50 kHz/250 kHz/500 kHz	16 weeks	5:2 75 % ER (650 kcal < 50 g carbohydrate) DER: 25 % ER	−5.5	36.0 (26.4, 41.3) %**	−3.8	29.3 (25.0–38.1) * %*
Time restricted eating								
Lin 2022 et al. ⁽¹⁷⁾	<i>n</i> 63 (All female) Age 50.1 ± 7.5 years BMI 25.9 ± 3.7 kg/m ²	Multi-frequency bioelectrical impedance 1, 5, 50, 250, 550, 1000 kHz	8 weeks	TRE: 1400 kcal (8 h eating window) DER: 1400 (no time restriction) Both groups 8 × 30 min exercise sessions	−1.6	12.5 %	−2.7	26.0 %

Table 1. (Continued)

Jamshed et al 2022 ⁽¹⁸⁾	n 90 (80 % female) Age 43.0 ± 10 years BMI 40.1 ± 6.6 kg/m ²	DEXA	14 weeks	TRE (500 kcal/d below their resting energy expenditure) + 8 h eating window DER – 500 kcal deficit diet no time restriction Both groups advised to exercise 75–150 min/week	–4.0	26.0 %	–6.3	22.0 %
Thomas et al 2022 ⁽¹⁹⁾	n 81 (85 % female) Age 38.0 ± 8.0 years BMI 34.1 ± 5.7 kg/m ²	DEXA	39 weeks	TRE (33 % ER + 10 h eating window within 3 hrs of waking DER: 33 % ER no time restriction Both groups advised to exercise 150 min/week	–5.1	21.6 %	–6.2	24.2 %
Ribiero et al 2022 ⁽²⁰⁾	n 24 (83 % female) Age 33.0 ± 8.7 years BMI 31.7 ± 8.6 kg/m ²	Multi-frequency bioelectrical impedance 20, 100 kHz	8 weeks	TRE (20 % ER + 8 h eating window) DER (20 % ER no time restriction) Both groups active: 3 x 20 min/week aerobics exercises + resistance training + exercises to balance and proprioception.	–6.3	22.2 %	–5.7	0.0 %

FFM = fat free mass DEXA = Dual energy X-Ray absorptiometry ADF = alternate day fasting DER daily energy restriction IER intermittent energy restriction TRE = time restricted eating * mean (SD) ** mean (95 % CI).

women, and alongside more severe energy/protein restriction⁽³⁷⁾, and in older subjects⁽³⁸⁾. Weight loss studies with IER amongst women with overweight/obesity have shown that reductions of FFM align with the weight loss achieved with comparable reductions of FFM per kg body weight reduction to DER as summarised in Table 1^(17–19,39–44).

There are no data on the effects of IER on FFM in cohorts of healthy weight women who will be more susceptible to reductions in FFM than women with overweight or obesity. However, several studies report large reductions of FFM in groups of lean men and women exposed to ADF with alternate day 24-hour fasts. Heilbronn et al studied 8 women and 8 men for 22 d who experienced a weight loss of 2.5 %, of which 57 % was FFM⁽⁴⁸⁾. Likewise, Templeman et al compared the effects of this ADF with an energy matched DER over 3 weeks in 12 lean women and 12 lean men. Mean (SD) weight loss and % of weight loss as body fat with DER were –1.91 (0.99) and 92 % compared to –1.60 (1.06) and 46 % for IER⁽³⁴⁾. These data highlight a potentially lower FFM retention amongst healthy weight subjects undertaking IER.

There are few data of the effects of IER on FFM amongst postmenopausal women who are at higher risk of developing sarcopenia⁽⁴⁹⁾. In the absence of data, it is prudent to advise adequate protein and exercise alongside IER diets. Exercise is well known to attenuate loss of FFM with energy-restricted diets⁽³⁸⁾. This has been reported alongside an ADF diet (mean age 45 years, 81 women, 3 men)⁽⁵⁰⁾. Two relatively small studies have examined whether exercise attenuates reductions in FFM with IER. Cooke et al reported that 10 min of sprint exercise three times a week was not sufficient to attenuate FFM loss that occurred alongside an intermittent 5:2 diet (mean age 35 years, 8 women, 3 men)⁽⁵¹⁾. However a study young trained women (13 TRE 13 control diet aged 18–30 years), reported that TRE eating in a 7.5 h window did not compromise accretion of FFM alongside a resistance exercise and high protein diet compared to consumption of regular meals throughout the day⁽⁵²⁾.

Effects of intermittent energy restriction on bone health

Weight loss with energy-restricted diets in individuals with overweight or obesity can reduce bone quantity, bone density and bone quality. The latter is already compromised in subjects with obesity. Bone effects may be partly through reduced mechanical loading at a reduced weight and may also relate to increased bone marrow adipose tissue and associated cytokine production and adipokines and reduced osteoblast formation⁽⁵³⁾. Weight loss has been associated with reductions in total hip bone mineral density (BMD), but not lumbar spine BMD⁽⁵⁴⁾. BMD reductions of approximately 1–1.5 % in weight loss studies are comparable to annual losses in older women, which have been associated with a 10 % to 15 % increase in fracture risk⁽⁵⁴⁾.

The effects of 5:2, ADF and TRE on bone health are not known, nor whether they differ from those of DER. IER could exert detrimental effects on bone health alongside reduced physical activity during the energy-restricted spells of intermittent diets. In addition, elevated post-prandial insulin resistance in response to the first post fast meal consumed with IER has the potential to suppress concentrations of C-terminal telopeptide and osteocalcin. The bone effects of an IER regime will also relate to its nutritional adequacy for bone health (calcium, vitamin D intake)⁽⁵⁵⁾.

One of the few data on IER and bone reported that 6 months of ADF or DER both resulted in a weight loss of 8 % and that neither diet was associated with reductions in total body dual energy x-ray

absorptiometry (DXA) measured bone mineral density, or in circulating bone turnover markers osteocalcin, bone alkaline phosphatase or C-terminal telopeptide⁽²⁸⁾. This study has limitations and is likely to be underpowered for these bone measures, and it did not collect specific hip/spine BMD responses. Powered research of the bone effects of IER are required using validated bone end points, that is, hip/lumbar spine BMD, bone microstructure and fracture risk⁽⁵⁵⁾.

Reproductive hormone levels

Energy restriction may disturb regularity of the menstrual cycle and fertility. Menstrual cycle disturbances, that is, shortened luteal phase, anovulation and/or oligomenorrhea (cycle length 36–90 d, have been observed amongst healthy weight eumenorrheic women aged 18–30 years when exposed to daily energy deficits of between 470 and 810 kcal/day (22–42 % energy restriction) over 4 menstrual cycles⁽⁵⁶⁾. Energy restriction can suppress the menstrual cycle through inhibition of gonadotropin-releasing hormone (GnRH) pulsatility. There are few data on the effect of IER on reproductive hormones and menstrual cycle function. One study of a 5:2 diet amongst premenopausal women with overweight/obesity has shown average cycle length across the dietary intervention to be slightly longer with IER than with DER, 29.7 (3.8) days vs. 27.4 (2.7) days. However there were comparable increases in sex hormone binding globulin (SHBG) +14 % v. +6 % and reductions in free androgen index –6 % v. –10 %⁽⁴⁶⁾. The increased cycle length may reflect a slightly longer follicular phase when undertaking IER. However, the clinical significance of this observation is not known. For example research to date has not linked menstrual cycle length with risk of breast cancer⁽⁵⁷⁾.

Intermittent diets and eating disorders

Another frequently cited concern is whether IER could trigger or exacerbate disordered eating. These concerns are based on a theoretical risk and cross-sectional studies which report higher eating disorder scores amongst populations undertaking IER. For example, a recent survey amongst 2762 Canadian adolescents and young adults aged 16–30 years (1477 women, mean age 23, 40 % who consider themselves overweight) reported that 47 % had engaged in IER in the past 3 and 12 months⁽⁵⁸⁾. The majority of these had undertaken TRE (80 %), 9 % ADF and 11 % other IER regimes. The group reporting IER had higher scores for overeating, loss of control, binge eating, vomiting, laxative use, compulsive eating than those not undertaking IER⁽⁵⁸⁾. Likewise, a survey which included 40 women undertaking TRE (fasting for > or = 16 h/d, mean age 33, mean BMI 27 kg/m²) reported that those engaging with IER had higher eating disorder scores for binge eating, vomiting and laxative use than community and clinical norms⁽⁵⁹⁾. Approximately a third of this cohort reported eating disorder symptom scores above the diagnostic cut offs. These cross-sectional studies do not allow us to ascertain the direction of the relationship between IER and eating disorders. It is likely that IER is not a cause of these symptoms, but that those with eating disorder traits may be more likely to undertake IER. It is also possible that individuals with eating disorder traits may be attracted to volunteer for studies which focus on IER. For future studies, it would be prudent to conduct baseline screening using validated eating disorder scales⁽⁶⁰⁾ so these subjects can be excluded from studies and directed to appropriate support where relevant.

In contrast, prospective intervention studies of IER in those with overweight and obesity have shown either no change or reductions in eating disorder scores alongside weight loss. A systematic review of 4 studies of TRE (194 participants, 71 % women) concluded TRE had neither beneficial or adverse effects on disordered eating⁽⁶¹⁾. Whilst an 8 week ADF intervention amongst women with overweight and obesity reported reduced scores for depression, binge eating and concern about body size/shape assessed with the Multidimensional Assessment of Eating-Disorder Symptoms alongside mean (SD) % weight loss of 4.2 (0.3) %⁽⁶²⁾. The lack of effect or reductions in eating disorders reported with IER is consistent with reports with weight loss from other behavioural weight management programmes⁽⁶³⁾.

Intermittent energy restriction and sleep

Sleep quality is a key factor for health and well-being. Women may be more predisposed to disturbed sleep than men associated with fluctuating hormone levels. During the luteal phase women can experience more daytime sleepiness, decreased sleep efficiency and difficulty initiating sleep. Sleep disturbance is well documented during the menopause and reported by around 60 % of women. This hormonal predisposition can be exacerbated by social factors with many working women undertaking domestic tasks and childcare⁽⁶⁴⁾.

Intermittent diets have the potential to have either beneficial or adverse effects on sleep quality. TRE may have a beneficial effect on sleep if the eating window avoids bedtime eating /snacking which can disturb sleep⁽⁶⁵⁾. Alternatively, IER regimens with an overall energy restriction may disturb sleep if people experience hunger at bedtime and during the nighttime⁽⁶⁶⁾.

A recent review of TRE and sleep reported no effect of overall sleep quality assessed with the Pittsburgh Sleep Quality Index, but it included some reports of reduced sleep efficiency with both a late TRE (eating window 12.00–20.00) and an early TRE (eating window 07.00–15.00) and reduced sleep duration and sleep onset latency with an early TRE (eating window 07.00–15.00)⁽⁶⁷⁾. Studies of ADF in all female⁽⁶⁸⁾ and predominately female cohorts (81 %) have shown no impact on sleep quality assessed with the PSQI. This aligns to data from a range of behavioural diet and exercise weight loss interventions using DER showing no effects on sleep⁽⁷⁰⁾. Improved sleep with weight loss in these studies may not be seen since many participants already have good baseline sleep scores^(67–69). One study reported that TRE can reduce sleep disturbance amongst shift workers with circadian rhythm misalignment and disturbed sleep⁽⁷¹⁾. Although it did not impact on the other measurements of sleep quality, that is, latency, daytime dysfunction, efficiency, overall quality, need for medication. Future studies should focus on the effects of IER in those with circadian rhythm misalignment and explore whether TRE has potential adverse effects on sleep in those with good baseline sleeping patterns.

Intermittent diets and polycystic ovary syndrome

An estimated 3.5 % of women in the UK have polycystic ovary syndrome (PCOS), which is increasingly linked to increasing rates of overweight and obesity⁽⁷²⁾. Women with PCOS have hyper-androgenism, sub-fertility and are at increased risk of type-2 diabetes and cardiovascular events. First-line treatment involves diet, weight loss and exercise. Li et al tested the effects of 5 weeks of an 8 h TRE amongst 18 premenopausal women with overweight/obesity and PCOS⁽⁷³⁾. There was a weight loss of 2 % over 5 weeks

Table 2. Summary of research findings and gaps in research of intermittent energy restriction v. dDaily energy restriction for weight control and metabolic health for women

Outcome	Effects in women who are overweight and obese		Effects in women who are a healthy weight	
	Current evidence	Research gaps	Current evidence	Research gaps
Weight loss	ADF/ 5:2 diets are superior to DER in short-term studies (< 6 months) but equivalent in longer-term studies (≥ 6 months) Dietary adherence & weight loss success is reduced with ADF/ 5:2 diets and DER amongst younger women.	Need effective strategies for weight management with DER or IER for young women	No evidence	Are there potential benefits of IER for metabolic health or weight gain prevention?
Potential harms of IER				
Bone health	Limited data	Require powered research with validated bone end points, i.e. hip/lumbar spine BMD, bone microstructure and fracture risk	No evidence	What are the effects of IER on bone health in women who are a healthy weight?
Reproductive hormones	No evidence of harm but minimal data	Studies of IER should collect this information	No evidence	Does IER affect reproductive hormones in lean individuals who are more susceptible to the hormonal effects of energy restriction?
Eating disorders	Either no effect or beneficial effect amongst overweight/ obese alongside weight loss	Studies of IER should collect this information	No evidence	Does IER lead to disordered eating in healthy weight individuals?
Sleep quality	Some reports of decreased sleep efficiency and sleep onset latency with TRE. No effect on sleep with ADF which aligns with findings from DER weight loss interventions.	Need more evidence in those with circadian misalignment and those with current good sleep. Also, populations at high risk of disturbed sleep, i.e. menopausal women	No evidence	What are the effects of IER on sleep in women who are a healthy weight?
Benefits to specific women's health conditions	Weight loss with TRE improved PMS Weight loss with ADF improved PCOS These IER diets are feasible alternative weight loss diets in these populations. However, there is no evidence of weight independent effects of IER on these conditions The effects of IER on GDM is under study.	Need randomised trials to compare the effects of IER to DER for these conditions	No evidence	Require more research of the effects of IER on specific women's health conditions amongst women who are a healthy weight

PMS = Premenstrual syndrome PCOS = polycystic ovarian syndrome GDM = gestational diabetes IER= intermittent energy restriction DER = Daily energy restriction TRE = Time restricted eating ADF = Alternate day fasting BMD = Bone mineral density.

alongside reductions in testosterone (9 %) increased SHBG (2 %) and decreased free androgen index (26 %). These positive results provide some evidence that TRE is a potential weight loss diet for women with PCOS. However, the reported hormonal effects are in line with those expected alongside the weight loss seen and do not verify a weight independent effect of TRE for those with PCOS.

Intermittent diets and premenstrual syndrome

Weight control is a potential strategy for management of premenstrual syndrome⁽⁷⁴⁾. Hooshier et al conducted a small short-term trial assessed the impact of ADF (alternate days of 75 % energy restriction and consuming estimated energy requirements) compared to a matched DER on premenstrual symptoms amongst 60 women with overweight or obesity⁽⁷⁵⁾. The ADF diet group reported reductions in some premenstrual syndrome traits, that is, mood lability and expressed anger, but no global reduction in premenstrual syndrome scores. These beneficial changes were seen alongside greater percentage weight loss in the ADF group v. DER (−6.7 % v. −3.7 %). The study shows greater short-term weight loss

success with ADF v. DER in this population but is unable to conclude there is a weight independent effect of ADF on symptoms of premenstrual syndrome.

Intermittent diets and gestational diabetes

Gestational diabetes (GDM) occurs in between 1 and 25 % of pregnancies worldwide affecting an estimated 16 % of pregnancies in the UK⁽⁷⁶⁾. Rates are rising linked to higher levels of obesity and maternal age. GDM and poor glycaemic control lead to maternal and neonatal complications, that is, macrosomia, shoulder dystocia, neonatal hypoglycaemia and/or hyperbilirubinemia, preterm delivery, caesarean-sections, preeclampsia and stillbirth. Diet and exercise are first line therapies for the management of GDM⁽⁷⁷⁾. Metformin and/or insulin are utilised if blood glucose targets are not met by changes to diet and exercise. UK NICE guidance promotes healthy eating and promotes low glycaemic index foods and physical activity but has no specific guidance for energy restriction. Other countries advocate carbohydrate and energy restriction, and optimal timing of meals and many include

targets for limiting gestational weight gain (GWG). There are no specific targets for GWG for GDM. Many GDM guidelines follow the Institute of Medicine Guidelines (USA) for GWG which recommend lower weight amongst women with overweight or obesity, that is, e. healthy weight women weight (BMI 18.5–24.9 kg/m²) should gain 11.5–16 kg (25–35 pounds) during pregnancy. Overweight women (BMI 25–29.9 kg/m²) should gain 7–11.5 kg (15–25 pounds) and obese women (BMI greater than 30 kg/m²) should only put on 5–9 kg (11–20 pounds)⁽⁷⁷⁾. A range of dietary approaches have been studied for the management of GDM including low-GI diets (limiting refined and promoting complex carbohydrates), low carbohydrate diets or modest energy-restriction (1800 kcal/d)⁽⁷⁷⁾. However, there is currently no strong evidence to recommend one dietary regimen to improve outcomes in GDM. IER has potential utility for the management of GDM. GDM is a strongly associated with obesity and insulin resistance and is considered a form of evolving Type 2 diabetes. IER has been associated with greater or equivalent reductions in weight and hyperglycaemia in patients with Type 2 diabetes when compared to DER^(22,39), and so could potentially also have benefits for GDM. Dietary management of GDM typically requires a 10–12-week dietary intervention in the third trimester. Since 5:2 or ADF diets appear to have a greater adherence and weight loss than daily diets in the short-term (< 3 months)⁽¹³⁾, they could have superior adherence to DER in the required short-term treatment period.

Our team is currently undertaking a feasibility study (*n* 48) to test the safety, feasibility and acceptability of a 5:2 diet between diagnosis of GDM and delivery in our MIDDAS – GDM study (Trial Registration Number: NCT053440660). The 5:2 diet involves 2 non-consecutive days of 1000 kcal which includes 100 g low-GI carbohydrate and 70 g of protein and 5 d/week of the NICE healthy eating low-GI diet and physical activity recommended for the best NHS care group⁽⁴⁰⁾.

The benefits of a 5:2 diet *v.* DER have been tested over 12 months amongst non-pregnant women with a history of GDM. The 5:2 diet included 2 non-consecutive days of 500 kcal to include 45 g low-GI carbohydrate and 50 g of protein and 5 d/week of their habitual diet *vs.* a daily 25 % energy-restricted diet which typically included 1500 kcal (6000 kJ) per day (170 g carbohydrate, 110 g protein)⁽⁴¹⁾. There was an equivalent high drop-out (49 %) and modest weight loss both groups (IER, 5.0 ± 5.4 %; DER, 3.5 ± 5.6 %; *P* = 0.3). These results reflect the challenge of weight control within the GDM population and that this challenge was not overcome by those attempting to follow IER.

Conclusion

Research findings and gaps in research of IER and DER amongst women is summarised in Table 2. IER (5:2 or ADF) diets are an equivalent (but not superior) approach to DER for weight loss and managing weight-related metabolic conditions for women with overweight and obesity. Adherence to behavioural weight control interventions can be a particular challenge for younger women, which is seen with both DER and IER diets. IER has some theoretical harms for women affecting reproductive health, bone health, disordered eating and sleep, but there are few data to support or refute these.

Weight loss with ADF and TRE has respectively improved symptoms of polycystic ovarian syndrome and premenstrual syndrome, but there is no evidence of weight independent effects of IER on these conditions. Thus, IER is a possible weight loss strategy for these conditions, but it is not currently advocated as a strategy

for treating/managing these conditions. IER is a popular diet amongst women who are healthy weight, but there are few data to inform its potential benefit or harms which needs to be a focus of ongoing IER research.

Acknowledgements. Professor Tony Howell who was a key investigator on the Manchester Intermittent Diet studies. Research dietitians and project management teams who supported the Manchester Intermittent studies; Dr Mary Pegington, Cheryl Lombardelli, Sarah McDiarmid, Grace Cooper, Kath Sellers, Suzy Krizak and Helen Ruane. Also, co-investigators working on the MIDDAS-GDM (gestational diabetes) study; Basil Issa, Liz Dapre, Avni Vyas, Brian Mcmillan, Helen Morley and Emma Thorpe for proofreading the manuscript.

Authorship. M Harvie devised the concept of the paper. M Harvie and M Haiba wrote the original draft. Both authors approved the final manuscript.

Financial support. Michelle Harvie receives part funding from the NIHR Manchester Biomedical Research Centre award NIHR203308). Mai Haiba is undertaking a PhD funded by the NIHR Manchester Biomedical Research Centre (award NIHR203308). The funders had no role in the design, conduct, analysis or write up of this review paper.

Competing interests. Michelle Harvie has co-authored three self-help books for the public to follow intermittent diets. All author proceeds are paid directly to the charity Prevent Breast Cancer (Registered Charity Number 1109839) to fund breast cancer research.

References

- House of Commons Library (2023) Obesity Statistics. <https://researchbriefings.files.parliament.uk/documents/SN03336/SN03336.pdf> (accessed 23 September 2024).
- Department of Health and Social Care (2024) Obesity Profile. <https://fingertips.phe.org.uk/profile/national-child-measurement-programme> (accessed 23 September 2024).
- NHS Digital (2022) Health Survey England Additional Analyses, Ethnicity and Health, 2011–2019 Experimental Statistics. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-england-additional-analyses/ethnicity-and-health-2011-2019-experimental-statistics/overweight-and-obesity> (accessed 23 September 2024).
- National Institute of Clinical Excellence (2021) Obesity: Identification, Assessment and Management Clinical Guideline [CG189 National Institute of Clinical Excellence]. <https://www.nice.org.uk/guidance/cg189#:~:text=This%20guideline%20covers%20identifying,%20assessing%20and%20managing%20obesity%20in%20children> (accessed 23 September 2024).
- NHS Digital (2021) Health Survey England, Part 1. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/latest-statistics> (accessed 23 September 2024).
- Rehnan AG, Tyson M, Egger M, *et al.* (2008) Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 371, 569–578.
- Argyropoulou G, Dalamaga M, Spyrou N, *et al.* (2021) Gender differences in obesity-related cancers. *Curr Obes Rep* 10, 100–115.
- Harrison SM, Hughes LA & Chang S (2021) Obesity, cardiovascular disease, and diabetes: insights from recent research. *Lancet Diabetes Endocrinol* 9, 408–423.
- Shulman RJ, McMahon M & Newnham T (2021) The impact of obesity on osteoarthritis and reproductive health: recent evidence. *J Women's Health* 30, 1502–1511.
- Wing RR, Blair EH, Bononi P, *et al.* (1994) Caloric restriction per se is a significant factor in improvements in glycemic control and insulin sensitivity during weight loss in obese NIDDM patients. *Diabetes Care* 17, 30–36.
- Henry RR, Scheaffer L & Olefsky JM (1985) Glycemic effects of intensive caloric restriction and isocaloric refeeding in noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 61, 917–925.

12. International Food Information Council (2024) Food and Health Survey. <https://foodinsight.org/wp-content/uploads/2024/06/2024-IFIC-Food-Health-Survey.pdf> (accessed 28 August 2024).
13. He S, Wang J, Zhang J, *et al.* (2021) Intermittent v. continuous energy restriction for weight loss and metabolic improvement: a meta-analysis and systematic review. *Obesity (Silver Spring)* **29**, 108–115.
14. van Baak MA & Mariman ECM (2023) Obesity-induced and weight-loss-induced physiological factors affecting weight regain. *Nat Rev Endocrinol* **19**, 655–670.
15. Jefcoate PW, Robertson MD, Ogden J, *et al.* (2023) Exploring rates of adherence and barriers to time-restricted eating. *Nutrients* **15**, 2336.
16. de Oliveira Maranhão Puraiza IR, da Silva Junior AE, Silva Praxedes DR, *et al.* (2021) Effects of time-restricted feeding on body weight, body composition and vital signs in low-income women with obesity: a 12-month randomized clinical trial. *Clin Nutr* **40**, 759–766.
17. Lin YJ, Wang YT, Chan LC, *et al.* (2022) Effect of time-restricted feeding on body composition and cardio-metabolic risk in middle-aged women in Taiwan. *Nutrition* **93**, 111504.
18. Jamshed H, Steger FL, Bryan DR, *et al.* (2022) Effectiveness of early time-restricted eating for weight loss, fat loss, and cardiometabolic health in adults with obesity: a randomized clinical trial. *JAMA Intern Med* **182**, 953–962.
19. Thomas EA, Zaman A, Sloggett KJ, *et al.* (2022) Early time-restricted eating compared with daily caloric restriction: a randomized trial in adults with obesity. *Obesity (Silver Spring)* **30**, 1027–1038.
20. Ribeiro D, Santiago AF & Cesar de Abreu W (2021) Continuous energy restriction (CER) plus 16/8 time-restricted feeding improve body composition and metabolic parameters in overweight and obese, but no more than CER alone. *Nutr Healthy Aging* **6**, 147–156.
21. Schübel R, Nattenmüller J, Sookthai D, *et al.* (2018) Effects of intermittent and continuous calorie restriction on body weight and metabolism over 50 weeks: a randomized controlled trial. *Am J Clin Nutr* **108**, 933–945.
22. Cioffi I, Evangelista A, Ponzio V, *et al.* (2018) Intermittent v. continuous energy restriction on weight loss and cardiometabolic outcomes: a systematic review and meta-analysis of randomized controlled trials. *J Transl Med* **16**, 371.
23. Chang Y, Du T, Zhuang X, *et al.* (2024) Time-restricted eating improves health because of energy deficit and circadian rhythm: a systematic review and meta-analysis. *iScience* **27**, 109000.
24. Chopra S, Malhotra A, Ranjan P, *et al.* (2021) Predictors of successful weight loss outcomes amongst individuals with obesity undergoing lifestyle interventions: a systematic review. *Obes Rev* **22**, e13148.
25. Bramante CT, Thornton RLJ, Pilla SJ, *et al.* (2020) Parents lose less weight than nonparents in an intensive lifestyle intervention. *Obes Sci Pract* **6**, 494–506.
26. Burgess E, Hassmén P & Pumpa KL (2017) Determinants of adherence to lifestyle intervention in adults with obesity: a systematic review. *Clin Obes* **7**, 123–135.
27. Lin S, Lima Oliveira M, Gabel K, *et al.* (2021) Does the weight loss efficacy of alternate day fasting differ according to sex and menopausal status? *Nutr Metab Cardiovasc Dis* **31**, 641–649.
28. Barnosky A, Kroeger CM, Trepanowski JF, *et al.* (2017) Effect of alternate day fasting on markers of bone metabolism: an exploratory analysis of a 6-month randomized controlled trial. *Nutr Healthy Aging* **4**, 255–263.
29. Schroor MM, Joris PJ, Plat J, *et al.* (2024) Effects of intermittent energy restriction compared with those of continuous energy restriction on body composition and cardiometabolic risk markers - a systematic review and meta-analysis of randomized controlled trials in adults. *Adv Nutr* **15**, 100130.
30. Davidsen L, Vistisen B & Astrup A (2007) Impact of the menstrual cycle on determinants of energy balance: a putative role in weight loss attempts. *Int J Obes (Lond)* **31**, 1777–1785.
31. Geiker NR, Ritz C, Pedersen SD, *et al.* (2016) A weight-loss program adapted to the menstrual cycle increases weight loss in healthy, overweight, premenopausal women: a 6-month randomized controlled trial. *Am J Clin Nutr* **104**, 15–20.
32. Rius-Bonet J, Macip S, Closa D, *et al.* (2024) Intermittent fasting as a dietary intervention with potential sexually dimorphic health benefits. *Nutr Rev* **83**, e635–e648.
33. Willoughby D, Hewlings S & Kalman D (2018) Body composition changes in weight loss: strategies and supplementation for maintaining lean body mass, a brief review. *Nutrients* **10**, 1876.
34. Templeman I, Smith HA, Chowdhury E, *et al.* (2021) A randomized controlled trial to isolate the effects of fasting and energy restriction on weight loss and metabolic health in lean adults. *Sci Transl Med* **13**, eabd8034.
35. Lowe DA, Wu N, Rohdin-Bibby L, *et al.* (2020) Effects of time-restricted eating on weight loss and other metabolic parameters in women and men with overweight and obesity: the TREAT randomized clinical trial. *JAMA Intern Med* **180**, 1491–1499.
36. Areta JL, Burke LM, Ross ML, *et al.* (2013) Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J Physiol* **591**, 2319–2331.
37. Forbes GB (1987) Lean body mass-body fat interrelationships in humans. *Nutr Rev* **45**, 225–231.
38. Heymsfield SB, Gonzalez MC, Shen W, *et al.* (2014) Weight loss composition is one-fourth fat-free mass: a critical review and critique of this widely cited rule. *Obes Rev* **15**, 310–321.
39. Wang X, Li Q, Liu Y, *et al.* (2021) Intermittent fasting v. continuous energy-restricted diet for patients with type 2 diabetes mellitus and metabolic syndrome for glycemic control: a systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract* **179**, 109003.
40. Dapre E, Issa BG, Harvie M, *et al.* (2024) Manchester Intermittent Diet in Gestational Diabetes Acceptability Study (MIDDAS-GDM): a two-arm randomised feasibility protocol trial of an intermittent low-energy diet (ILED) in women with gestational diabetes and obesity in Greater Manchester. *BMJ Open* **14**, e078264.
41. Gray KL, Clifton PM & Keogh JB (2021) The effect of intermittent energy restriction on weight loss and diabetes risk markers in women with a history of gestational diabetes: a 12-month randomized control trial. *Am J Clin Nutr* **114**, 794–803.
42. Hutchison AT, Liu B, Wood RE, *et al.* (2019) Effects of intermittent v. continuous energy intakes on insulin sensitivity and metabolic risk in women with overweight. *Obesity (Silver Spring)* **27**, 50–58.
43. Beaulieu K, Casanova N, Oustric P, *et al.* (2020) Matched weight loss through intermittent or continuous energy restriction does not lead to compensatory increases in appetite and eating behavior in a randomized controlled trial in women with overweight and obesity. *J Nutr* **150**, 623–633.
44. Coutinho SR, Halset EH, Gåsbakk S, *et al.* (2018) Compensatory mechanisms activated with intermittent energy restriction: a randomized control trial. *Clin Nutr* **37**, 815–823.
45. Bowen J, Brindal E, James-Martin G, *et al.* (2018) Randomized trial of a high protein, partial meal replacement program with or without alternate day fasting: similar effects on weight loss, retention status, nutritional, metabolic, and behavioral outcomes. *Nutrients* **10**, 1145.
46. Harvie MN, Pegington M, Mattson MP, *et al.* (2011) The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes (Lond)* **35**, 714–727.
47. Harvie M, Wright C, Pegington M, *et al.* (2013) The effect of intermittent energy and carbohydrate restriction v. daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br J Nutr* **110**, 1534–1547.
48. Heilbronn LK, Smith SR, Martin CK, *et al.* (2005) Alternate-day fasting in nonobese subjects: effects on body weight, body composition, and energy metabolism. *Am J Clin Nutr* **81**, 69–73.
49. Messier V, Rabasa-Lhoret R, Barbat-Artigas S, *et al.* (2011) Menopause and sarcopenia: a potential role for sex hormones. *Maturitas* **68**, 331–336.
50. Bhutani S, Klempel MC, Kroeger CM, *et al.* (2013) Alternate day fasting and endurance exercise combine to reduce body weight and favorably alter plasma lipids in obese humans. *Obesity (Silver Spring)* **21**, 1370–1379.
51. Cooke MB, Deasy W, Ritenis EJ, *et al.* (2022) Effects of intermittent energy restriction alone and in combination with sprint interval training on body composition and cardiometabolic biomarkers in individuals with overweight and obesity. *Int J Environ Res Public Health* **19**, 7969.

52. Tinsley GM, Moore ML, Graybeal AJ, *et al.* (2019) Time-restricted feeding plus resistance training in active females: a randomized trial. *Am J Clin Nutr* **110**, 628–640.
53. Rosen CJ & Horowitz MC (2023) Nutrient regulation of bone marrow adipose tissue: skeletal implications of weight loss. *Nat Rev Endocrinol* **19**, 626–638.
54. Zibellini J, Seimon RV, Lee CM, *et al.* (2015) Does diet-induced weight loss lead to bone loss in overweight or obese adults? A systematic review and meta-analysis of clinical trials. *J Bone Miner Res* **30**, 2168–2178.
55. Clayton DJ, Varley I & Papageorgiou M (2023) Intermittent fasting and bone health: a bone of contention? *Br J Nutr* **130**, 1487–1499.
56. Williams NI, Leidy HJ, Hill BR, *et al.* (2015) Magnitude of daily energy deficit predicts frequency but not severity of menstrual disturbances associated with exercise and caloric restriction. *Am J Physiol Endocrinol Metab* **308**, E29–39.
57. Olsson HL & Olsson ML (2020) The menstrual cycle and risk of breast cancer: a review. *Front Oncol* **10**, 21.
58. Ganson KT, Cuccolo K, Hallward L, *et al.* (2022) Intermittent fasting: describing engagement and associations with eating disorder behaviors and psychopathology among Canadian adolescents and young adults. *Eat Behav* **47**, 101681.
59. Cuccolo K, Kramer R, Petros T, *et al.* (2022) Intermittent fasting implementation and association with eating disorder symptomatology. *Eat Disord* **30**, 471–491.
60. Schaefer LM, Crosby RD & Machado PPP (2021) A systematic review of instruments for the assessment of eating disorders among adults. *Curr Opin Psychiatry* **34**, 543–562.
61. Vizthum D, Katz SE & Pacanowski CR (2023) The impact of time restricted eating on appetite and disordered eating in adults: a mixed methods systematic review. *Appetite* **183**, 106452.
62. Hoddy KK, Kroeger CM, Trepanowski JF, *et al.* (2015) Safety of alternate day fasting and effect on disordered eating behaviors. *Nutr J* **14**, 44.
63. Jebeile H, Libesman S, Melville H, *et al.* (2023) Eating disorder risk during behavioral weight management in adults with overweight or obesity: a systematic review with meta-analysis. *Obes Rev* **24**, e13561.
64. Andersen ML, Hachul H, Ishikura IA, *et al.* (2023) Sleep in women: a narrative review of hormonal influences, sex differences and health implications. *Front Sleep* **2**, 1271827.
65. Simon SL, Blankenship J, Manoogian ENC, *et al.* (2022) The impact of a self-selected time restricted eating intervention on eating patterns, sleep, and late-night eating in individuals with obesity. *Front Nutr* **9**, 1007824.
66. Steger FL, Jamshed H, Bryan DR, *et al.* (2023) Early time-restricted eating affects weight, metabolic health, mood, and sleep in adherent completers: a secondary analysis. *Obesity (Silver Spring)* **31**, 96–107.
67. Bohlman C, McLaren C, Ezzati A, *et al.* (2024) The effects of time-restricted eating on sleep in adults: a systematic review of randomized controlled trials. *Front Nutr* **11**, 1419811.
68. Teong XT, Hutchison AT, Liu B, *et al.* (2021) Eight weeks of intermittent fasting v. calorie restriction does not alter eating behaviors, mood, sleep quality, quality of life and cognitive performance in women with overweight. *Nutr Res* **92**, 32–39.
69. Kalam F, Gabel K & Cienfuegos S (2021) Alternate day fasting combined with a low carbohydrate diet: effect on sleep quality, duration, insomnia severity and risk of obstructive sleep apnea in adults with obesity. *Nutrients* **13**, 211.
70. Knowlden AP, Ottati M, McCallum M, *et al.* (2024) The relationship between sleep quantity, sleep quality and weight loss in adults: a scoping review. *Clin Obes* **14**, e12634.
71. Manoogian ENC, Zadourian A, Lo HC, *et al.* Feasibility of time-restricted eating and impacts on cardiometabolic health in 24-h shift workers: the Healthy Heroes randomized control trial. *Cell Metab* 2022; **34**, 1442–1456.e7.
72. Berni TR, Morgan CL & Rees DA (2024) Rising incidence, health resource utilization and costs of Polycystic Ovary Syndrome in the United Kingdom. *J Clin Endocrinol Metab*, 1–10. Advance access publication 25 July 2024.
73. Li C, Xing C, Zhang J, *et al.* (2021) Eight-hour time-restricted feeding improves endocrine and metabolic profiles in women with anovulatory polycystic ovary syndrome. *J Transl Med* **19**, 148.
74. Bertone-Johnson ER, Hankinson SE, Willett WC, *et al.* (2010) Adiposity and the development of premenstrual syndrome. *J Womens Health (Larchmt)* **19**, 1955–1962.
75. Hooshiair SH, Yazdani A & Jafarnejad S (2023) Does an alternate-day modified fasting diet improve premenstrual syndrome symptoms and health-related quality of life in obese or overweight women with premenstrual syndrome? A randomized, controlled trial. *Front Nutr* **10**, 1298831.
76. Zhu Y & Zhang C (2016) Prevalence of gestational diabetes and risk of progression to type 2 diabetes: a Global Perspective. *Curr Diab Rep* **16**, 7.
77. Kapur K, Kapur A & Hod M (2021) Nutrition management of gestational diabetes mellitus. *Ann Nutr Metab* 1–13.