European Psychiatry S211

Conclusions: This study highlights the belief-in-change of negative emotions as a key mechanism underlying the impact of self-compassion, and has important implications for research and interventions aimed at promoting youth mental health.

Disclosure of Interest: None Declared

Mental Health Policies

EPP163

Deinstitutionalization in Georgia-why it is so slow

N. Zavradashvili¹* and G. Matiashvili¹

¹School of Health Sciences, The University of Georgia, Tbilisi, Georgia *Corresponding author.

doi: 10.1192/j.eurpsy.2025.491

Introduction: Mental health reform represents one of the most transformative changes in the field of healthcare, as it not only changes the forms of services, but also the nature of services offered. While many countries have successfully implemented such reforms, others, including Georgia, have struggled with a protracted and inconsistent process. Despite decades of advocacy by professionals for deinstitutionalization psychiatric hospital treatment continues to dominate in Georgia's mental healthcare system.

Objectives: The purpose of the review is to explore the concept of deinstitutionalization within the mental health landscape and assess its status in the context of Georgia. It aims to study the lessons learned from successful deinstitutionalization and illuminate achievements and challenges surrounding deinstitutionalization in Georgia's reality.

Methods: A qualitative analysis including desk review, in-depth interviews and focus group discussions was conducted. Proceeding from the research objectives we analyzed the existing legislation, strategic documents and clinical practices concerning individuals with mental disorders; Interviews were also conducted with key informants on the shortcomings and problems in deinstitutionalization practices

Results: The review findings reveal, that despite recent progress such as the development of community mobile teams and increased funding allocated for community services within mental health budget, a number of issues still remain a problem: there is no agreement among stakeholders on how to restructure existing hospital beds and financial provisions remains unresolved. The field of mental health in Georgia suffers from a lack of human resources. Attracting new personnel, ensuring regional distribution, and enhancing qualifications are necessary components of deinstitutionalization that require the involvement of all stakeholders, coordinated and time-planned action. The current mental healthcare system in Georgia is characterized by a lack of coordination and collaboration among its various components. Establishing patient care pathways between services with clear referral criteria is crucial for improving the efficiency of mental health services.

Conclusions: This research highlights that successful deinstitutionalization requires additional funds, time, and trained people. Institutions should have a long-term (3-5) year development plan, detailing the source of funding, activities to be implemented, and expected outcomes. In the absence of such a plan, progress remains sporadic, intermittent, uncoordinated, and less effective.

By addressing identified challenges and promoting coordination among mental health components, Georgia can guide a more effective course toward a community-based, patient-centered mental healthcare system.

Disclosure of Interest: None Declared

Obsessive-Compulsive Disorder

EPP164

The effects of DBS treatment on OCD symptoms in pediatric and young adult patients with myoclonusdystonia: a case series

X. Acosta-Villalobos¹∗, B. Pérez-Dueñas¹,

L. Dougherty-De Miguel¹, A. Bescós¹, J. A. Ramos-Quiroga¹, M. Loredo-Silos¹, I. Delgado-Álvarez¹ and G. Español-Martín¹

¹Hospital Universitari Vall d'Hebron, Barcelona, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.492

Introduction: Myoclonus-dystonia syndrome (MDS) is a rare childhood-onset movement disorder characterized by the combination of myoclonus and dystonia. It is caused by loss of function of the epsilon-sarcoglycan gene (*SGCE*) due to mutations, although the pathogenic mechanism is not fully understood. Deep brain stimulation is used in patients that are refractory to pharmacological treatment. A higher prevalence of psychiatric disorders has been observed in MDS patients, especially obsessive-compulsive disorder (OCD) (Timmers *et al.* Parkinsonism Relat Disord. 2019; 69 85-90). A dysfunction of cortico-basal ganglia-thalamo-cortical (CBGTC) loops is present in both conditions, but the mechanism by which they display high comorbidity is unknown.

Objectives: To determine the presence of OCD comorbidity in pediatric and young adult patients that are candidates for DBS treatment for motor disorders, and the effects that the treatment may have on OCD symptoms.

Methods: This study is part of an ongoing cohort (n = 34) of children and adolescents (at the time of diagnosis) receiving internal globus pallidus DBS as treatment for motor disorders in Vall d'Hebron Hospital. The candidates go through a psychiatric evaluation before the procedure, including the diagnostic interview K-SADS, to diagnose psychiatric comorbidity. Long-term evaluation is performed by subsequent visits to the specialist.

Results: Of the seven patients with *SGCE*+ MDS that underwent DBS treatment, four of them presented OCD symptoms before the procedure, two did not, and one did not receive an evaluation. Of the patients with OCD symptoms, all four of them showed significant improvement in motor function after DBS but had worsening of OCD symptoms. Some of the patients also had a worsening of other psychiatric symptoms, such as anxiety, mood and behavioral symptoms.

Conclusions: OCD symptoms are present in a significant proportion of our patients with *SGCE*+ MDS. OCD and other psychiatric symptoms worsened during follow-up in our patients, although it is not clear if this is due to DBS treatment or due to the natural course of the disorder. Further studies are needed to explore this observation.

Disclosure of Interest: None Declared