

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. **Tables** Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLE

Interventions for enhancing return to work in individuals with a common mental illness: systematic review and meta-analysis of randomized controlled trials

Nigatu YT, Liu Y, Uppal M, McKinney S, Rao S, Gillis K & Wang J 3263

ORIGINAL ARTICLES

Cognitive remediation for individuals with psychosis: efficacy and mechanisms of treatment effects

Fiszdon JM, Choi KH, Bell MD, Choi J & Silverstein SM 3275

Cost-effectiveness of focal psychodynamic therapy and enhanced cognitive-behavioural therapy in out-patients with anorexia nervosa

Egger N, Wild B, Zipfel S, Junne F, Konnopka A, Schmidt U, de Zwaan M, Herpertz S, Zeeck A, Löwe B, von Wietersheim J, Tagay S, Burgmer M, Dinkel A, Herzog W & König H-H 3291

Prevalence of serum anti-neuronal autoantibodies in patients admitted to acute psychiatric care

Schou M, Sæther SG, Borowski K, Teegen B, Kondziella D, Stoecker W, Vaaler A & Reitan SK 3303

Recent trends in primary-care antidepressant prescribing to children and young people: an e-cohort study

John A, Marchant AL, Fone DL, McGregor JL, Dennis MS, Tan JOA & Lloyd K 3315

Prospective associations between televiewing at toddlerhood and later self-reported social impairment at middle school in a Canadian longitudinal cohort born in 1997/1998

Pagani LS, Lévesque-Seck F & Fitzpatrick C 3329

Childhood trauma, psychosis liability and social stress reactivity: a virtual reality study

Veling W, Counotte J, Pot-Kolder R, van Os J & van der Gaag M 3339

Reactivity to uncertain threat as a familial vulnerability factor for alcohol use disorder

Gorka SM, Hee D, Lieberman L, Mittal VA, Phan KL & Shankman SA 3349

Network analysis of depression and anxiety symptom relationships in a psychiatric sample

Beard C, Millner AJ, Forgeard MJC, Fried EI, Hsu KJ, Treadway MT, Leonard CV, Kertz SJ & Björgvísson T 3359

Casting wider nets for anxiety and depression: disability-driven cross-diagnostic subtypes in a large cohort

Wanders RBK, van Loo HM, Vermunt JK, Meijer RR, Hartman CA, Schoevers RA, Wardenaar KJ & de Jonge P 3371

Salience attribution and its relationship to cannabis-induced psychotic symptoms

Bloomfield MAP, Mouchlianitis E, Morgan CJA, Freeman TP, Curran HV, Roiser JP & Howes OD 3383

Mental illness and suicide after self-harm among young adults: long-term follow-up of self-harm patients, admitted to hospital care, in a national cohort

Beckman K, Mittendorfer-Rutz E, Lichtenstein P, Larsson H, Almqvist C, Runeson B & Dahlin M 3397

Suicide risk linked with clinical consultation frequency, psychiatric diagnoses and psychotropic medication prescribing in a national study of primary-care patients

Windfuhr K, While D, Kapur N, Ashcroft DM, Kontopantelis E, Carr MJ, Shaw J, Appleby L & Webb RT 3407

Psychosocial therapy and causes of death after deliberate self-harm: a register-based, nationwide multicentre study using propensity score matching

Birkbak J, Stuart EA, Lind BD, Qin P, Stenager E, Larsen KJ, Wang AG, Nielsen AC, Pedersen CM, Winsløv J-H, Langhoff C, Mühlmann C, Nordentoft M & Erlangsen A 3419

Predictors of psychological improvement on non-professional suicide message boards: content analysis

Niederkrotenthaler T, Gould M, Sonneck G, Stack S & Till B 3429

The effect of sodium nitroprusside on psychotic symptoms and spatial working memory in patients with schizophrenia: a randomized, double-blind, placebo-controlled trial

Stone JM, Morrison PD, Koychev I, Gao F, Reilly TJ, Kolanko M, Mohammadinasab A, Kapur S & McGuire PK 3443

The effects of cognitive and behavioural therapies for anxiety disorders on depression: a meta-analysis

Cuijpers P, Cristea IA, Weitz E, Gentili C & Berking M 3451