

Article

Cite this article: Deleamar T, Zamora S (2024). Systematics, diversity and paleoecology of cyrtocrinids (Crinoidea, Echinodermata) from the Oxfordian sponge meadows of northeast Spain (Tosos, Zaragoza). *Journal of Paleontology* **98**, 1066–1086. <https://doi.org/10.1017/jpa.2024.59>

Received: 10 June 2024
Revised: 29 July 2024
Accepted: 20 August 2024
First published online: 19 March 2025

Handling Editor: Przemyslaw Gorzelak.

Corresponding author:
Samuel Zamora;
Email: s.zamora@igme.es

Systematics, diversity and paleoecology of cyrtocrinids (Crinoidea, Echinodermata) from the Oxfordian sponge meadows of northeast Spain (Tosos, Zaragoza)

Théo Deleamar¹ and Samuel Zamora^{2,3} 

¹UMR CNRS 6118 Géosciences, Laboratoire de Paléontologie, campus de Beaulieu, Université Rennes, 263, avenue du Général-Leclerc, 35042 Rennes, France

²Instituto Geológico y Minero de España (IGME-CSIC), Residencia CSIC, Campus Aula Dei, Av. Montañana 1005, E-50059 Zaragoza, Spain

³Grupo Aragosaurus-IUCA, Área de Paleontología, Facultad de Ciencias, Universidad de Zaragoza, Zaragoza, Spain

Abstract

Jurassic cyrtocrinids from Spain are first documented here as representing relatively diverse assemblages from the western Tethys. The species *Ascidicrinus pentagonus*, *Eugeniocrinites cariophilites*, *Gammarocrinites compressus*, *Pilocrinus moussoni*, *Sclerocrinus* cf. *S. strambergensis*, and *Tetracrinus moniliformis* are described from Oxfordian marl levels belonging to the Yátova Formation, around Tosos (Zaragoza, NE Spain). Although based on partially disarticulated material, these fossils preserve cups, stem columnals, attachment structures, and brachial plates. Based on the sedimentology and associated invertebrate fauna, cyrtocrinids from this area lived below storm wave action but eventually were affected by storms, as opposed to their modern counterparts that occupy deep water environments. Some specimens preserve traces of interactions with other organisms, such as predation marks or epibiontic colonization during life and post-mortem. Attachment structures suggest cyrtocrinids mostly attached on sponges. We note that diversity of cyrtocrinids changed in step with the abundance of sponge reefs in the Jurassic, suggesting that both groups probably had an important link in that period related with similar ecological requirements.

Non-technical Summary

Stalked crinoids belonging to various groups such as isocrinids, bourgetocrinids, and cyrtocrinids are restricted to deep water habitats today. In contrast, they were widespread in shallow marine environments during the Jurassic (about 150–200 million years ago). Herein we document a diversified assemblage of cyrtocrinids from Spain including six different species. The crinoids were found in the Yátova Formation, which has been interpreted as a shallow marine sponge-dominated unit sporadically affected by storm events. Specimens are disarticulated but preserve all anatomical elements including cups, columnals, brachial plates, and attachment structures. Some of the attachment structures suggest that cyrtocrinids were encrusters on sponges. A global database suggests cyrtocrinids and reefs followed a similar pattern of diversification, which supports the idea that both groups had similar requirements in the Jurassic.

Introduction

Cyrtocrinids are a well-supported clade of modern crinoids belonging to Articulata Zittel, 1879, and are a sister group of Hyocrinida (Rouse et al., 2013). Cyrtocrinids are characterized by the absence of, or very reduced basal plates, short or absent column without nodals and cirri, immovable articulation (symplectial to cryptosymplectial) between columnals, short arms with only one or two primibrachials, and cemented attachment disc (Hess and Messing, 2011). Modern representatives (i.e., *Holopus* d'Orbigny, 1837; *Neogymnocrinus* Hess, 2006) include only a few species that live in relatively deep water that can reach depths of 900 m (Bourseau et al., 1991; Hess, 1999; Donovan and Jakobsen, 2004; Syverson et al., 2015) and are commonly seen in association with sponges (Gorzelak and Salamon, 2023, fig. 12).

Cyrtocrinids are known from the Lower Jurassic (Hess and Thuy, 2018) but highly diversified faunas from this age suggest they originated earlier. Based on molecular data, Rouse et al. (2013) suggested that stem cyrtocrinids probably originated in the Late Triassic. Unfortunately, the Triassic has failed thus far to provide unquestionable cyrtocrinids (Hess and Thuy, 2018; but see alternative in Hess, 2006, or in Salamon et al., 2009). The group increased its diversity rapidly in the Pliensbachian (Lower Jurassic) with a peak in the Late Jurassic–Lower Cretaceous (Romano et al., 2016; Brom, 2019).

© The Author(s), 2025. Published by Cambridge University Press on behalf of Paleontological Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

During the Jurassic cyrtocrinids were abundant in a wide range of depths and environments (Hess and Spichiger, 2001; Charbonnier et al., 2007; Salamon and Gorzelak, 2007; Brom, 2019). Detailed systematics of Jurassic cyrtocrinids include material from different parts of the Tethys, including France, Poland, Italy, Germany, and Switzerland, among others (Goldfuss, 1826–1833; Manni and Nicosia, 1990; Hess, 2006; Charbonnier et al., 2007; Salamon and Gorzelak, 2010a), but the material from Spain, which was first mentioned in Zamora et al. (2018), remains undetailed.

This work aims to provide the first detailed description of cyrtocrinid material from the Spanish Jurassic collected in the village of Tosos (Iberian Cordillera, NE Spain), where the quality of fossil preservation allows a proper systematic approach. This locality has well-exposed outcrops of the Yátova Formation (Urresti, 1995; Aurell et al., 1999) with the development of sponge meadows. Based on abundant disarticulated material containing cups, brachials, columns, and holdfasts, we provide information about the paleoecology of the group. Finally, we compare the diversity of the sponge meadows with the diversity of cyrtocrinids to understand potential causes of diversity fluctuations.

Geological setting and stratigraphy

The village of Tosos lies in the Huerva valley, 50 km southwest of Zaragoza. Most of the town is surrounded by Cenozoic strata, but to the south of Tosos Mesozoic exposures of Cretaceous and Jurassic age crop out in an anticline structure (Fig. 1.2). The section studied here (TO.3), which is located a few kilometers south of Tosos (Fig. 1.2), consists of around 90 levels (Fig. 2), with alternating sponge limestones and marls (Urresti, 1996). This outcrop corresponds to the Yátova Formation, and calibrating with ammonoids suggests it expands from the beginning of the middle Oxfordian (Transversarium Biozone) to the upper Oxfordian (Bimammatum Biozone) (Urresti, 1995; Aurell et al., 1999).

The transition between the middle and upper Oxfordian is marked by the transition to the Hypselum Biozone (Meléndez et al., 2009). Other organisms are abundant in these layers (Fig. 2), such as brachiopods, echinoids, belemnites, sponges, bivalves, and serpulids (Aurell et al., 1999). Magnetostratigraphic studies have confirmed the dates provided by biostratigraphy (Juárez et al., 1994). Crinoids are common in 10 layers (27, 33, 39, 43, 47, 49, 65, 77, 79, 85) of marly interbeds along the Yátova Formation (Fig. 2; also see methodology).

During the middle Oxfordian, and more specifically during the Transversarium Biozone (Fig. 2), the different parts of the East Iberian carbonate platform evolved into a very wide and homogeneous carbonate ramp, resulting in several major deposits of carbonate sequences, including the Yátova Formation and its sponge limestones. This platform reached its maximum depth shortly after the boundary between the middle and upper Oxfordian (Meléndez et al., 2006). After this transgressive event, a regression began in the Hypselum biozone and became widespread in the Bimammatum biozone, extending into the Kimmeridgian (Meléndez et al., 2006; Ramajo, 2006).

Paleoenvironment

The Jurassic is marked by a global event of reef proliferation in the oceans, caused by a reconfiguration of global circulation (Leinfelder et al., 2002). During the Late Jurassic, coral diversity appears to have peaked, which explains the sharp increase in coral reefs. The abundance of siliceous sponges increased sharply

in the northern part of the Tethys during the Middle and Late Jurassic (Leinfelder et al., 2002). According to some authors (Veron, 1995; Leinfelder et al., 2002), the “Reef Age” of the Late Jurassic can be explained by the formation of a global east–west current system, which favored distribution and mixing of the larvae of these organisms. In northeastern Spain, during the Oxfordian, the Basque–Cantabrian Basin was connected to the Iberian Basin by a shallow carbonate platform known as the Soria seaway (Fig. 1.1). Numerous siliciclastic inputs were supplied to this platform and its surroundings by the Ebro Massif to the north and the Iberian Massif to the south (Strasser et al., 2005). This geographical configuration, and the generalized transgression during the Transversarium zone (where level TO.3/27 is found), contributed to the development of a facies of mudstone to packstone limestones with sponges and bioclasts around Tosos (Figs. 1.1, 2). Sedimentation rates, nutrient supply and energy from the environment were sufficiently stable to allow the formation of sponge meadows and other associated photo-dependent organisms (Leinfelder et al., 2002; Ramajo, 2006). However, the low height of the sponge meadows and the few levels of remobilization, interpreted as produced by the action of storms, suggest a bathymetry below the lower limit of storm wave action but eventually spreading out around that limit (Ramajo, 2006). This environment enabled numerous cyrtocrinid colonies to establish themselves and increase their diversity.

Materials, preservation, and methods

Preliminary work included sampling several marly levels from the Tosos 3 section (see above). About 10 kg of marls from 10 levels were sampled (levels indicated in gray in Fig. 1.3) and cleaned with hot water and hydrogen peroxide (H₂O₂). All levels were positive for echinoderm material and more than 500 first cyrtocrinid cups (or calyces) were discovered and preliminarily studied. However, different types of preservation were observed, with some levels preserving the material better than others. Some preliminary results were published in Zamora et al. (2018). Therefore, a second sampling was done in the level 27 (this study) from the same section (Fig. 1.4) because this level records the best-preserved specimens and a high diversity of cyrtocrinids. In all, 130 kg of marls were collected and prepared following the aforementioned procedure. Samples were sieved using mesh sizes of 5 mm, 2 mm, and 1 mm. Most cups and columnals remained in the 5-mm and 2-mm mesh size so only those sediment fractions were studied in detail. The 1-mm mesh size revealed mostly brachials and other small elements. Of the 3,647 crinoid remains found, there were around 1603 brachials, 1412 columnals, 47 attachment structures, and 585 cups, the latter representing number of individuals.

In this study, the cups are used mainly for systematic purposes because they offer a wider range of characters to study and allow better identification of specimens than the other skeletal parts of crinoids (except for rare forms). Attachment structures also were studied in order to extract paleoecological information, as well as certain columnals with marks of parasites or epizoids. When possible, some isolated columnals are assigned to certain cups. Most cyrtocrinid remains are disarticulated, with the exception of some columnals, a few roots and, more rarely, some cups.

Most of the collected specimens were prepared with potassium hydroxide (KOH), cleaned with water, and neutralized with acetic acid (10%). They were photographed using a Nikon D7100




Figure 1. (1) Paleogeographic map of northeastern Spain during the Oxfordian, with the location of the village of Tosos (modified after Ramajo and Aurell, 2008). (2) Simplified geological map of the area around the village of Tosos, with the TO.3 outcrop in red. Abbreviations: Fm. = Formation; Oxf. = Oxfordian; Kimm. = Kimmeridgian; Mid. Jur. = Middle Jurassic (modified after Lendinez et al., 1989). (3) Photograph of the outcrop of the Yátova Formation in Tosos (Zaragoza, Spain), showing the first sampling of marly levels. (4) Photograph of the outcrop during resampling of level 27. (5) Outcrop surface with cyrtocrinid material indicated by white arrows. (6) Example of associated fauna including complete tests of echinoids (*Plegiocidaris* sp.); cyrtocrinid material indicated by white arrow.

equipped with a 60 mm AF-S Micro NIKKOR lens. Before photography, the specimens were coated with ammonium chloride to increase contrast. Some specimens were photographed using SEM model JSM 6360-LV in the “Servicio General de Apoyo a la Investigación-SAI, Universidad de Zaragoza.”

The resulting diversity of cyrtocrinids was analyzed using a rarefaction curve produced with PAST software. Since only one level was studied, this curve was used to show the sampling effort and the diversity, rather than a comparison with other sites. The global diversity of Jurassic cyrtocrinids was explored

using a database referencing the total number of cyrtocrinid species and genera known in the literature, considering a single occurrence per species (or genus) and per stage, which then were linked in the intermediate stages, using a range-through approach according to Foote (2000) (see Supplementary Data). This was made in order to compare the diversity curve for cyrtocrinids in the Jurassic with the curves for reef domains and siliceous sponge reefs taken from Leinfelder et al. (2002). A simple correlation test and a cross-correlation between these data was carried out (Pearson coefficient) to explore the




Figure 2. Stratigraphic log of the "TO.3" outcrop at Tosos (Zaragoza, Spain), with details of lithology and fossil content at the Yátova Formation; levels with crinoid material are indicated in gray. Abbreviations: m = meter; M = mudstone; W = wackestone; P = packstone; Fm. = Formation; St. = Stage; Bz. = Biozone; Bimam. = Bimammatum (modified after Ramajo, 2006).

possible relationship between cyrtocrinid diversity and abundance of sponge reefs.

Repository and institutional abbreviation. All figured specimens and important non-figured specimens have a museum number and are deposited in the following institution: Museo de Ciencias Naturales de la Universidad de Zaragoza, Zaragoza, Spain, under the acronyms MPZ2024/97–204. There is also extra cyrtocrinid material that remains in the same collection.

Systematic paleontology

Classification and terminology. The classification and most of the terminology used in descriptions (Fig. 3) follow Hess and Messing (2011).

Class **Crinoidea** Miller, 1821

Subclass **Articulata** Zittel, 1879

Order **Cyrtocrinida** Sieverts-Doreck in Moore et al., 1952

Suborder **Cyrtocrinina** Sieverts-Doreck in Moore et al., 1952

Superfamily **Eugeniocrinitoidea** Roemer, 1855

Family **Eugeniocrinitidae** Roemer, 1855

Genus **Eugeniocrinites** Miller, 1821

Type species. *Encrinites cariophilites* von Schlotheim, 1813.

Diagnosis. Cup conical to funnel-shaped, rarely nearly cylindrical. Radial articular facets steep and wide in adults, with small aboral ligament fossa and deeply excavated adoral muscle fossae, separated from radial cavity by narrow ledges extending from interradial processes to median notch; interarticular ligaments indistinct. Articular facets separated by small interradial projections. Radial cavity wide and rather shallow. Aboral side of cup entirely occupied by almost flat, articular facet for column. First primibrachials very low. Axillary second primibrachials large, with compact, spearhead-shaped, median prolongation above small, lateral, articular facets for secundibrachials, and joined aborally by symplectial facets as protective covers over adoral side of cup when retracted. Arms small, may be coiled between second primibrachials. Columnals cylindrical to slightly conical, commonly high. Uppermost columnal with diameter equal to base of cup, commonly closely attached to cup. Articular facet of columnals with marginal granulation (after Hess and Messing, 2011; Hess, 2014b).

Eugeniocrinites cariophilites (von Schlotheim, 1813)

Figure 4

- 1996 *Eugeniocrinites cariophilites*; Klikushin, p. 103, pl. 1, figs. 3–22 (with previous synonymy).
- 1998 *Eugeniocrinites cariophilites*; Kaufmann, p. 10, text-fig. 7.5?.
- 2011 *Eugeniocrinites cariophilites*; Hess and Messing, p. T181, text-fig. 88.1.
- 2014b *Eugeniocrinites cariophilites*; Hess, p. 123, figs. 2–14.
- ?2018 *Eugeniocrinites cariophilites*; Zamora et al., p. 788, text-fig. 9a.
- 2020 *Eugeniocrinites cariophilites*; Krajewski et al., p. 70, text-fig. 5e, f.

Holotype. Cup figured by Knorr (1775, pl. 36, fig. 20) and refigured by von Schlotheim, 1813, from the Jurassic of Switzerland (according to Rasmussen, 1961, p. 223).

Emended diagnosis. Cup smooth, funnel-shaped, slightly conical or bowl-shaped; pentagonal in adoral view. Radial articular facet steep, large, and almost flat, muscle fossae deep, wide, and oval in adult, more rounded in juveniles. Interradial processes not very high, with a triangular apex, sometimes with a rounded angle. Aboral part of the cup circular or subcircular, flat or slightly concave, with marginal and irregular roughness made up of tubercles that are sometimes elongated and can be folded up. Columnals cylindrical, sometimes with concave sides, crenularium with the same roughness pattern as the aboral part of the cup.




Figure 3. Schematic cup of a cyrtocrinid of Eugeniocrinitidae Roemer, 1855, typical morphology (1) in aboral view and (2) in lateral view, describing the terminology used here.

Occurrence. Late Jurassic (Oxfordian) of Spain, Oxfordian and Kimmeridgian of France, Switzerland, Germany, and Early Cretaceous of France? (Rasmussen, 1961).

Description. Cups of varying sizes and shapes, slightly higher than wide, funnel-shaped, with a cylindrical aboral part (Fig. 4.2, 4.6). Other cups are wider than high, bowl-shaped (Fig. 4.14). Slight asymmetry sometimes visible. Cups smooth, sutures of radial plates visible, sometimes forming slight grooves in small cups. Clearly pentagonal in adoral view, radial cavity varies in width, moderately deep, adoral groove well marked, fairly deep, interradial grooves more or less distinct, sometimes with crenulated edges (Fig. 4.4, 4.5). Interradial processes of variable height and width, triangular or slightly angular apex (Fig. 4.6, 4.10, 4.14).

Radial articular facet not outward sloping, aboral articular facets slightly crescent-shaped to almost flat, aboral ligament pit small and shallowly depressed, axial canal round, very small, slightly more depressed. Adoral articular facets not very high, wide, sunken, muscular fossae deep, wide, oval (Fig. 4.6) to sub-rectangular (Fig. 4.14), separated by a deep adoral groove, varying in width. Aboral face circular, sometimes subcircular (Fig. 4.11), facet for the column slightly depressed, contour ornamented with irregular ridges in the crenularium, well-developed areola, central canal small and round. Columnals of variable size, generally short, cylindrical (Fig. 4.2) or with concave sides (Fig. 4.18). Articular facets circular or oval, areola developed, crenularium ornamented with irregular tubercles more or less elongated and branched, lumen small and rounded (Fig. 4.19). Axillary second primibrachial spearhead-shaped (Fig. 4.16), basal part forming an almost right angle with the distal part, articular facet for the secundibrachials small and laterally arranged (Fig. 4.17).

Materials. Several cups, some with columnals still attached (MPZ2024/103–104), others isolated (MPZ2024/97–100), columnals (MPZ2024/101), brachials (MPZ2024/102), and additional

material from the TO3/27 level middle Oxfordian, Yátova Formation.

Remarks. The funnel or bowl shape of the cups and the wide adoral articular facets and muscular fossae in adults (Fig. 4.1–4.14) are characteristic of the species *Eugeniocrinites cariophilites* (Hess, 2014b). Variation of the shape of the cups has already been demonstrated by various authors (Quenstedt, 1876; Hess, 2014b). Some small cups have slightly rounder and narrower muscular fossae, but this is a juvenile feature of the species because the muscular fossae follow the widening of the facets with growth (Fig. 4.6, 4.10, 4.14) (Hess, 2014b). The same applies to the interradial fossae, which are more or less distinct depending on the stage of development (Fig. 4.6, 4.10) (Hess, 2014b). The Tosos specimens are also variable in the shape of the interradial processes, which can be wider and higher.

The studied material differs from the morphologically related species *Lonchocrinus dumortieri* (de Loriol, 1882) in their interradial processes, which are generally still less high, less acuminate, and more curved inwards (Fig. 4.5, 4.9) (de Loriol, 1882–1884; Hess and Messing, 2011). In addition, *L. dumortieri* has more vertical and round muscular fossae than *E. cariophilites*, as well as outwardly sloping radial facets (de Loriol, 1882–1884). *Eugeniocrinites cariophilites* can be distinguished from *E. crenulatus* d'Orbigny, 1850, another Oxfordian species, by its facet to the column, which does not show the same ornamentation (de Loriol, 1882–1884). In fact, *E. cariophilites* columnals are cylindrical, with a fine lumen on the articular surface, and a broad areola surrounded by a marginal, irregular crenularium.

Family **Sclerocrinidae** Jaekel, 1918
Genus **Gammarocrinites** Quenstedt, 1858

Type species. *Eugeniocrinites compressus* (Goldfuss, 1829).




Figure 4. *Eugeniocrinites cariophilites* (von Schlotheim, 1813) from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1–3) Cup with first columnal still attached in adoral, lateral, and aboral views (MPZ2024/104); (4–7) cup with adoral groove crenulated in adoral, oblique, lateral, and aboral views (MPZ2024/97); (8–11) cup with very distinct interradial processes in adoral, oblique, lateral, and aboral views (MPZ2024/100); (12–15) cup with small interradial processes in adoral, oblique, lateral, and aboral views (MPZ2024/98); (16, 17) second primibrachial in adoral and lateral views (MPZ2024/102); (18, 19) columnal with serpulids in lateral and articular views (MPZ2024/101). All specimens photographed whitened with ammonium chloride sublimated; all scale bars = 1 mm.

Diagnosis. Cup low, compact, more or less oblique, with granular surface, radial cavity rather small. Facet to column wide, deep, and concave (after Hess and Messing, 2011).

Remarks. The genus name *Sclerocrinus* Jaekel, 1891, has no prevalence over *Gammarocrinites* (Hess, 2004; ICZN, 2006), so *Gammarocrinites* is used here. These two genera differ in shape.

The calyx is more massive in *Sclerocrinus* than in *Gammarocrinites* and the radial articular facets are rounded in *Sclerocrinus* (Hess and Messing, 2011) and rather flat in *Gammarocrinites*. The specimens studied here are low forms, with rather flat articular facets, thus suggesting inclusion in *Gammarocrinites*.

Gammarocrinites compressus (Goldfuss, 1829)

Figure 5

- 1829 *Eugeniocrinites compressus* Goldfuss, p. 164, pl. L, fig. 5.
- 1852 *Eugeniocrinites compressus*; Quenstedt, p. 615, pl. 53, figs. 42, 43.
- 1858 *Eugeniocrinites compressus*; Quenstedt, p. 654, pl. 80, figs. 70–78.
- 1876 *Eugeniocrinus compressus*; Quenstedt, p. 429, pl. 106, figs. 6–9.
- 1885 *Eugeniocrinites compressus*; Quenstedt, p. 936, pl. 74, figs. 35–38.
- 1891 *Sclerocrinus compressus*; Jaekel, p. 626, pl. XXXIV, figs. 1–8.
- 1961 *Sclerocrinus compressus*; Rasmussen, p. 217, pl. 32, fig. 5.
- 1974 *Sclerocrinus compressus*; Arendt, p. 109, pl. XI, figs. 1–8, 14, 19, text-fig. 14a–c.
- 1978 *Gammarocrinites compressus*; Rasmussen, p. T831, text-fig. 557.3.
- 1987 *Gammarocrinites compressus*; Głuchowski, p. 26, pl. 7, figs. 2–8, pl. 8, figs 1–7, text-fig. 14:8.
- ? *Sclerocrinus compressus*; Salamon and Gorzelak, p. 875, text-fig. 2H–J.
- 2010a *Gammarocrinites compressus*; Hess and Messing, p. T188, text-fig. 90.4.
- 2011 *Gammarocrinites* sp., Zamora et al., p. 788, text-fig. 9c.

Lectotype. Cup from the Jura limestone of Germany, Jurassic (Goldfuss, 1829, p. 164, pl. L, fig. 5; according to Rasmussen, 1961, p. 218).

Emended diagnosis. Cup rather low, wider than high, granulated, asymmetric. Adoral articular facet almost flat, muscle fossae rounded. Sides of radial plates in lateral view straight, slightly concave, or flared at base, with distinctive suture, very finely and geometrically sinuous. Aboral cavity large, delimited by fine edges.

Occurrence. Late Jurassic, Oxfordian from Spain, Oxfordian to Tithonian in Europe (Hess and Messing, 2011).

Description. Cup low, very or slightly asymmetrical, wider than high, pentagonal with rounded edges in adoral view. Highly variable aboral part, ranging from cylindrical (Fig. 5.4) to very flared, sometimes forming slight lobes (Fig. 5.17). Radial cavity highly variable in size, from small (Fig. 5.12) to moderate (Fig. 5.15, 5.18), funnel-shaped form, smooth cavity wall, interrarial grooves more or less visible. Interrarial processes of variable height, forming small spikes. Radial plate sutures clearly visible, with alternating rectangular lobes and hollows (zygosynostosis) at each end of the plate, becoming entangled at the sutures (Fig. 5.3, 5.5, 5.10, 5.16, 5.19). Marked, rather regular granulation on the cup surface, with small circular granules becoming larger and elongated aborally (Fig. 5.3), sometimes granulation is only visible on the

lower contour of the aboral articular facet or totally faded and absent.

Radial articular facets generally slightly sloping outwards, adoral facets fairly flat and shallowly depressed, with variable size and inclination, trapezoidal in lateral view, circular or subcircular muscle fossae. Distinct transverse ridge, sometimes protruding when unabraded. Aboral articular facet of variable size, sometimes higher than the adoral articular facet, sometimes lower, generally as wide as the base of the adoral facets, rounded. Axial canal circular, sometimes subcircular, fairly large, sunken, located above the transverse ridge. Aboral ligament pit large, oval, and deep. Aboral face subcircular (Fig. 5.4), sometimes more pentagonal (Fig. 5.11), moderately deep, large, and conical, with quite distinct suture and sometimes with the outer edges granulated. Facet to the column shows a small, more pronounced depression at the bottom of the cavity and sometimes shows a slight crenularium with irregular granulation (Fig. 5.14, 5.17).

Materials. Several cups (MPZ2024/105–131) and additional material from the TO3/27 level, middle Oxfordian, Yátova Formation.

Remarks. Specimens are included within the species *Gammarocrinites compressus* because of the very distinct granulation around the entire circumference of the section (Fig. 5.3, 5.19) or around the aboral facets, the flat adoral articular facet (Fig. 5.3, 5.10, 5.13, 5.19), the small radial cavity, whose diameter does not appear to exceed the radius of the total section, and the protruding transverse ridge (as described in Goldfuss, 1826–1833; Rasmussen, 1961; Hess and Messing, 2011). Given the large number of well-preserved samples we studied, we assume that *G. compressus* and *Cyrtocrinus nutans* (Goldfuss, 1829) have very similar cups, as already discussed in literature (de Lorient, 1882–1884; Hess and Messing, 2011; Hess, 2012). Indeed, granulation is sometimes absent (Fig. 5.7, 5.8), although it remains present on the majority of specimens.

As described by de Lorient (1882–1884), it seems that the diameter of the radial cavity is an important feature in the variation of several morphological characters in the adoral part, such as the size of the interrarial processes, or the orientation of the adoral articular facet. However, by observing calyces from other Spanish levels, or by looking at photographs of *Cyrtocrinus nutans* cups made available to us by the Museum national d'Histoire naturelle de Paris (lot MNHN.F.A08034, collection d'Orbigny 3865), certain differences become apparent in the aboral part or in the shape of the radial plates, thus separation of two species is justifiable. Indeed, in the *G. compressus* specimens presented here, the aboral cavity appears large and flared, possibly shallow, and surrounded by thin radial plates (Fig. 5.4), sometimes to the point of making the delineation of the aboral cavity indistinct (Fig. 5.11). In *C. nutans*, this cavity is deep, always well delimited by thick radial plates, making it narrower and more circular. Also, in *G. compressus*, in lateral view, the radial plates are straight (Fig. 5.3, 5.10), slightly concave, or convex at the base of the sides, showing the maximum width of these plates, most often in the aboral part (sometimes becoming small lobes, as in Fig. 5.16) or close to mid-height. Conversely, in *C. nutans*, the radial plates have strongly convex sides in lateral view, reaching their maximum width just below the aboral radial articular facet, the minimum width being at the base of the calyx.




Figure 5. *Gammarocrinites compressus* (Goldfuss, 1829) from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1–4) Cup with well-detailed sutures and granulations in adoral, oblique, lateral, and aboral views (MPZ2024/105); (5) suture of radial plate observed through a scanning electron microscope (MPZ2024/105); (6) microstructure and granulation observed through a scanning electron microscope (MPZ2024/105); (7, 8) slightly abraded cup in lateral and aboral views (MPZ2024/117); (9–11) medium-sized cup with detailed sutures and granulations in adoral, lateral, and aboral views (MPZ2024/123); (12–14) cup with slightly erased granulation in adoral, oblique, and aboral view (MPZ2024/112); (15–17) cup with a flared base in adoral, oblique, and aboral views (MPZ2024/114); (18–20) small cup with granulation in adoral, lateral, and aboral views (MPZ2024/115). All specimens photographed whitened with ammonium chloride sublimated with exception of 5 and 6 which are covered by gold under SEM; scale bars = 1 mm (1–4; 9–20); 500 μ m (5, 7, 8); 200 μ m (6).

The species studied here is easily distinguished from the Tithonian species *Ascidicrinus pentagonus* (Jaekel, 1891) by its discoidal base without distinct interrarial lobes. Material described here is distinguished from many other species by the particular simplex suture of its radial plates (Fig. 5.3). In literature, only some specimens of the species *Cyrtocrinus praenutans* Hess, 2012, appear to have similar sutures, but are still different in that they are composed of non-rectangular lobes and hollows (Hess, 2012). Finally, *G. compressus* appears to be the only species of the genus *Gammarocrinites* described from the Oxfordian of Europe.

Genus *Pilocrinus* Jaekel, 1907

Type species. *Eugeniocrinus moussoni* Desor, 1845.

Diagnosis. As for the species by monotypy.

Pilocrinus moussoni (Desor, 1845)

Figure 6

- 1845 *Eugeniocrinus Moussoni* Desor, p. 220.
- 1852 *Eugeniocrinites coronatus*, Quenstedt, p. 615, pl. LIII, fig. 45.
- 1858 *Eugeniocrinites coronatus*; Quenstedt, p. 654, pl. LXXX, fig. 79.
- 1867 *Eugeniocrinus coronatus*; Quenstedt, p. 733, pl. LXVII, fig. 45.
- 1876 *Eugeniocrinus coronatus*; Quenstedt, p. 425, pl. 106, figs. 1–5.
- 1879 *Eugeniocrinus moussoni*; de Loriol, p. 212, pl. XVIII, figs. 53–67.
- 1882–1884 *Eugeniocrinus moussoni*; de Loriol, p. 138, pl. 14, figs. 13–24.
- 1885 *Eugeniocrinus coronatus*; Quenstedt, p. 936, pl. 74, fig. 38.
- 1891 *Gymnocrinus moussoni*; Jaekel, p. 639, pl. XVIII, figs. 1, 2, text-fig. 17.
- 1961 *Pilocrinus moussoni*; Rasmussen, p. 226, pl. 32, fig. 6.
- 1974 *Pilocrinus moussoni*; Arendt, p. 113, pl. XII, figs. 5–8, pl. XIII, fig. 1.
- 1975 *Pilocrinus moussoni*; Hess, p. 73, pl. 23, figs. 5–10, 17–18, pl. 25, fig. 5.
- 1975 *Pilocrinus moussoni*; Rasmussen, p. 281, text-fig. 1.
- 1978 *Pilocrinus moussoni*; Rasmussen, p. T833, text-fig. 559.3.
- 1987 *Pilocrinus moussoni*; Gluchowski, p. 27, pl. 25, fig. 15, text-figs. 5, 17, 18.
- 1990 *Pilocrinus moussoni*; Kästle, p. 96, text-fig. 64.
- 1996 *Pilocrinus moussoni*; Klikushin, p. 106, pl. 2, figs. 1–15, text-figs. 2, 7.
- 1998 *Pilocrinus moussoni*; Kaufmann, p. 10, text-fig. 7.4?.
- 2008 *Pilocrinus moussoni*; Gorzelak, p. 228, text-fig. 2E.
- 2008 *Pilocrinus moussoni*; Salamon, p. 88, text-fig. 4J, K.
- 2011 *Pilocrinus moussoni*; Hess and Messing, p. T189; text-fig. 92.2.
- ?2018 *Pilocrinus* sp., Zamora et al., p. 786, text-fig. 9B.
- 2021 *Pilocrinus moussoni*; Salamon et al., text-fig. 3O.

Holotype. Rasmussen (1961) referred to *P. moussoni* as the type species of the genus *Pilocrinus*, but indicated that no holotype had been designated for *P. moussoni*.

Diagnosis. Wide, cylindrical or conical cup, with a more or less pronounced mid-height restriction, differentiating a cylindrical adoral part in lateral view and circular in adoral view, from an aboral part forming a more or less flared cap, with a pentagonal or subcircular shape in aboral view. Radial cavity deep and narrow surrounded by well-developed interrarial processes. Articular facets are wide and slightly inclined outwards, adoral facet as a sub-horizontal base. Aboral part of cup has a very deep, concave cavity, with a rather wide articular facet for column. Cylindrical to strongly convex columnals. Primibrachials 1 and 2 united by synostosis. Primibrachial 2 axillary with thickened sides, mostly united adorally and enclosing radial furrow. Secundibrachials commonly prolonged on one side but may also be united adorally (modified after Hess and Messing, 2011).

Occurrence. Late Jurassic (Oxfordian) from Spain. Middle Jurassic (Callovian) from Poland (Salamon, 2008) to Late Cretaceous (Campanian) from Europe (Rasmussen, 1975).

Description. Cup high, aboral part considerably wider (Fig. 6.7) to slightly wider (Fig. 6.2) than adoral part, constriction at mid-height of varying degree. Smooth cup surface, plate sutures slightly developed, sometimes finely sinuous (Fig. 6.5). Adoral part of cup rounded, aboral part with subcircular or pentagonal outline. Radial cavity deep, moderately wide, adoral grooves deep and distinct, interrarial grooves mostly indistinguishable. Thick interrarial processes, rather high. Radial articular facets sloping outwards, aboral facet small, more rounded than crescent-shaped, aboral ligament pit relatively large, oval and deep, axial canal rounded, clearly distinguishable on the other side of the transverse ridge. Adoral facet wide and high, with rounded corners, generally sunken, so that its base forms a plateau almost perpendicular to the aboral articular facet, muscle fossae subvertical, elliptical, relatively deep (Fig. 6.2, 6.5, 6.17). Aboral side wide, sometimes with very thick edge (Fig. 6.10), large and deep cavity, radial plate sutures more or less visible. Facet to column large, circular, with thin but clearly visible crenularium, which appears to be composed of long irregular tubercles in adults (Fig. 6.3, 6.10), and much shorter and regular tubercles in juveniles (Fig. 6.6). Lumen circular, moderately wide, large areola. Columnals barrel-shaped, with significant shape variation (Fig. 6.12, 6.14, 6.16), sometimes the same width throughout, sometimes much wider at mid-height, generally convex, articular surfaces with moderately wide lumen, thin crenularium, and large areola.

Materials. Several cups (MPZ2024/136–138, MPZ2024/142–144), some with columnals still attached (MPZ2024/145), several columnals (MPZ2024/139–141), and additional material from the TO3/27 level, middle Oxfordian, Yátova Formation.

Remarks. The constriction at mid-height (Fig. 6.2) and the deep, wide aboral part of the cup (Fig. 6.3, 6.10) are characteristic of the species *P. moussoni* (de Loriol, 1882–1884). The radial cavity of the specimens is moderately wide, but deep (Fig. 6.2, 6.10), as originally described by Desor (1845) and later considered by de Loriol (1882–1884). Other diagnostic features of the specimens are their large size in the adult stage, and their “cap-like appearance” (Desor, 1845). This appearance is explained by a wide, very deep aboral cavity. The shape and structure of the radial articular facet (Fig. 6.17) agree with descriptions by de Loriol (1882–1884). Specimens show an important variability in different morphological characteristics, such as the size of the cup, the form of the




Figure 6. *Pilocrinus moussoni* (Desor, 1845) from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1–3) High cylindrical cup in adoral, lateral, and aboral views (MPZ2024/137); (4–6) cup with first columnal still attached in adoral, lateral and aboral views (MPZ2024/145); (7, 8) cup with very wide basal part in lateral and aboral views (MPZ2024/136); (9, 10) thick cup in lateral and aboral views (MPZ2024/144); (11, 12) thick columnal in articular and lateral views (MPZ2024/139); (13, 14) strongly convex columnal in articular and lateral views (MPZ2024/141); (15, 16) cylindrical columnal in articular and lateral views (MPZ2024/140); (17) radial articular facet of a cup observed through a scanning electron microscope (MPZ2024/143); (18) stereom of a cup (MPZ2024/143). All specimens photographed whitened with ammonium chloride sublimated with exception of 17 and 18 which are covered by gold under SEM; scale bars = 1 mm (1–16); 200 μm (17); 50 μm (18).

aboral part (Fig. 6.3, 6.8, 6.10), the size of the interrarial processes (Fig. 6.2, 6.7), their asymmetry, and the thickness of the cup (Fig. 6.3, 6.10), contradicting the explanations of Jaekel (1891). This phenomenon could be linked to an occurrence in shallow water, which could indicate adaptations to hydrodynamic regimes or stronger predation pressures, but this requires further study on a larger number of specimens. Columnals show a variety of shapes, but most often barrel-shaped, sometimes very wide (Fig. 6.12), convex, and their articular surface has a wide lumen with marginal tubercles arranged in a circle, whose organization seems to vary according to specimen size (Fig. 6.11, 6.13, 6.15). These characteristics correspond to specimens figured in Salamon (2008) and Salamon et al. (2021).

Pilocrinus moussoni seems to be the only species of the genus *Pilocrinus* described from the Oxfordian and is very characteristic of this period (de Loriol, 1882–1884; Jaekel, 1891; Hess and Messing, 2011), but some columnals have been described from the Callovian (Salamon, 2008), and others from the Cretaceous (Rasmussen, 1961). A cup of this species also has been recorded from the Campanian of Germany (Rasmussen, 1975). For some reason, some authors seem not to integrate the later discovery into the stratigraphic extension of the species (Hess and Messing, 2011; Salamon et al., 2021).

Genus *Ascidicrinus* Hess, Salamon, and Gorzelak, 2011

Type species. *Ascidicrinus armatus* Hess, Salamon, and Gorzelak, 2011.

Diagnosis. Cups rather low, bulging or lobed interradially, especially in proximal view, commonly ornamented; radial articular facets broad, not sunken; radial cavity relatively deep, variable in diameter; facet for the column deep (after Hess et al., 2011).

Ascidicrinus pentagonus (Jaekel, 1891)

Figure 7.1–7.9

- 1891 *Sclerocrinus strambergensis* var. *pentagona* Jaekel, p. 624, pl. XXXVII, figs. 8, 9.
- 1974 *Sclerocrinus pentagonus*; Žitt, pl. 3, figs. 1–12, pl. 4, figs. 7–12.
- 1992 *Gammarocrinites bakonicus* Manni, Nicosia, and Szabó, pl. 1, figs. 3, 4, text-fig. 4.
- 2011 *Ascidicrinus pentagonus*; Hess et al., p. 124, text-fig. 2A.
- 2020 *Ascidicrinus pentagonus*; Krajewski et al., p. 70, text-fig. 7A, B.

Lectotype. According to Žitt (1974), cup from red marl of Štramberk, Czech Republic, early Lower Cretaceous (Jaekel, 1891, pl. XXXVII, fig. 8).

Emended diagnosis. Cup low, pentagonal, generally smooth or very finely granular, with well-developed and robust interrarial lobes in adoral part. Radial cavity narrow, moderately deep.

Occurrence. Late Jurassic, Oxfordian of Spain, Tithonian of Hungary, to Early Cretaceous (Berriasian–Valanginian) of France, Poland, Ukraine, and Czech Republic (Hess et al., 2011).

Description. Rather small, low cup, slightly restricted at mid-height (Fig. 7.3, 7.7), aboral part wider than adoral part. Clearly

pentagonal in oral and adoral views. Suture of radial plates generally slightly distinct but forms a very visible groove on the interrarial lobes (Fig. 7.8). Smooth or slightly granular surface. Interrarial lobes rather developed in basal part, very visible in lateral view, sometimes less indented in aboral view (Fig. 7.4). Interrarial processes poorly developed, surrounding a small, fairly deep, circular to slightly pentagonal radial cavity with clearly visible adoral grooves. Articular facets of radial plates shallowly depressed and sloping outwards, aboral facet small, rounded or crescent-shaped, with fairly wide, subcircular aboral ligament pit, axial canal very small, rarely visible (Fig. 7.3, 7.7). Adoral facet wider and higher, rather trapezoidal (Fig. 7.2, 7.6), muscle fossae small, round when visible (Fig. 7.2). Aboral cavity pentagonal, wide, fairly deep, sutures of radial plates clearly visible, sometimes toothed (Fig. 7.4), facet to the column small but deep with crenulated edges.

Materials. Several cups (MPZ2024/132–135) from the TO3/27 level, middle Oxfordian, Yátova Formation.

Remarks. The pentagonal shape and the fairly distinct interrarial lobes (Fig. 7.3, 7.7, 7.9) are the main characters that allow these specimens to be assigned to *Ascidicrinus pentagonus*. Other features such as the narrow radial cavity, aboral cavity, and general appearance of the specimen are in line with the description by Hess et al. (2011). Some specimens also correspond to the description of *A. pentagonus* from Manni et al. (1992), in that the aboral articular facets are subvertical, crescent-shaped, with a wide aboral ligament pit and a small axial canal (Fig. 7.9). The adoral articular facet, on the other hand, is wider, with subcircular or circular muscular pits (Fig. 7.3) (Manni et al., 1992). This morphology of the facets supports the species determination of these specimens, even though the interarticular lobes are not as well developed, a feature already discussed by Hess et al. (2011). Finally, the surface of the specimens studied here appears to have little or no ornamentation (Fig. 7.3, 7.7), unlike the specimen of Manni et al. (1992), but Hess et al. (2011) did not consider this as a characteristic of the species *A. pentagonus*. Moreover, *A. pentagonus* was also described as a small form, with no particular granulation by Jaekel (1891). The difference in ornamentation, added to the pentagonal shape of the specimen, easily distinguishes this species from *Ascidicrinus armatus* Hess, Salamon, and Gorzelak, 2011.

The specimens described here, and particularly the smaller forms, resemble certain individuals of *Gammarocrinites compressus* by their adoral part and the general shape of their articular facet. However, the presence of interrarial lobes in the aboral part is a diagnostic character of *A. pentagonus*. Thus, the attribution of these specimens to different species seems difficult. This discovery modifies the stratigraphic distribution of this species, making this the oldest known occurrence for *A. pentagonus*.

Genus *Sclerocrinus* Jaekel, 1891

Type species. *Sclerocrinus strambergensis* Jaekel, 1891.

Diagnosis. Cup mostly round, extremely massive; radial articular facet rounded, sloping strongly outward, with very small muscle fossae; radial cavity small. Facet to column large and deep, with differentiated inner part containing minute columnals (slightly modified after Hess and Messing, 2011).

Sclerocrinus cf. *S. strambergensis* Jaekel, 1891
Figure 7.10–7.13




Figure 7. Other Sclerocrinidae Jaekel, 1918, from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1–4) *Ascidicrinus pentagonus* (Jaekel, 1891) cup with serpulids in adoral, oblique, lateral, and aboral views (MPZ2024/133); (5–8) *A. pentagonus* cup slightly abraded in adoral, oblique, lateral, and aboral views (MPZ2024/134); (9) *A. pentagonus* complete cup observed through a scanning electron microscope in lateral view (MPZ2024/135). (10–13) *Sclerocrinus* cf. *S. strambergensis* Jaekel, 1891 cup with pits and scratches in adoral, oblique, lateral, and aboral views (MPZ2024/146). All specimens photographed whitened with ammonium chloride sublimate with exception of (9), which is covered by gold under SEM; all scale bars represent 1 mm.

Occurrences. Late Jurassic (Oxfordian) of Spain, Tithonian of Czech Republic (Jaekel, 1891), and Early Cretaceous (Valanginian) from Poland (Salamon and Gorzelak, 2010b).

Description. Fairly large, compact cup, very wide and not very high, slightly asymmetrical. Wide and thick, convex-sided radial plates with flat lower edge and concave upper edge in lateral view (Fig. 7.12). Smooth cup surface with well-marked, straight or sometimes finely striated interrarial sutures (Fig. 7.12, 7.13). Pentagonal adoral face with rounded rim, rather small radial cavity, a bit smaller than half the diameter of the cup, smooth rim. Deep, moderately narrow adoral grooves, interrarial grooves inconspicuous (Fig. 7.10). Fairly well-developed interrarial processes, surrounding a wide radial articular facet, whose edges meet the radial plate sutures. Aboral articular facet subvertical, crescent-shaped, rounded, not very high, sunken. Deep aboral ligament pit, moderately wide, subcircular or oval, more or less pronounced transverse ridge, thin, deep axial canal, slightly recessed, similar in size to the aboral ligament pit. Slightly high adoral articular facet, with a subhorizontal base, then, more vertical as the interrarial processes rise, trapezoidal in shape with more or less rounded sides, fairly deep (Fig. 7.12). Muscle fossae almost subvertical in lateral view (Fig. 7.12), oblong or oval in adoral view (Fig. 7.10). Thick-edged, pentagonal aboral face with no visible lobes, surrounding a circular or elliptical, fairly wide and deep cavity with rather visible, sometimes raised sutures.

Materials. One cup (MPZ2024/146) from the TO3/27 level, middle Oxfordian, Yátova Formation.

Remarks. The overall shape of the specimen combined with the thick radial plates that make it globular (Fig. 7.11, 7.12) are reminiscent of the *S. strambergensis* specimens described by Jaekel (1891) or Rasmussen (1961). However, there are a number of characteristics that make species assignments problematic, such as the radial cavity, which is not as small as that of previously described specimens (Jaekel, 1891; Rasmussen, 1961), the aboral face, which is not completely exposed (Fig. 7.13), making it impossible to observe the facets to the column. These features combined with the scarcity of material prevent a confident assignment to *S. strambergensis*.

This specimen differs from those of the *Cyrtocrinus nutans* and *Gammarocrinites compressus* in its large size, globular shape, the form of its radial articular facets, and its well-developed interrarial processes. It also differs from the *Pilocrinus moussoni* specimens in that its adoral part is not raised and has straight edges; also, there is no constriction at mid-height, just the opposite, with the greatest width at this point. It differs from the Tithonian species *Cyrtocrinus praenutans* by the shape of its radial plates and by the size or the shape of the aboral cavity. The sutures of the specimen are sometimes slightly sinuous, giving patterns similar to those of *G. compressus*, *C. praenutans*, and *P. moussoni*, perhaps indicating that this feature may be peculiar to the Sclerocrinitidae.

Superfamily **Plicatocrinoidea** Zittel, 1879

Family **Tetracrinidae** Nicosia, 1991

Genus **Tetracrinus** Münster, 1839

Type species. *Eugeniocrinites moniliformis* Münster in Goldfuss, 1826.

Diagnosis. Cup low, compact, with 3–6, commonly 4, radials, separated by a constriction and suture, more or less distinct

from compact basal element, which may be discoidal, lenticular, barrel-shaped, or low subconical, resembling a columnal. Radial cavity moderate, continued in shallow central depression in basal element, generally less than half diameter of cup. Radial articular facet large and broad, occupying entire distal edge of cup and meeting along interrarial sutures. Primibrachials 1 and 2 fused. Columnal articular facets with few crenulae or ridges, commonly in 4 groups (slightly modified after Hess and Messing, 2011).

Tetracrinus moniliformis (Münster in Goldfuss, 1826–1833)

Figure 8

- 1829 *Eugeniocrinites moniliformis* Münster in Goldfuss, p. 200, pl. LX, fig. 8.
- 1839 *Tetracrinus moniliformis*; Münster, p. 88, pl. XI, figs. 3a, b, 4.
- 1851–1852 *Tetracrinus moniliformis*; Bronn, p. 117, pl. XV, fig. 15a–d.
- 1852 *Eugeniocrinites moniliformis*; Quenstedt, p. 616, pl. LIII, fig. 52.
- 1852 *Tetracrinus moniliformis*; Quenstedt, p. 616, pl. LIII, figs. 49, 50.
- 1857 *Tetracrinus moniliformis*; Pictet, p. 335, pl. CII, fig. 2.
- 1858 *Tetracrinus moniliformis*; Quenstedt, p. 655, pl. LXXX, figs. 82–92.
- 1862 *Eugeniocrinites moniliformis* Goldfuss, p. 155, pl. LX, fig. 8a–i, k–m.
- 1876 *Tetracrinus moniliformis*; Quenstedt, p. 437, pl. 106, figs. 72–107.
- 1879 *Tetracrinus moniliformis*; de Loriol, p. 243, pl. XIX, figs. 37–48.
- 1882–1884 *Tetracrinus moniliformis*; de Loriol, p. 182, pl. 19, figs. 1–10.
- 1885 *Tetracrinus moniliformis*; Quenstedt, p. 937, pl. 74, figs. 41–44.
- 1889 *Tetracrinus moniliformis*; Bather, p. 167, pl. VI, fig. 16.
- 1892 *Tetracrinus moniliformis*; Jaekel, p. 647, pl. XXVII, figs. 1–18, pl. XXVIII, figs. 7–9.
- 1902 *Tetracrinus moniliformis*; Remès, p. 206, pl. XX, fig. 3a, b.
- 1974 *Tetracrinus moniliformis*; Arendt, p. 86, pl. I, fig. 1–9, pl. II, figs. 1–15, text-fig. 13a–c.
- 1975 *Tetracrinus moniliformis*; Hess, p. 75, pl. 23, fig. 20, pl. 26, figs. 1–11.
- 1978 *Tetracrinus moniliformis*; Rasmussen, p. T829, text-fig. 557.2.
- 2008 *Tetracrinus moniliformis*; Zatoń et al., p. 564, text-fig. 2F.
- 2008 *Tetracrinus moniliformis*; Salamon, p. 84, text-figs. 3E–H, 5A.
- 2011 *Tetracrinus moniliformis*; Hess and Messing, p. T196, text-fig. 96.1
- 2011 *Tetracrinus moniliformis*; Quereilhac and Guinot, p. 43, pl. 6, fig. 4.
- 2018 *Tetracrinus* sp., Zamora et al., p. 786, text-fig. 9E.
- 2021 *Tetracrinus moniliformis*; Salamon et al., p. 8, fig. 3P–T.

Lectotype. Basal circlet from the Jurassic of Streitberg (Oberfranken, Germany) (Münster in Goldfuss, 1826, p. 165, pl. LX, fig. 8l, m).

Diagnosis. Cup low, conical, well differentiated from the basal circlet. Basal circlet flat, strongly circular with a discoidal appearance and sometimes an enlarged top edge where granulation is located. Columnals varying in shape with fine and more or less regular granulation at mid-height, sometimes thorny. Aboral side of basal circlet and columnals with a large number of tubercles placed in a circle.

Occurrences. Middle Oxfordian of Spain, Callovian of Poland (Salamon and Feldman-Olszewska, 2018), and Kimmeridgian of Czech Republic (Arendt, 1974).

Description. Small cup, rectangular in adoral view, with 4 radials surrounding a fairly deep, subrectangular radial cavity, relatively small in diameter (Fig. 8.1, 8.4). Radial plates not very high, convex, and bear a flat, almost horizontal adoral articular facet, as well as more outwardly inclined aboral articular facet with small and round axial canal, a slightly larger, oval aboral ligament pit, and a filiform transverse ridge. Cup can be found still articulated with the basal circles (Fig. 8.2, 8.5), and the suture seems quite deep and clearly discernible. Basal circlet flat, strongly circular with a discoidal appearance, rounded or sometimes with raised edge in lateral view (Fig. 8.2, 8.5, 8.8, 8.11), ornamented or smooth. Four crests running from center to outside of the circle in adoral view, with fairly small, shallow cavity. On some specimens (Fig. 8.10) the four ribs do not reach the outer edge of the basal circlet and the articulation facets for the radials are small, revealing a wide areola between these zones and the edge of the basal circlet. With these types of ossicles, basal circlet is wider than the rest of the cup, revealing part of its adoral side (Fig. 8.2). Aboral face flat or slightly concave, facet to the column comprises a small lumen (Fig. 8.9, 8.12), surrounded by a slightly wide areola and a crenularium composed of elongated tubercles, alternating with grooves, arranged in a circle (Fig. 8.6, 8.9, 8.12). Columnals cylindrical, convex laterally, sometimes discoid with almost triangular sides (Fig. 8.2), sometimes as high as wide, granulation at mid-height with tubercles more or less aligned, most of the time forming distinct spines (Fig. 8.13, 8.14). Articular surface smooth on the edges, with facet in the center featuring a small lumen and a few laterally flattened and regular granules on its contour (Fig. 8.13).

Materials. Cup with basal circlet (MPZ2024/148) and sometimes first columnals (MPZ2024/150) still attached, isolated basal circlets (MPZ2024/147, MPZ2024/151), few columnals (MPZ2024/149, MPZ2024/152), and additional material from the TO3/27 level, middle Oxfordian; Yátova Formation.

Remarks. This species is particularly recognizable by a cup with four radials well separated from a flat basal circlet (Fig. 8.2, 8.5). Disarticulated basal circlets are flat, bearing four ribs, sometimes the surface is marked with granulations, recalling the description of de Loriol (1879). The material described here shows considerable variability in terms of basal circlets and columnals, and some specimens also vary from those in the literature. Basal circlet varies from laterally rounded (Fig. 8.5, 8.8) to flanged (Fig. 8.2, 8.11). In plan view, articulation facet also changes and is sometimes restricted to the whole width of the basal circlet (Fig. 8.7), having in other cases a relatively wide epifacet (Fig. 8.10). These differences also have been described in the literature (see Jaekel, 1892).

In terms of columnals, there is significant variation in ornamentation, with most of the columnals described here having

ornamentation composed of tubercles transformed into long spines (Fig. 8.13, 8.14), while others have less developed ornamentation (Fig. 8.15, 8.16) or lack ornamentation. The shape of the columnals is also variable, but the first columnal still attached to the section (see Fig. 8.2) has a single row of spines that is different from material described in Hess and Messing (2011, fig. 96, 1a), which is barrel-shaped. However, despite all these variations on several types of *T. moniliformis* ossicles, this species was described by Jaekel, 1892, as highly variable, reinforcing the idea that our specimens are included in this species. Further studies are still needed, with much more material.

All the specimens described here can be distinguished from *Tetracrinus galei* Hess, 2014a, by the numerous tubercles surrounding the facet for the column in aboral view (Fig. 8.3, 8.6, 8.9, 8.11), its well-defined suture between the cup and the basal circlet, and by its extensive granulation (Fig. 8.13, 8.14). *Tetracrinus moniliformis* also has finer, more regular granulation on its stem articles than *T. galei*, whose ornamentation is more tessellated and irregular (Hess, 2014a).

Discussion

Diversity and taphonomy of cyrtocrinids from the studied interval. Cyrtocrinids are important components in Jurassic sponge meadows from the Tethyan realm (Klikushin, 1996; Salamon, 2008; Zatoń et al., 2008; Hess et al., 2011) but they have been only figured in Iberia by Zamora et al. (2018). According to this new study, at least six species of cyrtocrinids are present in the middle Oxfordian of Tosos (level 27), for a total of 585 cups. Sampling of the level appears to have been statistically representative, since it seems to have captured the majority of the species present (Fig. 9).

The cyrtocrinids discovered in the Yátova Formation (Tosos, Zaragoza) appear to be an important parautochthonous component of the paleoenvironment of this locality during the Oxfordian. In fact, several remains are sometimes as well preserved as in certain Lagerstätten (Charbonnier et al., 2007) with complete cups (sometimes articulated to the proximal columnals) and other disarticulated elements (brachials, columnals, holdfasts) that even preserve traces of stereom (Figs. 5.6, 6.18). This degree of preservation is incompatible with long-distance and/or high-energy transport, suggesting parautochthony.

Cyrtocrinids can be classified as taphonomic type III echinoderms (Brett et al., 1997; Zatoń et al., 2008; Salamon et al., 2020), especially because they have cups composed with fussed radials. They may have remained on the seafloor for some time without being disarticulated (Brett et al., 1997), so they might have undergone some transport. The fact that some specimens are well preserved but contain epibionts points to some exposure time before final burial. Preservation varies from excellent to moderate (with some ossicles showing traces of abrasion), which is compatible with time-averaged communities that remain on the sea floor after death in environments with low sedimentation rates (Zamora, 2022).

The cyrtocrinids from Tosos appear to have lived in a shallow environment, which may have slightly exceeded the lower limit of storm wave action (Zamora et al., 2018), dominated by sponge meadows alongside many other organisms. The presence of a rich and diverse fauna of cyrtocrinids, sponges, brachiopods, and other suspension-feeding organisms, combined with geological and paleogeographical contexts that may have generated numerous siliciclastic deposits, indicates a nutrient-rich




Figure 8. *Tetracrinus moniliformis* (Münster, 1829, in Goldfuss, 1826–1833), specimens from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1–3) Cup with basal circler and first columnal still attached in adoral, lateral, and aboral views (MPZ2024/150); (4–6) cup with basal circler still attached in adoral, lateral, and aboral views (MPZ2024/148); (7–9) basal circler in adoral, lateral, and aboral views (MPZ2024/151); (10–12) basal circler in adoral, lateral, and aboral views (MPZ2024/147); (13, 14) columnal in articular and lateral views (MPZ2024/149); (15, 16) slightly abraded columnal in articular and lateral views (MPZ2024/152). All specimens photographed whitened with ammonium chloride sublimate; all scale bars represent 1 mm.

environment, but does not necessarily indicate a high rate of sedimentation. In fact, because remains from the environment contain a large number of epibionts (notably serpulids) and because cyrtocrinid larvae potentially need a time of lower sedimentation rate to attach themselves to the substrate (Zamora et al., 2018), this indicates a generally low or episodically interrupted sedimentation rates during deposition of the Yátova Formation. However, the hypothesis of occasionally interrupted sedimentation seems less appropriate in view of the stable conditions that are necessary for the development of sponge meadows (Leinfelder et al., 2002).

Concerning epizoans, it seems that the small sizes of cyrtocrinids do not make them a preferential substrate for many organisms, with the exception of serpulids (Klikushin, 1996). From the latter, only two forms seem to be present here, most of which are very thin serpulids, less than a millimeter in diameter, generally running along a single face of a cup or column, in the form of more or less pronounced zigzags (Fig. 10.5). These forms probably settled after the death of the animal because they are present on only one side and/or on articular facets (Figs. 10.6, 11.3), but sometimes they were able to encrust the animal during life (Klikushin, 1996). More rarely, serpulids are found in the form of a girdle (“girdling serpulids”) around a cyrtocrinid stem (Fig. 10.3), which may point to syn-vivo encrustation (Klikushin, 1996).

Finally, some cyrtocrinids appear to have been affected by parasites (Fig. 10.1, 10.4). In fact, some cyrtocrinid remains show

more or less deep cavities on their cups or columnals, having pierced their skeleton. The part of the cyrtocrinid stereom around these cavities appears abnormally swollen. This suggests that a foreign body entered or attached itself to the surface of the crinoid skeleton, causing a reaction (swelling). These galleries have been observed in other crinoids, especially in millericrinids (Klikushin, 1996; Wilson et al., 2014; Zamora, 2022). There are some occurrences of these pits on cyrtocrinid radials, without showing any particular swelling around the cavity (Hess and Thuy, 2018). Previous works have included these marks in the ichnogenus *Oichnus* Bromley, 1981, which may be caused by gastropods or octopods (Wilson et al., 2014; Zamora, 2022). Other authors have referenced similar traces, with a slit-shaped form when the parasite settles on a suture between cup and first columnal (Fig. 10.1 for example), or oval-shaped when it settles in a single columnal (Fig. 10.4 for example). According to some authors, they could be typical of myzostomids worms (Arendt, 1961; Klikushin, 1996). These traces are still uncommon compared with the number of columnals discovered here and require further research on modern cyrtocrinids.

Evidence of predation or grazing on some specimens (Figs. 10.7, 11.8) suggests that cyrtocrinids from Tosos were exposed to predators. Predator–prey relationships have been documented in other cyrtocrinids from different localities (Salamon and Gorzelak, 2010a; Zamora and López-Horgue, 2022). Considering that predation in the Mesozoic was more frequent in shallow areas than in deep areas (Oji, 1996; Baumiller and




Figure 9. (1) Rarefaction curve for Tosos cyrtocrinids (level 27), within a 95% confidence interval, rapidly reaching a plateau. (2) Cyrtocrinid species studied with their respective number of cups.

Gahn, 2013; Syverson et al., 2015), it is not surprising to find those signs in the Jurassic of Tosos. Predation in shallow environments has been considered the main reason for migration of cyrtocrinids to deep environments during the Mesozoic Marine Revolution (Gorzelak et al., 2012; Zamora and López-Horgue, 2022).

Substrate preferences. As obligate encrusters, cyrtocrinids preferred hard substrates (Hess, 2006; Charbonnier et al., 2007; Hess and Messing, 2011). Forty-seven holdfasts have been

documented from the material of Tosos, 18 of which were found on sponge fragments (Fig. 11.1–11.7), 27 are no longer attached to any substrate, one was found on a mollusk shell (Fig. 11.8, 11.9), and another on a cyrtocrinid stem (Fig. 11.10, 11.11). Because a large number of attachment structures were found on sponges, we suggest that they were preferential substrates (benthic islands) for cyrtocrinids, as has been documented previously in other assemblages (Zatoñ et al., 2008; Hess and Messing, 2011; Zamora and López-Horgue, 2022). The sponges probably also provided protection from predators and more




Figure 10. Traces of interactions on cyrtocrinids remains from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1, 2) *Eugeniocrinites cariophilites* with parasite on the suture between cup and first columnal (MPZ2024/103), observed through a scanning electron microscope in lateral and enlarged views; (3) serpulid surrounding a cyrtocrinid columnal (MPZ2024/153); (4) *Ooichnus* sp. in a columnal (MPZ2024/202); (5, 6) serpulids on one side of a *Pilocrinus moussoni* columnal in lateral and articular views (MPZ2024/204); (7) trace of predation on a columnal (MPZ2024/201). All specimens photographed whitened with ammonium chloride sublimated with exception of (1, 2, 7), which are covered by gold under SEM; scale bars = 1 mm (1, 3–7); 100 μ m (2).




Figure 11. Cyrtocrinid holdfasts from the middle Oxfordian of Tosos (Zaragoza, Spain), Yátova Formation, layer TO3/27. (1, 2) Root-like holdfast on a sponge in articular and lateral views (MPZ2024/155); (3, 4) disk-like holdfast on a sponge with serpulids and a few traces of predation in lateral and articular views (MPZ2024/158); (5) abraded holdfast on a sponge (MPZ2024/161); (6, 7) disk-like holdfast on a sponge in lateral and articular views (MPZ2024/156); (8, 9) disk-like holdfast on a mollusk shell with traces of predation and/or grazing, in lateral and articular views (MPZ2024/160); (10, 11) disk-like holdfast on a cyrtocrinid columnal in lateral and articular views (MPZ2024/157); (12, 13) cross-like holdfast with serpulid in lateral and articular views (MPZ2024/159). All specimens photographed whitened with ammonium chloride sublimate; all scale bars = 1 mm.




Figure 12. Abundance of siliceous sponge reefs classified in categories ranging from extremely rare (1) to abundant (5), abundance of reef domains and diversity of cyrtocrinid species and genera in the world through the Jurassic; *p*-value and *Cor* (Pearson's coefficient) are calculated between the abundance of reef domain and cyrtocrinid species diversity data using a simple correlation test (in black) or with a cross-correlation test (in red).

elevated areas for filter feeding. The presence of an attachment disc wrapped around a cyrtocrinid stem is notable and interesting. The stem is well preserved (Fig. 11.10, 11.11) and it cannot be ruled out that colonization occurred during life.

Possible link between sponges and cyrtocrinids in the Jurassic. The cyrtocrinids of Tosos developed in an environment shaped by sponge meadows and show a particular link with the latter because they attach to them (see above). To explore a link between cyrtocrinids and sponges, we developed a database of cyrtocrinids and sponge reefs in the Jurassic (see below). Despite a lack of paleontological data concerning the exact number of sponge taxa in the Jurassic (Leinfelder et al., 2002), studies of reefs and bioherms in general provide a general overview of their abundance and diversity.

The curves for the abundance of reef domains around the world in the Jurassic, the abundance of reefs built only by siliceous sponges, and the diversity of cyrtocrinids show a similar trend (Fig. 12). Diversity in the Early Jurassic was rather low (except for cyrtocrinid diversity), then increased slightly in the Middle Jurassic, although with the lowest peak during the Aalenian (Fig. 12), and increased drastically in the Late Jurassic (Oxfordian), followed by a more or less continuous decline from the Kimmeridgian onwards (Fig. 12). Similar trends were noted previously (Leinfelder et al., 2002), given that siliceous sponges are the least known reef builders in the Early Jurassic, except in northern Italy (e.g., Zempolich, 1993; Leinfelder et al., 2002), unlike in the Late Jurassic where they are abundant. It is also important to note that during the Oxfordian, the majority

of siliceous sponge reefs formed along a belt at the northern margin of the Tethys, passing through France, Spain, Switzerland, southern Germany, Poland, and Romania (Leinfelder et al., 2002) and that important cyrtocrinid faunas were described from rocks of this age in these countries (de Loriol, 1879, 1882–1884; Zatoń et al., 2008; Zamora et al., 2018). Taking such a link into account, the decrease in cyrtocrinid diversity in the Kimmeridgian could then be explained by a well-known fundamental change in the northern Tethys (Leinfelder et al., 2002), which is marked by an increased influx of siliciclastic material that led to a significant decrease in sponge reefs.

Finally, the difference between trajectories of the curves in the Pliensbachian (i.e., much greater diversity of cyrtocrinid than sponges; Fig. 12) could be explained by a sampling bias of sponge diversity in Lower Jurassic units (Leinfelder et al., 2002), or by the existence of the Arzo paleontological site (Switzerland), where many cyrtocrinid species are endemic (Hess, 2006). At the same time, the peak in abundance of siliceous sponge reefs in the Bajocian (Fig. 12), which is not observed in cyrtocrinid diversity, could be caused by scarcity of paleontological sites with Bajocian-age cyrtocrinids, which likely is the case for Spain.

Although a possible link between cyrtocrinids and sponges is observed, the statistical relationships between Jurassic reef domains and cyrtocrinid diversity are more complex. A simple correlation test confirms a moderate correlation (see *p*-value and Pearson coefficient in Fig. 12). However, a cross-correlation test does not provide enough confidence and the correlation itself is less significant (see *p*-value and Pearson coefficient in Fig. 12). Thus, the hypothesis of a potential link between sponges and

cyrtocrinids needs further analyses that include more data and likely encompass other geological periods.

Conclusions

The cyrtocrinids from Tosos, detailed for the first time in the Spanish Oxfordian, provide a rather significant diversity, including six species: *Ascidicrinus pentagonus*, *Eugeniocrinites cariophilites*, *Gammarocrinites compressus*, *Pilocrinus moussoni*, *Sclerocrinus* cf. *S. stramburgensis*, and *Tetracrinus moniliformis*. They display considerable intraspecific variation, making systematic work sometimes difficult, thus necessitating abundant and well-preserved material. The Yátova Formation around Tosos was deposited in a shallow marine environment, which expands the distribution of those taxa in such environments to the western Tethys. This research confirms that cyrtocrinids were able to live in shallow environments in the Jurassic period, and that their distribution today in deep water is probably a consequence of the Mesozoic Marine Revolution. In addition, the numerous attachment structures discovered at Tosos confirm a preference for hard substrates, and especially for sponges. The diversity of cyrtocrinids and sponge reefs in the Jurassic suggests there might be a link between these two groups, but additional data are necessary to confirm or reject this hypothesis.

Acknowledgments. This is a contribution to project E18_20 R “Aragosaurus: Recursos geológicos y paleoambientales” from the Government of Aragón. I. Pérez (Zaragoza) helped with photography. The authors would like to acknowledge the use of Servicio General de Apoyo a la Investigación-SAI, Universidad de Zaragoza and especially to M^a J. Marín. Reviewers M. Salamon and P. Gorzelak greatly improved the resulting manuscript with important comments. P. Gorzelak is also acknowledging for help during the editorial process.

Competing interests. The authors declare none.

Data availability statement. Data available from the Zenodo Digital Repository: <https://doi.org/10.5281/zenodo.13378362>.

References

- Arendt, Y.A., 1961, On sea-lilies damages caused by Schizoproscina: *Paleontological Journal of Moscow*, v. 2, p. 101–106.
- Arendt, Y.A., 1974, Morskie lilii. Tsirtokrinidy [Sealilies, cyrtocrinids]: *Academy of Sciences of the USSR Transactions of the Paleontological Institute*, v. 144, 252 p.
- Aurell, M., Bádenas, B., Bello, J., Delvene, G., Meléndez, G., Urresti, I.P., and Ramajo, J., 1999, The Callovian and the Upper Jurassic in the northeastern Iberian Chain and in its transition to the Catalan Ranges (Sierra de Arcos, Calanda, and Xerta-Paüls areas): *Cuadernos de Geología Ibérica*, v. 25, p. 73–110.
- Bather, P.A., 1889, *Trigonocrinus*, a new genus of Crinoidea, from the “Weisser Jura” of Bavaria; with the description of a new species, *T. liratus*—appendix. Sudden deviations from normal symmetry in Neocrinoidea: *Quarterly Journal of the Geological Society of London*, v. 45, p. 149–171.
- Baumiller, T.K., and Gahn, F.J., 2013, Reconstructing predation pressure on crinoids: estimating arm-loss rates from regenerating arms: *Paleobiology*, v. 39, p. 40–51.
- Bourseau, J.P., Amézière-Cominardi, N., and Avocat, R., 1991, Echinodermata: Les Crinoïdes pédonculés de Nouvelle-Calédonie, in Crosnier, A., ed., *Résultats des Campagnes Musorstom*: Mémoire du Muséum Nationale d'Histoire Naturelle, v. 8, p. 229–333.
- Brett, C.E., Moffat, H.A., and Taylor, W.L., 1997, Echinoderm taphonomy, taphofacies, and Lagerstätten, in Waters, J.A., and Maples, C.G., *Geobiology of Echinoderms: Paleontological Society Papers*, v. 3, p. 147–190.
- Brom, K.R., 2019, Body-size trends of cyrtocrinids (Crinoidea, Cyrtocrinida): *Annales de Paléontologie*, v. 105, p. 109–118.
- Bromley, R.G., 1981, Concepts in ichnotaxonomy illustrated by small round holes in shells: *Acta Geologica Hispanica*, v. 16, p. 55–64.
- Bronn, H.G., 1851–1852, *Lethaea Geognostica oder Abbildung und Beschreibung der für die Gebirgs-Formationen Bezeichnendsten Versteinerungen*: Stuttgart, E. Schweizerbart, v. 3, p. 1–120.
- Charbonnier S., Vannier, J., Gaillard, C., Bourseau J.P., and Hantzpergue, P., 2007, The La Voulte Lagerstätte (Callovian): evidence for a deep water setting from sponge and crinoid communities: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 250, p. 216–236.
- de Loriol, P., 1879, Monographie des crinoïdes fossiles de la Suisse: *Mémoires de la Société Paléontologique Suisse*, v. 6, p. 125–300.
- de Loriol, P., 1882–1884, *Paléontologie Française, Série 1, Animaux Invertébrés. Terrain Jurassique*, v. 11, Crinoïdes, pt. 1: Paris, G. Masson, 627 p.
- Desor, E., 1845, Résumé des études sur les crinoïdes fossiles de la Suisse: *Bulletin de la Société Neuchâteloise des Sciences Naturelles*, v. 1, p. 211–222.
- Donovan, S.K., and Jakobsen, S.L., 2004, An unusual crinoid–barnacle association in the type area of the Danian (Paleocene), Denmark: *Lethaia*, v. 37, p. 407–415.
- d’Orbigny, A., 1837, Mémoire sur une seconde espèce vivante de la famille des Crinoïdes ou Encrines, servant de type au nouveau genre Holope (*Holopus*): *Magasin de Zoologie (7ème année)*, v. 10, p. 1–8.
- d’Orbigny, A., 1850, *Prodrome de Paléontologie Stratigraphique Universelle des Animaux Mollusques et Rayonnés*: Paris, Masson, v. 1, 394 p.
- Foote, M., 2000, Origination and extinction components of taxonomic diversity: general problems: *Paleobiology*, v. 26, p. 74–102.
- Gluchowski, E., 1987, Jurassic and Lower Cretaceous articulate crinoids of the Pieniny Klippen Belt and the Tatras Mts., Poland: *Studia Geologica Polonica*, v. 94, p. 6–102.
- Goldfuss, G.A., 1826–1833, *Petrefacta Germaniae. Echinodermata: Düsseldorf, Arnz and Co.*, v. 1, p. 115–221.
- Goldfuss, G.A., 1862, *Petrefacta Germaniae. Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angrenzenden Länder*: Leipzig, List & Franke, 234 p.
- Gorzelak, P., 2008, New data on fossil cyrtocrinids (Cyrtocrinida, Crinoidea) from the lower Oxfordian of the Polish Jura Chain (southern Poland): *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, v. 248, p. 215–232.
- Gorzelak, P., and Salamon, M.A., 2023, Deep-sea exploration of the Caribbean Sea in the area of Roatán Island (Honduras) from the point of view of Polish palaeontologists – the present is the key to the past: *Przegląd Geologiczny*, v. 71, p. 71–81.
- Gorzelak, P., Salamon, M.A., and Baumiller, T.K., 2012, Predator-induced macroevolutionary trends in Mesozoic crinoids: *Proceedings of the National Academy of Sciences USA*, v. 109, p. 7005–7007.
- Hess, H., 1975, Die fossilen Echinodermen des Schweizer Juras: *Seesterne, Schlangensterne, Seelilien, Seeigel, Seewalzen*: Basel, Veröffentlichungen Naturhistorischen Museum Basel, v. 8, 130 p.
- Hess, H., 1999, Recent, in Hess, H., Ausich, W.I., Brett, C.E., and Simms, M.J., eds., *Fossil Crinoids*: Cambridge, UK, Cambridge University Press, p. 237–244.
- Hess, H., 2004, Case 3235. *Sclerocrinus* Jaekel, 1891 (Crinoidea, Cyrtocrinida): proposed precedence over *Gammarocrinites* Quenstedt, 1857: *The Bulletin of Zoological Nomenclature*, v. 61, p. 251–253.
- Hess, H., 2006, Crinoids (Echinodermata) from the Lower Jurassic (upper Pliensbachian) of Arzo, southern Switzerland: *Schweizerische Paläontologische Abhandlungen*, v. 126, p. 1–144.
- Hess, H., 2012, Crinoids from the Middle Jurassic (Bajocian–lower Callovian) of Ardèche, France: *Swiss Journal of Palaeontology*, v. 131, p. 211–253.
- Hess, H., 2014a, *Balanocrinus* and other crinoids from Late Jurassic mudstones of France and Switzerland: *Swiss Journal of Palaeontology*, v. 133, p. 47–75.
- Hess, H., 2014b, The crinoid *Eugeniocrinites cariophilites* from the Late Jurassic of southern Germany: babies, cripples and enigmatic wing plates: *Swiss Journal of Palaeontology*, v. 133, p. 121–140.
- Hess, H., and Messing, C.G., 2011, *Treatise on Invertebrate Paleontology, Part T, Echinodermata 2 (Revised), Crinoidea, Volume 3, Chapter 2*: Lawrence, Kansas, University of Kansas, Paleontological Institute, 261 p.
- Hess, H., and Spichiger, P.U., 2001, *Argoviocrinus rarissimus* n. g. n. sp., a new crinoid (Echinodermata) from the middle Oxfordian of Northern Switzerland: *Eclogae Geologicae Helvetiae*, v. 94, p. 489–494.

- Hess, H., and Thuy, B., 2018, Emergence and early radiation of cyrtocrinids with new species from a Lower to Middle Jurassic rock reef of Feugueroles (Normandy, France): *Swiss Journal of Palaeontology*, v. **137**, p. 133–158.
- Hess, H., Salamon, M., and Gorzelak, P., 2011, Late Jurassic–Early Cretaceous (Tithonian–Berriasian) cyrtocrinids from south-eastern Poland: *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, v. **260**, p. 119–128.
- ICZN (International Commission on Zoological Nomenclature), 2006, Opinion 2155 (Case 3235). *Sclerocrinus* Jaekel, 1891 (Crinoidea, Cyrtocrinida): not conserved; priority maintained for *Gammarocrinites* Quenstedt, 1857: *Bulletin of Zoological Nomenclature*, v. **63**, p. 150–151.
- Jaekel, O., 1891, Ueber Holopocriniden mit besonderer Berücksichtigung der Stramberger Formen: *Zeitschrift der Deutschen Geologischen Gesellschaft*, v. **43**, p. 557–670.
- Jaekel, O., 1892, Ueber Plicatocriniden, *Hyocrinus* und *Saccocoma*: *Zeitschrift der Deutschen Geologischen Gesellschaft*, v. **44**, p. 619–696.
- Jaekel, O., 1907, Ueber die Körperform der Holopocriniden: *Neues Jahrbuch für Mineralogie, Geologie, Paläontologie, Festband*, v. **1907**, p. 272–309.
- Jaekel, O., 1918, Phylogenie und System der Pelmatozoen: *Paläontologische Zeitschrift*, v. **3**, p. 1–128.
- Juárez, M.T., Osete, M.L., Meléndez, G., Langereis, C.G., and Zijdeveld, J.D.A., 1994, Oxfordian magnetostratigraphy of the Aguilón and Tosos sections (Iberian Range, Spain) and evidence of a pre-Oligocene overprint: *Physics of the Earth and Planetary Interiors*, v. **85**, p. 195–211.
- Kästle, B., 1990, Fauna und Fazies der kondensierten Sedimente des Dogger und Malm (Bajocium bis Oxfordium) im südlichen Frankenjura: *Münchener Geowissenschaftliche Abhandlungen: Reihe A, Geologie und Paläontologie*, v. **18**, 134 p.
- Kaufmann, A., 1998, Versteinerte Meerestiere von Aarburg: *Aarburger Neujahrsblatt*, v. **1998**, p. 7–10.
- Klikushin, V.G., 1996, Late Jurassic crinoid from Sudak environ (Crimea): *Palaeontographica Abteilung A*, v. **238**, p. 97–151.
- Knorr, G.W., 1775, Sammlung von Merckwürdigkeiten der Natur und Alterthümern des Erdbodens Welche Petrifizirte Körper Enthalten: *Nürnberg, printed by the author*, v. **1**, 177 p.
- Krajewski, M., Ferré, B., and Salamon, M.A., 2020, Cyrtocrinids (Cyrtocrinida, Crinoidea) and other associated crinoids from the Jurassic (Kimmeridgian–Tithonian)–Cretaceous (Berriasian–Barremian) of the Carpathian Foredeep basement (western Ukraine): *Geobios*, v. **60**, p. 61–77.
- Leinfelder, L.R., Schmid, D.U., Nose, M., and Werner, W., 2002, Jurassic reef patterns, the expression of a changing globe, in Flügel, E., Kiessling W., and Golonka, J., eds., *Phanerozoic Reef Patterns: Society for Sedimentary Geology, SEPM Special Publication*, v. **72**, p. 465–520.
- Lendinez, G.A., Fernandez, R., and Carls, P., 1989, *Mapa Geológico de España*: Azuara, Spain, Instituto Tecnológico GeoMinero de España, scale 1:50,000, 25 p.
- Manni, R., and Nicosia, U., 1990, New crinoids from Liassic sediments nearby Tivoli (central Italy): rearrangement in the systematics of eudesicrinids and cotyledermatids: *Geologica Romana*, v. **26**, p. 81–92.
- Manni, R., Nicosia, U., and Szabó, J., 1992, Late Jurassic crinoids from the Eperkés-hegy (Bakony Mts., Hungary): *Fragmenta Mineralogica et Palaeontologica*, v. **15**, p. 115–137.
- Meléndez, G., Atrops, F., Ramajo, J., Urresti, I.P., and Delvene, G., 2006, Upper Oxfordian to lower Kimmeridgian successions in the NE Iberian Range (E Spain): some new stratigraphical and palaeontological data: *Neues Jahrbuch für Geologie und Paläontologie*, v. **241**, p. 203–224.
- Meléndez, G., Atrops, F., Bello, J., Brochwicz-Lewinski, W., D'Arpa, C., Fözy, I., Urresti, I.P., Ramajo, J., and Sequieros, L., 2009, The Oxfordian ammonite genus *Passendorferia* Brochwicz-Lewinski and the Tethyan subfamily Passendorferiinae Meléndez: origin and palaeobiogeography: *Volumina Jurassica*, v. **7**, p. 113–134.
- Miller, J.S., 1821, A natural history of the Crinoidea or lily-shaped animals, with observations on the genera *Asteria*, *Euryale*, *Comatula*, and *Marsupites*: Bristol, UK, C. Frost, 150 p.
- Moore, R.C., Lalicker C.G., and Fischer, A.G., 1952, *Invertebrate Fossils*: New York, Toronto, London, McGraw-Hill, 766 p.
- Münster, G.G.Z., 1839, Beschreibung einiger neuer Crinoideen aus der Übergangsformation: *Beiträge zur Petrefactenkunde*, v. **1**, p. 1–167.
- Nicosia, U., 1991, Mesozoic crinoids from the north-western Turkey: *Estratto da Geologica Romana*, v. **27**, p. 389–436.
- Oji, T., 1996, Is predation intensity reduced with increasing depth? Evidence from the West Atlantic stalked crinoid *Endoxocrinus parrae* (Gervais) and implications for the Mesozoic Marine Revolution: *Paleobiology*, v. **22**, p. 339–351.
- Pictet, F.J., 1857, *Traité de Paléontologie: ou, Histoire Naturelle des Animaux Fossiles Considérés dans leurs Rapports Zoologiques et Géologiques (second edition)*: Paris, J-B Baillière et Fils, v. **4**, 768 p.
- Quenstedt, F.A. von, 1852, *Handbuch der Petrefaktenkunde*: Tübingen, H. Laupp, 755 p.
- Quenstedt, F.A. von, 1858, *Der Jura*: Tübingen, H. Laupp, 842 p.
- Quenstedt, F.A. von, 1867, *Handbuch der Petrefaktenkunde, 2nd edition*: Tübingen, H. Laupp, 982 p.
- Quenstedt, F.A. von, 1876, *Petrefaktenkunde Deutschlands, 1 abt. Echinodermen (Asteriden und Encriniden) und Atlas zu den Echiniden und Encriniden*: Leipzig, Fues's Verlag (R. Reisland), 724 p.
- Quenstedt, F.A. von, 1885, *Handbuch der Petrefaktenkunde, 3rd edition*: Tübingen, H. Laupp, 1239 p.
- Quereilhac, P., and Guinot, Y., 2011, Les “Marnes à theoi” de Pamproux (Deux-Sèvres, France), Sous-Zone à Antecedens (Oxfordien moyen, Zone à Plicatilis): diversité des faunes et découverte de nouvelles espèces d’ammonites: *Carnet de Géologie*, v. **11** p. 21–61.
- Ramajo, J., 2006, *Evolución Sedimentaria del Calloviense–Oxfordiense en el Sector Central de la Cordillera Ibérica [Ph.D. dissertation]*: Rama, Aragonese, University of Zaragoza, 405 p.
- Ramajo, J., and Aurell, M., 2008, Long-term Callovian–Oxfordian sea-level changes and sedimentation in the Iberian carbonate platform (Jurassic, Spain): possible eustatic implications: *Basin Research*, v. **20**, p. 163–184.
- Rasmussen, H.W., 1961, A Monograph on the Cretaceous Crinoidea: *Biologiske Skrifter udgivet af Det Kongelige Danske Videnskabskabes Selskab*, v. **12**, 428 p.
- Rasmussen, H.W., 1975, Neue crinoiden aus der Oberkreide bei Hannover: *Bericht der Naturhistorischen Gesellschaft zu Hannover*, v. **119**, p. 279–283.
- Rasmussen, H.W., 1978, Systematic description, Articulata, in Moore, R. C., and Teichert, C., eds., *Treatise on Invertebrate Paleontology, Pt. T, Echinodermata 2*: Lawrence, Kansas, Geological Society of America and University of Kansas Press, p. T813–T927.
- Remès, M., 1902, Nachträge zur Fauna von Stramberg: *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, v. **14**, p. 200–229.
- Roemer, F.A., 1855, Erste Periode, Kohlen-Gebirge, in Bronn, H.G., ed., *Leithaea Geognostica (third edition)*: Stuttgart, E. Schweizerbart, v. **2**, 788 p.
- Romano, M., Manni, R., and Nicosia, U., 2016, Phylogenetic analysis of cyrtocrinid crinoids and its influence on traditional classifications: *Journal of Mediterranean Earth Sciences*, v. **8**, p. 1–21.
- Rouse, G.W., Jermin, L.S., Wilson, N.G., Eeckhaut, I., Lanterbecq, D., et al., 2013, Fixed, free, and fixed: the fickle phylogeny of extant Crinoidea (Echinodermata) and their Permian–Triassic origin: *Molecular Phylogenetics and Evolution*, v. **66**, p. 161–181.
- Salamon, M.A., 2008, Jurassic cyrtocrinids (Cyrtocrinida, Crinoidea) from extra-Carpathian Poland: *Palaeontographica Abteilung A*, v. **285**, p. 77–99.
- Salamon, M.A., and Feldman-Olszewska, A., 2018, (Crinoidea, Echinodermata) from the Middle Jurassic (Callovian) of eastern Poland: a case study from the Żebrak IG 1 borehole: *Annales Societatis Geologorum Poloniae*, v. **88**, p. 273–283.
- Salamon, M.A., and Gorzelak, P., 2007, Evidence of shallow-water cyrtocrinids (Crinoidea) from the Callovian of Poland: *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, v. **244**, p. 257–260.
- Salamon, M.A., and Gorzelak, P., 2010a, Cyrtocrinids (Echinodermata, Crinoidea) from Upper Jurassic Stramberk-type limestones in southern Poland: *Palaeontology*, v. **53**, p. 869–885.
- Salamon, M.A., and Gorzelak, P., 2010b, Late Cretaceous crinoids (Crinoidea) from eastern Poland: *Palaeontographica Abteilung A*, v. **291**, p. 1–43.
- Salamon, M.A., Gorzelak, P., and Zatoń, M., 2009, In quest of cyrtocrinids origins: evidence from Late Triassic ossicles from the Tatra Mountains: *Acta Palaeontologica Polonica*, v. **54**, p. 171–174.
- Salamon M.A., Bubík M., Ferré B., Szydło A., Nescieruk P., Plachno B.J., Brachaniec T., and Pascza K., 2020, Additional data on

- post-Paleozoic sea-lilies (Crinoidea, Echinodermata) from the Outer Carpathians of the Czech Republic and Poland: *Carnets de Géologie*, v. **20**, p. 283–299.
- Salamon, M.A., Bubík, M., Ferré, B., Duda, P., and Plachno, B.J.**, 2021, *Hrabalicrinus zitti* gen. et sp. nov., and other Upper Jurassic crinoids (Echinodermata, Crinoidea) from the Brno area (Czech Republic): *Annales de Paléontologie*, v. **107**, 102482, <https://doi.org/10.1016/j.annpal.2021.102482>.
- Strasser, A., Aurell, M., Badenas, B., Meléndez, G., and Tomas, S.**, 2005, From platform basin to swell: orbital control on sedimentary sequences in the Oxfordian, Spain: *Terra Nova*, v. **17**, p. 407–413.
- Syverson, V.J., Messing, C.G., Stanley, K., and Baumiller, T.K.**, 2015, Growth, injury, and population dynamics in the extant cyrtocrinid *Holopus mikihe* (Crinoidea, Echinodermata) near Roatan, Honduras: *Bulletin of Marine Sciences*, v. **91**, p. 47–61.
- Urresti, I.P.**, 1995, *Estudio bioestratigráfico y paleontológico basado en Ammonoideos del Oxfordiense Superior en la Rama Aragonesa de la Cordillera Ibérica [M.Sc. thesis]*: Zaragoza, Spain, University Zaragoza, 102 p.
- Urresti, I.P.**, 1996, Upper Oxfordian ammonite successions from the north-eastern Iberian Chain: new biostratigraphic data: *Coloquios de Paleontología*, v. **48**, p. 125–145.
- Veron, J.E.N.**, 1995, *Corals in Space & Time. The Biogeography & Evolution of the Scleractinia*: Ithaca, NY, London, Cornell University Press, 321 p.
- von Schlotheim, E.F.**, 1813, Beiträge zur Naturgeschichte der Versteinerungen in geognostischer Hinsicht: *Mineralogisches Taschenbuch*, v. **7**, p. 3–134.
- Wilson, M.A., Reinthal, E.A., and Ausich, W.A.**, 2014, Parasitism of a new apiocrinitid crinoid species from the Middle Jurassic (Callovian) of southern Israel: *Journal of Paleontology*, v. **88**, p. 1212–1221.
- Zamora, S.**, 2022, Systematics, taphonomy, and paleoecology of Millericrinid (Millericrinida, Articulata, Crinoidea) from the late Jurassic of Spain: *Contributions from the Museum of Paleontology, University of Michigan*, v. **34**, p. 82–102.
- Zamora, S., and López-Horgue, M.A.** 2022, A shallow-water cyrtocrinid crinoid (Articulata) from the upper Albion of the Western Pyrenees, North Spain: *Cretaceous Research*, v. **134**, 105161, <https://doi.org/10.1016/j.cretres.2022.105161>.
- Zamora, S., Aurell, M., Veitch, M., Saulsbury, J., López-Horgue, M.A., Ferratges F.A., Arz, J.A., and Baumiller, T.K.**, 2018, Environmental distribution of post-Palaeozoic crinoids from the Iberian and south-Pyrenean basins, NE Spain: *Acta Palaeontologica Polonica*, v. **63**, p. 779–794.
- Zatoń, M., Salamon, M.A., and Kaźmierczak, J.**, 2008, Cyrtocrinids (Crinoidea) and associated stalked crinoids from the lower/middle Oxfordian (Upper Jurassic) shelfal deposits of southern Poland: *Geobios*, v. **41**, p. 559–569.
- Zempolich, W.G.**, 1993, The drowning succession in Jurassic carbonates of the Venetian Alps, Italy: a record of supercontinent breakup, gradual eustatic rise, and eutrophication of shallow-water environments, in Loucks, R.G., and Sarg, J.F., eds., *Carbonate Sequence Stratigraphy: Recent Developments and Applications*: American Association of Petroleum Geologists Memoir 57, p. 63–105.
- Žitt, J.**, 1974, *Sclerocrinus* Jaekel, 1891 and *Proholopus* Jaekel, 1907 (Crinoidea, Cyrtocrinida) from the Lower Cretaceous of Štramberk (Czech Republic): *Sborník Geologických věd, Paleontologie*, v. **16**, p. 7–34.
- Zittel, K.A. von**, 1876–1880, *Handbuch der Palaeontologie, Band 1, Palaeozoologie*: München and Leipzig, Germany, R. Oldenbourg, 765 p.