





Article

Nomenclature of the ancylite supergroup

Mineralogy, petrology and geochemistry of pegmatites: Alessandro Guastoni memorial issue

Yanjuan Wang^{1,2} , Fabrizio Nestola² , Zengqian Hou¹, Ritsuro Miyawaki³, Igor V. Pekov⁴, Xiangping Gu⁵ ,
Guochen Dong¹ and Kai Qu^{6,7} 

¹School of Earth Sciences and Resources, China University of Geosciences, Beijing 100083, China; ²Department of Geosciences, University of Padova, Padova 35131, Italy; ³Department of Geology, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba 305-0005, Japan; ⁴Faculty of Geology, Moscow State University, Vorobievsky Gory, Moscow 119991, Russia; ⁵School of Geosciences and Info-Physics, Central South University, Changsha 410083, Hunan, China; ⁶Tianjin Center, China Geological Survey, Tianjin 300170, China; and ⁷School of Earth Sciences and Engineering, Nanjing University, Nanjing 210023, China

Abstract

The ancylite supergroup has been approved by the Commission on New Minerals, Nomenclature and Classification of the International Mineralogical Association, with the general crystal chemical formula $(M_x^{3+}M_{2-x}^{2+})(\text{CO}_3)_2[(\text{OH})_x \cdot (2-x)\text{H}_2\text{O}]$ ($1 \leq x \leq 2$, $Z = 2$). The ancylite supergroup can be divided into two groups defined by different proportions of the M cation and hydroxyl anion and/or water molecule: the ancylite group is defined for $1 \leq x \leq 1.5$; the kozoite group is defined for $1.5 < x \leq 2$. The ancylite supergroup minerals are orthorhombic with space group $Pm\bar{c}n$, or monoclinic with space group $Pm11$, and have a crystal structure with species-defining trivalent and divalent M cations ($M = \text{La}^{3+}$, Ce^{3+} , Nd^{3+} , Ca^{2+} , Sr^{2+} and Pb^{2+}) which centre ten-vertex polyhedra formed by oxygen atoms at three independent O sites. Two vertices of the triangular $(\text{CO}_3)^{2-}$ anion are oxygen atoms, whereas the third one, O(3), is statistically filled with $(\text{OH})^-$ groups and H_2O molecules. The triangular faces of three oxygen atoms of MO_{10} coordination polyhedra join the chains of this ten-vertex polyhedron, which is extended along the c axis. The (CO_3) triangles connect chains in three dimensions. To date, eight valid mineral species with $M^{2+} = \text{Sr}^{2+}$, Ca^{2+} and Pb^{2+} belong to the ancylite group [ancylite-(La), ancylite-(Ce), calcioancylite-(La), calcioancylite-(Ce), calcioancylite-(Nd), gysinite-(La), gysinite-(Ce) and gysinite-(Nd)]. Two hydroxyl carbonates with only rare earth elements as species-defining cations, kozoite-(La) and kozoite-(Nd) are members of the kozoite group.

Keywords: ancylite supergroup; ancylite; calcioancylite; gysinite; kozoite; nomenclature; classification

(Received 9 December 2023; accepted 10 February 2024; Accepted Manuscript published online: 19 February 2024)

Introduction

Ancylite-type minerals, which are hydrous/hydroxyl carbonates containing rare earth elements (REEs), Ca^{2+} , Sr^{2+} and Pb^{2+} as major cations, commonly occur in alkaline rocks as late accessories or, in some types of carbonatites, as rock-forming minerals and important concentrators of light rare earth elements (LREE) and strontium (Fig. 1).

Prior to this work, neither ancylite group nor supergroup were formally approved by the Commission on New Minerals, Nomenclature and Classification of the International Mineralogical Association (IMA–CNMNC). However, the term ‘ancylite group’ appeared in the Fleischer’s Glossary of Mineral Species in 2008 (Back and Mandarino, 2008), has been repeated in subsequent editions, and is now a common term in literature.

Corresponding author: Yanjuan Wang; Email: wangyanjuan_cugb@foxmail.com

This paper is part of a thematic set on pegmatites in memory of Alessandro Guastoni
Guest Editor: Simone Molinari

Cite this article: Wang Y., Nestola F., Hou Z., Miyawaki R., Pekov I.V., Gu X., Dong G. and Qu K. (2024) Nomenclature of the ancylite supergroup. *Mineralogical Magazine* 88, 677–681. <https://doi.org/10.1180/mgm.2024.8>

The general chemical formula for ancylite-group minerals was previously described as: $M_x^{3+}M_{2-x}^{2+}(\text{CO}_3)_2(\text{OH})_x \cdot (2-x)\text{H}_2\text{O}$ ($Z = 2$), where $1 < x \leq 2$, $M^{3+} = \text{REE}^{3+}$ such as La^{3+} , Ce^{3+} and Nd^{3+} , $M^{2+} = \text{Ca}^{2+}$, Sr^{2+} and Pb^{2+} (Dal Negro *et al.*, 1975; Sarp and Bertrand, 1985; Miyawaki *et al.*, 2000). Note that there are significant differences in the formulae given in the original descriptions of valid mineral species belonging to the ancylite group, e.g. ancylite-(La): $\text{Sr}(\text{La,Ce})(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O}$ (Yakovenchuk *et al.*, 1997); calcioancylite-(Ce): $(\text{Ce,Ca,Sr})(\text{CO}_3)(\text{OH},\text{H}_2\text{O})$ (Belovitskaya *et al.*, 2013); calcioancylite-(Nd): $\text{Nd}_{2.8}\text{Ca}_{1.2}(\text{CO}_3)_4(\text{OH})_3 \cdot \text{H}_2\text{O}$ (Orlandi *et al.*, 1990). Without detailed knowledge of the crystal chemistry of this group, it is not easy to understand whether trivalent REEs and the other divalent cations occupy the same crystallographic site or independent sites. For example: chemically, formulae of some REE minerals can be written as $\text{La}_2\text{Sr}(\text{CO}_3)_3(\text{OH})_2 \cdot \text{H}_2\text{O}$ or $\text{La}_3\text{Sr}(\text{CO}_3)_4(\text{OH})_3 \cdot \text{H}_2\text{O}$, the ratio of La to Sr is 2:1 or 3:1, which is different from the ideal formula of ancylite-(La) $[\text{LaSr}(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O}]$ in the current IMA mineral list (Pasero, 2024). This suggests potentially two new mineral species, however they are intermediate between the end-members, $[\text{LaSr}(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]]$ (revised ideal formula, this work)




Figure 1. The localities of ancylium supergroup minerals worldwide, following the literature records on [Mindat.org](https://www.mindat.org) (Accessed November 2023). TL = Type Locality.

[ancylium-(La)] and $\text{La}_2(\text{CO}_3)_2(\text{OH})_2$ [kozyte-(La)]. This ambiguity is potentially confusing, not only for novices but also for expert crystallographers. The situation is additionally complicated by the specific behaviour of the $M^{3+}:M^{2+}$ ratio in the solid-solution series between ancylium-(Ce) and calcioancylium-(Ce) (Pekov *et al.*, 1997).

These problems of inconsistent chemical formulae, the complex chemical variability of the ancylium supergroup minerals, and vague boundaries between different hierarchies, have prompted us to formally establish the ancylium supergroup and to recommend updating the ideal formulae of ancylium supergroup members in the IMA–CNMNC official mineral list (Table 1). The proposal for the “*Nomenclature of the ancylium supergroup*” has been approved by the IMA–CNMNC (Bosi *et al.*, 2024). The present work aims to not only identify and classify complex ancylium supergroup minerals but to also convey important chemical information for mineralogists and ore geologists who are interested in alkaline complexes and REE behaviours.

Crystal structure

In the structure of the ancylium supergroup minerals, trivalent and divalent cations (M cations), such as La^{3+} , Ce^{3+} , Nd^{3+} , Ca^{2+} , Sr^{2+} and Pb^{2+} , centre ten-vertex polyhedra, formed by oxygen atoms at three independent O sites. Two of them, at the apices of triangular $(\text{CO}_3)^{2-}$ anion, are occupied by oxygen atoms, whereas the third one, expressed as O(3) in Table 2, is statistically filled with $(\text{OH})^-$ groups and H_2O molecules. The chains of this ten-vertex polyhedral are stretched along the c axis and connected by sharing the triangular faces of three oxygen atoms of MO_{10} coordination polyhedra. The chains are interconnected into a three-dimensional framework via (CO_3) triangles (Fig. 2a).

The crystal structure of ancylium-(Ce) was solved for the first time by Dal Negro *et al.* (1975). Belovitskaya *et al.* (2002) performed the structure refinement for two specimens of ancylium-(Ce) within the space groups $Pm\bar{c}n$ and $Pm\bar{c}2_1$. Belovitskaya *et al.* (2013) studied the crystal structure of

Table 1. Formulae of ancylium supergroup minerals.

Name	Formula in the current IMA list	Simplified crystal chemical formula	Recommended ideal formula
Ancylium-(La) ¹	$\text{LaSr}(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{La,Sr})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{LaSr})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Ancylium-(Ce) ²	$\text{CeSr}(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{Ce,Sr})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{CeSr})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Calcioancylium-(La) ³	$(\text{La,Ca})(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{La,Ca})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{La,Ca})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Calcioancylium-(Ce) ⁴	$(\text{Ce,Ca,Sr})(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})$	$(\text{Ce,Ca})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{Ce,Ca})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Calcioancylium-(Nd) ⁵	$\text{Nd}_{2.8}\text{Ca}_{1.2}(\text{CO}_3)_4(\text{OH})_3\cdot\text{H}_2\text{O}$	$(\text{Nd,Ca})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{Nd,Ca})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Gysinite-(La) ⁶	$\text{PbLa}(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{La,Pb})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{La,Pb})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Gysinite-(Ce) ⁷	$\text{PbCe}(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{Ce,Pb})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{Ce,Pb})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Gysinite-(Nd) ⁸	$\text{PbNd}(\text{CO}_3)_2(\text{OH})\cdot\text{H}_2\text{O}$	$(\text{Nd,Pb})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$	$(\text{Nd,Pb})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, $Z = 2$
Kozyte-(La) ⁹	$\text{La}_2(\text{CO}_3)_2(\text{OH})$	$\text{La}_2(\text{CO}_3)_2(\text{OH})_2$	$\text{La}(\text{CO}_3)(\text{OH})$, $Z = 4$
Kozyte-(Nd) ¹⁰	$\text{Nd}(\text{CO}_3)(\text{OH})$	$\text{Nd}_2(\text{CO}_3)_2(\text{OH})_2$	$\text{Nd}(\text{CO}_3)(\text{OH})$, $Z = 4$

References: ¹ Yakovenchuk *et al.* (1997); Petersen *et al.* (2001); ² Dal Negro *et al.* (1975); Belovitskaya *et al.* (2002); ³ Wang *et al.* (2023); ⁴ Belovitskaya *et al.* (2013); ⁵ Orlandi *et al.* (1990); ⁶ Wu *et al.* (2023); ⁷ Kampf *et al.* (2023); ⁸ Chabot and Sarp (1985); ⁹ Miyawaki *et al.* (2003); ¹⁰ Miyawaki *et al.* (2000).

Table 2. Characteristics of the ancylite supergroup described by the general formula $(M_x^{3+}M_{2-x}^{2+})(\text{CO}_3)_2[(\text{OH})_x(2-x)\text{H}_2\text{O}]$, with $Z=2$.

Group	Root name	Range	Species	Essential <i>M</i> cation	Essential O(3) anion/mol.	Crystal system	Space group	<i>a</i> (Å)	<i>b</i> (Å)	<i>c</i> (Å)	α (°)
Ancylite	ancylite	$1 \leq x \leq 1.5$	ancylite-(La)	La ³⁺ , Sr ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.044	8.541	7.292	90
			ancylite-(Ce)	Ce ³⁺ , Sr ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.03	8.53	7.29	90
	calcioancylite		calcioancylite-(La)	La ³⁺ , Ca ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.0253	8.5152	7.2717	90
			calcioancylite-(Ce)	Ce ³⁺ , Ca ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.0095	8.5006	7.2670	90
	gysinite		calcioancylite-(Nd)	Nd ³⁺ , Ca ²⁺	(OH) ⁻ , H ₂ O	Mon.	<i>Pm</i> 11	4.976	8.468	7.212	90.04
			gysinite-(La)	La ³⁺ , Pb ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	4.986	8.513	7.227	90
			gysinite-(Ce)	Ce ³⁺ , Pb ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.0780	8.6689	7.3255	90
			gysinite-(Nd)	Nd ³⁺ , Pb ²⁺	(OH) ⁻ , H ₂ O	Orth.	<i>Pm</i> <i>cn</i>	5.0028	8.555	7.2392	90
Kozoite	kozoite	$1.5 < x \leq 2$	kozoite-(La)	La ³⁺	(OH) ⁻	Orth.	<i>Pm</i> <i>cn</i>	4.986	8.513	7.227	90
			kozoite-(Nd)	Nd ³⁺	(OH) ⁻	Orth.	<i>Pm</i> <i>cn</i>	4.9829	8.5188	7.2570	90

Note: The references are the same as in Table 1 for each mineral, respectively.

calcioancylite-(Ce) using the Rietveld method and showed that most minerals of the ancylite group were regarded as orthorhombic, and that the whole structure of ancylite group minerals can be derived from orthorhombic carbonates by adding hydroxyl groups that are positioned on the mirror planes and bonded to the M cations. It should be noted that calcioancylite-(Nd) from Baveno, Italy, described by Orlandi *et al.* (1990) has revealed that M^{3+} and M^{2+} cations order into four M sites in the ancylite-type structure to lower the symmetry from orthorhombic (*Pm**cn*) to monoclinic (*Pm*11) with a minor deviation of α from 90° to $90.04(3)^\circ$ (Belovitskaya *et al.*, 2013). It is the only approved valid mineral species with monoclinic symmetry in the ancylite supergroup so far (Table 2). However, if the crystal structures of the polymorphs have essentially the same topology, differing only in terms of a structural distortion or in the order–disorder

relationship of some of the atoms comprising the structure, such polymorphs are not regarded as separate species (Nickel and Grice, 1998). The recommendation is that both of the ordered monoclinic and disordered orthorhombic phases be classified into a mineral species. The difference in symmetry owing to the order–disorder can be indicated by a suffix, as is the polytype distinction, e.g. calcioancylite-(Nd)-*M* and calcioancylite-(Nd)-*O*.

Nomenclature and classification of the ancylite supergroup

In the frame of the present IMA-approved nomenclature (Mills *et al.*, 2009), ancylite-type minerals are divided into two groups: the ancylite group and the kozoite group. These mineral groups compose the ancylite supergroup (named after the first adequately characterised mineral). In addition, the members are classified




Figure 2. (a) The crystal structure of ancylite supergroup minerals viewed along [001]; (b) quaternary diagram, showing the boundaries between kozoite $\text{REE}(\text{CO}_3)(\text{OH})$, ancylite $(\text{REESr})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, calcioancylite $(\text{REECa})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$, and gysinite $(\text{REEPb})(\text{CO}_3)_2[(\text{OH})(\text{H}_2\text{O})]$. Figure 2a drawn using Vesta software (Momma and Izumi, 2011).

into root names corresponding to the essential *M* cations: ancylite ($REE^{3+}Sr^{2+}$), calcioancylite ($REE^{3+}Ca^{2+}$), gysinite ($REE^{3+}Pb^{2+}$) and kozoite (REE^{3+}).

The general chemical formula of ancylite supergroup minerals can be defined as $(M^{3+}_x M^{2+}_{2-x})(CO_3)_2[(OH)_x \cdot (2-x)H_2O]$ ($1 \leq x \leq 2$, $Z=2$), and different combinations of M^{3+} and M^{2+} constituents should be regarded as separate mineral species. Both M^{3+} and M^{2+} cations generally occupy the same crystallographic site, and the formula is charge-balanced through the following substitution mechanism: $M^{3+} + OH^- \rightleftharpoons M^{2+} + H_2O$. The excess positive charge of M^{3+} is compensated by the incorporation of $(OH)^-$, and the number of hydroxyl ions and water molecules are equivalent to those of M^{3+} and M^{2+} , respectively (Wang *et al.*, 2023). Taking into account the recently defined species, the boundaries between different hierarchies of the ancylite supergroup is as follows (Fig. 2b):

- (1) $1 \leq x \leq 1.5$, REE^{3+} dominant for M^{3+} : ancylite group
 - (a) Sr^{2+} dominant for M^{2+} : ancylite (root name)
 - (b) Ca^{2+} dominant for M^{2+} : calcioancylite (root name)
 - (c) Pb^{2+} dominant for M^{2+} : gysinite (root name)
- (2) $1.5 < x \leq 2$, REE^{3+} dominant for M^{3+} : kozoite group

Each distinct mineral species within the ancylite supergroup has a hyphenated suffix between parentheses, the Levinson modifier (Levinson, 1966), indicating the dominant *REE* constituent (Hatert and Burke, 2008; Hatert *et al.*, 2013). To date, the supergroup includes ten valid mineral species: ancylite-(La), ancylite-(Ce), calcioancylite-(La), calcioancylite-(Ce), calcioancylite-(Nd), gysinite-(La), gysinite-(Ce), gysinite-(Nd), kozoite-(La) and kozoite-(Nd) (Table 2).

Historical synopsis of the ancylite supergroup minerals

The name ancylite first appeared in literature in 1898, when this mineral was discovered in Narsaarsuk, Southern Greenland. The name is taken from the Greek word ἀγκυλός (ankylos) for ‘curved’, in allusion to the planes of the crystals usually rounded and distorted. The formula was given by Flink (1898, 1901) as $Ce_4Sr_3(CO_3)_7(OH)_4 \cdot 3H_2O$. The La-dominant ancylite species, ancylite-(La), was discovered by Yakovenchuk *et al.* (1997) from a hydrothermal vein at the Kukisvumchorr mountain, Khibiny alkaline complex, Kola peninsula, Russia.

A brief but interesting review of the history of the discoveries and early studies of calcioancylite-(Ce) has been given by Pekov *et al.* (1997): Chernik (1904) described a Ca-dominant analogue of ancylite but without a clear locality, with the address written as “Western land of the Russian Empire” (Pekov *et al.*, 1997); the name ‘calcio-ancylite’ in literature was first mentioned by Chernik (1923), and it has been renamed to calcioancylite-(Ce) according to the revised nomenclature for the *REE*-bearing minerals (Nickel and Mandarino, 1987). The Nd-dominant calcioancylite was found in miarolitic cavities of Baveno granite, Italy (Orlandi *et al.*, 1990). At the same time, Orlandi *et al.* (1990) re-examined the type samples identified as ‘weibyeite’ by Artini (1915) from Baveno, and their powder X-ray diffraction and chemical analyses indicated that the material was actually a calcioancylite-(Ce). It should be noted that the term ‘weibyeite’ was first introduced by Brögger (1890) to describe a *REE* carbonate phase in Langesundsfjorden, Norway. If it is the same material as the ‘weibyeite’ from Baveno, it should be the first finding of the

ancylite-type mineral, but, unfortunately, Saebø (1963) discredited Broegger’s ‘weibyeite’ and the result shows that it is just a bastnäsite-(Ce) pseudomorph. Recently, the La-dominant calcioancylite species, calcioancylite-(La), was discovered from Gejiu nepheline syenite, Yunnan Province, China (Wang *et al.*, 2023).

Gysinite-(Nd), the first Pb-dominant member of the ancylite group, was discovered by Sarp and Bertrand (1985) in a specimen from the mineral collection at the Geneva Natural History Museum. The sample was originally labelled ‘schuilingite’ from Shinkolobwe, Shaba, Zaïre (now the Democratic Republic of the Congo). The mineral is named after the late Professor Marcel Gysin (1891–1974) from the University of Geneva. Almost four decades later, the La-dominant mineral species, gysinite-(La), was discovered in Iujavrite from the Saima alkaline complex, Liaoning Province, China (Wu *et al.*, 2023). Gysinite-(Ce), the mineral species with Ce prevailing among *REEs*, has recently been found in Abendröthe Mine, located in Sankt Andreasberg, Braunlage, Goslar District, Lower Saxony, Germany (Kampf *et al.*, 2023).

In 1997, a kind of pale pink *REE* carbonate was collected by Koichi Takeuchi from alkali olivine basalt from Hizen-cho, Saga Prefecture, Japan. Miyawaki *et al.* (1998) found that it was different from other ancylites because it was characterised by very low Ca and Sr and almost absent Pb. Several years later, two new minerals and mineral names were approved by the IMA Commission on New Mineral and Mineral Names (IMA–CNMMN, the predecessor of the IMA–CNMNC, as kozoite-(Nd) and kozoite-(La), respectively (Miyawaki *et al.*, 2000; 2003). The minerals are named in honour of the late Kozo Nagashima (1925–1985), a chemist and pioneer in the study of the crystal chemistry of rare earth minerals in Japan.

Note that if considering the extension of the solid solution of the ancylite series beyond the point $M^{2+}/M^{3+} < 0.5$, there probably exists the potential compound $Ca(CO_3)_2H_2O$ (for x close to 0 in the general formula) (Sarp and Bertrand, 1985). However, monohydrocalcite, as a mineral with the chemical formula $Ca(CO_3)(H_2O)$ that occurs in nature (Semenov, 1964), has a topology and symmetry differing from the ancylite-type series (Effenberger, 1981; Swainson, 2008). While it is not clear if that composition could be stable with the ancylite topology, previous studies that reviewed the reported chemical compositions of ‘ancylites’ and ‘calcioancylites’ from several localities showed that the value ‘ x ’ in the general formula, $(M^{3+}_x M^{2+}_{2-x})(CO_3)_2[(OH)_x \cdot (2-x)H_2O]$, always exceeded 1 (Bulakh *et al.*, 1998; Pekov *et al.*, 1997). This indicates that ancylite supergroup minerals are not intermediate solid solutions between $M^{3+}(CO_3)(OH)$ and $M^{2+}(CO_3)(H_2O)$. The end-members of this group are $M^{3+}(CO_3)(OH)$ and $(M^{3+}M^{2+})(CO_3)_2[(OH) \cdot H_2O]$ (Sarp and Bertrand, 1985; Miyawaki *et al.*, 2000). Taking into account the above reasons and the fact that no minerals having an ancylite-type topology and a chemical composition with x close to 0 (even < 0.5) have been found in Nature so far, the potential ancylite-type topology $M^{2+}(CO_3)(H_2O)$ phases are not included in this nomenclature.

Acknowledgements. The helpful comments from three anonymous reviewers are greatly appreciated. We are grateful to Stuart Mills and CNMNC members for their valuable suggestions regarding the nomenclature. This study was supported by the Natural Science Foundation of China (NSFC) (Grant: 92062217, 92062220, and 42072054 for GD, KQ, and XG, respectively). YW and KQ acknowledge financial support from the China Scholarship Council (CSC) (Grant: 202106400047, 202108575009, respectively).

Competing interests. One of the co-authors is a guest member of the editorial board of Mineralogical Magazine for the special issue “Mineralogy, petrology and geochemistry of pegmatites”. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

References

- Artini E. (1915) Due minerali di Baveno contenenti terra rare – weibeyite e bazzite. *Rendiconti dell'Accademia Nazionale dei Lincei*, **24**, 313–319.
- Back M.E and Mandarino J.A. (2008) *Fleischer's Glossary of Mineral Species 2008*. The Mineralogical Record Inc., Tucson, USA, pp.275.
- Belovitskaya Y.V., Pekov I.V., Gobetchia E.R., Yamnova N.A., Kabalov Yu.K., Chukanov N.V. and Schneider J. (2002) Crystal structures of two ancylite modifications. *Crystallography Reports*, **47**, 223–228.
- Belovitskaya Y.V., Pekov I.V., Gobetchia E.R. and Kabalov Y.K. (2013) Refinement of the crystal structure of calcioancylite-(Ce) by the Rietveld method. *Crystallography Reports*, **58**, 216–219.
- Bosi F., Hatert F., Pasero M. and Mills S.J. (2024) Establishment of the ancylite supergroup, CNMNC Newsletter 78. *Mineralogical Magazine*, **88**, 345–349. <https://doi.org/10.1180/mgm.2024.23>
- Brögger W.C. (1890) Die Mineralien der Syentipegmatitgänge der südnorwegischen Augit- und Nephelinsyenite. 71. Weibeyit und 72. Parisit. *Zeitschrift für Kristallographie*, **16**, 650–654.
- Bulakh A.G., Zaitsev A.N., Le Bas M.J. and Wall F. (1998) Ancylite-bearing carbonatites of the Sebyavr massif, Kola Peninsula, Russia. *Neues Jahrbuch für Mineralogie, Monatshefte*, **4**, 171–192.
- Chabot B. and Sarp H. (1985) Structure refinement of Gysinite $\text{La}_{0.16}\text{Nd}_{1.18}\text{Pb}_{0.66}(\text{CO}_3)_2(\text{OH})_{1.34} \cdot 0.66\text{H}_2\text{O}$. *Zeitschrift für Kristallographie*, **171**, 155–158.
- Chernik G.P. (1904) A nature and chemical composition of one a new cerian ancylite-like mineral. *Zapiski Imperatorskogo Mineralogicheskogo Obshchestva*, **71**, 43–54 [in Russian].
- Chernik G.P. (1923) Chemical examination of calcian ancylite and associated minerals from Khibina tundra. *Izvestiya Rossiyskoy Akademii Nauk*, Ser.VI, **17**, 81–94 [in Russian].
- Dal Negro A., Rossi G. and Tazzoli V. (1975) The crystal structure of ancylite, $(\text{RE})_x(\text{Ca},\text{Sr})_{2-x}(\text{CO}_3)_2(\text{OH})_x \cdot (2-x)\text{H}_2\text{O}$. *American Mineralogist*, **60**, 280–284.
- Effenberger H. (1981) Kristallstruktur und Infrarot-Absorptionsspektrum von synthetischem Monohydrocalcit, $\text{CaCO}_3 \cdot \text{H}_2\text{O}$. *Monatshefte für Chemie*, **112**, 899–909.
- Flink G. (1898) Berättelse om en Mineralogisk Resa i Syd-Grönland sommaren 1897. 4. Gult anataslikt, *Meddelelser om Grönland*, **14**, 235–235.
- Flink G. (1901) On the minerals from Narsarsuk on the Firth of Tunugdliarfik in Southern Greenland. 11 Ancylite, *Meddelelser om Grönland*, **24**, 49–56.
- Hatert F. and Burke E.A. (2008) The IMA–CNMNC dominant-constituent rule revisited and extended. *The Canadian Mineralogist*, **46**, 717–728.
- Hatert F., Mills S.J., Pasero M. and Williams P.A. (2013) CNMNC guidelines for the use of suffixes and prefixes in mineral nomenclature, and for the preservation of historical names. *European Journal of Mineralogy*, **25**, 113–115.
- Kampf A.R., Möhn G., Ma C., Désor J. and Groß M. (2023) Gysinite-(Ce), IMA 2023-035, CNMNC Newsletter 74. *Mineralogical Magazine*, **87**, 783–787. <https://doi.org/10.1180/mgm.2023.54>
- Levinson A.A. (1966) A system of nomenclature for rare-earth minerals. *American Mineralogist*, **51**, 152–158.
- Mills S.J., Hatert F., Nickel E.H. and Ferraris G. (2009) The standardisation of mineral group hierarchies: application to recent nomenclature proposals. *European Journal of Mineralogy*, **21**, 1073–1080.
- Miyawaki R., Matsubara S., Yokoyama K. and Takeuchi, K. (1998) An ancylite group mineral in alkali olivine basalt from Saga Prefecture, Japan. *Memoirs of the National Science Museum*, **31**, 49–56.
- Miyawaki R., Matsubara S., Yokoyama K., Takeuchi K., Nakai I. and Terada Y. (2000) Kozoite-(Nd), $\text{Nd}(\text{CO}_3)(\text{OH})$, a new mineral in an alkali olivine basalt from Hizen-cho, Saga Prefecture, Japan. *American Mineralogist*, **85**, 1076–1081.
- Miyawaki R., Matsubara S., Yokoyama K., Iwano S., Hamasaki K. and Yukinori I. (2003) Kozoite-(La), $\text{La}(\text{CO}_3)(\text{OH})$, a new mineral from Mitsukoshi, Hizen-cho, Saga Prefecture, Japan. *Journal of Mineralogical and Petrological Sciences*, **98**, 137–141.
- Momma K. and Izumi F. (2011) VESTA 3 for three-dimensional visualization of crystal, volumetric and morphology data. *Journal of Applied Crystallography*, **44**, 1272–1276.
- Nickel E.H. and Grice J.D. (1998) The IMA Commission on New Minerals and Mineral Names: procedures and guidelines on mineral nomenclature, 1998. *Mineralogy and Petrology*, **64**, 237–263.
- Nickel E.H. and Mandarino J.A. (1987) Procedures involving the IMA commission on New Minerals and Mineral Names, and guidelines on mineral nomenclature. *The Canadian Mineralogist*, **25**, 353–377.
- Orlandi P., Pasero M. and Vezzolini G. (1990) Calcio-ancylite-(Nd), a new REE-carbonate from Baveno, Italy. *European Journal of Mineralogy*, **2**, 413–418.
- Pasero M. (2024) *The New IMA List of Minerals – A Work in Progress – Updated: January 2024*. The Commission on New Minerals, Nomenclature and Classification (CNMNC) of the International Mineralogical Association (IMA). <http://cnmnc.units.it/>.
- Pekov I.V., Petersen O.V. and Voloshin A.V. (1997) Calcio-ancylite-(Ce) from Ilimaussaq and Narssarsuk, Greenland, Kola peninsula and Polar Urals, Russia; ancylite-(Ce) – calcio-ancylite-(Ce) an isomorphous series. *Neues Jahrbuch für Mineralogie, Abhandlungen*, **171**, 309–322.
- Petersen O.V., Niedermayr G., Gault R.A., Brandstetter F. and Giester G. (2001) Ancylite-(La) from the Ilimaussaq alkaline complex, South Greenland: Contribution to the mineralogy of Ilimaussaq, no. 106. *Neues Jahrbuch für Mineralogie - Monatshefte*, **11**, 493–504.
- Saebø P.C. (1963) Contributions to the mineralogy of Norway. No. 20. The identity of weibeyite. *Norsk Geologisk Tidsskrift*, **43**, 441–443.
- Sarp H. and Bertrand J. (1985) Gysinite, $\text{Pb}(\text{Nd},\text{La})(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O}$, a new lead, rare-earth carbonate from Shinkolobwe, Shaba, Zaïre and its relationship to ancylite. *American Mineralogist*, **70**, 1314–1317.
- Semenov E.I. (1964) Hydrated carbonates of sodium and calcium. *Kristallografiya*, **9**, 109–110 [in Russian].
- Swainson I.P. (2008) The structure of monohydrocalcite and the phase composition of the beachrock deposits of Lake Butler and Lake Fellmongery, South Australia. *American Mineralogist*, **93**, 1014–1018.
- Wang Y., Gu X., Dong G., Hou Z., Nestola F., Yang Z., Fan G., Wang Y. and Qu K. (2023) Calcioancylite-(La), $(\text{La},\text{Ca})_2(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O})_2$, a new member of the ancylite group from Gejiu nepheline syenite, Yunnan Province, China. *Mineralogical Magazine*, **87**, 554–560.
- Wu B., Gu X., Rao C., Wang R., Xing X., Wan J. and Zhong, F. (2023) Gysinite-(La), $\text{PbLa}(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O}$, a new rare earth mineral of the ancylite group from the Saima alkaline complex, Liaoning Province, China. *Mineralogical Magazine*, **87**, 143–150.
- Yakovenchuk V.N., Menshikov Y.P., Pakhomovsky Y.A. and Ivanyuk G.Y. (1997) Ancylite-(La) $\text{Sr}(\text{La},\text{Ce})(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O}$ – a new carbonate from hydrothermal vein at the Kukisvumchorr mountain (Khibiny massif) and its comparison with ancylite-(Ce). *Zapiski Vserossiyskogo Mineralogicheskogo Obshchestva*, **126**, 96–108 [in Russian].