

NOTE ON NON-COMMUTATIVE SEMI-LOCAL RINGS

YUKITOSHI HINOHARA

Our aim in this note is to generalize some topological results of commutative noetherian rings to non-commutative rings. As a supplemental remark of [2] we prove in §1 that any right ideal of a complete right semi-local ring is closed, and that

$$\bigcap_{s=1}^{\infty} MJ^s = (0)$$

for any finitely generated right module M over a complete right semi-local ring A where J is the Jacobson radical of A .

In §2 we are concerned with the flatness of modules. C. Lech gave in [7] an ideal theoretical criterion of the flatness of modules over a commutative ring. We notice that his criterion of the flatness is valid for non-commutative rings.

§1. Non-commutative semi-local rings

DEFINITION. *Let A be a ring with a unit element 1 and J its Jacobson radical; A is said to be right semi-local if the following conditions are satisfied:*

- (a) $\bigcap_{s=1}^{\infty} J^s = 0$,
- (b) A is right noetherian,
- (c) A/J satisfies the minimum condition on right ideals.

This definition is due to E. H. Batho who studied the basic property of this class of rings in [2].

By virtue of the condition (a), we may introduce a Hausdorff topology (called the J -adic topology) in A and construct the completion \hat{A} of A with respect to this topology.

For brevity, we call an ideal I of A a nucleus if $\bigcap_{s=1}^{\infty} I^s = 0$ and denote the Jacobson radical of the ring A by $J(A)$.

LEMMA 1. *Let A be a right semi-local ring and \hat{A} the completion of A with respect to the $J(A)$ -adic topology. Then we have*

Received May 18, 1960.

$$J(\hat{A})^s = J(A)^s \hat{A}, \quad \bigcap_{s=1}^{\infty} J(\hat{A})^s = 0,$$

$$J(\hat{A})^s \cap A = J(A)^s, \quad A/J(A)^s \cong \hat{A}/J(\hat{A})^s$$

for any positive integer s .

For the proof we refer to [Theorem 2.3 of 2] and [Theorem 2 of 3].

LEMMA 2. *Let A be a ring with a unit element 1. Then we have the relation $NJ(A) \cong N$ for any finitely generated right A -module $N (\neq 0)$.*

This is [Proposition 2 of 8, p. 200].

LEMMA 3. *Let A be a ring and J its Jacobson radical. Assume the following conditions for A :*

- (a) A/J satisfies the minimum condition on right ideals,
- (b) J is a nucleus and has a finite number of right A -basis,
- (c) A is complete with respect to the J -adic topology.

Then we have the relation

$$\bigcap_{s=1}^{\infty} (M + FJ^s) = M$$

for any finitely generated A -submodule M of a free right A -module F .

We notice that $\bigcap_{s=1}^{\infty} FJ^s = (0)$ since F is a free A -module and J is a nucleus. Therefore we can define a Hausdorff topology in F by taking F, FJ, FJ^2, \dots to be neighbourhoods of zero. Then the closure \bar{N} of any submodule N of F is equal to $\bigcap_{s=1}^{\infty} (N + FJ^s)$.

Before proving Lemma 3 we prove

LEMMA 4. *If a submodule M of F is finitely generated we have $\bar{M} = M + \bar{M}J$.*

Proof. We consider the residue class module F/MJ of which $(\bar{M} + MJ)/MJ$ is a submodule. Since $\bar{M}J \subseteq MJ$, we have $((\bar{M} + MJ)/MJ)J = (0)$. Let \bar{m} be any element of \bar{M} . Then \bar{m} can be written in the following form, for any positive integer t ,

$$\bar{m} = \sum_{i=1}^n m_i \lambda_i + j_t$$

where $\{m_1, \dots, m_n\}$ is a A -basis of M , $\lambda_i \in A$ and $j_t \in FJ^t$. Now $m = (\bar{m}A + M + MJ)/\bar{M}J$ is a finitely generated module and $mJ = 0$. Therefore m is considered

as a finitely generated A/J -module. Since A/J satisfies the minimum condition on right ideals, the module \mathfrak{m} satisfies the minimum condition on submodules. Consider the descending sequence of submodules

$$\mathfrak{m} \supseteq \mathfrak{m} \cap (FJ + MJ) / \bar{M}J \supseteq \mathfrak{m} \cap (FJ^2 + MJ) / MJ \supseteq \dots,$$

and there exists an integer u such that

$$\mathfrak{m} \cap (FJ^u + MJ) / MJ = \mathfrak{m} \cap (FJ^{u+1} + MJ) / MJ = \dots.$$

The fact that MJ is closed implies

$$\mathfrak{m} \cap (FJ^u + MJ) / MJ = (0).$$

Let $\bar{m} = \sum_i m_i \lambda_i' + j_u$, $j_u \in FJ^u$. Then we have

$$j_u = \bar{m} - \sum_i m_i \lambda_i' \in FJ^u \cap (\bar{m}A + M).$$

Therefore $j_u \in MJ$, i.e. $\bar{m} \in M + \bar{M}J$. Thus we have $\bar{M} \subseteq M + MJ$. The converse inclusion is obvious and we complete the proof of Lemma 4.

Proof of Lemma 3. Since M is a finitely generated module and the two-sided ideal J is finitely generated as a right ideal, MJ^s is finitely generated for any positive integer s . Thus we have

$$MJ^i = MJ^i + MJ^{i+1}.$$

Now we are in a position to prove $\bar{M} = M$. Let \bar{m} be any element of \bar{M} . Then we have, by virtue of the above relation of submodules,

$$\begin{aligned} \bar{m} &= \sum m_i \lambda_i^{(0)} + \bar{m}', \quad \lambda_i^{(0)} \in A, \quad \bar{m}' \in MJ \subseteq FJ, \\ \bar{m}' &= \sum m_i \lambda_i^{(1)} + \bar{m}'', \quad \lambda_i^{(1)} \in J, \quad \bar{m}'' \in \bar{M}J^2 \subseteq FJ^2, \\ \bar{m}'' &= \dots \end{aligned}$$

Let $\bar{\lambda}_i = \sum_{j=0}^{\infty} \lambda_i^{(j)}$. Then we have $\bar{m} = \sum_i m_i \bar{\lambda}_i \in M$. This completes the proof of Lemma 3.

As an immediate consequence of Lemma 1 and Lemma 3, we have

THEOREM 1. *Let A be a right semi-local ring and J its Jacobson radical. Then any finitely generated right ideal of the completion \hat{A} of A is closed, and therefore there holds the relation $I\hat{A} \cap A = \bar{I}$ where \bar{I} is the closure in A of a right ideal I of A .*

THEOREM 2. *Let A be a complete right semi-local ring. Then any right ideal of A is closed. Further, for any finitely generated right A -module M , we have*

$\bigcap_{s=1}^{\infty} MJ^s = 0$ where $J = J(A)$.

Proof. There exists an exact sequence of right A -modules

$$0 \rightarrow N \rightarrow F \rightarrow M \rightarrow 0$$

where F is a finitely generated free A -module. Since A is right noetherian, N is finitely generated. Thus we deduce $\bigcap_{s=1}^{\infty} (N + FJ^s) = N$, i.e. $\bigcap_{s=1}^{\infty} (F/N)J^s = 0$. This is the required result since $F/N \cong M$.

By combining this theorem with [Remark 2 of § 4 in 9] and [Theorem 3.4 of 2], we have

COROLLARY. *A complete right semi-local ring A is linearly compact as a right A -module in the discrete topology.*

Finally we have the following result:

THEOREM 3. *Let A be a right noetherian ring with a unit element 1, and Q a two-sided ideal of A which is a nucleus. If any right ideal I of A is closed with respect to the Q -adic topology, then we have $\bigcap_{s=1}^{\infty} MQ^s = (0)$ for any finitely generated right A -module M .*

Proof. We assume that $\bigcap_{s=1}^{\infty} MQ^s \neq 0$ and deduce a contradiction. We consider the set \mathfrak{S} of all submodules S such that $\bigcap_{i=1}^{\infty} (M/S)Q^i \neq 0$. Let N' be a maximal element of \mathfrak{S} . Then by assumption we have $N = \bigcap_{s=1}^{\infty} (N' + MQ^s) \neq N'$. Let M' be any submodule of M properly containing N' . Then $M' \supseteq N$ by the maximality of N' . Let \bar{m} be any element of M/N' . Then we have $\bar{m}A \cong A/0(\bar{m})$ where $0(\bar{m}) = \{\lambda \in A \mid \bar{m}\lambda = 0\}$. Since any submodule of $\bar{m}A$ contains the unique minimal submodule N/N' of M/N' (this implies that M/N' is sub-directly irreducible), $A/0(\bar{m})$ is subdirectly irreducible. Therefore there exists a positive integer t such that $Q^t \subseteq 0(\bar{m})$ since $0(\bar{m})$ is closed by assumption. This implies that $\bar{m}Q^t = 0$. Therefore there exists an integer s such that $(M/N')Q^s = 0$ since M/N' is finitely generated. This contradicts our assumption $N \neq N'$.

§ 2. Flatness of modules

Let A be a ring with a unit element 1 and M a (unitary) left A -module. Then the module M is said to be A -flat if $\text{Tor}_n^A(C, M) = 0$ for all right A -modules C and all $n > 0$.

Let λ be an element of A , I a right ideal of A and M a left A -module. Then we use the following notations:

- Z for the ring of all integers,
- IM for the Z -submodule of M generated by the set IM ,
- $(I : \lambda)$ for $\{\mu \in A \mid \lambda\mu \in I\}$ and
- $(IM : \lambda)_M$ for $\{m \in M \mid \lambda m \in IM\}$.

THEOREM 4. *Let A be a ring with a unit element 1. Then, for each left A -module M , the following conditions are equivalent to each other:*

- (a) M is A -flat,
- (b) $\text{Tor}_1^A(A/I, M) = 0$ for each right ideal I of A ,
- (c) i) For any right ideals I_1 and I_2 of A , there holds the relation $(I_1 \cap I_2)M = I_1M \cap I_2M$, and
 ii) For each element λ of A , there holds the relation $(0 : \lambda)M = (0 : \lambda)_M$.¹⁾
- (d) $(I : \lambda)M = (IM : \lambda)_M$ for each right ideal I and each element λ of A .

The equivalence of the conditions (a) and (b) is an exercise of 4 (see p. 123 of [4]) and the implication (d) \Rightarrow (a) is proved by the same way as in [7], so we prove only the implications (a) \Rightarrow (c) \Rightarrow (d).

To deduce i) of (c) from (a), it suffices to prove

LEMMA 5. *Let I_1 and I_2 be right ideals of A and M a left A -module. If $\text{Tor}_1^A(A/I_1 + I_2, M) = 0$ we have the relation $(I_1 \cap I_2)M = I_1M \cap I_2M$.²⁾*

This lemma follows immediately from the exact sequence:

$$\begin{array}{ccccccc}
 (I_1 \cap I_2) \otimes M & \longrightarrow & I_1 \otimes M & \longrightarrow & I_1/I_1 \cap I_2 \otimes M & \longrightarrow & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \\
 I_2 \otimes M & \longrightarrow & A \otimes M & \longrightarrow & A/I_2 \otimes M & \longrightarrow & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \\
 0 \rightarrow I_2/I_1 \cap I_2 \otimes M & \rightarrow & A/I_1 \otimes M & \rightarrow & A/I_1 + I_2 \otimes M & \rightarrow & 0
 \end{array}$$

where \otimes means \otimes_A .

Proof of the implication (a) \Rightarrow ii) of (c). From the flatness of the module M , we deduce a commutative exact diagram;

¹⁾ A. Hattori called this property of a module torsion-free in [6].

²⁾ The proof of this lemma is a formal generalization of those of [Theorems 5 and 6 in 1, p. 111].

$$\begin{array}{ccccccc}
 0 & \rightarrow & (0 : \lambda) \otimes M & \rightarrow & A \otimes M & \rightarrow & A / (0 : \lambda) \otimes M \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & (0 : \lambda)_M & \rightarrow & M & \longrightarrow & \lambda M \rightarrow 0
 \end{array}$$

where \otimes means \otimes_Δ . From this diagram we have the required result $(0 : \lambda)M = (0 : \lambda)_M$ by virtue of the well known “five lemma”.

Proof of the implication (c) \Rightarrow (d). It suffices to prove $(I : \lambda)M \supseteq (IM : \lambda)_M$. Let m be any element of $(IM : \lambda)_M$. Then we have $\lambda m \in \lambda(IM : \lambda)_M = IM \cap \lambda M = (I \cap \lambda A)M = \lambda(I : \lambda)M$. Therefore there exists an element $m' \in (I : \lambda)M$ such that

$$\lambda m = \lambda m', \quad \text{i.e.} \quad \lambda(m - m') = 0.$$

This implies that $m - m' \in (0 : \lambda)_M = (0 : \lambda)M \subseteq (I : \lambda)M$. Thus we have $m \in (I : \lambda)M$, and this completes the proof of Theorem 4.

Remark. As an immediate consequence of Theorem 4, we have the following corollary by combining [Theorem 2 or Corollary of 5]:

A commutative integral domain A is a Prüfer ring if and only if there holds the relation

$$(I_1 \cap I_2)I = I_1 I \cap I_2 I,$$

for any ideals I_1, I_2 , and I of A .³⁾

REFERENCES

[1] Y. Akizuki and M. Nagata, Modern algebra (in Japanese), Ser. of Modern Math., Kyoritsu-Shuppan Co., Tokyo (1957).
 [2] E. H. Batho, Non-commutative semi-local and local rings, Duke Math. J., **24** (1957), pp. 163-172.
 [3] I. S. Cohen, On the structure and ideal theory of complete local rings, Trans. of the Amer. Math. Soc., **59** (1946), pp. 54-106.
 [4] H. Cartan and S. Eilenberg, Homological algebra, Princeton Univ., (1956).
 [5] A. Hattori, On Prüfer rings, Jour. Math. Soc. Japan, **9** (1957), pp. 381-385.
 [6] A. Hattori, A foundation of torsion theory for modules over general rings, Nagoya Math. J., this issue.
 [7] C. Lech, Note on multiplicities of ideals, Arkiv för Math., B. 4, H. 1 (1960), pp. 63-86.
 [8] N. Jacobson, Structure of rings, Amer. Math. Soc. Colloq. Pub. XXXVII (1956).
 [9] D. Zelinsky, Linearly compact modules and rings, Amer. J. of Math., **76** (1953), pp. 79-90.

Tokyo Metropolitan University

³⁾ This result was suggested to the writer by T. Ishikawa.