
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Winches Farm Field Station,
395 Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Translation products of mRNA from infective larvae of <i>Trichinella spiralis</i> include an antigenic polypeptide. K. SUGANE, T. MATSUURA and H. MAEKAWA	1-8
The involvement of eggs and soluble egg antigens in resistance to <i>Schistosoma japonicum</i> . N. A. MOLONEY, A. J. F. LUTY, P. HINCHCLIFFE and G. WEBBE	9-17
<i>Brugia pahangi</i> in nude mice: protective immunity to infective larvae is Thy 1.2 ⁺ cell dependent and cyclosporin A resistant. A. C. VICKERY and J. K. NAYAR	19-27
<i>Heterophyes</i> species in Greece: record of <i>H. heterophyes</i> , <i>H. aequalis</i> and <i>H. dispar</i> from the first intermediate host, <i>Pirenella conica</i> . H. TARASCHEWSKI and A. NICOLAIDOU	28-32
Experiments on habitat selection of <i>Heterophyes</i> species in different definitive hosts. H. TARASCHEWSKI	33-42
Chromosomes of <i>Onchocerca volvulus</i> (Spirurida: Onchocercidae): a comparative study between Nigeria and Guatemala. H. HIRAI, I. TADA, H. TAKAHASHI, B. E. B. NWOKE and G. O. UFOMADU	43-46
A comparison of two procedures for labelling the surface of the hydatid disease organism, <i>Echinococcus granulosus</i> , with ¹²⁵ I. D. P. MCMANUS, D. J. MCLAREN, N. W. T. CLARK and R. M. E. PARKHOUSE	47-52
An evaluation of the efficacy of monensin against <i>Fasciola hepatica</i> in the albino rat. P. J. SKUCE, I. FAIRWEATHER and J. W. M. KENNY	53-58
Histochemical studies on eggshell formation in <i>Paramphistomum cervi</i> (Digenea: Paramphistomatidae). B. C. GUPTA, V. R. PARSHAD and S. S. GURAYA	59-64
Field experiments on the ability of <i>Arthrotrys oligospora</i> (Hyphomycetales) to reduce the number of larvae of <i>Cooperia oncophora</i> (Trichostrongylidae) in cow pats and surrounding grass. J. GRØNVOLD, J. WOLSTRUP, S. A. HENRIKSEN and P. NANSEN	65-71
<i>Protostrongylus rufescens</i> : a cytogenetic study. C. CUTILLAS, D. C. GUEVARA, A. VALERO and C. ARIZA	72-76
Helminth parasites of Jamaican anoles (Reptilia: Iguanidae): a comparison of the helminth fauna of 6 <i>Anolis</i> species. D. A. P. BUNDY, P. VOGEL and E. A. HARRIS	77-83
Amino acid metabolism in <i>Ancylostoma ceylanicum</i> and <i>Nippostrongylus brasiliensis</i> . G. SINGH, S. GUPTA, J. C. KATIYAR and V. M. L. SRIVASTAVA	84-88
Genetic variability of <i>Schistosoma bovis</i> cercarial production according to miracidial dose. A. MOUAHID and C. COMBES	89-94
Book review	18
Announcement (Filariasis: Ciba Foundation Symposium 127)	46