

Original Article

Cite this article: Yuksel C *et al* (2024). REM disruption and REM vagal activity predict extinction recall in trauma-exposed individuals. *Psychological Medicine* **54**, 4623–4634. <https://doi.org/10.1017/S0033291724002757>

Received: 19 April 2024
Revised: 8 September 2024
Accepted: 7 October 2024
First published online: 9 December 2024

Keywords:

PTSD; fear extinction; memory consolidation; vagal activity; heart rate variability; sleep; REM

Corresponding author:

Cagri Yuksel;
Email: ayuksel@mgb.org

REM disruption and REM vagal activity predict extinction recall in trauma-exposed individuals

Cagri Yuksel^{1,2} , Lauren Watford¹, Monami Muranaka¹, Carolina Daffre^{3,4}, Emma McCoy¹, Hannah Lax^{1,4}, Augustus Kram Mendelsohn^{3,4}, Katelyn I. Oliver^{3,4}, Alexis Acosta^{3,4}, Abigail Vidrin^{3,4}, Uriel Martinez^{3,4}, Natasha Lasko^{2,5}, Scott Orr^{2,5} and Edward F. Pace-Schott^{2,3,4}

¹Schizophrenia and Bipolar Disorder Program, McLean Hospital, Belmont, MA, USA; ²Department of Psychiatry, Harvard Medical School, Boston, MA, USA; ³Athinoula A. Martinos Center for Biomedical Imaging, Charlestown, MA, USA; ⁴Department of Psychiatry, Massachusetts General Hospital, Charlestown, MA, USA and ⁵Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA

Abstract

Background. Accumulating evidence suggests that rapid eye movement sleep (REM) supports the consolidation of extinction memory. REM is disrupted in posttraumatic stress disorder (PTSD), and REM abnormalities after traumatic events increase the risk of developing PTSD. Therefore, it was hypothesized that abnormal REM in trauma-exposed individuals may pave the way for PTSD by interfering with the processing of extinction memory. In addition, PTSD patients display reduced vagal activity. Vagal activity contributes to the strengthening of memories, including fear extinction memory, and recent studies show that the role of vagus in memory processing extends to memory consolidation during sleep. Therefore, it is plausible that reduced vagal activity during sleep in trauma-exposed individuals may be an additional mechanism that impairs extinction memory consolidation. However, to date, the contribution of sleep vagal activity to the consolidation of extinction memory or any emotional memory has not been investigated.

Methods. Trauma-exposed individuals ($n = 113$) underwent a 2-day fear conditioning and extinction protocol. Conditioning and extinction learning phases were followed by extinction recall 24 h later. The association of extinction recall with REM characteristics and REM vagal activity (indexed as heart rate variability) during the intervening consolidation night was examined.

Results. Consistent with our hypotheses, REM disruption was associated with poorer physiological and explicit extinction memory. Furthermore, higher vagal activity during REM was associated with better explicit extinction memory, and physiological extinction memory in males.

Conclusions. These findings support the notion that abnormal REM, including reduced REM vagal activity, may contribute to PTSD by impairing the consolidation of extinction memory.

Introduction

Posttraumatic stress disorder (PTSD) is characterized by excessive fear responses to cues associated with a traumatic event. It is hypothesized that this is due to dysfunction in fear extinction, the process whereby fear response is gradually reduced after multiple exposures to a conditioned stimulus (CS) without reinforcement (Milad & Quirk, 2012; Zuj, Palmer, Lommen, & Felmingham, 2016). After a traumatic event, fear extinction is achieved by repeated exposure to reminders of the traumatic event without the feared outcome. This mechanism is suggested to be the basis of exposure therapy for PTSD and other anxiety-related disorders (Craske, Treanor, Conway, Zbozinek, & Vervliet, 2014).

Fear extinction is not simply an erasure of fear but relies on new learning and memory. While the preponderance of studies does not show impaired extinction learning in PTSD (reviewed in Bottary, Straus, & Pace-Schott, 2023; Pace-Schott, Seo, & Bottary, 2023), accumulating evidence suggests that impaired retention of extinction memory, acquired after trauma exposure (Milad *et al.*, 2008), may be a critical mechanism that leads to PTSD (Helpman *et al.*, 2016; Milad *et al.*, 2008, 2009; Milad & Quirk, 2012; Shvil *et al.*, 2014; Suarez-Jimenez *et al.*, 2020; Wicking *et al.*, 2016). Sleep supports the consolidation of different types of memories, which stabilizes and integrates them, and enhances their retrieval (Klinzing, Niethard, & Born, 2019). Recent evidence suggests that this extends to extinction memory, with rapid eye movement sleep (REM) implicated as an important sleep stage involved in its consolidation (Davidson & Pace-Schott, 2020). In healthy people, more REM has been shown to be associated with better extinction recall (Bottary *et al.*, 2020; Menz, Rihm, & Buchel, 2016; Pace-Schott *et al.*, 2014; Spoormaker *et al.*, 2010), and REM deprivation leads to impaired extinction memory (Spoormaker *et al.*, 2012), suggesting the possibility of a causal role for

this sleep stage. Several studies and meta-analyses find REM abnormalities in PTSD (Kobayashi, Boarts, & Delahanty, 2007; Richards, Kanady, & Neylan, 2020; Zhang et al., 2019), which also have been shown to be associated with the risk of developing PTSD after trauma (Mellman, Bustamante, Fins, Pigeon, & Nolan, 2002; Mellman, Knorr, Pigeon, Leiter, & Akay, 2004; Mellman, Pigeon, Nowell, & Nolan, 2007). Therefore, it is postulated that sleep disruption, specifically of REM, may predispose traumatized individuals to develop PTSD symptoms by interfering with extinction memory (Colvonen, Straus, Acheson, & Gehrman, 2019; Pace-Schott et al., 2023). Supporting this hypothesis, in a sample overlapping that of the current report, we showed that prefrontal activations during extinction recall were associated with REM density during the night before extinction recall (Seo et al., 2022). However, much remains to be learned regarding the association between REM and extinction memory in trauma-exposed individuals. One preliminary study (Straus, Norman, Risbrough, Acheson, & Drummond, 2018) did not find any association of REM with extinction recall in PTSD; however, the small sample size ($n = 13$) may explain this negative finding. This study's sample was also limited to those with PTSD diagnoses and, therefore, did not provide insight into this association in individuals with subthreshold PTSD symptoms. To address this gap in knowledge, we examine here the association of REM measures on the night following extinction learning with extinction recall the following day, in a large sample of trauma-exposed individuals. We hypothesized that disruptions in REM sleep would be associated with impaired extinction recall (Hypothesis 1).

PTSD is characterized by abnormal autonomic nervous system function, including reduced parasympathetic nervous system (PNS) activity (Seligowski, Webber, Marvar, Ressler, & Philip, 2022). PNS activity can be estimated by measuring heart rate variability (HRV), the variation in the interval between successive heartbeats. Specific HRV measures in the frequency (high frequency; HF-HRV) and time (root mean square of successive differences in the R-R interval; RMSSD) domains reflect control of heart rate by the vagus nerve, the main component of PNS outflow from the CNS (Laborde, Mosley, & Thayer, 2017; Shaffer & Ginsberg, 2017). Meta-analyses show that patients with PTSD display reduced vagal activity, as indexed by HF-HRV and RMSSD, during wakefulness (Chalmers, Quintana, Abbott, & Kemp, 2014; Ge, Yuan, Li, & Zhang, 2020; Schneider & Schwerdtfeger, 2020), and in one study, lower baseline HF-HRV predicted the severity of later post-traumatic stress symptoms (Pyne et al., 2016). Similar to wakefulness, a small number of recent studies in sleep find reduced vagal activity in PTSD (Kobayashi, Lavelle, & Mellman, 2014), including in specific sleep stages (Daffre et al., 2023; Ulmer, Hall, Dennis, Beckham, & Germain, 2018). Additionally, in one study, HF-HRV across actigraphically measured sleep was associated with PTSD symptom severity (Rissling et al., 2016).

Vagally mediated HRV is considered to reflect the efficiency of the medial prefrontal cortex (mPFC) in inhibiting limbic subcortical structures (Thayer, Hansen, Saus-Rose, & Johnsen, 2009), a process also critical for fear extinction (Giustino & Maren, 2015). Furthermore, converging evidence indicates that the vagus nerve contributes to emotional regulation and memory via its afferents to the brainstem nuclei, which in turn modulate the activity of multiple neurotransmitter systems and relevant brain regions within the central autonomic network, including the hippocampus, amygdala, and mPFC (Broncel, Bocian, Klos-Wojtczak, Kulbat-Warycha, & Konopacki, 2020; McGaugh,

2013). Consistent with these, high vagal tone is associated with better extinction learning in humans (Jenness, Miller, Rosen, & McLaughlin, 2019; Pappens et al., 2014; Wendt, Neubert, Koenig, Thayer, & Hamm, 2015), and enhancing vagal activity improves extinction learning and retention of extinction memory in both rodents (Noble et al., 2017, 2019b; Pena et al., 2014; Pena, Engineer, & McIntyre, 2013; Souza et al., 2022) and humans (Burger et al., 2016, 2019; Szeska, Richter, Wendt, Weymar, & Hamm, 2020). In addition, a separate line of recent studies shows that vagal activity during sleep, as indexed by HF-HRV, independently contributes to memory consolidation over and above contributions by sleep architecture and sleep-related neural oscillations (Whitehurst, Cellini, McDevitt, Duggan, & Mednick, 2016; Whitehurst, Subramoniam, Krystal, & Prather, 2022). However, to date, no study has examined the role of vagal activity in the sleep-dependent consolidation of extinction memory in any population. REM is characterized by activation in the aforementioned regions that regulate both vagal activity and fear extinction (Pace-Schott et al., 2023). We thus hypothesized that lower HF-HRV in REM during the night following extinction learning would independently predict lower extinction recall (i.e. consolidation of extinction memory) in trauma-exposed individuals (Hypothesis 2). If this hypothesis is supported, it would further underscore vagal activity as a promising target for enhancing extinction memory through readily accessible interventions (Jung, Yang, & Lee, 2024; Siepmann, Weidner, Petrowski, & Siepmann, 2022) in the aftermath of trauma exposure.

Methods

Participants

A total of 139 participants, aged between 18 and 40, were recruited from the greater Boston metropolitan area using online and posted advertisements. All participants reported experiencing a DSM-5 criterion-A traumatic event ('index trauma') in the past two years, except within the last one month. Participants were allowed with a concurrent anxiety disorder, dysthymia, major depressive disorder (remitted or on a case-by-case basis), or if on a stable dose of an antidepressant (≥ 8 weeks). Exclusion criteria included: history of chronic childhood abuse or neglect; PTSD diagnosis preceding traumatic event indexed at study interview; neurological disorder or injury; major medical disorders; psychotic, bipolar, autism spectrum, or other neurodevelopmental disorders; current drug or alcohol abuse or dependence; history of sleep disorder other than insomnia or nightmare disorder; current use of hypnotic or recently adjusted psychiatric medications; shift work; and any contraindication to MRI scans.

Twenty-six participants were excluded from all analyses because of missing or unusable EEG or ECG data (final $n = 113$). Psychiatric history was ascertained using the Structured Clinical Interview for DSM-IV-TR for Non-Patients (SCID-I/NP) (First, Gibbon, Spitzer, & Williams, 2007). The Clinician-Administered PTSD Scale for DSM-5 (CAPS-5; [Weathers et al., 2018]) was used to assess post-traumatic stress symptom severity (see online Supplementary Fig. S1 for its distribution in the sample). This study followed a Research Domain Criteria (RDoC) (Insel et al., 2010) design in which dimensional rather than categorical measures were targeted, and PTSD diagnoses were established post-hoc from diagnostic evaluations rather than by assigning subjects to PTSD and non-PTSD groups. 49.6% ($n = 56$) of participants met diagnostic criteria for PTSD. The Quick Inventory of Depressive

Symptomatology, Self-Report (QIDS-SR) (Rush et al., 2003), and the Pittsburg Structured Clinical Interview for Sleep Disorders (SCID-SLD) (Stocker, Khan, Henry, & Germain, 2017) were administered to evaluate depressive and sleep disorder symptoms, respectively. Demographic and clinical characteristics of the final sample are displayed in Table 1. All participants provided written consent to participate in the study and were paid for their participation. All procedures were approved by the Partners Healthcare Institutional Review Board.

Procedures

Timeline of the study procedures are included in online Supplementary Methods (Fig. S2). Participants completed an

approximately 2-week sleep assessment period during which they wore an actigraph (Actiwatch-2, Philips Respironics) and filled out a daily sleep and nightmare diary. Approximately midway through the 14-day period, they underwent a combined sleep-disorders-screening and acclimation night of ambulatory polysomnography (PSG) recording. This was followed by a 'baseline' overnight PSG recording. Starting in the evening immediately after the baseline night, participants completed a 2-day fear conditioning and extinction protocol with simultaneous fMRI recordings (Milad et al., 2013). Between fMRI sessions, participants completed a third ('consolidation') night of ambulatory PSG.

Ambulatory PSG

Ambulatory PSG was recorded on three nights using the Somte-PSG ambulatory sleep monitor (Compumedics USA, Charlotte, NC, USA). Sampling rate was 256 Hz. EEG data were acquired using six EEG channels (F3, F4, C3, C4, O1, O2; positioned according to the 10–20 system). Additional electrodes were placed on bilateral mastoids, above the right and below the left eye (EOG), under the chin (EMG), and below the right clavicle and in the left fifth intercostal space (ECG). Participants returned home to sleep after being instrumented. During the acclimation/screening (first) PSG night, additional channels for pulse-oximeter, respiration transducer belts, nasal cannula, and tibialis movement sensors were added to screen for obstructive sleep apnea (OSA) and periodic limb movement disorder (PLMD). No participant met criteria for clinically significant OSA or PLMD. All sleep records were scored by an experienced, registered polysomnographic technologist according to American Academy of Sleep Medicine criteria (Berry, Gamaldo, Harding, & Brooks, 2015).

Fear conditioning, extinction learning, and extinction recall

A validated 2-day paradigm (Milad et al., 2007, 2013) was used to probe fear conditioning and extinction during ongoing fMRI recording. This protocol consisted of four phases, with habituation, fear conditioning, and extinction learning phases taking place on the first day and extinction recall 24 h later. During each phase, images of a colored desk lamp (red, yellow, or blue) appearing in a contextual background (office for conditioning context and conference room for extinction context) served as conditioned stimuli (CS). Context images were presented for nine seconds, with three seconds with the lamp off and six seconds with the lamp on (red, yellow, or blue). The unconditioned stimulus (US) was a mild (0.8–4.0 mA), 500 msec electric shock delivered to the index and middle fingers of participants' right hand using a Coulbourn Transcutaneous Aversive Finger Stimulator (Coulbourn Instruments, Allentown, PA). Prior to entering the scanner, participants were administered increasing intensities of shock and they each selected a level that they perceived as 'highly annoying but not painful'. Prior to entering the scanner, MRI-safe 11-mm, Ag/AgCl skin conductance monitoring electrodes were attached to the participant's left palm. Skin conductance level (SCL) was continuously monitored at 37.5 Hz using the MP150 system with Acqknowledge 4.3 (BIOPAC Systems, Inc., Goleta, CA) software.

During habituation, all six possible combinations of lamp colors and contexts were presented across six trials. During the following fear conditioning phase, two of the three colored lamps (CS+) were each presented eight times paired with the US at stimulus offset, on a partial reinforcement schedule (five out of

Table 1. Demographic and clinical characteristics of the participants

Age	24.0 ± 4.8 (18–39)
Sex (%female)	69.9%
Race (%)	
American Indian or Alaskan Native	2.7%
Asian	9.8%
Black or African-American	16.8%
More than one race	6.2%
Unknown/unreported	1.8%
White	62.9%
Ethnicity	
Hispanic or Latino	9.7%
Not Hispanic or Latino	85%
Unknown/unreported	3.5%
Type of trauma	
Transportation accident	26.5%
Violent assault	19.5%
Rape or sexual assault	17.7%
Mass shooting	2.7%
Sudden loss of family or friend	4.5%
Combat incident	2.7%
Multiple or other	26.4%
PTSS Severity	
CAPS-5	21.8 ± 13.0 (0–53)
Depression Severity	
QIDS	7.4 ± 4.4 (0–18)
Months since trauma	13.0 ± 6.8 (1–28)
Medications	
Antidepressants	18.6%
Benzodiazepines	3.5%
Beta-blockers	0%
Mood stabilizers	0%
Antipsychotics	0%

CAPS-5, Clinician-Administered PTSD Scale for DSM-5; PTSS, Post-traumatic stress symptom; QIDS, Quick Inventory of Depressive Symptomatology. Some of the data is displayed as mean ± standard deviation (minimum–maximum).

eight presentations were paired with US). The third lamp color, which was never paired with US (CS-), was interspersed among the CS + s for a total of 16 presentations. Fear conditioning was followed by extinction learning, during which one CS + (CS + E) was presented in the extinction context 16 times without the US along with 16 interspersed presentations of the CS-. The other CS+ remained conditioned but unextinguished (CS + U). During extinction recall, which took place 24 h later, each CS+ was presented eight times in the extinction context, with no US, along with 16 interspersed CS-.

During each phase, physiologic reactivity for each trial was indexed using skin conductance response (SCR), a measure of sympathetic activity (Dawson, Schell, & Filion, 2007), calculated as the mean SCL in microSiemens (μ S) during the last 2 s of context presentation subtracted from the maximum SCL during the 6 s of CS presentation. Negative SCRs were coded as zero (Lonsdorf et al., 2017), and then all values were square root transformed. 'non-conditioners' were defined as those who exhibited less than two non-square-root transformed SCR responses to either of the two CS + s that were equal to or exceeding 0.05 μ S during the Fear Conditioning phase (Orr et al., 2000). Thirty-one non-conditioners were further excluded from SCR analyses based on these criteria. In addition, SCR data for three participants were lost.

Immediately following each phase except habituation, participants verbally reported shock expectancy for the first and last presentations of each CS (i.e. colored light) appearing in that phase on a scale from 1 ('not expecting a shock at all') to 5 ('expecting a shock very much'). Ratings were obtained at this frequency with the goal of capturing the shift in expectancy from baseline to the end of the corresponding session. Expectancy ratings for seven participants were not included in the analyses, because they did not complete the experiment or because of missing data.

Extinction recall variables

To examine the association of physiologically expressed and subjective extinction recall with sleep and HRV measures, we used an extinction retention index (ERI) (Lonsdorf, Merz, & Fullana, 2019) and a subjective extinction retention index (sERI) (Bottary et al., 2020), respectively. ERI was calculated as: $([\text{Average of the SCRs of the first 4 CS + E presentations at extinction recall phase}/\text{maximum SCR to the 'to-be' CS + E during the fear conditioning phase}] \times 100)$, which is one of the most commonly used formulae in the literature (Lonsdorf et al., 2019). Only the first four CS + E trials from the extinction recall phase were included in this calculation in order to avoid confounding recalled extinction with new extinction learning. Higher ERI reflected lower extinction memory. sERI was calculated as: $\text{Expectancy to the first CS + E in Extinction Recall}/\text{mean expectancy for the last of each CS + during Fear conditioning} \times 100$. Larger sERI indicated lower extinction memory. Both ERI and sERI were formulated to quantify the degree of dominance for the putatively competing memories of extinction and fear (Lonsdorf et al., 2019).

Sleep measures

The PSG data analyzed in the present study were recorded during the consolidation night (the night between the extinction learning and extinction recall phases). For regression analyses, we selected the REM measures that have been associated with PTSD, including %REM (Zhang et al., 2019), REM density (REMD)

(Kobayashi et al., 2007), REM latency (REML) (Mellman, Kobayashi, Lavelle, Wilson, & Hall Brown, 2014), and REM fragmentation (REMF) (Breslau et al., 2004; Habukawa et al., 2018; Insana, Kolko, & Germain, 2012; Lipinska & Thomas, 2019; Mellman et al., 2002; Saguin, Gomez-Merino, Sauvet, Leger, & Chennaoui, 2021). Percent time spent in each sleep stage (%N1–N3, %REM) was computed as a percentage of total sleep time (TST). REMD is the number of rapid eye movements per minute of REM sleep and was calculated using an automatic algorithm (Yetton et al., 2016). REML was calculated as the number of minutes occurring after sleep onset before the first REM epoch. REMF was calculated as the average duration of REM segments (Mellman et al., 2002). REM segments were defined as continuous REM from the start of at least 1 min of REM to the onset of at least 1 min of non-REM or wake (Mellman et al., 2002).

Heart rate variability

HRV was calculated using Kubios HRV Premium software (Kubios Oy, Kuopio, Finland) in continuous REM ECG segments lasting at least 5 min during the entire night's sleep. Prior to analysis, ECG traces were visually inspected for physiological and technical artifacts (Laborde et al., 2017). Misplaced or ectopic beats were manually corrected if possible. Very low to low, or occasionally medium, filter was applied, if necessary, after correction (<https://www.kubios.com/hrv-preprocessing/>). Segments that did not provide reliable estimates due to excessive artifacts were removed. Twenty-three participants were excluded because of artifacts, not having ≥ 5 min REM segments, or because their records were missing. High frequency (HF; 0.15–0.4 Hz) absolute power ($\text{HF}[\text{ms}^2]$) was used as a predictor variable in the primary analyses because it was shown to be associated with sleep-dependent memory processing (Whitehurst et al., 2016). We also carried out the same analyses with root mean square of the successive differences in the R–R interval (RMSSD), another measure of vagal activity (Shaffer & Ginsberg, 2017). For frequency domain analyses, an autoregressive method was used, with model order set at 16 (Laborde et al., 2017). Data were transformed by their natural logarithm (Laborde et al., 2017).

Statistical analyses

Associations were explored between extinction recall indices (ERI and sERI), and age, sex, months since index trauma, scores of symptom scales (CAPS-5 and QIDS), and medication use as a binary variable. For correlations, Pearson and Spearman's correlations were used depending on the distribution of variables. Student's *t* tests were used for comparisons between sexes and between medication users and non-users.

We used linear regression models to test whether REM sleep variables predicted extinction recall (ERI and sERI; Hypothesis 1). The models included %REM, REMD, REML, and REMF as predictor variables. Sex and medication use were also included as predictors as these factors have been shown to modulate fear extinction and its retention (Bouton, Kenney, & Rosengard, 1990; Gunduz-Cinar et al., 2016; Milad et al., 2006).

Next, hierarchical regression analyses were used to test the hypothesis that vagal activity would improve the predictions of ERI and sERI above and beyond REM sleep measures alone (Hypothesis 2). For this purpose, two models were built. Model 1 included sex, medication use, and REM measures. $\text{HF}[\text{ms}^2]$ was added as an additional predictor in Model 2. Visual inspection of the data suggested that sex may be moderating the effect

of HF[ms²] on ERI. Therefore, sex $\times$ HF[ms²] interaction was also added to the model for ERI. Because missing HF[ms²] data further limited the number of participants that could be included in the analysis, only REM variables that were significant predictors in the regression analysis that tested Hypothesis 1 were included. Results for the models that included all REM variables and RMSSD instead of HF[ms²] were also reported for completeness.

In all regression analyses, partial regression plots and a plot of studentized residuals against the predicted values indicated that assumptions of linear relationship were met. Variance inflation factors were within acceptable ranges (1.1–1.4), indicating that there was no confounding multicollinearity. Little's Missing Completely At Random (MCAR) test indicated that data was missing at random in our sample ($\chi^2 = 84.253$, $DF = 71$, $Sig. = 0.135$). For ERI, the distribution histogram and P–P plot of standardized residuals showed a non-normal distribution. In addition, plotting standardized residuals and predictors revealed heteroscedasticity. Therefore, ERI was log-transformed. After transformation, normal distributions of residuals and homoscedasticity were confirmed.

For all variables, outliers ± 3 standard deviations from the mean were removed. All analyses were two-tailed. The significance level for hypothesis testing (α) was set at 0.05.

Results

Association of extinction recall indexes with demographic and clinical measures

Age was inversely correlated with sERI ($r_s = -0.27$, $p = 0.006$), while it was not significantly correlated with ERI ($r_s = 0.12$, $p = 0.305$). There were no differences between sexes in ERI or sERI ($t(76) = 0.233$, $p = 0.817$ and $t(104) = 0.116$, $p = 0.250$, respectively). There were no significant correlations between either extinction recall index and the number of months since index trauma (ERI: $r_s = -0.03$, $p = 0.796$; sERI: $r_s = -0.04$, $p = 0.659$), or CAPS-5 (ERI: $r_s = 0.11$, $p = 0.333$; sERI: $r_s = -0.02$, $p = 0.831$) or QIDS (ERI: $r_s = 0.13$, $p = 0.350$; sERI: $r_s = -0.04$, $p = 0.692$) scores.

Regression analyses predicting ERI and sERI

Measures of REM characteristics and sleep architecture are displayed in Table 2. The multiple regression model with ERI as the dependent variable included sex, medication use, %REM, REMD, REML, and REMF as predictor variables ($N = 67$). The overall model explained a significant proportion of the variance (Adj. $R^2 = 0.14$, $p = 0.021$), with sex, %REM, REMD, and REML emerging as significant predictors (Table 3; online Supplementary Fig. S3). Poorer physiological extinction recall was associated with the male sex ($B[s.e.] = -0.71[0.34]$, $p = 0.041$), less time in REM sleep (%REM; $B[s.e.] = -0.06[0.03]$, $p = 0.018$), increased REMD ($B[s.e.] = 0.14[0.05]$, $p = 0.009$), and shorter REML ($B[s.e.] = -0.01[0.003]$, $p = 0.002$).

A similar multiple regression model for sERI was carried out ($N = 94$). In our sample, age was associated with sERI (see above), therefore it was also included in the model. This model also explained a significant proportion of the variance (Adj. $R^2 = 0.09$, $p = 0.031$). In this model, REMD was the only significant predictor (Table 4; online Supplementary Fig. S4), and as with ERI, increased REMD was associated with poorer subjective extinction recall ($B[s.e.] = 2.80[0.94]$, $p = 0.004$).

Table 2. Sleep and HRV measures

REMD	5.1 $\pm$ 3.1 (0.62–13.0)
REML	102.3 $\pm$ 45.9 (0–238.0)
REMF	9.8 $\pm$ 4.0 (2.8–23.4)
%N1	5.7 $\pm$ 3.8 (0.8–27.0)
%N2	54.5 $\pm$ 9.3 (26–81.7)
%N3	21.7 $\pm$ 10.6 (1.7–63.3)
%REM	18.3 $\pm$ 6.5 (2.0–35.0)
HF[ms ²]	1198.0 $\pm$ 1069.5 (39.2–4788.6)
RMSSD	54.3 $\pm$ 26.8 (13.0–128.3)

HF [ms²], absolute power of high frequency heart rate variability; %N1-REM, proportion of the sleep stage to the total sleep time; REMD, REM density; REMF, REM fragmentation; REML, REM latency; RMSSD, root mean square of the successive differences in the R–R interval. Data is displayed as mean $\pm$ standard deviation (minimum–maximum).

Hierarchical regressions with HF HRV predicting ERI and sERI

Average HRV values are displayed in Table 2. For the hierarchical regression analyses with ERI as the dependent variable ($N = 61$), sex, medication use, and REM sleep measures (%REM, REMD, and REML) were included in the first model. HF [ms²] and sex $\times$ HF [ms²] interaction were included as additional predictors in the second model. The final model was significant (Adj. $R^2 = 0.17$, $p = 0.018$; Table 5). There was a trend for an increase in the variance accounted for by the second model ($\Delta R^2 = 0.08$, $p = 0.074$). HF [ms²] and the sex $\times$ HF [ms²] interaction were significant predictors ($B[s.e.] = -0.42[0.20]$, $p = 0.042$ and $B[s.e.] = 0.65[0.29]$, $p = 0.030$), respectively. We therefore probed this interaction by examining the simple slopes, which showed that the effect of HF[ms²] on ERI was significant for males ($B[s.e.] = -0.42[0.20]$, $p = 0.042$) but not for females ($B[s.e.] = 0.23[0.21]$, $p = 0.263$), and the effects were different between sexes ($p = 0.031$).

A similar hierarchical regression analysis was carried out for sERI ($N = 85$). The first model included age, sex, medication use, and REMD. HF[ms²] was added to the predictors in the second model. The addition of HF [ms²] led to a significant increase in the variance accounted for by the model ($\Delta R^2 = 0.06$, $p = 0.018$), indicating that vagal activity contributed to the prediction of subjective extinction recall above and beyond REM sleep measures. The final model was significant (Adj. $R^2 = 0.13$, $p = 0.006$), and medication use, REMD, and HF[ms²] emerged as significant predictors. Poorer subjective extinction recall was associated with not using medications ($B[s.e.] = -18.81[8.19]$, $p = 0.024$), higher REMD ($B[s.e.] = 2.64[0.95]$, $p = 0.007$), and lower HF [ms²] ($B[s.e.] = -7.05[2.92]$, $p = 0.018$) (Table 6, online Supplementary Fig. S4).

For completeness, we repeated the same hierarchical regression analyses by adding all REM variables. The results were similar (online Supplementary Tables S1 and S2). Using RMSSD instead of HF [ms²] also did not change the results (online Supplementary Tables S3 and S4).

Results of regression analyses after the removal of non-contributory predictors are included in the Supplementary Methods (online Supplementary Tables S5–S8).

Discussion

We investigated the association of REM measures with the consolidation of physiological and subjective extinction memory in

Table 3. Linear regression analysis for physiological extinction recall (ERI; $N = 67$)

Predictors	B	S.E.	β	t	p	95% CI	Adj. R^2	ANOVA	
								F	p
Sex	−0.714	0.341	−0.279	−2.092	0.041	−1.397	−0.031	0.135	2.716
Medications	0.113	0.372	0.038	0.304	0.762	−0.630	0.856		
%REM	−0.061	0.025	−0.326	−2.427	0.018	−0.111	−0.011		
REMD	0.141	0.052	0.365	2.701	0.009	0.036	0.245		
REML	−0.011	0.003	−0.421	−3.167	0.002	−0.018	−0.004		
REMF	0.036	0.031	0.153	1.155	0.253	−0.026	0.098		

%REM, proportion of REM sleep to the total sleep time; REMD, REM density; REMF, REM fragmentation; REML, REM latency. Sex, %REM, REMD and REML were significant predictors. Note that smaller ERI denotes better extinction recall.

a large sample of trauma-exposed individuals. Confirming our first hypothesis, we found that less time spent in REM sleep, shorter REML, and higher REMD independently predicted poorer physiological extinction recall. Similarly, poorer subjective extinction recall was predicted by higher REMD. Furthermore, our second hypothesis, that higher HF HRV during REM would predict greater extinction recall above and beyond REM measures, was supported for subjective extinction recall and partially supported for physiological extinction recall. To the best of our knowledge, this is the first study to show that features of REM are associated with the consolidation of extinction memory in trauma-exposed individuals. In addition, we show here for the first time that vagal activity during a specific sleep stage contributes to the consolidation of an emotional memory.

REM measures predicting extinction recall is in agreement with previous research in healthy individuals (Davidson & Pace-Schott, 2020; Pace-Schott, Germain, & Milad, 2015a) and in insomnia disorder (Bottary et al., 2020). In healthy individuals, better extinction recall was associated with the presence of REM during a nap (Spoormaker et al., 2010), higher overnight % REM (Pace-Schott et al., 2014), less fragmented REM, and increased REM theta power (Bottary et al., 2020). In addition, late-night (REM-rich) sleep but not early-night, NREM stage 3-rich sleep benefited extinction memory (Menz et al., 2016), and selective REM, but not NREM, deprivation impaired extinction recall (Spoormaker et al., 2012). In contrast, among individuals with insomnia disorder (Bottary et al., 2020), better

extinction recall was associated with less %REM, shorter REM bouts, and longer REML. Findings are also in agreement with the general notion that REM is involved in processing emotional memories (Goldstein & Walker, 2014), although this assertion is not consistently supported by experimental evidence (Davidson, Jonsson, Carlsson, & Pace-Schott, 2021).

We further showed that specific REM features were independently associated with extinction recall. There is a sizable literature that implicates both REML and REMD in a variety of psychiatric disorders, including PTSD (Baglioni et al., 2016). However, in PTSD, alterations in these variables are not consistently found, and the direction of change showed variability across studies, possibly due to the heterogeneity in samples and the settings in which data were collected (Baglioni et al., 2016; Kobayashi et al., 2007; Zhang et al., 2019). It was also proposed that two competing processes may be at play simultaneously, REM dysregulation and the resulting pressure to achieve REM (Mellman, 1997), which would explain the contradictory findings across studies (e.g. shorter or longer REML, depending on which of these processes predominate). Nonetheless, shortened REML and increased REMD have emerged as characteristics of PTSD in meta-analyses (Baglioni et al., 2016; Kobayashi et al., 2007; Zhang et al., 2019). Our findings advance insight into the significance of these REM alterations and indicate that they are associated with impaired extinction memory, a mechanism that is considered central to the development of PTSD (Bottary et al., 2023).

Table 4. Linear regression analysis for subjective extinction recall (sERI; $N = 94$)

Predictors	B	S.E.	β	t	p	95% CI	Adj. R^2	ANOVA	
								F	p
Age	−0.512	0.629	−0.084	−0.814	0.418	−1.763	0.739	0.092	2.345
Sex	−2.903	6.842	−0.046	−0.424	0.672	−16.503	10.698		
Medications	−12.574	7.632	−0.175	−1.648	0.103	−27.747	2.598		
%REM	0.238	0.527	0.052	0.452	0.653	−0.810	1.286		
REMD	2.801	0.936	0.331	2.994	0.004	0.942	4.661		
REML	0.087	0.072	0.136	1.209	0.230	−0.056	0.229		
REMF	0.528	0.653	0.090	0.808	0.421	−0.771	1.827		

%REM, Proportion of REM sleep to the total sleep time; REMD, REM density; REMF, REM fragmentation; REML, REM latency. REMD was the only significant predictor. Note that smaller sERI denotes better extinction recall.

Table 5. Hierarchical regression analysis for physiological extinction recall (ERI; $N = 61$)

Model	Predictors	<i>B</i>	S.E.	β	<i>t</i>	<i>p</i>	95% CI						
1	Sex	−0.531	0.345	−0.213	−1.539	0.130	−1.221	0.160					
	Medications	0.017	0.386	0.006	0.045	0.964	−0.757	0.791					
	%REM	−0.043	0.025	−0.234	−1.763	0.083	−0.093	0.006					
	REMD	0.143	0.053	0.379	2.718	0.009	0.037	0.248					
	REML	−0.008	0.003	−0.323	−2.481	0.016	−0.015	−0.002					
									Change		ANOVA		
									ΔF	<i>p</i>	Adj. R^2	<i>F</i>	<i>p</i>
2	Sex	−0.591	0.338	−0.237	−1.750	0.086	−1.269	0.086	2.738	0.074	0.165	2.700	0.018
	Medications	0.143	0.385	0.048	0.370	0.713	−0.630	0.915					
	%REM	−0.051	0.024	−0.275	−2.113	0.039	−0.100	−0.003					
	REMD	0.156	0.052	0.414	2.976	0.004	0.051	0.261					
	REML	−0.010	0.003	−0.397	−3.021	0.004	−0.017	−0.003					
	HF[ms ²]	−0.416	0.199	−0.358	−2.089	0.042	−0.816	−0.016					
	HF[ms ²] × Sex	0.651	0.293	0.401	2.223	0.030	0.064	1.238					

HF [ms²], absolute power of high frequency heart rate variability; %REM, proportion of REM sleep to the total sleep time; REMD, REM density; REML, REM latency. HF [ms²] and HF [ms²] × sex interaction were significant predictors. Note that smaller ERI denotes better extinction recall.

Table 6. Hierarchical regression analysis for subjective extinction recall (sERI; $N = 85$)

Model	Predictors	<i>B</i>	S.E.	β	<i>t</i>	<i>p</i>	95% CI						
1	Age	−0.506	0.630	−0.085	−0.803	0.424	−1.761	0.748					
	Sex	−0.224	7.153	−0.004	−0.031	0.975	−14.458	14.010					
	Medications	−14.982	8.278	−0.203	−1.810	0.074	−31.456	1.492					
	REMD	2.917	0.970	0.348	3.006	0.004	0.986	4.849					
									Change		ANOVA		
									ΔF	<i>p</i>	Adj. R^2	<i>F</i>	<i>p</i>
2	Age	−0.678	0.616	−0.114	−1.100	0.274	−1.905	0.549	5.834	0.018	0.133	3.566	0.006
	Sex	0.830	6.960	0.013	0.119	0.905	−13.022	14.683					
	Medications	−18.809	8.194	−0.255	−2.296	0.024	−35.118	−2.500					
	REMD	2.638	0.950	0.314	2.778	0.007	0.748	4.528					
	HF[ms ²]	−7.048	2.918	−0.255	−2.415	0.018	−12.857	−1.240					

HF [ms²], absolute power of high frequency heart rate variability; REMD, REM density. Addition of HF [ms²] significantly increased the proportion of variance explained by the model. Note that smaller sERI denotes better extinction recall.

The mechanisms underlying REM alterations in PTSD and, by extension, how they might be associated with impaired extinction recall are not clear. However, it was proposed that hyperarousal, characterized by impairment in the inhibitory control of amygdala activity by the medial PFC (mPFC) with a concomitant increase in noradrenergic activity, contributes to REM dysregulation (Cabrera et al., 2024; Germain, Buysse, & Nofzinger, 2008; Pace-Schott et al., 2023; Pace-Schott, Germain, & Milad, 2015b). Indeed, increased REMD may be a direct manifestation of hyperarousal in PTSD (Barbato, 2023). Rapid eye movements are associated with activation in the limbic and paralimbic structures (Andrillon, Nir, Cirelli, Tononi, & Fried, 2015; Calvo & Fernandez-Guardiola, 1984; Corsi-Cabrera et al., 2016; Ioannides et al., 2004; Miyauchi, Misaki, Kan, Fukunaga, & Koike, 2009; Wehrle et al., 2007), as well as heart rate surges (Rowe et al., 1999) and nightmares concurrent with autonomic arousal (Paul, Alpers, Reinhard, & Schredl, 2019). Furthermore, in an overlapping sample, we recently showed that REMD was one of the predictors of self-reported hyperarousal symptoms in trauma-exposed individuals (Daffre et al., 2023). Therefore, our results suggest that hyperarousal during REM interferes with the consolidation of extinction memory. REML has been considered an indicator of REM pressure, and the shortening of REML in PTSD has been attributed to an 'unmet need' (Mellman, 1997). This interpretation would suggest that a history of insufficient REM or a latent factor associated with a REM deficit may impair the consolidation of extinction memory. Alternatively, this association may be due to an adaptive REM enhancement to facilitate emotional processing (Mellman, 1997). Yet another possibility is that shortened REML may reflect depression (Palagini, Baglioni, Ciapparelli, Gemignani, & Riemann, 2013). However, self-reported depressive symptoms were not correlated with REML ($r_s = 0.05$, $p = 0.65$) or ERI ($r_s = 0.02$, $p = 0.86$).

Vagal activity during REM, indexed as HF HRV, was a significant predictor of subjective extinction recall. This novel finding is consistent with the role of the vagal nerve in supporting memory formation (McGaugh, 2013) and emotional regulation (Thayer & Lane, 2000), as well as the growing body of evidence indicating that vagal activity is causally involved in fear processing (Alvarez-Dieppa, Griffin, Cavalier, & McIntyre, 2016; Pena et al., 2013, 2014). Vagal activity in response to cognitive and emotional demands and vagally mediated HRV are considered to reflect the tonic inhibitory control of the amygdala by the mPFC (Thayer & Lane, 2009; Thayer, Ahs, Fredrikson, Sollers, & Wager, 2012), circuitry that is also critical for fear extinction (Milad & Quirk, 2012). Consistent with this, high vagal activity during wake was repeatedly shown to be associated with better fear extinction in humans (Jenness et al., 2019; Pappens et al., 2014; Wendt et al., 2015, 2019). Furthermore, vagal nerve stimulation (VNS) facilitated plasticity in ventromedial PFC-amygdala connectivity in rodents (Pena et al., 2014), and improved extinction learning and recall in both rodents (Alvarez-Dieppa et al., 2016; Noble et al., 2017; Noble, Chuah, Callahan, Souza, & McIntyre, 2019a; Pena et al., 2013, 2014; Souza et al., 2019, 2020, 2021a, 2021b, 2022) and humans (Burger et al., 2016, 2017, 2019; Szeska et al., 2020). Our results suggest that the contribution of vagal activity continues beyond extinction learning to its consolidation during sleep and supports the potential utility of interventions that can enhance sleep vagal activity in the prevention and treatment of fear-related disorders (Noble, Souza, & McIntyre, 2019c). The association of HRV with physiological extinction recall in males should be treated as a preliminary

finding, because of the small number of male participants in our sample ($N = 34$). Previous studies reported that sex and sex hormones were associated with differential activations in PFC and amygdala during extinction learning and recall (Benson et al., 2014; Merz et al., 2012; Zeidan et al., 2011). Therefore, the moderating effect of sex for the association of HRV with physiological extinction recall in our study may be due to the sex differences in the engagement of overlapping circuitry that processes fear extinction and regulates vagal activity.

Our study had several limitations. First, we had to exclude a substantial number of participants because of lost or unusable sleep data. Lack of oversight during ambulatory recordings can lead to more artifacts than in laboratory studies. Second, there is significant variability across studies in fear conditioning/extinction protocols and methods in analyzing generated data (Lonsdorf et al., 2017). Nonetheless, we used the widely employed Milad et al., 2007 fear conditioning and extinction protocol (Milad & Quirk, 2012), and we attempted to be consistent with our own and others' previously used metrics to index extinction recall. Third, a portion of our sample did not achieve physiological fear conditioning and was therefore excluded from the analysis. Nonetheless, the proportion of 'non-conditioners' in our sample (49/139, 35%) was smaller than in many previous studies (Alexandra Kredlow, Orr, & Otto, 2018). Lastly, the models for ERI and sERI revealed distinct predictors. A substantial body of research suggests that explicit and implicit fear conditioning, as well as extinction, are separate processes that engage overlapping but distinct neural circuits (Grillon, 2009; Knight, Nguyen, & Bandettini, 2003; Knight, Waters, & Bandettini, 2009; Siegel & Weinberger, 2012). This dissociation is also evident in our sample, where physiological and subjective extinction recall indexes (ERI and sERI) are not correlated (spearman's $\rho = 0.07$, $p = 0.52$). Therefore, it is not surprising that different predictors emerged for ERI and sERI.

Conclusions

Abnormalities in REM have repeatedly been reported in individuals diagnosed with PTSD and have been shown to be associated with an increased risk of developing PTSD after a traumatic event. Results of this study further our insight into the role of REM disruptions and indicate that they are associated with impaired consolidation of extinction memory, a mechanism proposed to be critical in the pathogenesis of PTSD.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291724002757>

Acknowledgments. The authors would like to thank Karen Gannon, RPSGT for scoring sleep recordings and Catherine Bostian for adapting the Yetton et al. REM density program.

Author contributions. Cagri Yuksel and Edward F. Pace-Schott conceptualized the study. Lauren Watford, Monami Muranaka, Emma McCoy, Augustus Kram Mendelsohn, Katelyn I. Oliver, Carolina Daffre, Hannah Lax, and Uriel Martinez, collected and reduced data. Cagri Yuksel, Lauren Watford, Monami Muranaka, Hannah Lax, Alexis Acosta, and Abigail Vidrin analyzed the data. Natasha B. Lasko interviewed participants. Carolina Daffre and Scott Orr provided consultation on statistical analyses. Cagri Yuksel and Edward F. Pace-Schott authored the manuscript draft. Carolina Daffre and Scott Orr provided feedback on the manuscript draft.

Funding statement. This project was primarily supported by the National Institute of Mental Health R01MH109638 and R21MH121832 to E.P.-S. and

K23MH119322 to C.Y. Praxis Precision Medicines also provided partial salary support to E.P.-S. Funding bodies did not have any role in the design of the study, collection, and analysis of data, and decision to publish.

Competing interests. The authors declare no competing interests.

References

- Alexandra Kredlow, M., Orr, S. P., & Otto, M. W. (2018). Who is studied in de novo fear conditioning paradigms? An examination of demographic and stimulus characteristics predicting fear learning. *International Journal of Psychophysiology*, 130, 21–28. <https://doi.org/10.1016/j.ijpsycho.2018.05.008>
- Alvarez-Dieppa, A. C., Griffin, K., Cavalier, S., & McIntyre, C. K. (2016). Vagus nerve stimulation enhances extinction of conditioned fear in rats and modulates arc protein, CaMKII, and GluN2B-containing NMDA receptors in the basolateral amygdala. *Neural Plasticity*, 2016, 4273280. <https://doi.org/10.1155/2016/4273280>
- Andrillon, T., Nir, Y., Cirelli, C., Tononi, G., & Fried, I. (2015). Single-neuron activity and eye movements during human REM sleep and awake vision. *Nature Communications*, 6, 7884. <https://doi.org/10.1038/ncomms8884>
- Baglioni, C., Nanovska, S., Regen, W., Spiegelhalter, K., Feige, B., Nissen, C., ... Riemann, D. (2016). Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychological Bulletin*, 142(9), 969–990. <https://doi.org/10.1037/bul0000053>
- Barbato, G. (2023). Is REM density a measure of arousal during sleep? *Brain Sciences*, 13(3), 378. <https://doi.org/10.3390/brainsci13030378>
- Benson, S., Kattoor, J., Kullmann, J. S., Hofmann, S., Engler, H., Forsting, M., ... Elsenbruch, S. (2014). Towards understanding sex differences in visceral pain: Enhanced reactivation of classically-conditioned fear in healthy women. *Neurobiology of Learning and Memory*, 109, 113–121. <https://doi.org/10.1016/j.nlm.2013.12.014>
- Berry, R. B., Gamaldo, C. E., Harding, S. M., & Brooks, R. (2015). *The AASM manual for the scoring of sleep and associated events: Rules, terminology and technical specifications*. Darian, IL: American Academy of Sleep Medicine.
- Bottary, R., Seo, J., Daffre, C., Gazecki, S., Moore, K. N., Kopotiyenko, K., ... Pace-Schott, E. F. (2020). Fear extinction memory is negatively associated with REM sleep in insomnia disorder. *Sleep*, 43(7). <https://doi.org/10.1093/sleep/zsaa007>
- Bottary, R., Straus, L. D., & Pace-Schott, E. F. (2023). The impact of sleep on fear extinction. *Current Topics in Behavioral Neurosciences*, 64, 133–156. https://doi.org/10.1007/7854_2023_431
- Bouton, M. E., Kenney, F. A., & Rosengard, C. (1990). State-dependent fear extinction with two benzodiazepine tranquilizers. *Behavioral Neuroscience*, 104(1), 44–55. <https://doi.org/10.1037//0735-7044.104.1.44>
- Breslau, N., Roth, T., Burduvali, E., Kapke, A., Schultz, L., & Roehrs, T. (2004). Sleep in lifetime posttraumatic stress disorder: A community-based polysomnographic study. *Archives of General Psychiatry*, 61(5), 508–516. <https://doi.org/10.1001/archpsyc.61.5.508>
- Broncel, A., Bocian, R., Klos-Wojtczak, P., Kulbat-Warycha, K., & Konopacki, J. (2020). Vagal nerve stimulation as a promising tool in the improvement of cognitive disorders. *Brain Research Bulletin*, 155, 37–47. <https://doi.org/10.1016/j.brainresbull.2019.11.011>
- Burger, A. M., Verkuil, B., Van Diest, I., Van der Does, W., Thayer, J. F., & Brosschot, J. F. (2016). The effects of transcutaneous vagus nerve stimulation on conditioned fear extinction in humans. *Neurobiology of Learning and Memory*, 132, 49–56. <https://doi.org/10.1016/j.nlm.2016.05.007>
- Burger, A. M., Verkuil, B., Fenlon, H., Thijs, L., Cools, L., Miller, H. C., ... Van Diest, I. (2017). Mixed evidence for the potential of non-invasive transcutaneous vagal nerve stimulation to improve the extinction and retention of fear. *Behaviour Research and Therapy*, 97, 64–74. <https://doi.org/10.1016/j.brat.2017.07.005>
- Burger, A. M., Van Diest, I., Van der Does, W., Korbee, J. N., Waziri, N., Brosschot, J. F., & Verkuil, B. (2019). The effect of transcutaneous vagus nerve stimulation on fear generalization and subsequent fear extinction. *Neurobiology of Learning and Memory*, 161, 192–201. <https://doi.org/10.1016/j.nlm.2019.04.006>
- Cabrera, Y., Koymans, K. J., Poe, G. R., Kessels, H. W., Van Someren, E. J. W., & Wassing, R. (2024). Overnight neuronal plasticity and adaptation to emotional distress. *Nature Reviews Neuroscience*, 25(4), 253–271. <https://doi.org/10.1038/s41583-024-00799-w>
- Calvo, J. M., & Fernandez-Guardiola, A. (1984). Phasic activity of the basolateral amygdala, cingulate gyrus, and hippocampus during REM sleep in the cat. *Sleep*, 7(3), 202–210. <https://doi.org/10.1093/sleep/7.3.202>
- Chalmers, J. A., Quintana, D. S., Abbott, M. J., & Kemp, A. H. (2014). Anxiety disorders are associated with reduced heart rate variability: A meta-analysis. *Frontiers in Psychiatry*, 5, 80. <https://doi.org/10.3389/fpsy.2014.00080>
- Colvonen, P. J., Straus, L. D., Acheson, D., & Gehrman, P. (2019). A review of the relationship between emotional learning and memory, sleep, and PTSD. *Current Psychiatry Reports*, 21(1), 2. <https://doi.org/10.1007/s11920-019-0987-2>
- Corsi-Cabrera, M., Velasco, F., Del Rio-Portilla, Y., Armony, J. L., Trejo-Martinez, D., Guevara, M. A., & Velasco, A. L. (2016). Human amygdala activation during rapid eye movements of rapid eye movement sleep: An intracranial study. *Journal of Sleep Research*, 25(5), 576–582. <https://doi.org/10.1111/jsr.12415>
- Craske, M. G., Treanor, M., Conway, C. C., Zbozinek, T., & Vervliet, B. (2014). Maximizing exposure therapy: An inhibitory learning approach. *Behaviour Research and Therapy*, 58, 10–23. <https://doi.org/10.1016/j.brat.2014.04.006>
- Daffre, C., Oliver, K. I., Nazareno, J. R. S., Mader, T., Seo, J., Dominguez, J. P., ... Pace-Schott, E. F. (2023). Rapid eye movement sleep parasympathetic activity predicts wake hyperarousal symptoms following a traumatic event. *Journal of Sleep Research*, 32(1), e13685. <https://doi.org/10.1111/jsr.13685>
- Davidson, P., & Pace-Schott, E. (2020). The role of sleep in fear learning and memory. *Current Opinion in Psychology*, 34, 32–36. <https://doi.org/10.1016/j.copsyc.2019.08.016>
- Davidson, P., Jonsson, P., Carlsson, I., & Pace-Schott, E. (2021). Does sleep selectively strengthen certain memories over others based on emotion and perceived future relevance? *Nature and Science of Sleep*, 13, 1257–1306. <https://doi.org/10.2147/NSS.S286701>
- Dawson, M. E., Schell, A. M., & Filion, D. L. (2007). The electrodermal system. In J. T. Cacioppo, L. G. Tassinary & G. G. Berntson (Eds.), *Handbook of psychophysiology* (3rd ed., pp. 159–181). New York, NY: Cambridge University Press.
- First, M., Gibbon, M., Spitzer, R., & Williams, J. (2007). *Structured clinical interview for DSM-IV-TR axis I disorders–NonPatient edition (SCID-I/NP)*. New York, NY: Biometrics Research Department, New York State Psychiatric Institute.
- Ge, F., Yuan, M., Li, Y., & Zhang, W. (2020). Posttraumatic stress disorder and alterations in resting heart rate variability: A systematic review and meta-analysis. *Psychiatry Investigation*, 17(1), 9–20. <https://doi.org/10.30773/pi.2019.0112>
- Germain, A., Buysse, D. J., & Nofzinger, E. (2008). Sleep-specific mechanisms underlying posttraumatic stress disorder: Integrative review and neurobiological hypotheses. *Sleep Medicine Reviews*, 12(3), 185–195. <https://doi.org/10.1016/j.smrv.2007.09.003>
- Giustino, T. F., & Maren, S. (2015). The role of the medial prefrontal cortex in the conditioning and extinction of fear. *Frontiers in Behavioral Neuroscience*, 9, 298. <https://doi.org/10.3389/fnbeh.2015.00298>
- Goldstein, A. N., & Walker, M. P. (2014). The role of sleep in emotional brain function. *Annual Review of Clinical Psychology*, 10, 679–708. <https://doi.org/10.1146/annurev-clinpsy-032813-153716>
- Grillon, C. (2009). D-cycloserine facilitation of fear extinction and exposure-based therapy might rely on lower-level, automatic mechanisms. *Biological Psychiatry*, 66(7), 636–641. <https://doi.org/10.1016/j.biopsych.2009.04.017>
- Gunduz-Cinar, O., Flynn, S., Brockway, E., Kaugars, K., Baldi, R., Ramikie, T. S., ... Holmes, A. (2016). Fluoxetine facilitates fear extinction through amygdala endocannabinoids. *Neuropsychopharmacology*, 41(6), 1598–1609. <https://doi.org/10.1038/npp.2015.318>
- Habukawa, M., Uchimura, N., Maeda, M., Ogi, K., Hiejima, H., & Kakuma, T. (2018). Differences in rapid eye movement (REM) sleep abnormalities between posttraumatic stress disorder (PTSD) and major depressive disorder patients: REM interruption correlated with nightmare complaints in PTSD. *Sleep Medicine*, 43, 34–39. <https://doi.org/10.1016/j.sleep.2017.10.012>
- Helpman, L., Marin, M. F., Papini, S., Zhu, X., Sullivan, G. M., Schneier, F., ... Neria, Y. (2016). Neural changes in extinction recall following prolonged

- exposure treatment for PTSD: A longitudinal fMRI study. *Neuroimage. Clinical*, 12, 715–723. <https://doi.org/10.1016/j.nicl.2016.10.007>
- Insana, S. P., Kolk, D. J., & Germain, A. (2012). Early-life trauma is associated with rapid eye movement sleep fragmentation among military veterans. *Biological Psychology*, 89(3), 570–579. <https://doi.org/10.1016/j.biopsycho.2012.01.001>
- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., ... Wang, P. (2010). Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*, 167(7), 748–751. <https://doi.org/10.1176/appi.ajp.2010.09091379>
- Ioannides, A. A., Corsi-Cabrera, M., Fenwick, P. B., del Rio Portilla, Y., Laskaris, N. A., Khurshudyan, A., ... Kostopoulos, G. K. (2004). MEG tomography of human cortex and brainstem activity in waking and REM sleep saccades. *Cerebral Cortex (New York, N.Y.: 1991)*, 14(1), 56–72. <https://doi.org/10.1093/cercor/bhg091>
- Jenness, J. L., Miller, A. B., Rosen, M. L., & McLaughlin, K. A. (2019). Extinction learning as a potential mechanism linking high vagal tone with lower PTSD symptoms among abused youth. *Journal of Abnormal Child Psychology*, 47(4), 659–670. <https://doi.org/10.1007/s10802-018-0464-0>
- Jung, B., Yang, C., & Lee, S. H. (2024). Vagus nerves stimulation: Clinical implication and practical issue as a neuropsychiatric treatment. *Clinical Psychopharmacology and Neuroscience*, 22(1), 13–22. <https://doi.org/10.9758/cpn.23.1101>
- Klinzing, J. G., Niethard, N., & Born, J. (2019). Mechanisms of systems memory consolidation during sleep. *Nature Neuroscience*, 22(10), 1598–1610. <https://doi.org/10.1038/s41593-019-0467-3>
- Knight, D. C., Nguyen, H. T., & Bandettini, P. A. (2003). Expression of conditional fear with and without awareness. *Proceedings of the National Academy of Sciences of the United States of America*, 100(25), 15280–15283. <https://doi.org/10.1073/pnas.2535780100>
- Knight, D. C., Waters, N. S., & Bandettini, P. A. (2009). Neural substrates of explicit and implicit fear memory. *Neuroimage*, 45(1), 208–214. <https://doi.org/10.1016/j.neuroimage.2008.11.015>
- Kobayashi, I., Boarts, J. M., & Delahanty, D. L. (2007). Polysomnographically measured sleep abnormalities in PTSD: A meta-analytic review. *Psychophysiology*, 44(4), 660–669. <https://doi.org/10.1111/j.1469-8986.2007.037.x>
- Kobayashi, I., Lavelle, J., & Mellman, T. A. (2014). Nocturnal autonomic balance and sleep in PTSD and resilience. *Journal of Traumatic Stress*, 27(6), 712–716. <https://doi.org/10.1002/jts.21973>
- Laborde, S., Mosley, E., & Thayer, J. F. (2017). Heart rate variability and cardiac vagal tone in psychophysiological research – recommendations for experiment planning, data analysis, and data reporting. *Frontiers in Psychology*, 8, 213. <https://doi.org/10.3389/fpsyg.2017.00213>
- Lipinska, G., & Thomas, K. G. F. (2019). Rapid eye movement fragmentation, not slow-wave sleep, predicts neutral declarative memory consolidation in posttraumatic stress disorder. *Journal of Sleep Research*, 28(6), e12846. <https://doi.org/10.1111/jsr.12846>
- Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J., ... Merz, C. J. (2017). Don't fear 'fear conditioning': Methodological considerations for the design and analysis of studies on human fear acquisition, extinction, and return of fear. *Neuroscience & Biobehavioral Reviews*, 77, 247–285. <https://doi.org/10.1016/j.neubiorev.2017.02.026>
- Lonsdorf, T. B., Merz, C. J., & Fullana, M. A. (2019). Fear extinction retention: Is it what we think it is? *Biological Psychiatry*, 85(12), 1074–1082. <https://doi.org/10.1016/j.biopsycho.2019.02.011>
- McGaugh, J. L. (2013). Making lasting memories: Remembering the significant. *Proceedings of the National Academy of Sciences of the United States of America*, 110(Suppl 2), 10402–10407. <https://doi.org/10.1073/pnas.1301209110>
- Mellman, T. A. (1997). Psychobiology of sleep disturbances in posttraumatic stress disorder. *Annals of the New York Academy of Sciences*, 821, 142–149. <https://doi.org/10.1111/j.1749-6632.1997.tb48275.x>
- Mellman, T. A., Bustamante, V., Fins, A. I., Pigeon, W. R., & Nolan, B. (2002). REM sleep and the early development of posttraumatic stress disorder. *American Journal of Psychiatry*, 159(10), 1696–1701. <https://doi.org/10.1176/appi.ajp.159.10.1696>
- Mellman, T. A., Knorr, B. R., Pigeon, W. R., Leiter, J. C., & Akay, M. (2004). Heart rate variability during sleep and the early development of post-traumatic stress disorder. *Biological Psychiatry*, 55(9), 953–956. <https://doi.org/10.1016/j.biopsycho.2003.12.018>
- Mellman, T. A., Pigeon, W. R., Nowell, P. D., & Nolan, B. (2007). Relationships between REM sleep findings and PTSD symptoms during the early aftermath of trauma. *Journal of Traumatic Stress*, 20(5), 893–901. <https://doi.org/10.1002/jts.20246>
- Mellman, T. A., Kobayashi, I., Lavelle, J., Wilson, B., & Hall Brown, T. S. (2014). A relationship between REM sleep measures and the duration of post-traumatic stress disorder in a young adult urban minority population. *Sleep*, 37(8), 1321–1326. <https://doi.org/10.5665/sleep.3922>
- Menz, M. M., Rihm, J. S., & Buchel, C. (2016). REM sleep is causal to successful consolidation of dangerous and safety stimuli and reduces return of fear after extinction. *Journal of Neuroscience*, 36(7), 2148–2160. <https://doi.org/10.1523/JNEUROSCI.3083-15.2016>
- Merz, C. J., Tabbert, K., Schweckendiek, J., Klucken, T., Vaitl, D., Stark, R., & Wolf, O. T. (2012). Neuronal correlates of extinction learning are modulated by sex hormones. *Social Cognitive and Affective Neuroscience*, 7(7), 819–830. <https://doi.org/10.1093/scan/nsr063>
- Milad, M. R., & Quirk, G. J. (2012). Fear extinction as a model for translational neuroscience: Ten years of progress. *Annual Review of Psychology*, 63, 129–151. <https://doi.org/10.1146/annurev.psych.121208.131631>
- Milad, M. R., Goldstein, J. M., Orr, S. P., Wedig, M. M., Klitsch, A., Pitman, R. K., & Rauch, S. L. (2006). Fear conditioning and extinction: Influence of sex and menstrual cycle in healthy humans. *Behavioral Neuroscience*, 120(6), 1196–1203. <https://doi.org/10.1037/0735-7044.120.5.1196>
- Milad, M. R., Wright, C. I., Orr, S. P., Pitman, R. K., Quirk, G. J., & Rauch, S. L. (2007). Recall of fear extinction in humans activates the ventromedial prefrontal cortex and hippocampus in concert. *Biological Psychiatry*, 62(5), 446–454. <https://doi.org/10.1016/j.biopsycho.2006.10.011>
- Milad, M. R., Orr, S. P., Lasko, N. B., Chang, Y., Rauch, S. L., & Pitman, R. K. (2008). Presence and acquired origin of reduced recall for fear extinction in PTSD: Results of a twin study. *Journal of Psychiatric Research*, 42(7), 515–520. <https://doi.org/10.1016/j.jpsychires.2008.01.017>
- Milad, M. R., Pitman, R. K., Ellis, C. B., Gold, A. L., Shin, L. M., Lasko, N. B., ... Rauch, S. L. (2009). Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biological Psychiatry*, 66(12), 1075–1082. <https://doi.org/10.1016/j.biopsycho.2009.06.026>
- Milad, M. R., Furtak, S. C., Greenberg, J. L., Keshaviah, A., Im, J. J., Falkenstein, M. J., ... Wilhelm, S. (2013). Deficits in conditioned fear extinction in obsessive-compulsive disorder and neurobiological changes in the fear circuit. *JAMA Psychiatry*, 70(6), 608–618, quiz 554. <https://doi.org/10.1001/jamapsychiatry.2013.914>
- Miyauchi, S., Misaki, M., Kan, S., Fukunaga, T., & Koike, T. (2009). Human brain activity time-locked to rapid eye movements during REM sleep. *Experimental Brain Research*, 192(4), 657–667. <https://doi.org/10.1007/s00221-008-1579-2>
- Noble, L. J., Gonzalez, I. J., Meruva, V. B., Callahan, K. A., Belfort, B. D., Ramanathan, K. R., ... McIntyre, C. K. (2017). Effects of vagus nerve stimulation on extinction of conditioned fear and post-traumatic stress disorder symptoms in rats. *Translational Psychiatry*, 7(8), e1217. <https://doi.org/10.1038/tp.2017.191>
- Noble, L. J., Chuah, A., Callahan, K. K., Souza, R. R., & McIntyre, C. K. (2019a). Peripheral effects of vagus nerve stimulation on anxiety and extinction of conditioned fear in rats. *Learning & Memory (Cold Spring Harbor, N.Y.)*, 26(7), 245–251. <https://doi.org/10.1101/lm.048447.118>
- Noble, L. J., Meruva, V. B., Hays, S. A., Rennaker, R. L., Kilgord, M. P., & McIntyre, C. K. (2019b). Vagus nerve stimulation promotes generalization of conditioned fear extinction and reduces anxiety in rats. *Brain Stimulation*, 12(1), 9–18. <https://doi.org/10.1016/j.brs.2018.09.013>
- Noble, L. J., Souza, R. R., & McIntyre, C. K. (2019c). Vagus nerve stimulation as a tool for enhancing extinction in exposure-based therapies. *Psychopharmacology (Berl)*, 236(1), 355–367. <https://doi.org/10.1007/s00213-018-4994-5>
- Orr, S. P., Metzger, L. J., Lasko, N. B., Macklin, M. L., Peri, T., & Pitman, R. K. (2000). De novo conditioning in trauma-exposed individuals with and without posttraumatic stress disorder. *Journal of Abnormal Psychology*, 109(2), 290–298.

- Pace-Schott, E. F., Tracy, L. E., Rubin, Z., Mollica, A. G., Ellenbogen, J. M., Bianchi, M. T., ... Orr, S. P. (2014). Interactions of time of day and sleep with between-session habituation and extinction memory in young adult males. *Experimental Brain Research*, 232(5), 1443–1458. <https://doi.org/10.1007/s00221-014-3829-9>
- Pace-Schott, E. F., Germain, A., & Milad, M. R. (2015a). Effects of sleep on memory for conditioned fear and fear extinction. *Psychological Bulletin*, 141(4), 835–857. <https://doi.org/10.1037/bul0000014>
- Pace-Schott, E. F., Germain, A., & Milad, M. R. (2015b). Sleep and REM sleep disturbance in the pathophysiology of PTSD: The role of extinction memory. *Biology of Mood & Anxiety Disorders*, 5, 3. <https://doi.org/10.1186/s13587-015-0018-9>
- Pace-Schott, E. F., Seo, J., & Bottary, R. (2023). The influence of sleep on fear extinction in trauma-related disorders. *Neurobiology of Stress*, 22, 100500. <https://doi.org/10.1016/j.ynstr.2022.100500>
- Palagini, L., Baglioni, C., Ciapparelli, A., Gemignani, A., & Riemann, D. (2013). REM sleep dysregulation in depression: State of the art. *Sleep Medicine Reviews*, 17(5), 377–390. <https://doi.org/10.1016/j.smrv.2012.11.001>
- Pappens, M., Schroijen, M., Sutterlin, S., Smets, E., Van den Bergh, O., Thayer, J. F., & Van Diest, I. (2014). Resting heart rate variability predicts safety learning and fear extinction in an interoceptive fear conditioning paradigm. *PLoS One*, 9(9), e105054. <https://doi.org/10.1371/journal.pone.0105054>
- Paul, F., Alpers, G. W., Reinhard, I., & Schredl, M. (2019). Nightmares do result in psychophysiological arousal: A multimeasure ambulatory assessment study. *Psychophysiology*, 56(7), e13366. <https://doi.org/10.1111/psyp.13366>
- Pena, D. F., Engineer, N. D., & McIntyre, C. K. (2013). Rapid remission of conditioned fear expression with extinction training paired with vagus nerve stimulation. *Biological Psychiatry*, 73(11), 1071–1077. <https://doi.org/10.1016/j.biopsych.2012.10.021>
- Pena, D. F., Childs, J. E., Willett, S., Vital, A., McIntyre, C. K., & Kroener, S. (2014). Vagus nerve stimulation enhances extinction of conditioned fear and modulates plasticity in the pathway from the ventromedial prefrontal cortex to the amygdala. *Frontiers in Behavioral Neuroscience*, 8, 327. <https://doi.org/10.3389/fnbeh.2014.00327>
- Pyne, J. M., Constans, J. L., Wiederhold, M. D., Gibson, D. P., Kimbrell, T., Kramer, T. L., ... McCune, T. R. (2016). Heart rate variability: Pre-deployment predictor of post-deployment PTSD symptoms. *Biological Psychology*, 121(Pt A), 91–98. <https://doi.org/10.1016/j.biopsycho.2016.10.008>
- Richards, A., Kanady, J. C., & Neylan, T. C. (2020). Sleep disturbance in PTSD and other anxiety-related disorders: An updated review of clinical features, physiological characteristics, and psychological and neurobiological mechanisms. *Neuropsychopharmacology*, 45(1), 55–73. <https://doi.org/10.1038/s41386-019-0486-5>
- Risling, M. B., Dennis, P. A., Watkins, L. L., Calhoun, P. S., Dennis, M. F., Beckham, J. C., ... Ulmer, C. S. (2016). Circadian contrasts in heart rate variability associated with posttraumatic stress disorder symptoms in a young adult cohort. *Journal of Traumatic Stress*, 29(5), 415–421. <https://doi.org/10.1002/jts.22125>
- Rowe, K., Moreno, R., Lau, T. R., Wallooppillai, U., Nearing, B. D., Kocsis, B., ... Verrier, R. L. (1999). Heart rate surges during REM sleep are associated with theta rhythm and PGO activity in cats. *American Journal of Physiology*, 277(3), R843–R849. <https://doi.org/10.1152/ajpregu.1999.277.3.R843>
- Rush, A. J., Trivedi, M. H., Ibrahim, H. M., Carmody, T. J., Arnow, B., Klein, D. N., ... Keller, M. B. (2003). The 16-item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): A psychometric evaluation in patients with chronic major depression. *Biological Psychiatry*, 54(5), 573–583. [https://doi.org/10.1016/S0006-3223\(02\)01866-8](https://doi.org/10.1016/S0006-3223(02)01866-8)
- Saguin, E., Gomez-Merino, D., Sauvet, F., Leger, D., & Chennaoui, M. (2021). Sleep and PTSD in the military forces: A reciprocal relationship and a psychiatric approach. *Brain Sciences*, 11(10), 1310. <https://doi.org/10.3390/brainsci11101310>
- Schneider, M., & Schwerdtfeger, A. (2020). Autonomic dysfunction in post-traumatic stress disorder indexed by heart rate variability: A meta-analysis. *Psychological Medicine*, 50(12), 1937–1948. <https://doi.org/10.1017/S003329172000207X>
- Seligowski, A. V., Webber, T. K., Marvar, P. J., Ressler, K. J., & Philip, N. S. (2022). Involvement of the brain-heart axis in the link between PTSD and cardiovascular disease. *Depression and Anxiety*, 39(10–11), 663–674. <https://doi.org/10.1002/da.23271>
- Seo, J., Oliver, K. I., Daffre, C., Moore, K. N., Gazecki, S., Lasko, N. B., ... Pace-Schott, E. F. (2022). Associations of sleep measures with neural activations accompanying fear conditioning and extinction learning and memory in trauma-exposed individuals. *Sleep*, 45(3), zsab261. <https://doi.org/10.1093/sleep/zsab261>
- Shaffer, F., & Ginsberg, J. P. (2017). An overview of heart rate variability metrics and norms. *Frontiers in Public Health*, 5, 258. <https://doi.org/10.3389/fpubh.2017.00258>
- Shvil, E., Sullivan, G. M., Schafer, S., Markowitz, J. C., Campeas, M., Wager, T. D., ... Neria, Y. (2014). Sex differences in extinction recall in posttraumatic stress disorder: A pilot fMRI study. *Neurobiology of Learning and Memory*, 113, 101–108. <https://doi.org/10.1016/j.nlm.2014.02.003>
- Siegel, P., & Weinberger, J. (2012). Less is more: The effects of very brief versus clearly visible exposure. *Emotion (Washington, D.C.)*, 12(2), 394–402. <https://doi.org/10.1037/a0026806>
- Siepmann, M., Weidner, K., Petrowski, K., & Siepmann, T. (2022). Heart rate variability: A measure of cardiovascular health and possible therapeutic target in dysautonomic mental and neurological disorders. *Applied Psychophysiology and Biofeedback*, 47(4), 273–287. <https://doi.org/10.1007/s10484-022-09572-0>
- Souza, R. R., Robertson, N. M., Pruitt, D. T., Gonzales, P. A., Hays, S. A., Rennaker, R. L., ... McIntyre, C. K. (2019). Vagus nerve stimulation reverses the extinction impairments in a model of PTSD with prolonged and repeated trauma. *Stress (Amsterdam, Netherlands)*, 22(4), 509–520. <https://doi.org/10.1080/10253890.2019.1602604>
- Souza, R. R., Robertson, N. M., Mathew, E., Tabet, M. N., Bucksot, J. E., Pruitt, D. T., ... Kilgard, M. P. (2020). Efficient parameters of vagus nerve stimulation to enhance extinction learning in an extinction-resistant rat model of PTSD. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 99, 109848. <https://doi.org/10.1016/j.pnpbp.2019.109848>
- Souza, R. R., Oleksiak, C. R., Tabet, M. N., Rennaker, R. L., Hays, S. A., Kilgard, M. P., & McIntyre, C. K. (2021a). Vagus nerve stimulation promotes extinction generalization across sensory modalities. *Neurobiology of Learning and Memory*, 181, 107425. <https://doi.org/10.1016/j.nlm.2021.107425>
- Souza, R. R., Robertson, N. M., McIntyre, C. K., Rennaker, R. L., Hays, S. A., & Kilgard, M. P. (2021b). Vagus nerve stimulation enhances fear extinction as an inverted-U function of stimulation intensity. *Experimental Neurology*, 341, 113718. <https://doi.org/10.1016/j.expneurol.2021.113718>
- Souza, R. R., Powers, M. B., Rennaker, R. L., McIntyre, C. K., Hays, S. A., & Kilgard, M. P. (2022). Timing of vagus nerve stimulation during fear extinction determines efficacy in a rat model of PTSD. *Scientific Reports*, 12(1), 16526. <https://doi.org/10.1038/s41598-022-20301-9>
- Spoormaker, V. I., Sturm, A., Andrade, K. C., Schroter, M. S., Goya-Maldonado, R., Holsboer, F., ... Czisch, M. (2010). The neural correlates and temporal sequence of the relationship between shock exposure, disturbed sleep and impaired consolidation of fear extinction. *Journal of Psychiatric Research*, 44(16), 1121–1128. <https://doi.org/10.1016/j.jpsychires.2010.04.017>
- Spoormaker, V. I., Schroter, M. S., Andrade, K. C., Dresler, M., Kiem, S. A., Goya-Maldonado, R., ... Czisch, M. (2012). Effects of rapid eye movement sleep deprivation on fear extinction recall and prediction error signaling. *Human Brain Mapping*, 33(10), 2362–2376. <https://doi.org/10.1002/hbm.21369>
- Stocker, R. P. J., Khan, H., Henry, L., & Germain, A. (2017). Effects of sleep loss on subjective complaints and objective neurocognitive performance as measured by the immediate post-concussion assessment and cognitive testing. *Archives of Clinical Neuropsychology*, 32(3), 349–368. <https://doi.org/10.1093/arclin/acx003>
- Straus, L. D., Norman, S. B., Risbrough, V. B., Acheson, D. T., & Drummond, S. P. A. (2018). REM sleep and safety signal learning in posttraumatic stress disorder: A preliminary study in military veterans. *Neurobiology of Stress*, 9, 22–28. <https://doi.org/10.1016/j.ynstr.2018.07.001>
- Suarez-Jimenez, B., Albajes-Eizaguirre, A., Lazarov, A., Zhu, X., Harrison, B. J., Radua, J., ... Fullana, M. A. (2020). Neural signatures of conditioning,

- extinction learning, and extinction recall in posttraumatic stress disorder: A meta-analysis of functional magnetic resonance imaging studies. *Psychological Medicine*, 50(9), 1442–1451. <https://doi.org/10.1017/S0033291719001387>
- Szeska, C., Richter, J., Wendt, J., Weymar, M., & Hamm, A. O. (2020). Promoting long-term inhibition of human fear responses by non-invasive transcutaneous vagus nerve stimulation during extinction training. *Scientific Reports*, 10(1), 1529. <https://doi.org/10.1038/s41598-020-58412-w>
- Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *Journal of Affective Disorders*, 61(3), 201–216. [https://doi.org/10.1016/S0165-0327\(00\)00338-4](https://doi.org/10.1016/S0165-0327(00)00338-4)
- Thayer, J. F., & Lane, R. D. (2009). Claude bernard and the heart-brain connection: Further elaboration of a model of neurovisceral integration. *Neuroscience & Biobehavioral Reviews*, 33(2), 81–88. <https://doi.org/10.1016/j.neubiorev.2008.08.004>
- Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart rate variability, prefrontal neural function, and cognitive performance: The neurovisceral integration perspective on self-regulation, adaptation, and health. *Annals of Behavioral Medicine*, 37(2), 141–153. <https://doi.org/10.1007/s12160-009-9101-z>
- Thayer, J. F., Ahs, F., Fredrikson, M., Sollers III J. J., & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews*, 36(2), 747–756. <https://doi.org/10.1016/j.neubiorev.2011.11.009>
- Ulmer, C. S., Hall, M. H., Dennis, P. A., Beckham, J. C., & Germain, A. (2018). Posttraumatic stress disorder diagnosis is associated with reduced parasympathetic activity during sleep in US veterans and military service members of the Iraq and Afghanistan wars. *Sleep*, 41(12), zsy174. <https://doi.org/10.1093/sleep/zsy174>
- Weathers, F. W., Bovin, M. J., Lee, D. J., Sloan, D. M., Schnurr, P. P., Kaloupek, D. G., ... Marx, B. P. (2018). The clinician-administered PTSD scale for DSM-5 (CAPS-5): Development and initial psychometric evaluation in military veterans. *Psychological Assessment*, 30(3), 383–395. <https://doi.org/10.1037/pas0000486>
- Wehrle, R., Kaufmann, C., Wetter, T. C., Holsboer, F., Auer, D. P., Pollmacher, T., & Cizisch, M. (2007). Functional microstates within human REM sleep: First evidence from fMRI of a thalamocortical network specific for phasic REM periods. *European Journal of Neuroscience*, 25(3), 863–871. <https://doi.org/10.1111/j.1460-9568.2007.05314.x>
- Wendt, J., Neubert, J., Koenig, J., Thayer, J. F., & Hamm, A. O. (2015). Resting heart rate variability is associated with inhibition of conditioned fear. *Psychophysiology*, 52(9), 1161–1166. <https://doi.org/10.1111/psyp.12456>
- Wendt, J., Konig, J., Hufenbach, M. C., Koenig, J., Thayer, J. F., & Hamm, A. O. (2019). Vagally mediated heart rate variability and safety learning: Effects of instructions and number of extinction trials. *Psychophysiology*, 56(10), e13404. <https://doi.org/10.1111/psyp.13404>
- Whitehurst, L. N., Cellini, N., McDevitt, E. A., Duggan, K. A., & Mednick, S. C. (2016). Autonomic activity during sleep predicts memory consolidation in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 113(26), 7272–7277. <https://doi.org/10.1073/pnas.1518202113>
- Whitehurst, L. N., Subramoniam, A., Krystal, A., & Prather, A. A. (2022). Links between the brain and body during sleep: Implications for memory processing. *Trends in Neurosciences*, 45(3), 212–223. <https://doi.org/10.1016/j.tins.2021.12.007>
- Wicking, M., Steiger, F., Nees, F., Diener, S. J., Grimm, O., Ruttorf, M., ... Flor, H. (2016). Deficient fear extinction memory in posttraumatic stress disorder. *Neurobiology of Learning and Memory*, 136, 116–126. <https://doi.org/10.1016/j.nlm.2016.09.016>
- Yetton, B. D., Niknazar, M., Duggan, K. A., McDevitt, E. A., Whitehurst, L. N., Sattari, N., & Mednick, S. C. (2016). Automatic detection of rapid eye movements (REMs): A machine learning approach. *Journal of Neuroscience Methods*, 259, 72–82. <https://doi.org/10.1016/j.jneumeth.2015.11.015>
- Zeidan, M. A., Igoe, S. A., Linnman, C., Vitalo, A., Levine, J. B., Klibanski, A., ... Milad, M. R. (2011). Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biological Psychiatry*, 70(10), 920–927. <https://doi.org/10.1016/j.biopsych.2011.05.016>
- Zhang, Y., Ren, R., Sanford, L. D., Yang, L., Zhou, J., Zhang, J., ... Tang, X. (2019). Sleep in posttraumatic stress disorder: A systematic review and meta-analysis of polysomnographic findings. *Sleep Medicine Reviews*, 48, 101210. <https://doi.org/10.1016/j.smr.2019.08.004>
- Zuj, D. V., Palmer, M. A., Lommen, M. J., & Felmingham, K. L. (2016). The centrality of fear extinction in linking risk factors to PTSD: A narrative review. *Neuroscience & Biobehavioral Reviews*, 69, 15–35. <https://doi.org/10.1016/j.neubiorev.2016.07.014>