



On smooth and peaked travelling waves in a local model for shallow water waves

Spencer Locke¹ and Dmitry E. Pelinovsky^{2,3,†}

¹Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA

²Department of Mathematics and Statistics, McMaster University, Hamilton, ON, Canada L8S 4K1

³Department of Applied Mathematics, Nizhny Novgorod State Technical University, 24 Minin street, 603950 Nizhny Novgorod, Russia

(Received 25 May 2024; revised 12 November 2024; accepted 20 November 2024)

We introduce a new model equation for Stokes gravity waves based on conformal transformations of Euler's equations. The local version of the model equation is relevant for the dynamics of shallow water waves. It allows us to characterize the travelling periodic waves both in the case of smooth and peaked waves and to solve the existence problem exactly, albeit not in elementary functions. Spectral stability of smooth waves with respect to co-periodic perturbations is proven analytically based on the exact count of eigenvalues in a constrained spectral problem.

Key words: Hamiltonian theory, surface gravity waves, wave breaking

1. Introduction

An irrotational motion of incompressible two-dimensional surface water waves can be fully described by means of evolution equations for two canonical variables in one spatial coordinate. This formalism was originated by Zakharov (1968) in the context of the stability of travelling periodic waves.

One approach to developing this formalism systematically is based on the Dirichlet-to-Neumann (D-N) operator (Craig & Sulem 1993). The two nonlinear evolution equations closed with the D-N operator have been studied in many works on water waves, including the recent study of modulational instability of travelling periodic waves (Berti, Maspero & Ventura 2022). See also Creedon & Deconinck (2023), Hur & Yang (2023) and Nguyen & Strauss (2023) for other recent works where three more methods have been explored in the same context.

† Email address for correspondence: pelinod@mcmaster.ca

Another approach to obtaining a closed system of two nonlinear evolution equations for water waves is based on a conformal transformation which maps the fluid domain with a variable surface profile to a fixed rectangular domain. This formalism was introduced in Babenko (1987) and Tanveer (1991) and has been explored in the context of travelling periodic waves in Dyachenko *et al.* (1996), Zakharov & Dyachenko (1996), Choi & Camassa (1999) and more recently in Dyachenko, Lushnikov & Korotkevich (2016), Dyachenko & Semenova (2023a,b), Korotkevich *et al.* (2023) and Lushnikov, Dyachenko & Silantyev (2017). Our work contributes to the analysis of the nonlinear evolution equations obtained in the latter approach.

The approach based on conformal transformations has been used to tackle many mathematical problems related to water waves such as the existence of standing waves (Wilkening 2020, 2021) and bifurcations of quasi-periodic wave solutions from the standing periodic waves (Wilkening & Zhao 2021, 2023a,b). Holomorphic coordinates were used for analysis of the well posedness of the water wave equations (Hunter, Ifrim & Tataru 2016; Harrop–Griffiths, Ifrim & Tataru 2017). The particular problem addressed in our work is the coexistence of smooth and peaked travelling periodic waves for different intervals of wave speeds as well as the linear stability of waves with smooth profiles.

1.1. A new model equation

The purpose of this paper is to introduce a new model equation which shares the same solutions as the travelling wave reduction of Euler’s equations in Babenko (1987) but simplifies the time evolution and, particularly, the linear stability analysis of the travelling periodic waves. This model equation can be written in the following non-local form:

$$2cT_h^{-1}\partial_t\eta = (c^2K_h - 1)\eta - \eta K_h\eta - \frac{1}{2}K_h\eta^2, \tag{1.1}$$

where $\eta = \eta(u, t) \in \mathbb{R}$ is the surface elevation in the reference frame moving with the constant wave speed $c > 0$, $t \in \mathbb{R}$ is time and u is the spatial coordinate defined on the periodic domain $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$. The spatial coordinate u arises after the conformal transformation of the fluid domain with variable surface elevation η to a rectangle $[-\pi, \pi] \times [-h, 0]$, where $h > 0$ is the fluid depth. The linear skew-adjoint operator T_h^{-1} in $L^2(\mathbb{T})$ is defined by the Fourier symbol

$$\widehat{(T_h^{-1})}_n = \begin{cases} -i \coth(hn), & n \in \mathbb{Z} \setminus \{0\}, \\ 0, & n = 0, \end{cases} \tag{1.2}$$

whereas the linear, self-adjoint, positive operator $K_h = T_h^{-1}\partial_u$ in $L^2(\mathbb{T})$ is defined by the Fourier symbol

$$\widehat{(K_h)}_n = \begin{cases} n \coth(hn), & n \in \mathbb{Z} \setminus \{0\}, \\ 0, & n = 0. \end{cases} \tag{1.3}$$

Appendix A explains how the non-local evolution equation (1.1) arises in the context of the original Euler’s equations.

Let us obtain the conserved quantities for the non-local model (1.1). Taking the mean value of (1.1), we get the constraint

$$\oint \eta(1 + K_h\eta) du = 0, \tag{1.4}$$

which represents the zero-mean constraint for the surface elevation η in the physical spatial coordinate. Furthermore, differentiating (1.1) in u , multiplying by η and integrating over

the period of \mathbb{T} yields

$$\begin{aligned} c \frac{d}{dt} \oint \eta K_h \eta \, du &= \oint (c^2 \eta K_h \eta_u - \eta \eta_u - \eta \eta_u K_h \eta - \eta^2 K_h \eta_u - \eta K_h \eta \eta_u) \, du \\ &= \oint \partial_u \left(\frac{1}{2} c^2 \eta K_h \eta - \frac{1}{2} \eta^2 - \eta^2 K_h \eta \right) \, du = 0, \end{aligned} \tag{1.5}$$

where we have used self-adjointness of K_h in $L^2(\mathbb{T})$ for every solution with $\eta, \eta_u, \eta \eta_u$ in the domain of K_h . It follows from (1.4) and (1.5) that the non-local evolution equation (1.1) admits two conserved quantities

$$\oint \eta \, du \quad \text{and} \quad \oint \eta K_h \eta \, du. \tag{1.6a,b}$$

In addition, the evolution equation (1.1) can be written in the Hamiltonian form

$$2cT_h^{-1} \partial_t \eta = \Lambda'_c(\eta), \quad \text{with } \Lambda_c(\eta) := \frac{1}{2} \oint [c^2 \eta K_h \eta - \eta^2 (1 + K_h \eta)] \, du, \tag{1.7}$$

where $\Lambda_c(\eta)$ is the action related to the conserved energy of the fluid. Critical points of Λ_c in the corresponding energy space satisfy the Euler–Lagrange equation

$$(c^2 K_h - 1)\eta = \frac{1}{2} K_h \eta^2 + \eta K_h \eta, \tag{1.8}$$

which is known as Babenko’s equation because it coincides with the travelling wave reduction of Euler’s equations after the conformal transformation (Babenko 1987). In the context of the evolution equation (1.1) with u defined in the reference frame moving with the wave speed c , solutions of (1.8) correspond to the time-independent solutions of (1.1).

1.2. Local model and main results

In the deep water limit ($h \rightarrow \infty$), we have from (1.2) and (1.3) that

$$\lim_{h \rightarrow \infty} T_h^{-1} = -\mathcal{H} \quad \text{and} \quad \lim_{h \rightarrow \infty} K_h = -\mathcal{H} \partial_u, \tag{1.9a,b}$$

where \mathcal{H} is the periodic Hilbert transform defined by the Fourier symbol

$$\hat{\mathcal{H}}_n = i \operatorname{sgn}(n), \quad n \in \mathbb{Z}. \tag{1.10}$$

This work explores the shallow water limit ($h \rightarrow 0$), when we replace T_h^{-1} and K_h by $-\partial_u$ and $-\partial_u^2$, respectively. In other words, we study herein the local evolution equation

$$2c \partial_u \partial_t \eta = (c^2 - 2\eta) \partial_u^2 \eta - (\partial_u \eta)^2 + \eta. \tag{1.11}$$

Appendix B describes how the local model (1.11) arises from T_h^{-1} and K_h as $h \rightarrow 0$ and compares it with other phenomenological models for fluid dynamics.

It is important to emphasize that (1.11) is not the asymptotic reduction of (1.1) as $h \rightarrow 0$ but rather a toy model to understand the existence and linear stability of travelling periodic waves in the shallow water limit.

The local equation (1.11) without the last term was derived in Hunter & Saxton (1991) in a different (geometric) context and has been referred to as the Hunter–Saxton equation (Hunter & Zheng 1994). The same equation (1.11) with the last term was also discussed in Alber *et al.* (1995, 1999) in connection to the high-frequency limit of the Camassa–Holm

equation, one of the toy models for the physics of fluids with smooth and peaked waves. Integrability of (1.11) was established in Hone, Novikov & Wang (2018) together with other peaked wave equations such as the reduced Ostrovsky and short-pulse equations. Some travelling wave solutions of this and similar equations were studied with Hirota's bilinear method in Matsuno (2020).

Next, we discuss the time evolution and the conserved quantities for the local model (1.11). Taking the mean value of (1.11) for smooth 2π -periodic solutions and integrating by parts yields the constraint

$$\oint [\eta + (\partial_u \eta)^2] du = 0, \tag{1.12}$$

which corresponds to (1.4) also after integration by parts. Let $\Pi_0 : L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})|_{\{1\}^T}$ be a projection operator to the periodic functions with zero mean. The evolution equation (1.11) can be written in the form

$$2c\partial_t \eta = (c^2 - 2\eta)\partial_u \eta + \Pi_0 \partial_u^{-1} \Pi_0 [(\partial_u \eta)^2 + \eta], \tag{1.13}$$

where $\Pi_0 \partial_u^{-1} \Pi_0$ is uniquely defined on the zero-mean functions in $L^2(\mathbb{T})$ with the zero-mean constraint. The evolution equation (1.13) is a non-local version of the inviscid Burgers equation. The initial-value problem for the inviscid Burgers equation is locally well posed in $H^1_{per}(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$. Since the mapping

$$\Pi_0 \partial_u^{-1} \Pi_0 [(\partial_u \eta)^2 + \eta] : H^1_{per}(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T}) \rightarrow H^1_{per}(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T}) \tag{1.14}$$

is bounded on every bounded subset, there exists a unique local solution of the evolution equation (1.13) for every initial data in $H^1_{per}(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$.

To get the conserved quantities, we multiply (1.11) by $\partial_u \eta$ and integrate over the period for smooth 2π -periodic solutions η . This implies the conservation of

$$Q(\eta) := \oint (\partial_u \eta)^2 du, \tag{1.15}$$

and, in view of the constraint (1.12), the conservation of

$$M(\eta) := \oint \eta du. \tag{1.16}$$

The conserved quantities (1.15) and (1.16) correspond to (1.6a,b). Furthermore, similar to (1.7), we can write (1.13) in the Hamiltonian form

$$2c\partial_t \eta = -\Pi_0 \partial_u^{-1} \Pi_0 \left[\frac{1}{2} c^2 Q'(\eta) - \frac{1}{2} H'(\eta) \right], \tag{1.17}$$

where H is the third conserved quantity given by

$$H(\eta) := \oint [\eta^2 + 2\eta(\partial_u \eta)^2] du. \tag{1.18}$$

The existence of travelling periodic waves in the local model (1.11) is defined by the second-order equation

$$(c^2 - 2\eta)\eta'' - (\eta')^2 + \eta = 0, \quad u \in \mathbb{T}, \tag{1.19}$$

where $\eta = \eta(u)$ is the 2π -periodic wave profile satisfying the constraint (1.12). The linear stability of the travelling wave with the profile η is defined by the linearized equation

$$2c\partial_t \hat{\eta} = -\Pi_0 \partial_u^{-1} \Pi_0 \mathcal{L} \hat{\eta}, \quad \mathcal{L} := -\partial_u (c^2 - 2\eta)\partial_u - 1 + 2\eta'', \tag{1.20a,b}$$

where $\hat{\eta} = \hat{\eta}(u, t)$ is the perturbation to the travelling wave with the profile $\eta = \eta(u)$ satisfying the orthogonality condition $\langle 1 - 2\eta'', \hat{\eta} \rangle = 0$ with the standard inner product in




Figure 1. Profiles in the family of smooth periodic waves (a) and singular periodic waves (b) vs u for different values of wave speeds $c \in (1, c_*)$ and $c \in (c_*, c_\infty)$, respectively. The dashed line shows the profile of the peaked periodic wave (1.23).

$L^2(\mathbb{T})$. The orthogonality condition $\langle 1 - 2\eta'', \hat{\eta} \rangle = 0$ follows by expanding the nonlinear constraint (1.12) near η .

The following two theorems describe the main results of this work. Our results are formulated for the single-lobe periodic solutions with an even profile η which possesses a single maximum on \mathbb{T} placed at $u = 0$. Such single-lobe periodic solutions are often referred to as Stokes waves.

THEOREM 1.1. *There exist $c_* := \pi/(2\sqrt{2})$ and $c_\infty \in (c_*, \infty)$ such that the stationary equation (1.19) admits a unique single-lobe solution with the profile $\eta \in C_{per}^\infty(\mathbb{T})$ for every $c \in (1, c_*)$ such that*

$$\|\eta\|_{L^\infty} \rightarrow 0 \quad \text{as } c \rightarrow 1, \tag{1.21}$$

and a single-lobe solution with the profile $\eta \in C_{per}^0(\mathbb{T})$ for every $c \in (c_, c_\infty)$ satisfying*

$$\eta(u) = \frac{c^2}{2} - A(c)|u|^{2/3} + O(|u|^{4/3}) \quad \text{as } u \rightarrow 0, \tag{1.22}$$

for some constant $A(c) > 0$. At $c = c_$, there exists a unique single-lobe solution with the profile $\eta \in C_{per}^0(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$ given explicitly as*

$$\eta(u) = \frac{1}{16}(\pi^2 - 4\pi|u| + 2u^2), \quad u \in [-\pi, \pi], \tag{1.23}$$

and extended as a 2π -periodic function on \mathbb{T} .

REMARK 1.1. *Figure 1 shows profiles of the periodic waves of Theorem 1.1. The profiles were obtained numerically by using solutions of the second-order equation (1.19).*

REMARK 1.2. *There exists another single-lobe solution with the singular behaviour (1.22) for every $c \in (0, c_\infty)$ which is not included in the statement of Theorem 1.1 as it does not bifurcate from the zero solution as $c \rightarrow 1$ compared with (1.21). See figure 2 for the bifurcation diagram of all single-lobe solutions of (1.19).*

REMARK 1.3. *The special solution (1.23) has a peaked profile with a finite jump of the first derivative. It is usually referred to as the peaked periodic wave. Such peaked periodic waves are commonly known in other fluid models such as the reduced Ostrovsky equation (Geyer & Pelinovsky 2019, 2020; Bruell & Dhara 2021), the Camassa–Holm equation (Madiyeva & Pelinovsky 2021) and the Degasperis–Procesi equation (Geyer & Pelinovsky 2024).*




Figure 2. Dependence of $E := \|\eta\|_{L^\infty}$ vs wave speed c for periodic solutions of (1.19). The smooth solutions of Theorem 1.1 exist between the black dots. The singular solutions of Theorem 1.1 exist between the rightmost black and red dots. The singular solutions which are not included in the statement of Theorem 1.1 exist between the red dots.

REMARK 1.4. The singular behaviour (1.22) corresponds to the singularity of the limiting Stokes wave with a 120° angle in the physical coordinate after the conformal transformation. The behaviour was rigorously proven for the original Euler’s equation in Amick, Fraenkel & Toland (1982), Plotnikov (2002) and Toland (1978), with many asymptotic results known in the literature (see Lushnikov (2016) and references therein). Note that

$$\max_{u \in \mathbb{T}} \eta(u) = \eta(0) = \frac{c^2}{2} \tag{1.24}$$

holds for every limiting Stokes wave for which the horizontal velocity at the wave height coincides with the wave speed c , see the second equation in system (A3) of Appendix A.

THEOREM 1.2. Consider the unique single-lobe solution with the profile $\eta \in C^\infty_{per}(\mathbb{T})$ in Theorem 1.1 for $c \in (1, c_*)$. For every initial data $\hat{\eta}_0 \in H^1_{per}(\mathbb{T})$ satisfying

$$\langle 1, \hat{\eta}_0 \rangle = 0 \quad \text{and} \quad \langle \eta'', \hat{\eta}_0 \rangle = 0, \tag{1.25a,b}$$

there exists a unique solution $\hat{\eta} \in C^0(\mathbb{R}, H^1_{per}(\mathbb{T}))$ of the linearized equation (1.20a,b) with $\hat{\eta}|_{t=0} = \hat{\eta}_0$ and a unique $a \in C^1(\mathbb{R}, \mathbb{R})$ such that

$$\|\hat{\eta}(\cdot, t) - a(t)\eta'\|_{H^1_{per}} \leq C\|\hat{\eta}_0\|_{H^1_{per}}, \quad |a'(t)| \leq C\|\hat{\eta}_0\|_{H^1_{per}}, \quad t \in \mathbb{R}, \tag{1.26a,b}$$

where $C > 0$ is independent of $\hat{\eta}_0$.

REMARK 1.5. Constraints (1.25a,b) are preserved in the time evolution of the linearized equation (1.20a,b) because they are linearizations of the conserved quantities (1.15) and (1.16). In view of the constraint $\langle 1 - 2\eta'', \hat{\eta} \rangle = 0$ imposed on solutions of the linearized equation (1.20a,b), only one constraint in (1.25a,b) is linearly independent. Imposing $\langle \eta'', \hat{\eta}_0 \rangle = 0$ is equivalent to the requirement that the perturbation $\hat{\eta}$ does not change

the conserved quantity Q in (1.15) up to the linear approximation. The bound (1.26a,b) expresses the concept of linear orbital stability of the orbit $\{\eta(\cdot + \mathbf{u})\}_{\mathbf{u} \in \mathbb{T}}$ of the travelling periodic wave with the profile η .

REMARK 1.6. The linear orbital stability of Theorem 1.2 implies spectral stability of the travelling periodic wave in the sense that the spectrum of the associated linearized operator $\partial_u^{-1} \mathcal{L}$ in $L^2(\mathbb{T})$ belongs to $i\mathbb{R}$. It is also worthwhile to point out that an eigenfunction $\hat{\eta}_0$ of the spectral stability problem

$$\partial_u^{-1} \mathcal{L} \hat{\eta}_0 = \lambda_0 \hat{\eta}_0, \quad \eta_0 \in H_{per}^1(\mathbb{T}) \tag{1.27}$$

for every non-zero eigenvalue $\lambda_0 \in \mathbb{C} \setminus \{0\}$ must satisfy the two constraints in (1.25a,b). The spectral stability problem in the form (1.27) was also considered in Stanislavova & Stefanov (2016).

REMARK 1.7. The proof of Theorem 1.2 relies on the construction of the coercive quadratic form $\langle \mathcal{L} \hat{\eta}, \hat{\eta} \rangle$ for $\hat{\eta} \in H_{per}^1(\mathbb{T})$ under the three constraints

$$\langle 1, \hat{\eta} \rangle = \langle \eta', \hat{\eta} \rangle = \langle \eta'', \hat{\eta} \rangle = 0. \tag{1.28}$$

The quadratic form is invariant in the time evolution of the linearized equation (1.20a,b). This yields the energetic stability of the travelling periodic wave, ensuring that the periodic wave with the profile η is a local minimizer of the energy H subject to fixed Q and M in $H_{per}^1(\mathbb{T})$. If local well posedness of the nonlinear evolution equation (1.13) can be shown in $H_{per}^1(\mathbb{T})$, then the energetic stability implies the nonlinear orbital stability of the travelling periodic wave. However, the local well posedness of (1.13) holds only in $H_{per}^1(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$ and the control of $\|\partial_u \hat{\eta}\|_{L^\infty}$ for the perturbation $\hat{\eta}$ does not follow from the conserved quantities (1.15), (1.16) and (1.18).

1.3. Discussion

The local model (1.11) shows a pattern of the existence and stability of travelling periodic waves parameterized by the wave speed c . There is a continuum of wave speeds for the smooth waves with profile $\eta \in C^\infty(\mathbb{T})$ bifurcating from the linear limit in (1.21) and a continuum of wave speeds for the cusped waves with the profile $\eta \in C^0(\mathbb{T})$ satisfying (1.22). The two continuous families are connected together at a particular value of the wave speed $c = c_*$ for which the wave is peaked with the profile $\eta \in C^0(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$. The same phenomenon is observed in the Camassa–Holm equation (Lenells 2005b; Geyer *et al.* 2022) and the Degasperis–Procesi equation (Lenells 2005a; Geyer & Pelinovsky 2024).

It is rather remarkable that exactly the same $|u|^{2/3}$ singularity in the limiting wave profile with

$$\max_{u \in \mathbb{T}} \eta(u) = \eta(0) = \frac{c^2}{2} \tag{1.29}$$

is recovered by the local model (1.11) as predicted by the full model for any depth h (Plotnikov 2002). After the conformal transformation, this singularity yields the limiting Stokes wave with the 120° angle in the physical coordinates. More precise details of such singular profiles are beyond the current capacities of the asymptotic (Lushnikov 2016) or numerical (Dyachenko *et al.* 2016; Lushnikov *et al.* 2017) methods. The local model (1.11) gives precise conclusions that the $|u|^{2/3}$ singularity is obtained in a range of wave speeds c

and that the borderline wave profile $\eta \in H_{per}^1(\mathbb{T}) \cap W^{1,\infty}(\mathbb{T})$ has a peaked profile at $c = c_*$ before getting the $|u|^{2/3}$ singularity for $c > c_*$. This might be an artefact of the local model (1.11) since the non-local models typically predict only the $|u|^{2/3}$ singularity in the fluid models, see Locke & Pelinovsky (2025).

Stability of the travelling periodic waves with singular profiles is a complicated problem, which is out of reach in the current analytical and numerical methods in the non-local models (Dyachenko & Semanova 2023a; Korotkevich *et al.* 2023). The local model (1.11) is a promising candidate for showing linear instability of the peaked wave (based on a similar analysis in Geyer & Pelinovsky 2019, 2020; Madiyeva & Pelinovsky 2021) and for attacking linear instability of the cusped wave with the $|u|^{2/3}$ singularity.

The remainder of this paper is organized as follows. Section 2 contains the proof of Theorem 1.1 on the existence of smooth and peaked periodic waves. Section 3 gives the proof of Theorem 1.2 on linear stability of the smooth periodic waves. Appendix A reviews the Euler equations after the conformal transformation and discusses how the non-local model (1.1) arises. Appendix B describes the local model (1.11) in the context of other phenomenological models for dynamics of fluid surfaces.

2. Existence of smooth and peaked travelling periodic waves

We consider the single-lobe periodic solutions of the second-order equation (1.19). Recall that a single-lobe periodic solution has the even profile η with a single maximum on \mathbb{T} placed at $u = 0$. Theorem 1.1 is proven by using the period function for the planar Hamiltonian systems used in a similar context in Geyer *et al.* (2022), Geyer & Pelinovsky (2017, 2024) and Long & Liu (2023).

We start with the first-order invariant of the second-order equation (1.19) given by the following lemma.

LEMMA 2.1. *For every solution $\eta \in C^2(a, b)$ of the second-order equation (1.19) with $-\infty \leq a < b \leq \infty$, the following function:*

$$E(\eta, \eta') := \frac{1}{2}(c^2 - 2\eta)(\eta')^2 + \frac{1}{2}\eta^2 \tag{2.1}$$

is constant for $u \in (a, b)$.

Proof. It is based on the elementary computation

$$\begin{aligned} \frac{d}{du}E(\eta, \eta') &= (c^2 - 2\eta)\eta'\eta'' - (\eta')^3 + \eta\eta' \\ &= \eta'[(c^2 - 2\eta)\eta'' - (\eta')^2 + \eta] \\ &= 0, \end{aligned} \tag{2.2}$$

since $\eta \in C^2(a, b)$ satisfies (1.19).

The next lemma explores the phase portrait for the second-order equation (1.19) on the phase plane (η, η') obtained from the level curves of $E(\eta, \eta')$ in (2.1). We obtain the existence of smooth and singular solutions in terms of the level \mathcal{E} of $E(\eta, \eta')$.




Figure 3. Phase portrait from the level curves of $E(\eta, \eta') = \mathcal{E}$ for $c = 1$.

LEMMA 2.2. For every $c > 0$, there exists $\mathcal{E}_c := c^4/8$ such that every periodic solution to (1.19) with profile $\eta \in C^\infty(\mathbb{R})$ belongs to $E(\eta, \eta') = \mathcal{E}$ with $\mathcal{E} \in (0, \mathcal{E}_c)$. For $E(\eta, \eta') = \mathcal{E}_c$, the only solution to (1.19) is the parabola

$$\eta(u) = -\frac{c^2}{2} + \frac{(u - u_0)^2}{8}, \tag{2.3}$$

with arbitrary $u_0 \in \mathbb{R}$. For $E(\eta, \eta') = \mathcal{E}$ with $\mathcal{E} \in (\mathcal{E}_c, \infty)$, there exist no bounded solutions to (1.19) with profile $\eta \in C^\infty(\mathbb{R})$.

Proof. The only equilibrium point of (1.19) is the centre $(\eta, \eta') = (0, 0)$, which corresponds to the minimum of $E(\eta, \eta')$ if $c > 0$. Every periodic solution to (1.19) belongs to the period annulus, which is the largest punctured neighbourhood of the centre $(0, 0)$ consisting entirely of periodic orbits.

The phase portrait from the level curves of $E(\eta, \eta') = \mathcal{E}$ is shown on figure 3. Each level curve defines the profile η from integration of

$$\left(\frac{d\eta}{du}\right)^2 = \frac{2\mathcal{E} - \eta^2}{c^2 - 2\eta}. \tag{2.4}$$

The vertical line corresponds to $\eta = c^2/2$ and divides the phase plane into two half-planes. For $\eta > c^2/2$, the level curves of $E(\eta, \eta') = \mathcal{E}$ contain no bounded solutions. For $\eta < c^2/2$, bounded level curves exist for $\mathcal{E} \in (0, \mathcal{E}_c)$ with $\mathcal{E}_c := c^4/8$ and contain periodic solutions with profile $\eta \in C^\infty(\mathbb{R})$. For $\mathcal{E} = \mathcal{E}_c$, all solutions are given by integrating

$$\left(\frac{d\eta}{du}\right)^2 = \frac{2\mathcal{E}_c - \eta^2}{c^2 - 2\eta} = \frac{1}{4}(c^2 + 2\eta). \tag{2.5}$$

Differentiating in u gives $\eta''(u) = \frac{1}{4}$. Integrating twice and using (2.5) yields (2.3) with profile $\eta \in C^\infty(\mathbb{R})$. Finally, for $\mathcal{E} > \mathcal{E}_c$, the level curve reaches $\eta = c^2/2$ with the singularity of η' , which rules out the existence of bounded solutions with profile $\eta \in C^\infty(\mathbb{R})$.

The next lemma clarifies how the solutions for $\eta < c^2/2$ reach the singularity line $\eta = c^2/2$.

LEMMA 2.3. *Let $\eta \in C^0(u_-, u_+)$ be a solution for the level curve $E(\eta, \eta') = \mathcal{E}$ for $\mathcal{E} \in (\mathcal{E}_c, \infty)$ such that $\eta(u) \rightarrow \pm c^2/2$ as $u \rightarrow u_{\pm}$. Then, $-\infty < u_- < u_+ < \infty$ and the solution satisfies*

$$\eta(u) = \frac{c^2}{2} - \frac{3^{2/3}}{2^{2/3}}(\mathcal{E} - \mathcal{E}_c)^{1/3}|u - u_{\pm}|^{2/3} + O(|u - u_{\pm}|^{4/3}) \quad \text{as } u \rightarrow u_{\pm}. \quad (2.6)$$

Proof. We consider the level curve $E(\eta, \eta') = \mathcal{E}$ with $\mathcal{E} \in (\mathcal{E}_c, \infty)$ for $\eta < c^2/2$. Then, we have

$$\begin{aligned} \left(\frac{d\eta}{du}\right)^2 &= \frac{2\mathcal{E} - \eta^2}{c^2 - 2\eta} \\ &= \frac{2(\mathcal{E} - \mathcal{E}_c)}{c^2 - 2\eta} + \frac{1}{2}c^2 + \frac{1}{4}(2\eta - c^2) \\ &= \frac{2(\mathcal{E} - \mathcal{E}_c)}{c^2 - 2\eta} + O(1) \quad \text{as } \eta \rightarrow \frac{c^2}{2}. \end{aligned} \quad (2.7)$$

This yields

$$\sqrt{c^2 - 2\eta}[1 + O(|c^2 - 2\eta|)]\frac{d\eta}{du} = \pm\sqrt{2(\mathcal{E} - \mathcal{E}_c)} \quad \text{as } u \rightarrow u_{\pm} \mp 0. \quad (2.8)$$

Integrating in u yields

$$\sqrt{(c^2 - 2\eta)^3}[1 + O(|c^2 - 2\eta|)] = \mp 3\sqrt{2(\mathcal{E} - \mathcal{E}_c)}(u - u_{\pm}) \quad \text{as } u \rightarrow u_{\pm} \mp 0, \quad (2.9)$$

which results in the expansion (2.6). It remains to prove that $[u_-, u_+]$ is compact. This follows from the bounds

$$\begin{aligned} u_+ - u_- &= 2 \int_{-\sqrt{2\mathcal{E}}}^{c^2/2} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} d\eta \\ &\leq 2\sqrt{c^2 + 2\sqrt{2\mathcal{E}}} \int_{-\sqrt{2\mathcal{E}}}^{c^2/2} \frac{d\eta}{\sqrt{2\mathcal{E} - \eta^2}} < \infty, \end{aligned} \quad (2.10)$$

since the integral in the upper bound is finite.

In order to analyse bounded periodic solutions in Lemma 2.2, we introduce the following period function associated with (2.1) and (2.4):

$$T(\mathcal{E}, c) := 2 \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} d\eta, \quad \mathcal{E} \in (0, \mathcal{E}_c). \quad (2.11)$$

For the singular solutions in Lemma 2.3, we augment the period function for $\mathcal{E} \geq \mathcal{E}_c$ as

$$T(\mathcal{E}, c) := 2 \int_{-\sqrt{2\mathcal{E}}}^{c^2/2} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} d\eta, \quad \mathcal{E} \in [\mathcal{E}_c, \infty). \quad (2.12)$$

The next result describes properties of the period function $(0, \infty) \ni \mathcal{E} \mapsto T(\mathcal{E}, c)$ for every fixed $c > 0$.

LEMMA 2.4. For every $c > 0$, there exist \mathcal{E}_* , $\mathcal{E}_{**} \in (\mathcal{E}_c, \infty)$ that depend on c such that

$$\frac{\partial}{\partial \mathcal{E}} T(\mathcal{E}, c) < 0, \quad \mathcal{E} \in (0, \mathcal{E}_*), \tag{2.13}$$

and

$$\frac{\partial}{\partial \mathcal{E}} T(\mathcal{E}, c) > 0, \quad \mathcal{E} \in (\mathcal{E}_{**}, \infty), \tag{2.14}$$

with

$$\left. \begin{aligned} T(\mathcal{E}, c) &\rightarrow 2\pi c \quad \text{as } \mathcal{E} \rightarrow 0, \\ T(\mathcal{E}, c) &\rightarrow 4\sqrt{2}c \quad \text{as } \mathcal{E} \rightarrow \mathcal{E}_c, \end{aligned} \right\} \tag{2.15}$$

and $T(\mathcal{E}, c) \rightarrow \infty$ as $\mathcal{E} \rightarrow \infty$. In addition, we have

$$\frac{\partial}{\partial c} T(\mathcal{E}, c) > 0, \quad \mathcal{E} \in (0, \infty), \quad c \in (0, \infty). \tag{2.16}$$

Proof. For the smooth periodic solutions, we use the change of variables $\eta = \sqrt{2\mathcal{E}}x$ in (2.11) and obtain

$$T(\mathcal{E}, c) = 2 \int_{-1}^1 \frac{\sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}}{\sqrt{1-x^2}} dx, \quad \mathcal{E} \in (0, \mathcal{E}_c). \tag{2.17}$$

Since the weak singularity is independent of \mathcal{E} , we can differentiate under the integration sign and obtain

$$\frac{\partial}{\partial \mathcal{E}} T(\mathcal{E}, c) = -\frac{2}{\sqrt{2\mathcal{E}}} \int_{-1}^1 \frac{x dx}{\sqrt{1-x^2} \sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}}, \quad \mathcal{E} \in (0, \mathcal{E}_c). \tag{2.18}$$

The result is strictly negative since

$$\frac{|x|}{\sqrt{1-x^2} \sqrt{c^2 + 2\sqrt{2\mathcal{E}}|x|}} < \frac{x}{\sqrt{1-x^2} \sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}}, \quad x \in (0, 1). \tag{2.19}$$

This proves (2.13) for $\mathcal{E} \in (0, \mathcal{E}_c)$. We also obtain from the same representation

$$T(\mathcal{E}, c) \rightarrow 2c \int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} = 2\pi c \quad \text{as } \mathcal{E} \rightarrow 0, \tag{2.20}$$

and

$$T(\mathcal{E}, c) \rightarrow 2c \int_{-1}^1 \frac{\sqrt{1-x}}{\sqrt{1-x^2}} dx = 4\sqrt{2}c \quad \text{as } \mathcal{E} \rightarrow \mathcal{E}_c. \tag{2.21}$$

Monotonicity (2.16) follows from the positive derivative of (2.11) in c .

For the singular solutions, we break (2.12) into the sum of two terms and use the same change of variables only in the first term

$$T(\mathcal{E}, c) = 2 \int_{-1}^{-c^2/2\sqrt{2\mathcal{E}}} \frac{\sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}}{\sqrt{1 - x^2}} dx + 2 \int_{-c^2/2}^{c^2/2} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} d\eta, \quad \mathcal{E} \in [\mathcal{E}_c, \infty). \tag{2.22}$$

Since $8\mathcal{E} > c^4$, both terms are differentiable under the integration sign and we obtain

$$\frac{\partial}{\partial \mathcal{E}} T(\mathcal{E}, c) = -\frac{2}{\sqrt{2\mathcal{E}}} \int_{-1}^{-c^2/2\sqrt{2\mathcal{E}}} \frac{x dx}{\sqrt{1 - x^2}\sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}} - 2 \int_{-c^2/2}^{c^2/2} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{(2\mathcal{E} - \eta^2)^3}} d\eta, \tag{2.23}$$

where the first term is positive and the second term is negative. The first term is zero as $\mathcal{E} \rightarrow \mathcal{E}_c$, monotonically increasing for $\mathcal{E} \gtrsim \mathcal{E}_c$ and monotonically decreasing as $\mathcal{E} \rightarrow \infty$. The second term is strictly negative as $\mathcal{E} \rightarrow \mathcal{E}_c$ and is monotonically increasing towards 0 for $\mathcal{E} > \mathcal{E}_c$. Since

$$\int_{-1}^{-c^2/2\sqrt{2\mathcal{E}}} \frac{\sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}}{\sqrt{1 - x^2}} dx \sim (8\mathcal{E})^{1/4} \int_{-1}^0 \frac{\sqrt{|x|} dx}{\sqrt{1 - x^2}} \quad \text{as } \mathcal{E} \rightarrow \infty, \tag{2.24}$$

and

$$\int_{-c^2/2}^{c^2/2} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} d\eta \sim (2\mathcal{E})^{-1/2} \int_{-c^2/2}^{c^2/2} \sqrt{c^2 - 2\eta} d\eta \quad \text{as } \mathcal{E} \rightarrow \infty, \tag{2.25}$$

the first term in the decomposition (2.23) is larger than the second term at infinity and we have $T(\mathcal{E}, c) = O(\mathcal{E}^{1/4})$ as $\mathcal{E} \rightarrow \infty$ so that

$$T(\mathcal{E}, c) \rightarrow \infty \quad \text{as } \mathcal{E} \rightarrow \infty. \tag{2.26}$$

At the same time, the first term in the decomposition (2.23) is zero at $\mathcal{E} = \mathcal{E}_c$, hence there exist \mathcal{E}_* , $\mathcal{E}_{**} \in (\mathcal{E}_c, \infty)$ such that (2.13) and (2.14) hold. Monotonicity (2.16) follows from the positive derivative of (2.12) in c .

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. We consider the family of solutions of Lemmas 2.2 and 2.3 for $\mathcal{E} \in (0, \infty)$ and $\eta < c^2/2$. For every $c > 0$, we select the intersection of the period function $T(\mathcal{E}, c)$ of Lemma 2.4 with the 2π period on \mathbb{T} . It follows from (2.16) that the period function is monotonically increasing in c for every $\mathcal{E} \in (0, \infty)$.

For the smooth periodic solutions with the period function (2.11), there exists only one root $\mathcal{E} \in (0, \mathcal{E}_c)$ of $T(\mathcal{E}, c) = 2\pi$ for every $c \in (1, c_*)$ with $c_* = \pi/(2\sqrt{2})$ due to monotonicity (2.13) and the limiting values $T(0, c) = 2\pi c$ and $T(\mathcal{E}_c, c) = 4\sqrt{2}c$. This gives the first assertion of the theorem with the limit (1.21) since the smooth periodic solution shrinks to the centre point $(0, 0)$ on the (η, η') plane as $\mathcal{E} \rightarrow 0$. At $c = c_*$, we have $\mathcal{E} = \mathcal{E}_{c_*}$ for the root of $T(\mathcal{E}, c) = 2\pi$. The unique single-lobe solution (1.23) follows from the unique solution (2.3) at $\mathcal{E} = \mathcal{E}_c$ from Lemma 2.2 by the translation $u_0 = \pi$ for $u \in [0, \pi]$ and an even reflection on $[-\pi, 0]$.

For the singular solutions with the period function (2.12), we have a root $\mathcal{E} \in (\mathcal{E}_c, \infty)$ of $T(\mathcal{E}, c) = 2\pi$ for every $c \in (c_*, c_\infty)$ due to monotonicity (2.13). The value c_∞ is obtained




Figure 4. Period function T vs \mathcal{E} for $c = 1.05$ (solid blue) and $c = 1.1$ (dashed blue). Black dots show the values $T(0, c) = 2\pi c$ and $T(\mathcal{E}_c, c) = 4\sqrt{2}c$. The red dashed line gives the level $T(\mathcal{E}, c) = 2\pi$ for periodic solutions on \mathbb{T} .

from the intersection of $T(\mathcal{E}_*, c)$ with 2π . The asymptotic expansion (1.22) of the singular single-lobe solutions follows from the expansion (2.6) with the translation $u_- = 0$ of the solution in Lemma 2.3 for $u \in [0, \pi]$ with $\eta(\pi) = -\sqrt{2\mathcal{E}}$ and $\eta'(\pi) = 0$ and an even reflection on $[-\pi, 0]$.

Figure 4 illustrates the result of Lemma 2.4 and the proof of Theorem 1.1. The period functions (2.11) and (2.12) can be computed in terms of complete elliptic integrals by using 3.141 (integrals 2 and 9) in Gradshteyn & Ryzhik (2007)

$$T(\mathcal{E}, c) = 4\sqrt{c^2 + 2\sqrt{2\mathcal{E}}} \left(\sqrt{\frac{4\sqrt{2\mathcal{E}}}{c^2 + 2\sqrt{2\mathcal{E}}}} \right), \quad \mathcal{E} \in (0, \mathcal{E}_c), \quad (2.27)$$

and

$$T(\mathcal{E}, c) = 4(2\mathcal{E})^{1/4} \left[2E \left(\sqrt{\frac{c^2 + 2\sqrt{2\mathcal{E}}}{4\sqrt{2\mathcal{E}}}} \right) + \left(\frac{c^2}{2\sqrt{2\mathcal{E}}} - 1 \right) K \left(\sqrt{\frac{c^2 + 2\sqrt{2\mathcal{E}}}{4\sqrt{2\mathcal{E}}}} \right) \right], \quad \mathcal{E} \in (\mathcal{E}_c, \infty), \quad (2.28)$$

where $K(k)$ and $E(k)$ are complete elliptic integrals of the first and second kind, respectively. The two definitions agree at $T(\mathcal{E}_c, c) = 4\sqrt{2}c$, where $\mathcal{E}_c = c^4/8$ shown by the black dot in figure 4. We can also see from figure 4 that the monotonicity results (2.13), (2.14) and (2.16) hold true with $\mathcal{E}_* = \mathcal{E}_{**}$. The periodic solutions of Theorem 1.1 on \mathbb{T} are obtained by the intersection of the plot of the period function with the level $T(\mathcal{E}, c) = 2\pi$.

3. Stability of smooth travelling periodic waves

We consider the unique single-lobe solution with the profile $\eta \in C_{per}^\infty(\mathbb{T})$ in Theorem 1.1 which exists in the second-order equation (1.19) for $c \in (1, c_*)$ with $c_* = \pi/(2\sqrt{2})$. Theorem 1.2 is proven by showing that the periodic solution is a constrained minimizer of the quadratic form $\langle \mathcal{L}\hat{\eta}, \hat{\eta} \rangle$ associated with the linear operator

$$\mathcal{L} = -\partial_u(c^2 - 2\eta)\partial_u - 1 + 2\eta'', \tag{3.1}$$

in the function space $H_{per}^1(\mathbb{T}) \cap \mathcal{X}_c$, where

$$\mathcal{X}_c := \{\hat{\eta} \in L^2(\mathbb{T}) : \langle 1, \hat{\eta} \rangle = \langle \eta', \hat{\eta} \rangle = 0\}. \tag{3.2}$$

The minimizer is only degenerate due to the translational symmetry which results in the one-dimensional kernel $\text{Ker}(\mathcal{L}) = \text{span}(\eta')$ with $\eta' \in \mathcal{X}_c$.

We start with the count of the negative eigenvalues of $\mathcal{L} : H_{per}^2(\mathbb{T}) \subset L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})$ in the following lemma.

LEMMA 3.1. *Let $\eta \in C_{per}^\infty(\mathbb{T})$ be the profile of the single-lobe solution in Theorem 1.1 for $c \in (1, c_*)$ and $\mathcal{L} : H_{per}^2(\mathbb{T}) \subset L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})$ be the linear operator given by (3.1). Then, \mathcal{L} has two simple negative eigenvalues and a simple zero eigenvalue, with the rest of its spectrum bounded away from zero.*

Proof. Since $\mathcal{L} : H_{per}^2(\mathbb{T}) \subset L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})$ is a self-adjoint Sturm–Liouville operator with $\eta'' \in L^\infty(\mathbb{T})$ and $c^2 - 2\eta(u) > 0$ for every $u \in \mathbb{T}$, its spectrum consists of isolated eigenvalues located on the real line.

For fixed $c \in (1, c_*)$, differentiating (1.19) in u yields $\mathcal{L}\eta' = 0$ with $\eta' \in H_{per}^2(\mathbb{T})$. Hence, \mathcal{L} admits a zero eigenvalue with the spatially odd eigenfunction η' . To consider the second linearly independent solution of $\mathcal{L}\hat{\eta} = 0$, we define the family $\{\eta(u; \mathcal{E})\}_{\mathcal{E} \in (0, \mathcal{E}_c)}$ of spatially even, smooth periodic solutions of the second-order equation (1.19) with

$$\eta(0; \mathcal{E}) = \sqrt{2\mathcal{E}}, \quad \partial_u \eta(0; \mathcal{E}) = 0, \tag{3.3a,b}$$

and

$$\eta(T(\mathcal{E}, c); \mathcal{E}) = \sqrt{2\mathcal{E}}, \quad \partial_u \eta(T(\mathcal{E}, c); \mathcal{E}) = 0, \tag{3.4a,b}$$

where $T(\mathcal{E}, c)$ is the period function (2.11) satisfying (2.13). Let $\mathcal{E}(c)$ be the root of the period function $T(\mathcal{E}, c) = 2\pi$ in the proof of Theorem 1.1 such that $\eta(u) = \eta(u; \mathcal{E}(c)) \in C_{per}^\infty(\mathbb{T})$. Differentiating (1.19) in \mathcal{E} along the family $\{\eta(u; \mathcal{E})\}_{\mathcal{E} \in (0, \mathcal{E}_c)}$ and setting $\mathcal{E} = \mathcal{E}(c)$ yields the second linearly independent solution of $\mathcal{L}\hat{\eta} = 0$. The solution is given by the spatially even function $\partial_{\mathcal{E}}\eta(\cdot; \mathcal{E}(c))$ which satisfies

$$\partial_{\mathcal{E}}\eta(0; \mathcal{E}) = \frac{1}{\sqrt{2\mathcal{E}}}, \quad \partial_u \partial_{\mathcal{E}}\eta(0; \mathcal{E}) = 0, \tag{3.5a,b}$$

and

$$\partial_{\mathcal{E}}\eta(T(\mathcal{E}, c); \mathcal{E}) = \frac{1}{\sqrt{2\mathcal{E}}}, \quad \partial_u \partial_{\mathcal{E}}\eta(T(\mathcal{E}, c); \mathcal{E}) = \partial_{\mathcal{E}}T(\mathcal{E}, c) \frac{\sqrt{2\mathcal{E}}}{c^2 - 2\sqrt{2\mathcal{E}}}, \tag{3.6a,b}$$

where we have used (1.19) at $u = T(\mathcal{E}, c)$ which implies

$$\partial_u^2 \eta(T(\mathcal{E}, c); \mathcal{E}) = -\frac{\sqrt{2\mathcal{E}}}{c^2 - 2\sqrt{2\mathcal{E}}}. \tag{3.7}$$

We thus have $\partial_{\mathcal{E}}\eta(\cdot; \mathcal{E}(c)) \in H_{per}^2(\mathbb{T})$ if and only if $\partial_{\mathcal{E}}T(\mathcal{E}(c), c) = 0$, which is impossible due to monotonicity (2.13). Hence, 0 is a simple eigenvalue of \mathcal{L} bounded away from the rest of its spectrum in $L^2(\mathbb{T})$.

To prove that there exist two negative eigenvalues of \mathcal{L} below the zero eigenvalue, we use Proposition 1 in Geyer *et al.* (2022) and construct the following two normalized solutions:

$$\eta_1(u) := \sqrt{2\mathcal{E}}\partial_{\mathcal{E}}\eta(u; \mathcal{E}) \quad \text{and} \quad \eta_2(u) := \frac{\eta'(u)}{\eta''(0)}, \tag{3.8a,b}$$

where $\eta_2(u + 2\pi) = \eta_2(u)$ and $\eta_1(u + 2\pi) = \eta_1(u) + \theta\eta_2(u)$ with

$$\theta := \partial_{\mathcal{E}}T(\mathcal{E}(c), c) \frac{\sqrt{2\mathcal{E}(c)}}{c^2 - 2\sqrt{2\mathcal{E}(c)}} < 0, \tag{3.9}$$

due to monotonicity (2.13). By Proposition 1 in Geyer *et al.* (2022), 0 is the third eigenvalue of \mathcal{L} with two simple negative eigenvalues below 0.

REMARK 3.1. *One can prove the assertion of Lemma 3.1 by using small-amplitude expansions. The periodic solution with the profile $\eta \in C_{per}^{\infty}(\mathbb{T})$ is expanded near the trivial solution, see (1.21), as*

$$\eta = a \cos(u) - a^2 \sin^2(u) + O(a^3), \quad c^2 = 1 + \frac{1}{2}a^2 + O(a^4), \tag{3.10a,b}$$

where $a > 0$ is a small parameter. Then, \mathcal{L} along the solution family has one negative eigenvalue $-1 + O(a^2)$ and a small negative eigenvalue $-a^2 + O(a^4)$ with 0 being the third eigenvalue. Since $\text{Ker}(\mathcal{L}) = \text{span}(\eta')$ along the family of smooth periodic solutions for $c \in (1, c_*)$, the inertia index of \mathcal{L} remains the same for every $c \in (1, c_*)$.

The next lemma specifies the criterion for the constrained linear operator $\mathcal{L}|_{\mathcal{X}_c}$ to be positive, where $\mathcal{L}|_{\mathcal{X}_c} = \mathcal{L}|_{\{1, \eta''\}^{\perp}}$ is defined by the two constraints in (3.2).

LEMMA 3.2. *Let $\mathcal{L} : H_{per}^2(\mathbb{T}) \subset L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T})$ be given by (3.1) as in Lemma 3.1 and \mathcal{X}_c be the constrained subspace of $L^2(\mathbb{T})$ given by (3.2). Then, $\mathcal{L}|_{\mathcal{X}_c}$ has a simple zero eigenvalue and no negative eigenvalues, with the rest of its spectrum being bounded away from zero, if and only if the mapping*

$$(1, c_*) \ni c \mapsto \mathcal{M}(c) := \oint \eta \, du \tag{3.11}$$

is monotonically decreasing for $c \in (1, c_*)$.

Proof. By Proposition 2 in Geyer *et al.* (2022), we construct the 2-by-2 matrix related to the two constraints in \mathcal{X}_c

$$A := \begin{bmatrix} \langle \mathcal{L}^{-1}1, 1 \rangle & \langle \mathcal{L}^{-1}1, \eta'' \rangle \\ \langle \mathcal{L}^{-1}\eta'', 1 \rangle & \langle \mathcal{L}^{-1}\eta'', \eta'' \rangle \end{bmatrix}. \tag{3.12}$$

The inverse operator \mathcal{L}^{-1} on $\text{span}(1, \eta'')$ is well defined since

$$\text{Ker}(\mathcal{L}) = \text{span}(\eta') \perp \text{span}(1, \eta''). \tag{3.13}$$

Differentiating (1.19) in c yields

$$\mathcal{L}\partial_c\eta = 2c\eta'', \tag{3.14}$$

where $\partial_c\eta$ is defined along the family of 2π -periodic solutions $\{\eta\}_{c \in (1, c_*)}$. The family is smooth in c since $T(\mathcal{E}, c)$ is C^1 on $(0, \mathcal{E}_c) \times (0, c_*)$ and $\mathcal{E}(c)$ is C^1 on $(0, c_*)$ due to the

implicit function theorem for $T(\mathcal{E}(c), c) = 2\pi$ with $\partial_{\mathcal{E}}T(\mathcal{E}(c), c) < 0$. In addition, we have $\mathcal{L}1 = 2\eta'' - 1$ so that

$$\mathcal{L}(c^{-1}\partial_c\eta - 1) = 1. \tag{3.15}$$

By using (3.14) and (3.15), we compute

$$A = \begin{bmatrix} c^{-1}\langle\partial_c\eta, 1\rangle - 2\pi & c^{-1}\langle\partial_c\eta, \eta''\rangle \\ (2c)^{-1}\langle\partial_c\eta, 1\rangle & (2c)^{-1}\langle\partial_c\eta, \eta''\rangle \end{bmatrix}. \tag{3.16}$$

Since A is symmetric, we have $\langle\partial_c\eta, 1\rangle = 2\langle\partial_c\eta, \eta''\rangle$. Furthermore,

$$\det(A) = -\frac{\pi}{2c}\langle\partial_c\eta, 1\rangle = -\frac{\pi}{2c}\mathcal{M}'(c), \tag{3.17}$$

where $\mathcal{M}(c)$ is given by (3.11). Since $c > 0$, we have the following trichotomy from Proposition 2 in Geyer *et al.* (2022):

- (i) If $\mathcal{M}'(c) > 0$, then $\det(A) < 0$, hence A has one negative and one positive eigenvalue so that $\mathcal{L}|_{\mathcal{X}_c}$ admits one simple negative and a simple zero eigenvalue.
- (ii) If $\mathcal{M}'(c) = 0$, then $\det(A) = 0$ but $\text{tr}(A) < 0$, hence A has one negative and one zero eigenvalue so that $\mathcal{L}|_{\mathcal{X}_c}$ admits a double zero eigenvalue and no negative eigenvalues.
- (iii) If $\mathcal{M}'(c) < 0$, then $\det(A) > 0$, hence A has two negative eigenvalues so that $\mathcal{L}|_{\mathcal{X}_c}$ admits a simple zero eigenvalue and no negative eigenvalues.

The last case yields the assertion of the lemma.

REMARK 3.2. *Due to constraint (1.12), we have*

$$M(\eta) = \oint \eta \, du = -\oint (\partial_u\eta)^2 \, du = -Q(\eta), \tag{3.18}$$

so that the criterion in Lemma 3.2 is equivalent to the mapping of

$$(1, c_*) \ni c \mapsto Q(c) := \oint (\eta')^2 \, du \tag{3.19}$$

being monotonically increasing for $c \in (1, c_*)$.

Since $\mathcal{M}(c) \rightarrow 0$ as $c \rightarrow 1$ by (1.21) and $\mathcal{M}(c) < 0$ by (3.18), it is clear that $\mathcal{M}'(c) < 0$ for $c \gtrsim 1$. The next lemma asserts that $\mathcal{M}'(c) < 0$ for every $c \in (1, c_*)$.

LEMMA 3.3. *The mapping (3.11) is monotonically decreasing for every $c \in (1, c_*)$.*

Proof. Differentiating $T(\mathcal{E}(c), c) = 2\pi$ in c yields

$$\partial_c T(\mathcal{E}(c), c) + \mathcal{E}'(c) \partial_{\mathcal{E}} T(\mathcal{E}(c), c) = 0, \tag{3.20}$$

where

$$\left. \begin{aligned} \partial_c T(\mathcal{E}, c) &= 2c \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}}, \\ \partial_{\mathcal{E}} T(\mathcal{E}, c) &= -\mathcal{E}^{-1} \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}}, \end{aligned} \right\} \tag{3.21}$$

and we have used (2.18) with the substitution $\eta = \sqrt{2\mathcal{E}x}$. By using the same substitution, we define $\mathcal{M}(c) \equiv \mathcal{M}(\mathcal{E}(c), c)$ with

$$\mathcal{M}(\mathcal{E}, c) := 2 \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta \sqrt{c^2 - 2\eta} d\eta}{\sqrt{2\mathcal{E} - \eta^2}} = 2\sqrt{2\mathcal{E}} \int_{-1}^1 \frac{x \sqrt{c^2 - 2\sqrt{2\mathcal{E}x}} dx}{\sqrt{1 - x^2}}, \tag{3.22}$$

from which we obtain

$$\left. \begin{aligned} \partial_c \mathcal{M}(\mathcal{E}, c) &= 2c \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}}, \\ \partial_{\mathcal{E}} \mathcal{M}(\mathcal{E}, c) &= \mathcal{E}^{-1} \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta \sqrt{c^2 - 2\eta} d\eta}{\sqrt{2\mathcal{E} - \eta^2}} - \mathcal{E}^{-1} \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta^2 d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}}. \end{aligned} \right\} \tag{3.23}$$

Since $\partial_{\mathcal{E}} T(\mathcal{E}(c), c) < 0$, we obtain

$$\begin{aligned} \mathcal{M}'(c) &= \partial_c \mathcal{M}(\mathcal{E}(c), c) + \mathcal{E}'(c) \partial_{\mathcal{E}} \mathcal{M}(\mathcal{E}(c), c) \\ &= \frac{2c}{\mathcal{E}(c) |\partial_{\mathcal{E}} T(\mathcal{E}(c), c)|} \Delta(\mathcal{E}(c), c), \end{aligned} \tag{3.24}$$

where

$$\begin{aligned} \Delta(\mathcal{E}, c) &:= \left(\int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}} \right)^2 \\ &+ \left(\int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}} \right) \left(\int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta(c^2 - 3\eta) d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}} \right). \end{aligned} \tag{3.25}$$

We show that $\Delta(\mathcal{E}, c) < 0$, which implies that $\mathcal{M}'(c) < 0$. Indeed, since

$$\frac{\eta \sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E} - \eta^2}} < \frac{|\eta| \sqrt{c^2 + 2|\eta|}}{\sqrt{2\mathcal{E} - \eta^2}}, \quad \eta \in (0, \sqrt{2\mathcal{E}}), \tag{3.26}$$

we have

$$\left(\int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{d\eta}{\sqrt{2\mathcal{E} - \eta^2} \sqrt{c^2 - 2\eta}} \right) \left(\int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\eta \sqrt{c^2 - 2\eta} d\eta}{\sqrt{2\mathcal{E} - \eta^2}} \right) < 0. \tag{3.27}$$

The remaining part of $\Delta(\mathcal{E}, c)$ is also negative since

$$\begin{aligned} & \left(\int \frac{\eta \, d\eta}{\sqrt{2\mathcal{E} - \eta^2 \sqrt{c^2 - 2\eta}}} \right)^2 - \left(\int \frac{d\eta}{\sqrt{2\mathcal{E} - \eta^2 \sqrt{c^2 - 2\eta}}} \right) \left(\int \frac{\eta^2 \, d\eta}{\sqrt{2\mathcal{E} - \eta^2 \sqrt{c^2 - 2\eta}}} \right) \\ &= \int \int \frac{\eta_1 \eta_2 - \eta_2^2}{\sqrt{2\mathcal{E} - \eta_1^2 \sqrt{c^2 - 2\eta_1}} \sqrt{2\mathcal{E} - \eta_2^2 \sqrt{c^2 - 2\eta_2}}} \, d\eta_1 \, d\eta_2 \\ &= -\frac{1}{2} \int \int \frac{(\eta_1 - \eta_2)^2}{\sqrt{2\mathcal{E} - \eta_1^2 \sqrt{c^2 - 2\eta_1}} \sqrt{2\mathcal{E} - \eta_2^2 \sqrt{c^2 - 2\eta_2}}} \, d\eta_1 \, d\eta_2 < 0, \end{aligned} \tag{3.28}$$

where the integrations are defined on $[-\sqrt{2\mathcal{E}}, \sqrt{2\mathcal{E}}]$. Hence, $\Delta(\mathcal{E}(c), c) < 0$ and the assertion of the lemma has been proven.

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. First, we prove that the two constraints (1.25a,b) are preserved in the time evolution of the linearized equation (1.20a,b). Since $\Pi_0 \partial_u^{-1} \Pi_0$ is defined on zero-mean functions with the zero-mean constraint, taking the mean value of (1.20a,b) yields

$$2c \frac{d}{dt} \langle 1, \hat{\eta} \rangle = 0. \tag{3.29}$$

Multiplying (1.20a,b) by η'' and integrating by parts, we obtain for any solution $\hat{\eta} \in C^0(\mathbb{R}, H_{per}^1(\mathbb{T}))$

$$\begin{aligned} 2c \frac{d}{dt} \langle \eta'', \hat{\eta} \rangle &= \langle (c^2 - 2\eta)\eta'', \partial_u \hat{\eta} \rangle - \langle (1 - 2\eta'')\eta', \hat{\eta} \rangle \\ &= \langle (\eta')^2 - \eta, \partial_u \hat{\eta} \rangle - \langle (1 - 2\eta'')\eta', \hat{\eta} \rangle \\ &= \langle \eta' - 2\eta' \eta'', \hat{\eta} \rangle - \langle (1 - 2\eta'')\eta', \hat{\eta} \rangle \\ &= 0, \end{aligned} \tag{3.30}$$

where (1.19) has been used with $\eta \in C_{per}^\infty(\mathbb{T})$. Hence, the two constraints (1.25a,b) are preserved in time and the solution $\hat{\eta} \in C^0(\mathbb{R}, H_{per}^1(\mathbb{T}))$ of the linearized equation (1.20a,b) with $\hat{\eta}(\cdot, 0) = \hat{\eta}_0$ and $\hat{\eta}_0 \in \mathcal{X}_c$ satisfies $\hat{\eta}(\cdot, t) \in \mathcal{X}_c$ for every $t \in \mathbb{R}$. Thus, we have

$$\langle 1, \hat{\eta}(\cdot, t) \rangle = 0, \quad \langle \eta'', \hat{\eta}(\cdot, t) \rangle = 0, \quad t \in \mathbb{R}. \tag{3.31a,b}$$

If we further decompose

$$\hat{\eta}(\cdot, t) = a(t)\eta' + w(\cdot, t), \quad t \in \mathbb{R}, \tag{3.32}$$

then $w(\cdot, t) \in H_{per}^1(\mathbb{T}) \cap \mathcal{X}_c$ for $t \in \mathbb{R}$ satisfies the additional constraint

$$\langle \eta', w(\cdot, t) \rangle = 0, \quad t \in \mathbb{R}. \tag{3.33}$$

Next, the existence and uniqueness of solutions $\hat{\eta} \in C^0(\mathbb{R}, H_{per}^1(\mathbb{T}))$ of the linearized equation (1.20a,b) such that $\hat{\eta}(\cdot, 0) = \hat{\eta}_0$ follow by the energy method (Renardy & Rogers 2004). The energy quadratic form $\langle \mathcal{L}\hat{\eta}, \hat{\eta} \rangle$ is bounded and conserved for the

Travelling waves in a local model for shallow water waves




Figure 5. Dependence of \mathcal{E} (a) and \mathcal{M} (b) on c along the family of solutions of $T(\mathcal{E}(c), c) = 2\pi$. The black dots show the values $\mathcal{E}(1) = 0$, $\mathcal{M}(1) = 0$ and $\mathcal{E}(c_*) = \pi^4/512$, $\mathcal{M}(c_*) = -(\pi^3/24)$.

solution $\hat{\eta} \in C^0(\mathbb{R}, H^1_{per}(\mathbb{T}))$ of the linearized equation (1.20a,b). Since $\mathcal{L}\eta' = 0$, we get

$$\langle \mathcal{L}w(\cdot, t), w(\cdot, t) \rangle = \langle \mathcal{L}\hat{\eta}(\cdot, t), \hat{\eta}(\cdot, t) \rangle = \langle \mathcal{L}\hat{\eta}_0, \hat{\eta}_0 \rangle \leq \beta \|\hat{\eta}_0\|_{H^1_{per}}^2, \quad (3.34)$$

for some fixed $\beta > 0$. By Lemmas 3.2 and 3.3, $\langle \mathcal{L}w(\cdot, t), w(\cdot, t) \rangle$ is coercive for $w(\cdot, t) \in H^1_{per}(\mathbb{T}) \cap \mathcal{X}_c$ and is non-degenerate if $w(\cdot, t)$ is orthogonal to η' . Hence, we get the lower bound with some fixed $\alpha > 0$

$$\alpha \|w(\cdot, t)\|_{H^1_{per}}^2 \leq \langle \mathcal{L}w(\cdot, t), w(\cdot, t) \rangle \leq \beta \|\hat{\eta}_0\|_{H^1_{per}}^2, \quad (3.35)$$

which implies the first estimate (1.26a). In addition, we get from (1.20a,b) due to $\mathcal{L}\eta' = 0$

$$2cd'(t)\eta' + 2c\partial_t w = -\Pi_0 \partial_u^{-1} \Pi_0 \mathcal{L}w, \quad (3.36)$$

which allows us to control the unique $a \in C^1(\mathbb{R}, \mathbb{R})$ from the bound

$$\begin{aligned} 2cd'(t)\|\eta'\|_{L^2}^2 &= \langle (c^2 - 2\eta)\eta', \partial_u w(\cdot, t) \rangle - \langle (1 - 2\eta'')\eta, w(\cdot, t) \rangle \\ &\leq \gamma \|w(\cdot, t)\|_{H^1_{per}}, \end{aligned} \quad (3.37)$$

for some fixed $\gamma > 0$, which yields the second estimate (1.26b).

Figure 5 displays the dependence of \mathcal{E} and \mathcal{M} on c for $c \in (1, c_*)$ computed along the family of solutions of $T(\mathcal{E}(c), c) = 2\pi$. The mass integral can be computed in terms of

complete elliptic integrals by using 3.141 (integral 20) in Gradshteyn & Ryzhik (2007)

$$\begin{aligned}
 \mathcal{M}(\mathcal{E}, c) &= -2 \int_{-\sqrt{2\mathcal{E}}}^{\sqrt{2\mathcal{E}}} \frac{\sqrt{2\mathcal{E} - \eta^2}}{\sqrt{c^2 - 2\eta}} d\eta \\
 &= -4\mathcal{E} \int_{-1}^1 \frac{\sqrt{1 - x^2}}{\sqrt{c^2 - 2\sqrt{2\mathcal{E}}x}} dx \\
 &= -\frac{2}{3} \sqrt{c^2 + 2\sqrt{2\mathcal{E}}} \left[c^2 E \left(\sqrt{\frac{4\sqrt{2\mathcal{E}}}{c^2 + 2\sqrt{2\mathcal{E}}}} \right) \right. \\
 &\quad \left. - (c^2 - 2\sqrt{2\mathcal{E}}) K \left(\sqrt{\frac{4\sqrt{2\mathcal{E}}}{c^2 + 2\sqrt{2\mathcal{E}}}} \right) \right], \tag{3.38}
 \end{aligned}$$

where $K(k)$ and $E(k)$ are complete elliptic integrals of the first and second kind, respectively. The values of $\mathcal{E} = \mathcal{E}(c)$ are computed numerically from $T(\mathcal{E}_c, c) = 2\pi$ by a root-finding algorithm. Figure 5 illustrates the monotonicity result of Lemma 3.3. Since $\mathcal{E}(c) \rightarrow 0$ as $c \rightarrow 1$ follows from (1.21), we have $\mathcal{M}(c) \rightarrow 0$ as $c \rightarrow 1$. On the other hand, $\mathcal{E}(c_*) \rightarrow c_*^4/8 = \pi^4/512$ as $c \rightarrow c_*$ follows by Lemma 2.4 and we compute from (1.23) that

$$\mathcal{M}(c_*) = \frac{1}{8} \int_0^\pi (\pi^2 - 4\pi u + 2u^2) du = -\frac{\pi^3}{24}, \tag{3.39}$$

which agrees well with the numerical data in figure 5.

Numerical methods. The numerical data in figure 2 are an extended version of figure 5(a), where all roots of $T(\mathcal{E}, c) = 2\pi$ have been computed numerically from a bisection method, see figure 4. The numerical data in figure 1 were obtained from finding roots of the implicit function

$$|u| = \int_\eta^{\eta_{max}} \frac{\sqrt{c^2 - 2\eta}}{\sqrt{2\mathcal{E}(c) - \eta^2}} d\eta, \tag{3.40}$$

where $\eta_{max} := \sqrt{2\mathcal{E}(c)}$ for smooth profiles (panel a) and $\eta_{max} := c^2/2$ for singular profiles (panel b) and $\mathcal{E}(c)$ is a root of $T(\mathcal{E}(c), c) = 2\pi$ obtained on the lower part of the bifurcation diagram in figure 2.

Supplementary material. The data that support the findings of this study are available upon request from the authors.

Acknowledgements. This work is a part of the undergraduate thesis of S. Locke at McMaster University (2023–2024). The authors thank S. Dyachenko, P. Lushnikov, J. Weber and X. Zhao for many discussions related to the content of this work. Figure 1 was prepared by S. Wang as a part of an undergraduate summer project (2024).

Funding. D.E.P. acknowledges the funding of this study provided by the grant no. FSWE-2023-0004 through the State task program in the sphere of scientific activity of the Ministry of Science and Higher Education of the Russian Federation and grant no. NSH-70.2022.1.5 for the State support of leading Scientific Schools of the Russian Federation.

Declaration of interests. The authors report no conflict of interest.

Author ORCID.

 Dmitry E. Pelinovsky <https://orcid.org/0000-0001-5812-440X>.

Appendix A. Euler equations after a conformal transformation

Let $y = \eta(x, t)$ be the profile for the free surface of an incompressible and irrotational fluid in the 2π -periodic domain and assume a flat bottom at $y = -h_0$, where the vertical velocity vanishes. For a proper definition of the fluid depth h_0 , we add the zero-mean constraint on the free surface, that is

$$\oint \eta(x, t) \, dx = 0, \tag{A1}$$

which is invariant in the time evolution of Euler’s equations.

Let $\varphi(x, y, t)$ be the velocity potential, which satisfies the Laplace equation in the time-dependent spatial domain

$$D_\eta := \{(x, y) \in \mathbb{R}^2 : -\pi \leq x \leq \pi, -h_0 \leq y \leq \eta(x, t)\}, \tag{A2}$$

subject to the periodic boundary conditions at $x = \pm\pi$ and the Neumann boundary condition at $y = -h_0$. The formulation of the water wave problem is completed by two additional (kinematic and dynamic) conditions at the free surface $y = \eta(x, t)$

$$\left. \begin{aligned} \eta_t + \varphi_x \eta_x - \varphi_y &= 0, \\ \varphi_t + \frac{1}{2}(\varphi_x)^2 + \frac{1}{2}(\varphi_y)^2 + \eta &= 0, \end{aligned} \right\} \text{ at } y = \eta(x, t), \tag{A3}$$

where the gravitational constant g is set to unity for convenience.

The method of conformal transformations is used to map the spatial domain D_η to the flat domain

$$\mathcal{D} := \{(u, v) \in \mathbb{R}^2 : -\pi \leq u \leq \pi, -h \leq v \leq 0\}, \tag{A4}$$

where h may be different from h_0 . The transformation is based on the conformal mapping $x + iy = z(u + iv, t)$, where $w := u + iv$ is a new complex variable and $z \in C^\omega(\mathcal{D})$ is a holomorphic function, the real and imaginary parts of which satisfy the Cauchy–Riemann equations

$$\frac{\partial x}{\partial u} = \frac{\partial y}{\partial v}, \quad \frac{\partial x}{\partial v} = -\frac{\partial y}{\partial u}. \tag{A5a,b}$$

To preserve the flat bottom $y = -h_0$ at $v = -h$, one needs to add the Neumann condition $\partial_v x|_{v=-h} = 0$, which ensures that $y(u, -h, t) = -h_0$ is u -independent. In addition, we require $x - u$ and $y - v$ be 2π -periodic functions of $u \in \mathbb{T} := \mathbb{R} \setminus (2\pi\mathbb{Z})$ to ensure that $x(\pi, v, t) - x(-\pi, v, t) = 2\pi$.

Abusing notations we refer to $x = x(u, t)$ and $y = \eta(u, t)$ at the top boundary of \mathcal{D}

$$x(u, t) + i\eta(u, t) = z(u, t), \quad \text{at } v = 0. \tag{A6}$$

Similarly, we abuse notations for the velocity potential $\varphi(u, v, t)$ and define

$$\xi(u, t) := \varphi(u, v = 0, t), \tag{A7}$$

on the flat top boundary of \mathcal{D} . Since the conformal transformation preserves the periodic boundary conditions and the zero vertical velocity condition at $v = -h$, the Laplace

equation can be solved with the following Fourier series:

$$\varphi(u, v, t) = \sum_{n \in \mathbb{Z}} \hat{\xi}_n(t) e^{inu} \frac{\cosh(n(v+h))}{\cosh(nh)}, \tag{A8}$$

where $\hat{\xi}_n(t)$ is the Fourier coefficient for $\xi(u, t) = \varphi(u, v = 0, t)$. Similarly, we obtain

$$\left. \begin{aligned} x(u, v, t) &= u + \sum_{n \in \mathbb{Z}} \hat{x}_n(t) e^{inu} \frac{\cosh(n(v+h))}{\cosh(nh)}, \\ y(u, v, t) &= v + \hat{\eta}_0 + \sum_{n \in \mathbb{Z}} \hat{x}_n(t) e^{inu} \frac{\sinh(n(v+h))}{\cosh(nh)}. \end{aligned} \right\} \tag{A9}$$

It follows from $y(u, -h, t) = -h_0$ that $\hat{\eta}_0 = h - h_0$. If $\hat{\eta}_0(t) = (1/2\pi) \oint \eta(u, t) du$ depends on time t , so does $h(t)$ which satisfies $\partial_t \eta(u, -h, t) - h'(t) \partial_v \eta(u, -h, t) = 0$ for all $u \in \mathbb{T}$.

Reducing the Fourier series for $x(u, v, t)$ and $y(u, v, t)$ on $v = 0$ yields

$$\left. \begin{aligned} x(u, t) &= u + \hat{x}_0(t) + \sum_{n \in \mathbb{Z} \setminus \{0\}} \hat{\eta}_n(t) e^{inu} (-i) \coth(nh), \\ \eta(u, t) &= \hat{\eta}_0(t) + \sum_{n \in \mathbb{Z} \setminus \{0\}} \hat{\eta}_n(t) e^{inu}, \end{aligned} \right\} \tag{A10}$$

with the correspondence $\hat{\eta}_n(t) = i \tanh(nh) \hat{x}_n(t)$ for $n \in \mathbb{Z} \setminus \{0\}$.

Let us introduce the non-local operator T_h with the Fourier symbol given by

$$\widehat{(T_h)}_n = i \tanh(hn), \quad n \in \mathbb{Z}, \tag{A11}$$

so that $\eta = \hat{\eta}_0 + T_h(x - u)$. The inverse of T_h is only defined on the zero-mean functions with the Fourier symbol given by

$$\widehat{(T_h^{-1})}_n = \begin{cases} -i \coth(hn), & n \in \mathbb{Z} \setminus \{0\}, \\ 0, & n = 0. \end{cases} \tag{A12}$$

Inverting $\eta = \hat{\eta}_0 + T_h(x - u)$ yields $x = u + \hat{x}_0 + T_h^{-1} \eta$ and

$$x_u = 1 + K_h \eta, \tag{A13}$$

where $K_h := T_h^{-1} \partial_u$ is a linear, self-adjoint, positive operator on $L^2(\mathbb{T})$. We set $\hat{x}_0 = 0$ in $x = u + \hat{x}_0 + T_h^{-1} \eta$ without loss of generality.

The equations of motion can be derived from the following Lagrangian (see Dyachenko *et al.* (1996) and Dyachenko *et al.* (2016), Appendix A for $h = \infty$):

$$\begin{aligned} \mathcal{L}(\xi, \eta, x) &:= \oint \xi (\eta_t x_u - \eta_u x_t) du + \frac{1}{2} \oint \xi T_h \xi_u du - \frac{1}{2} \oint \eta^2 x_u du \\ &+ \oint f(\eta - T_h(x - u)) du, \end{aligned} \tag{A14}$$

where f is the Lagrange multiplier satisfying $\oint f du = 0$. Variation of \mathcal{L} in ξ, η and x yields the system of equations

$$\left. \begin{aligned} \eta_t x_u - \eta_u x_t + T_h \xi_u &= 0, \\ -\xi_t x_u + \xi_u x_t - \eta x_u + f &= 0, \\ \xi_t \eta_u - \xi_u \eta_t + \eta \eta_u + T_h f &= 0, \end{aligned} \right\} \tag{A15}$$

with the additional constraint due to the reduction $h = h_0 + (1/2\pi) \oint \eta \, du$

$$\frac{1}{2} \oint \xi (\partial_h T_h) \xi_u \, du - \oint f (\partial_h T_h) (x - u) \, du = 0. \tag{A16}$$

Taking mean values in each equation of system (A15) and integrating by parts yields three conserved quantities

$$M_1(\eta) = \oint \eta x_u \, du = \oint \eta (1 + K_h \eta) \, du = 0, \tag{A17}$$

$$M_2(\xi, \eta) = \oint \xi x_u \, du = \oint \xi (1 + K_h \eta) \, du, \tag{A18}$$

$$M_3(\xi, \eta) = \oint \xi \eta_u \, du, \tag{A19}$$

where the constraint $M_1(\eta) = 0$ follows from the zero-mean constraint (A1) in physical coordinates. We express f from the second equation of system (A15), substitute it into the third equation and invert T_h on the periodic functions with zero mean. This transforms the system (A15) to the following system of two equations for ξ and η :

$$\left. \begin{aligned} \eta_t x_u - \eta_u x_t + T_h \xi_u &= 0, \\ \xi_t x_u - \xi_u x_t + \eta (1 + K_h \eta) + T_h^{-1} (\xi_t \eta_u - \xi_u \eta_t + \eta \eta_u) &= 0. \end{aligned} \right\} \tag{A20}$$

The constraint (A16) is rewritten in the equivalent form

$$\frac{1}{2} \oint \xi (\partial_h T_h) \xi_u \, du + \oint T_h^{-1} (\xi_t \eta_u - \xi_u \eta_t + \eta \eta_u) (\partial_h T_h) (T_h^{-1} \eta) \, du = 0. \tag{A21}$$

The conserved energy of system (A20) is given by

$$H(\xi, \eta) = \oint [\xi T_h \xi_u - \eta^2 (1 + K_h \eta)] \, du. \tag{A22}$$

To derive (A22), we multiply the first equation of system (A20) by ξ_t and the second equation by η_t , integrate over the period and subtract one equation from another. After integration by parts, we get $(d/dt)H(\xi, \eta) = 0$ if and only if

$$\begin{aligned} & -\frac{1}{2} h'(t) \oint \xi (\partial_h T_h) \xi_u \, du - h'(t) \oint \eta \eta_u (\partial_h T_h^{-1}) \eta \, du \\ & - \oint (\xi_t \eta_u - \xi_u \eta_t) (x_t - T_h^{-1} \eta_t) \, du = 0. \end{aligned} \tag{A23}$$

Since $x_t - T_h^{-1} \eta_t = h'(t) (\partial_h T_h^{-1}) \eta$, $\partial_h T_h^{-1} = -T_h^{-1} (\partial_h T_h) T_h^{-1}$ and T_h^{-1} is skew-adjoint, the last constraint is identical to the constraint (A21) for every $h'(t)$. This proves the conservation of $H(\xi, \eta)$.

The conserved quantities (A17), (A18), (A19) and (A22) coincide with the conserved quantities for Euler's equation in physical coordinates, see Benjamin & Olver (1982) and Dyachenko *et al.* (1996) for $h = \infty$.

In order to introduce the scalar model (1.1), we rewrite (A20) in the reference frame moving with the wave speed c

$$\left. \begin{aligned} \eta_t x_u - \eta_u x_t + T_h \xi_u - c \eta_u &= 0, \\ \xi_t x_u - \xi_u x_t + \eta(1 + K_h \eta) - c \xi_u + T_h^{-1}(\xi_t \eta_u - \xi_u \eta_t + \eta \eta_u) &= 0, \end{aligned} \right\} \quad (\text{A24})$$

where u now stands for $u - ct$ and we have used the chain rule with

$$\left. \begin{aligned} \xi_t &\rightarrow \xi_t - c \xi_u, \\ \eta_t &\rightarrow \eta_t - c \eta_u, \\ x_t &\rightarrow x_t + c - c x_u. \end{aligned} \right\} \quad (\text{A25})$$

We introduce a change of variables by

$$\xi = c T_h^{-1} \eta + \zeta, \quad (\text{A26})$$

after which the system (A24) can be rewritten in the form

$$\left. \begin{aligned} \eta_t x_u - \eta_u x_t + T_h \zeta_u &= 0, \\ \zeta_t x_u - \zeta_u x_t + \eta(1 + K_h \eta) - c \zeta_u + c x_t - c^2 K_h \eta \\ + T_h^{-1}(\zeta_t \eta_u - \zeta_u \eta_t + \eta \eta_u + c \eta_u x_t - c \eta_t K_h \eta) &= 0. \end{aligned} \right\} \quad (\text{A27})$$

Substituting $-c \zeta_u = c T_h^{-1}(\eta_t x_u - \eta_u x_t)$ to the second equation of system (A27) and taking the derivative of $x = u + T_h^{-1} \eta$ in t yields

$$\begin{aligned} \zeta_t x_u - \zeta_u x_t + T_h^{-1}(\zeta_t \eta_u - \zeta_u \eta_t) + h'(t)(\partial_h T_h^{-1}) \eta \\ + 2c T_h^{-1} \eta_t - c^2 K_h \eta + \eta(1 + K_h \eta) + \frac{1}{2} K_h \eta^2 = 0. \end{aligned} \quad (\text{A28})$$

The scalar model (1.1) follows by ignoring the constraint (A21) and the first equation of system (A27) and by setting $\zeta \equiv 0$ and $h'(t) \equiv 0$ in (A28). Babenko's equation (1.8), which is the exact equation for travelling waves, see Babenko (1987), corresponds to the time-independent solutions of (A27) and (A28) with $\zeta \equiv 0$ and $h'(t) \equiv 0$ since u in (1.1) stands for $u - ct$.

Appendix B. Introducing the local model

One popular model for fluid motion is the intermediate long-wave (ILW) equation written in the form

$$\partial_t \eta + \eta \partial_u \eta = \mathcal{K}_h(\partial_u \eta), \quad (\text{B1})$$

where \mathcal{K}_h is defined by the Fourier symbol

$$\widehat{(\mathcal{K}_h)}_n = \begin{cases} n \coth(hn), & n \in \mathbb{Z} \setminus \{0\}, \\ h^{-1}, & n = 0. \end{cases} \quad (\text{B2})$$

In comparison with (1.3), we have the correspondence

$$\mathcal{K}_h = K_h + \frac{1}{2\pi h} \oint \cdot du. \quad (\text{B3})$$

The ILW equation (B1) is integrable by inverse scattering and many results on well posedness and the dynamics of nonlinear waves have been obtained for this fluid model,

see review in Saut (2019). In the shallow water limit $h \rightarrow 0$, the scaling transformation

$$\eta(u, t) := h\tilde{\eta}(\tilde{u}, \tilde{t}), \quad \tilde{u} := u + h^{-1}t, \quad \tilde{t} := ht, \tag{B4}$$

recovers formally the Korteweg–de Vries (KdV) equation

$$\partial_{\tilde{t}}\tilde{\eta} + \tilde{\eta}\partial_{\tilde{u}}\tilde{\eta} + \frac{1}{3}\partial_{\tilde{u}}^3\tilde{\eta} = 0, \tag{B5}$$

due to the asymptotic expansion

$$\mathcal{K}_h = \frac{1}{h} - \frac{1}{3}h\partial_u^2 + O(h^3). \tag{B6}$$

For completeness, in the deep water limit $h \rightarrow \infty$, the ILW equation (B1) becomes the Benjamin–Ono (BO) equation

$$\partial_t\eta + \eta\partial_u\eta + \mathcal{H}(\partial_u^2\eta) = 0, \tag{B7}$$

where \mathcal{H} is the periodic Hilbert transform defined by (1.10). Both the KdV and BO equations are also integrable by inverse scattering.

To obtain the local evolution equation (1.11) from the non-local model (1.1), we replace K_h given by (1.3) with

$$\tilde{K}_h = K_h + \frac{1}{2\pi h} \oint \cdot du - \frac{1}{h} = \mathcal{K}_h - \frac{1}{h}. \tag{B8}$$

The difference between \tilde{K}_h and K_h appears in the local term $1/h$. It is removed from the mean term in the definition of K_h and from all Fourier modes in the definition of \tilde{K}_h . Since $K_h = T_h^{-1}\partial_u$, we can similarly define $\tilde{K}_h = \tilde{T}_h^{-1}\partial_u$ and expand asymptotically as $h \rightarrow 0$

$$\tilde{K}_h = -\frac{1}{3}h\partial_u^2 + O(h^3), \quad \tilde{T}_h^{-1} = -\frac{1}{3}h\partial_u + O(h^3). \tag{B9a,b}$$

By using the scaling transformation

$$\eta(u, t) := h^{-1}\tilde{\eta}(\tilde{u}, \tilde{t}), \quad u := 3^{-1/2}\tilde{u}, \quad t := 3^{-1/2}h^{1/2}\tilde{t}, \quad c = h^{-1/2}\tilde{c}, \tag{B10a–d}$$

we obtain the formal limit of the non-local model (1.1) as $h \rightarrow 0$ in the form

$$-2\tilde{c}\partial_{\tilde{u}}\partial_{\tilde{t}}\tilde{\eta} = (-\tilde{c}^2\partial_{\tilde{u}}^2 - 1)\tilde{\eta} + \tilde{\eta}\partial_{\tilde{u}}^2\tilde{\eta} + \frac{1}{2}\partial_{\tilde{u}}^2\tilde{\eta}^2. \tag{B11}$$

Expanding the derivatives, changing the sign and removing the tilde notations yields the local model (1.11).

REFERENCES

- ALBER, M., CAMASSA, R., FEDOROV, Y.N., HOLM, D.D. & MARSDEN, J.E. 1999 On billiard solutions of nonlinear PDEs. *Phys. Lett. A* **264**, 171–178.
- ALBER, M., CAMASSA, R., HOLM, D.D. & MARSDEN, J.E. 1995 On the link between umbilic geodesics and soliton solutions of nonlinear PDEs. *Proc. R. Soc. Lond. A* **450**, 677–692.
- AMICK, C.J., FRAENKEL, L.E. & TOLAND, J.F. 1982 On the Stokes conjecture for the wave of extreme form. *Acta Mathematica* **148**, 193–214.
- BABENKO, K.I. 1987 Some remarks on the theory of surface waves of finite amplitude. *Sov. Math. Dokl.* **35**, 599–603.
- BENJAMIN, T.B. & OLVER, P.J. 1982 Hamiltonian structure, symmetries and conservation laws for water waves. *J. Fluid Mech.* **125**, 137–185.
- BERTI, M., MASPERO, A. & VENTURA, P. 2022 Full description of Benjamin–Feir instability of Stokes waves in deep water. *Invent. Math.* **230**, 651–711.

- BRUELL, G. & DHARA, R.N. 2021 Waves of maximal height for a class of nonlocal equations with homogeneous symbols. *Indiana Univ. Math. J.* **70**, 711–742.
- CHOI, W. & CAMASSA, R. 1999 Exact evolution equations for surface waves. *J. Engng Maths* **125**, 756–760.
- CRAIG, W. & SULEM, C. 1993 Numerical simulation of gravity waves. *J. Comput. Phys.* **108**, 73–83.
- CREEDON, R.P. & DECONINCK, B. 2023 A high-order asymptotic analysis of the Benjamin–Feir instability spectrum in arbitrary depth. *J. Fluid Mech.* **956**, A29.
- DYACHENKO, A.I., KUZNETSOV, E.A., SPECTOR, M.D. & ZAKHAROV, V.E. 1996 Analytical description of the free surface dynamics of an ideal fluid (canonical formalism and conformal mapping). *Phys. Lett. A* **221**, 73–79.
- DYACHENKO, S.A., LUSHNIKOV, P.M. & KOROTKEVICH, A.O. 2016 Branch cuts of Stokes wave on deep water. Part I. Numerical solution and Padé approximation. *Stud. Appl. Maths* **137**, 419–472.
- DYACHENKO, S.A. & SEMENOVA, A. 2023a Quasiperiodic perturbations of Stokes waves: Secondary bifurcations and stability. *J. Comput. Phys.* **492**, 112411.
- DYACHENKO, S.A. & SEMENOVA, A. 2023b Canonical conformal variables based method for stability of Stokes waves. *Stud. Appl. Maths* **150**, 705–715.
- GEYER, A., MARTINS, R.H., NATALI, F. & PELINOVSKY, D.E. 2022 Stability of smooth periodic traveling waves in the Camassa–Holm equation. *Stud. Appl. Maths* **148**, 27–61.
- GEYER, A. & PELINOVSKY, D.E. 2017 Spectral stability of periodic waves in the generalized reduced Ostrovsky equation. *Lett. Math. Phys.* **107**, 1293–1314.
- GEYER, A. & PELINOVSKY, D.E. 2019 Linear instability and uniqueness of the peaked periodic wave in the reduced Ostrovsky equation. *SIAM J. Math. Anal.* **51**, 1188–1208.
- GEYER, A. & PELINOVSKY, D.E. 2020 Spectral instability of the peaked periodic wave in the reduced Ostrovsky equation. *Proc. Am. Math. Soc.* **148**, 5109–5125.
- GEYER, A. & PELINOVSKY, D.E. 2024 Stability of smooth periodic traveling waves in the Degasperis–Procesi equation. *J. Differ. Equ.* **404**, 354–390.
- GRADSHTEYN, I.S. & RYZHIK, I.M. 2007 *Table of Integrals, Series and Products*. Elsevier.
- HARROP–GRIFFITHS, B., IFRIM, M. & TATARU, D. 2017 Finite depth gravity water waves in holomorphic coordinates. *Ann. PDE* **3**, 1–102.
- HONE, A.N.W., NOVIKOV, V. & WANG, J.P. 2018 Generalizations of the short pulse equation. *Lett. Math. Phys.* **108**, 927–947.
- HUNTER, J.K., IFRIM, M. & TATARU, D. 2016 Two-dimensional water waves in holomorphic coordinates. *Commun. Math. Phys.* **346**, 483–552.
- HUNTER, J.K. & SAXTON, R. 1991 Dynamics of director fields. *SIAM J. Appl. Maths* **51**, 1498–1521.
- HUNTER, J.K. & ZHENG, Y. 1994 On a completely integrable nonlinear hyperbolic variational equation. *Physica D* **79**, 361–386.
- HUR, V.M. & YANG, Z. 2023 Unstable Stokes waves. *Arch. Rat. Mech. Anal.* **247**, 62.
- KOROTKEVICH, A.O., LUSHNIKOV, P.M., SEMENOVA, A. & DYACHENKO, S.A. 2023 Superharmonic instability of Stokes waves. *Stud. Appl. Maths* **150**, 119–134.
- LENELLS, J. 2005a Traveling wave solutions of the Degasperis–Procesi equation. *J. Math. Anal. Appl.* **306**, 72–82.
- LENELLS, J. 2005b Traveling wave solutions of the Camassa–Holm equation. *J. Differ. Equ.* **217**, 393–430.
- LOCKE, S. & PELINOVSKY, D.E. 2025 Peaked Stokes waves as solutions of Babenko’s equation. *Appl. Math. Lett.* **161**, 109359.
- LONG, T. & LIU, C. 2023 Orbital stability of smooth solitary waves for the b -family of Camassa–Holm equations. *Physica D* **446**, 133680.
- LUSHNIKOV, P.M. 2016 Structure and location of branch point singularities for Stokes waves on deep water. *J. Fluid Mech.* **800**, 557–594.
- LUSHNIKOV, P.M., DYACHENKO, S.A. & SILANTYEV, D.A. 2017 New conformal mapping for adaptive resolving of the complex singularities of Stokes wave. *Proc. R. Soc. A* **473**, 20170198.
- MADIYEVA, A. & PELINOVSKY, D.E. 2021 Growth of perturbations to the peaked periodic waves in the Camassa–Holm equation. *SIAM J. Math. Anal.* **53**, 3016–3039.
- MATSUNO, Y. 2020 Parametric solutions of the generalized short pulse equations. *J. Phys. A* **53**, 105202.
- NGUYEN, H.Q. & STRAUSS, W.A. 2023 Proof of modulational instability of Stokes waves in deep water. *Commun. Pure Appl. Maths* **76**, 1035–1084.
- PLOTNIKOV, P. 2002 A proof of the Stokes conjecture in the theory of surface waves. *Stud. Appl. Maths* **108**, 217–244.
- RENARDY, M. & ROGERS, R.C. 2004 *An Introduction to Partial Differential Equations*. Springer.
- SAUT, J.C. 2019 Benjamin–Ono and intermediate long wave equations: modeling, IST and PDE. In *Nonlinear Dispersive Partial Differential Equations and Inverse Scattering* (ed. P. Miller, P. Perry, J.C. Saut, & C. Sulem), Fields Institute Communications, vol. 83, pp. 95–160. Springer.

Travelling waves in a local model for shallow water waves

- STANISLAVOVA, M. & STEFANOV, A. 2016 On the spectral problem $Lu = \lambda u'$ and applications. *Commun. Math. Phys.* **343**, 361–391.
- TANVEER, S. 1991 Singularities in water waves and Rayleigh–Taylor instability. *Proc. R. Soc. Lond. A* **435**, 137–158.
- TOLAND, J.F. 1978 On the existence of a wave of greatest height and Stokes’s conjecture. *Proc. R. Soc. Lond. A* **363**, 469–485.
- WILKENING, J. 2020 Harmonic stability of standing water waves. *Q. Appl. Maths* **78**, 219–260.
- WILKENING, J. 2021 Traveling-standing water waves. *Fluids* **6**, 187.
- WILKENING, J. & ZHAO, X. 2021 Quasi-periodic travelling gravity-capillary waves. *J. Fluid Mech.* **915**, A7.
- WILKENING, J. & ZHAO, X. 2023a Spatially quasi-periodic bifurcations from periodic traveling water waves and a method for detecting bifurcations using signed singular values. *J. Comput. Phys.* **478**, 111954.
- WILKENING, J. & ZHAO, X. 2023b Spatially quasi-periodic water waves of finite depth. *Proc. R. Soc. A* **479**, 20230019.
- ZAKHAROV, V.E. 1968 Stability of periodic waves of finite amplitude on a surface. *J. Appl. Mech. Tech. Phys.* **9**, 190–194.
- ZAKHAROV, V.E. & DYACHENKO, A.I. 1996 High-Jacobian approximation in the free surface dynamics of an ideal fluid. *Physica D* **98**, 652–664.