

ASD, and 41 were suspected as having ASD. In total, 39% of service users had, or were suspected of having, ASD. The prevalence was higher in female service users than males with a total of 48% of female service users having, or being suspected of having ASD, compared with 22% of males. Comparative data demonstrated that autistic/suspected autistic young people presenting to the service were more likely than their neurotypical counterparts to be: over 13 years old, have a longer symptom duration before presentation, have an Educational Health Care Plan, report friendship difficulties, have a family history of neurodiversity, report sensory difficulties, and have sleep difficulties. RCADS scores found that the ASD group were more likely than the neurotypical group to have clinical levels of anxiety (58.3% vs 15.3%) and depression (80.6% vs 58.3%).

Conclusion. Our audit suggests that there is a higher prevalence of young people with ASD/ASD traits presenting to a paediatric fatigue service than found in the general population. Reasons for this may relate to undiagnosed ASD presenting as severe fatigue due to the energy draining nature of camouflaging, as well as sensory overload, known as autistic burnout. Do we need to develop a specialist treatment pathway which is better adapted to these young people's needs? We are planning a follow up study and focus groups to explore this complexity further.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Psychiatric Emergency Bleep Documentation Enhancement Audit

Dr Anastasija Davidova*, Dr Scott Young, Dr Amal Al Sayegh and Dr Jude Halford

NHS Lothian, West Lothian, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.553

Aims. West Lothian Psychiatry operates in a district general hospital, fostering a close working relationship between medical and psychiatric practitioners including the Psychiatric 2222 call (akin to medical emergency/cardiac arrest response). No other team like this has been identified in Scotland. Whilst there is a range of scenarios where this is used, there is no 'gold standard' for defining a psychiatric emergency or how to document these. Preliminary data gathered between August and November 2022 revealed concerns regarding call appropriateness, medical staff proficiency in de-escalation and restraint on medical wards, inadequate handovers, and poor documentation. This prompted a collaborative quality improvement project, undertaken by psychiatric and medical team leaders. Part of this initiative was an audit to improve the documentation of psychiatric emergencies to achieve a 90% compliance rate using a new checklist.

Methods. Cycle 1 of the audit (December 2022 to April 2023) identified patients through the 2222 calls to switchboard (n = 54). TrakCare notes were reviewed to assess call rationale and outcomes, focusing on documentation by the attending psychiatric team. A documentation checklist within the electronic records system was designed and introduced in July 2023, for completion by the junior doctor. Cycle 2 (November 2023 to January 2024, n = 47) aimed to assess improvements by comparing results with the previous cycle.

Results. There was a significant improvement in documentation rates with the checklist (44% to 90%). Indirect enhancements were observed in ward nursing documentation (65% to 83%) and medical ward doctor documentation (39% to 57%).

Appropriateness of emergency calls increased from 65% to 74%, with attending doctors' participation in emergencies longer than 10 minutes rising to 68% from 47%. The initial audit revealed a lack of awareness among senior medical staff regarding overnight psychiatric emergency calls, especially in cases of repeated calls for the same individual. The improved documentation played a pivotal role in addressing this issue, facilitating effective information sharing and changes in patient management plans, reducing further emergency calls.

Conclusion. The documentation checklist significantly improved junior doctor documentation, positively impacting patient care and communication among staff. This successful intervention serves as a promising model that can be replicated in other documentation domains. Moreover, this project has set the stage for broader initiatives within a larger Quality Improvement framework. The ongoing efforts are directed towards establishing a shared model for the psychiatric emergency bleep, optimising staffing resources for restraint procedures and improving staff de-escalation skills.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

An Audit of the Prescribing and Monitoring of Antipsychotic Medication in an Older Adult Inpatient Psychiatric Ward Using NICE Guidance [CG178] Psychosis and Schizophrenia in Adults: Prevention and Management

Dr Alaa Shabaka, Dr Thomas Parry, Dr Mary Ugah and Dr Stephen De Souza*

Somerset NHS Foundation Trust, Taunton, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.554

Aims. The National Institute for Health and Care Excellence (NICE) offers guidance for prescribing and monitoring of antipsychotic medications. In this audit we sought to investigate if our unit was compliant with this guidance.

Methods. The audit was carried out on a 28 bedded older adult inpatient psychiatric unit. The notes of all patients admitted to this ward on 27/11/2023 were reviewed. Any patient on an antipsychotic was included in the audit. Four standards reflecting the prescribing and monitoring of antipsychotics were identified. These were:

1.3.5.1 The choice of antipsychotic medication should be made by the service user and healthcare professional together, taking into account the views of the carer if the service user agrees.

1.3.6.1 Before starting antipsychotic medication, undertake and record the baseline investigations.

1.3.6.2 Before starting antipsychotic medication, offer the person with psychosis or schizophrenia an electrocardiogram (ECG).

1.3.6.3 Treatment with antipsychotic medication should be considered an explicit individual therapeutic trial.

1.3.6.4 Monitor and record the following (response to treatment – side effects – adherence – physical health) regularly and systematically throughout treatment.

These five areas of guidance were broken down into 22 domains which are outlined in results below.

Results. Of 28 patients admitted to the ward, 22 were on antipsychotic medication.