

## **$(H, C)$ -GROUPS WITH POSITIVE LINE BUNDLES**

YUKITAKA ABE

### **§ 0. Introduction**

Let  $G$  be a connected complex Lie group. Then there exists the smallest closed complex subgroup  $G^0$  of  $G$  such that  $G/G^0$  is a Stein group (Morimoto [8]). Moreover  $G^0$  is a connected abelian Lie group and every holomorphic function on  $G^0$  is a constant.  $G^0$  is called an  $(H, C)$ -group or a toroidal group. Every connected complex abelian Lie group is isomorphic to the direct product  $G^0 \times C^m \times C^{*n}$ , where  $G^0$  is an  $(H, C)$ -group ([7], [9]).

Recently, several interesting results with respect to  $(H, C)$ -groups have been obtained (Kazama [5], Kazama and Umeno [6], Vogt [13], [14] and [15]). The set of  $(H, C)$ -groups includes the set of complex tori. A complex torus is called an abelian variety if it satisfies Riemann condition. The definition of quasi-abelian variety for  $(H, C)$ -groups was given in [2]. In this paper we shall show that the concept of quasi-abelian variety is a natural generalization of abelian variety. Throughout this paper, we assume that  $\dim H^1(X, \mathcal{O}) < \infty$  for  $(H, C)$ -groups  $X$ . Our main result is the following.

Let  $X = C^n/\Gamma$  be an  $(H, C)$ -group. The following statements are equivalent:

- (1)  $X$  has a positive line bundle;
- (2)  $X$  is a quasi-abelian variety;
- (3)  $X$  is a covering space on an abelian variety;
- (4)  $X$  is embedded in a complex projective space as a locally closed submanifold.

The above result is well-known for complex tori. For the proof we use the theory of weakly 1-complete manifolds and results of Vogt. We note that implications (2)  $\Rightarrow$  (3) and (2)  $\Rightarrow$  (4) were obtained by Gherardelli

---

Received April 12, 1985.

Revised September 18, 1986.

and Andreotti [2]. Combining with a result of Gherardelli and Andreotti [2], we get the affirmative answer to a problem of the structure of weakly 1-complete manifolds in the case of  $(H, C)$ -groups (see § 6).

The author was inspired from dissertations of Pothering [12] and Vogt [13]. He is very grateful to Professor H. Kazama who told him the existence of their dissertations.

### § 1. Preliminaries

Let  $G$  be an  $n$ -dimensional connected complex Lie group without nonconstant holomorphic functions. Such a Lie group  $G$  is said to be an  $(H, C)$ -group or a toroidal group ([7], [8]). We recall that  $G$  is abelian and then  $G$  is isomorphic onto  $\mathbb{C}^n/\Gamma$  for some discrete subgroup  $\Gamma$  of  $\mathbb{C}^n$  as a Lie group ([8]). If  $\mathbb{C}^n/\Gamma$  is an  $(H, C)$ -group, then the generators of  $\Gamma$  contains  $n$  vectors linearly independent over  $\mathbb{C}$  and  $\text{rank } \Gamma = n + q$  ( $1 \leq q \leq n$ ). When  $\Gamma$  is generated by  $p_1, \dots, p_{n+q} \in \mathbb{C}^n$ , we write

$$P = (p_1, \dots, p_{n+q}),$$

and we call it a period basis of  $\Gamma$ , or also of  $X = \mathbb{C}^n/\Gamma$ . Two period bases  $P$  and  $P'$  are equivalent if and only if there exist a non-singular matrix  $S$  and a unimodular matrix  $M$  such that

$$P' = SPM.$$

A period basis  $P$  is always equivalent to the following standard form

$$(I_n V),$$

where  $V = (v_1, \dots, v_q)$ ,  $v_j = {}^t(v_{1j}, \dots, v_{nj})$ ,  $\det(\text{Im } v_{ij}; 1 \leq i, j \leq q) \neq 0$  and  $I_n$  is the  $(n, n)$  unit matrix.

It is well-known that  $\mathbb{C}^n/\Gamma$  is an  $(H, C)$ -group if and only if

$$\max \left\{ \left| \sum_{k=1}^n v_{kj} m_k - m_{n+j} \right|; 1 \leq j \leq q \right\} > 0$$

for all  $m = (m_1, \dots, m_n, m_{n+1}, \dots, m_{n+q}) \in \mathbb{Z}^{n+q} \setminus \{0\}$  (Kopfermann [7] and Morimoto [8]).

A discrete subgroup  $\Gamma$  of rank  $n + q$  in  $\mathbb{C}^n$  generates an  $(n + q)$ -dimensional real linear subspace  $\mathbb{R}_\Gamma^{n+q}$  of  $\mathbb{C}^n$ .  $\mathbb{R}_\Gamma^{n+q}$  contains the  $q$ -dimensional complex linear subspace  $\mathbb{C}_\Gamma^q$  which is the maximal complex linear subspace contained in  $\mathbb{R}_\Gamma^{n+q}$ . If we take the standard form for the period basis of  $\Gamma$ , then  $\text{Im } v_1, \dots, \text{Im } v_q$  generate  $\mathbb{C}_\Gamma^q$ .

§ 2. Factors of automorphy

We introduce some results of Vogt ([13] and [14]). For the details, we refer the reader to [13].

Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$ ,  $X = C^n/\Gamma$  and  $\pi: C^n \rightarrow X$  be the projection. If  $p: L \rightarrow X$  is a holomorphic line bundle, then its pull-back  $\pi^*L$  is given by the following fibre product

$$\pi^*L = C^n \times_X L = \{(z, v) \in C^n \times L; \pi(z) = p(v)\}.$$

Since any holomorphic Line bundle over  $C^n$  is analytically trivial, we have a trivialization

$$\varphi: \pi^*L \longrightarrow C^n \times C, (z, v) \longmapsto (z, \varphi_z(v))$$

of  $\pi^*L$ . We define

$$\alpha: \Gamma \times C^n \longrightarrow C^*, \alpha(\gamma, z) := \varphi_{z+\gamma}\varphi_z^{-1}.$$

Then  $\alpha$  satisfies the following conditions:

- (a)  $\alpha_\gamma(z) := \alpha(\gamma, z)$  is holomorphic for all  $\gamma \in \Gamma$ ;
- (b)  $\alpha(0, z) = 1$  for all  $z \in C^n$ ;
- (c)  $\alpha(\gamma + \gamma', z) = \alpha(\gamma, z + \gamma')\alpha(\gamma', z)$  for all  $\gamma, \gamma' \in \Gamma$  and  $z \in C^n$ .

DEFINITION. A map  $\alpha: \Gamma \times C^n \rightarrow C^*$  is called a factor of automorphy for  $\Gamma$  on  $C^n$  if it satisfies the above conditions (a), (b) and (c).

Conversely, if  $\alpha: \Gamma \times C^n \rightarrow C^*$  is a factor of automorphy, then we get a line bundle  $L$  over  $C^n/\Gamma$  defined as the quotient of  $C^n \times C$  by the following action of  $\Gamma$ :

$$\gamma(z, v) := (z + \gamma, \alpha(\gamma, z)v) \text{ for } \gamma \in \Gamma, z \in C^n, v \in C.$$

DEFINITION. Two factors of automorphy  $\alpha, \beta$  are said to be equivalent if there exists a holomorphic function  $h: C^n \rightarrow C^*$  such that

$$\beta(\gamma, z) = h(z + \gamma)\alpha(\gamma, z)h^{-1}(z)$$

for all  $\gamma \in \Gamma$  and  $z \in C^n$ .

PROPOSITION 1 (Vogt [13] and [14]). *The equivalent classes of factors of automorphy for  $\Gamma$  on  $C^n$  correspond one-to-one to the isomorphism classes of holomorphic line bundles over  $C^n/\Gamma$ .*

PROPOSITION 2 (Vogt [13]). *Let  $L_1$  and  $L_2$  be holomorphic line bundles over  $C^n/\Gamma$ . If factors of automorphy  $\alpha_1$  and  $\alpha_2$  give  $L_1$  and  $L_2$  respectively,*

then  $L_1 \otimes L_2$  is given by the factor of automorphy

$$\alpha_1 \alpha_2: \Gamma \times \mathbb{C}^n \longrightarrow \mathbb{C}^*, \quad \alpha_1 \alpha_2(\gamma, z) := \alpha_1(\gamma, z) \alpha_2(\gamma, z)$$

for all  $\gamma \in \Gamma$  and  $z \in \mathbb{C}^n$ .

**DEFINITION.** A map  $a: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}$  is called a summand of automorphy, if

- (a)  $a_\gamma(z) := a(\gamma, z)$  is holomorphic for all  $\gamma \in \Gamma$ ;
- (b)  $a(0, z) = 0$  for all  $z \in \mathbb{C}^n$ ;
- (c)  $a(\gamma + \gamma', z) = a(\gamma, z + \gamma') + a(\gamma', z)$  for all  $\gamma, \gamma' \in \Gamma$

and  $z \in \mathbb{C}^n$ .

Let  $a: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}$  be a summand of automorphy. We set  $\alpha(\gamma, z) := \exp(a(\gamma, z))$ . Then  $\alpha$  is a factor of automorphy.

**LEMMA 1** (Vogt [13]). *Let  $\Gamma$  be a discrete subgroup of rank  $r$  in  $\mathbb{C}^n$  with a basis  $\{\gamma_1, \dots, \gamma_r\}$ . If a map  $b: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}$  satisfies the following properties:*

- (a)  $b_j(z) := b(\gamma_j, z)$  is holomorphic for all  $j = 1, \dots, r$ ;
- (b)  $b(\gamma_i, z + \gamma_j) + b(\gamma_j, z) = b(\gamma_j, z + \gamma_i) + b(\gamma_i, z)$  for all  $i, j = 1, \dots, r$ ,

then there exists a summand of automorphy such that

$$a(\gamma_i, z) = b(\gamma_i, z) \quad \text{for all } i = 1, \dots, r.$$

**DEFINITION.** A factor of automorphy  $\alpha: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}^*$  is called a theta factor for  $\Gamma$  on  $\mathbb{C}^n$ , if it is expressed as follows,

$$\alpha(\gamma, z) = \exp 2\pi\sqrt{-1}(\mathcal{L}_\gamma(z) + c(\gamma)),$$

where  $\mathcal{L}_\gamma(z)$  is a linear polynomial and  $c(\gamma)$  is a constant for all  $\gamma \in \Gamma$ .

The following proposition was mentioned in [15], and its proof is hidden in the proof of Theorem in [14].

**PROPOSITION 3.** *Let  $\mathbb{C}^n/\Gamma$  be an  $(H, \mathbb{C})$ -group. For every line bundle  $L$  over  $\mathbb{C}^n/\Gamma$ , there exist a topologically trivial line bundle  $L_0$  and a line bundle  $L_1$  given by a theta factor such that  $L \cong L_0 \otimes L_1$ .*

*Proof.* The proof is along the argument of the implication 2)  $\Rightarrow$  1) of Theorem in [14]. Let  $\alpha: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}^*$  be a factor of automorphy which gives  $L$ . There exists a map  $a: \Gamma \times \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $\alpha(\gamma, z) = \exp a(\gamma, z)$ . We have

$$a(\gamma, z + \gamma') + a(\gamma', z) = a(\gamma', z + \gamma) + a(\gamma, z) + 2\pi\sqrt{-1}n(\gamma, \gamma'), \quad n(\gamma, \gamma') \in \mathbb{Z},$$

for all  $\gamma, \gamma' \in \Gamma$  and  $z \in C^n$ . We may assume that  $P = (I_n V)$  is a period basis of  $\Gamma$ . Putting

$$\ell(e_j, z) := \pi\sqrt{-1} \sum_{k=1}^n n(e_j, e_k)z_k, \quad j = 1, \dots, n,$$

$$\ell(v, e_j) := \ell(e_j, v) + 2\pi\sqrt{-1}n(v, e_j), \quad v \in V, j = 1, \dots, n,$$

we get a map  $\ell: \Gamma \times C^n \rightarrow C$  such that  $\ell(\gamma, \cdot): C^n \rightarrow C$  is  $C$ -linear and

$$(*) \quad \ell(\gamma, \gamma') = \ell(\gamma', \gamma) + 2\pi\sqrt{-1}n(\gamma, \gamma') \quad \text{for all } \gamma, \gamma' \in P.$$

We define

$$b(\gamma, z) := a(\gamma, z) - \ell(\gamma, z) \quad \text{for all } \gamma \in P.$$

We rewrite  $P = (I_n V) = (\gamma_1, \dots, \gamma_{n+q})$ . By (\*) we have

$$b(\gamma_i, z + \gamma_j) + b(\gamma_j, z) = b(\gamma_j, z + \gamma_i) + b(\gamma_i, z)$$

for all  $i, j = 1, \dots, n + q$ . By Lemma 1 there exists a summand of automorphy  $\tilde{b}$  such that

$$\tilde{b}(\gamma_i, z) = b(\gamma_i, z), \quad i = 1, \dots, n + q, z \in C^n.$$

Now we show the following (\*\*).

(\*\*) For any  $\gamma \in \Gamma$  there exists a constant  $c(\gamma) \in C$  such that

$$\tilde{b}(\gamma, z) = a(\gamma, z) - \ell(\gamma, z) + c(\gamma).$$

We use the proof of Lemma 1. Every  $\gamma \in \Gamma$  can be expressed uniquely as

$$\gamma = \sum_{i=1}^{n+q} t_i \gamma_i, \quad t_i \in \mathbf{Z}.$$

Let  $|\gamma| = \sum_{i=1}^{n+q} |t_i|$ . We show (\*\*) by induction on  $|\gamma|$ . For  $\gamma = 0$ , we have

$$a(0, z) - \ell(0, z) = a(0, z) = 2\pi\sqrt{-1}n_0, \quad n_0 \in \mathbf{Z}.$$

Then we set  $c(0) = -2\pi\sqrt{-1}n_0$ . When  $|\gamma| = 1$ ,  $\gamma = \pm\gamma_i$ . Since  $\tilde{b}(\gamma_i, z) = b(\gamma_i, z)$ , (\*\*) holds for  $\gamma = \gamma_i$  with  $c(\gamma_i) = 0$ . As in the proof of Lemma 1 ([13]),  $\tilde{b}(-\gamma_i, z)$  is defined by

$$\tilde{b}(-\gamma_i, z) = -\tilde{b}(\gamma_i, z - \gamma_i).$$

Hence it holds that

$$\tilde{b}(-\gamma_i, z) = -a(\gamma_i, z - \gamma_i) + \ell(\gamma_i, z - \gamma_i).$$

And we have

$$-a(\gamma_i, z - \gamma_i) = a(-\gamma_i, z) + 2\pi\sqrt{-1}\{n(\gamma_i, -\gamma_i) - n_0\}.$$

Setting  $c(-\gamma_i) = -\ell(\gamma_i, \gamma_i) + 2\pi\sqrt{-1}\{n(\gamma_i, -\gamma_i) - n_0\}$ , we obtain (\*\*). Assume that (\*\*) holds for  $|\gamma| \leq N$ . Let  $|\gamma| = N + 1$ . There exist  $\gamma', \gamma'' \in \Gamma$  such that  $\gamma = \gamma' \oplus \gamma''$ ,  $|\gamma'|, |\gamma''| \leq N$ . By the definition of  $\tilde{b}$  it holds that

$$\tilde{b}(\gamma, z) = \tilde{b}(\gamma', z + \gamma'') + \tilde{b}(\gamma'', z).$$

Then we obtain

$$\begin{aligned} \tilde{b}(\gamma, z) &= a(\gamma' + \gamma'', z) - 2\pi\sqrt{-1}n(\gamma', \gamma'') - \ell(\gamma' + \gamma'', z) - \ell(\gamma', \gamma'') \\ &\quad + c(\gamma') + c(\gamma''). \end{aligned}$$

Thus (\*\*) holds with

$$c(\gamma) = -2\pi\sqrt{-1}n(\gamma', \gamma'') - \ell(\gamma', \gamma'') + c(\gamma') + c(\gamma'').$$

We define the factor of automorphy by  $\tilde{\beta} := \exp \tilde{b}$ . Then the line bundle  $L_{\tilde{\beta}}$  given by  $\tilde{\beta}$  is topologically trivial (Vogt [13] and [14]). Put

$$\rho(\gamma, z) := \exp(\ell(\gamma, z) - c(\gamma)).$$

Then  $\rho$  is a theta factor and  $\alpha(\gamma, z) = \tilde{\beta}(\gamma, z)\rho(\gamma, z)$ . By Proposition 2 we obtain  $L \cong L_{\tilde{\beta}} \otimes L_{\rho}$ , where  $L_{\rho}$  is the line bundle given by  $\rho$ .

### § 3. Theta functions

**DEFINITION.** Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$ . A holomorphic function  $\theta(z)$  on  $C^n$  is called a theta function with theta factor  $\rho(\gamma, z)$  if it satisfies

$$\theta(z + \gamma) = \rho(\gamma, z)\theta(z), \quad \text{for all } \gamma \in \Gamma \text{ and } z \in C^n.$$

**PROPOSITION 4 (Kopfermann [7]).** *Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$  and  $\rho(\gamma, z)$  be a theta factor for  $\Gamma$  on  $C^n$ . Then there exist a hermitian form  $\mathcal{H}: C^n \times C^n \rightarrow C$  with  $\mathcal{A} := \text{Im } \mathcal{H}$   $\mathbf{Z}$ -valued on  $\Gamma \times \Gamma$ , a  $C$ -bilinear symmetric form  $\mathcal{Q}$ , a  $C$ -linear form  $\mathcal{L}$  and a semi-character  $\psi$  of  $\Gamma$  associated with  $\mathcal{A}|_{\Gamma \times \Gamma}$  such that*

$$\rho(\gamma, z) = \psi(\gamma) \exp 2\pi\sqrt{-1} \left[ \frac{1}{2\sqrt{-1}}(\mathcal{H} + \mathcal{Q})(\gamma, z) + \frac{1}{4\sqrt{-1}}(\mathcal{H} + \mathcal{Q})(\gamma, \gamma) + \mathcal{L}(\gamma) \right]$$

for all  $\gamma \in \Gamma$  and  $z \in C^n$ . If rank  $\Gamma = 2n$ , then this expression is unique.

A theta factor  $\rho$  with the expression as in Proposition 4 is called a

theta factor of type  $(\mathcal{H}, \psi, \mathcal{Q}, \mathcal{L})$ . A theta function with theta factor  $\rho$  of type  $(\mathcal{H}, \psi, \mathcal{Q}, \mathcal{L})$  is called a theta function of type  $(\mathcal{H}, \psi, \mathcal{Q}, \mathcal{L})$ . A trivial theta function is a theta function of type  $(0, 1, \mathcal{Q}, \mathcal{L})$ . A theta function of type  $(\mathcal{H}, \psi, 0, 0)$  is said to be reduced. Every theta function can be expressed as the product of a reduced theta function and a trivial theta function.

Let  $\mathcal{H}$  be a hermitian form on  $C^n \times C^n$ . We set

$$\text{Ker}(\mathcal{H}) := \{z \in C^n; \mathcal{H}(z', z) = 0 \text{ for all } z' \in C^n\}.$$

PROPOSITION 5 (Kopfermann [7]). *Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$ . If  $\theta(z)$  is a theta function for  $\Gamma$  of type  $(\mathcal{H}, \psi, \mathcal{Q}, \mathcal{L})$  and  $\theta(z) \not\equiv 0$ , then*

- (1)  $\mathcal{H}$  is positive semi-definite on  $C^n \times C^n$ ,
- (2)  $\theta$  is constant on  $\text{Ker}(\mathcal{H})$ , if  $\theta$  is reduced.

#### § 4. Quasi-abelian varieties

Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$  and  $C^n/\Gamma$  be an  $(H, C)$ -group. We consider the following condition.

- (R) There exists a hermitian form  $\mathcal{H}$  on  $C^n \times C^n$  such that
  - (1)  $\text{Im } \mathcal{H}$  is  $\mathbb{Z}$ -valued on  $\Gamma \times \Gamma$ ;
  - (2)  $\mathcal{H}$  is positive definite on  $C^n \times C^n$ .

When  $\text{rank } \Gamma = 2n$ , it is well-known that  $C^n/\Gamma$  is an abelian variety if and only if the above condition (R) is satisfied. The following definition is due to Gherardelli and Andreotti [2].

DEFINITION. An  $(H, C)$ -group  $C^n/\Gamma$  is called a quasi-abelian variety if it satisfies the condition (R).

PROPOSITION 6 (Gherardelli and Andreotti [2]). *Let  $\Gamma$  be a discrete subgroup of rank  $n + q$  in  $C^n$ . Suppose that  $\mathcal{H}$  is a hermitian form on  $C^n \times C^n$  satisfying the following properties:*

- (a)  $\text{Im } \mathcal{H}$  is  $\mathbb{Z}$ -valued on  $\Gamma \times \Gamma$ ,
- (b)  $\mathcal{H}$  is positive definite on  $C^n \times C^n$ .

*Then there exist  $\gamma \in C^n$  and a hermitian form  $\mathcal{Q}$  symmetric on  $\mathbb{R}^{n+q} \times \mathbb{R}^{n+q}$  such that*

- (1)  $\Gamma_1 = \Gamma + \mathbb{Z}\gamma$  is a discrete subgroup of rank  $n + q + 1$  in  $C^n$ ,
- (2)  $\text{Im}(\mathcal{H} + \mathcal{Q})$  is  $\mathbb{Z}$ -valued on  $\Gamma_1 \times \Gamma_1$ ,
- (3)  $\mathcal{H} + \mathcal{Q}$  is positive definite on  $C^n \times C^n$ .

Let  $C^n/\Gamma$  be a quasi-abelian variety with a discrete subgroup  $\Gamma$  of

rank  $n + q$ . Using Proposition 6 successively, we obtain a discrete subgroup  $\tilde{\Gamma}$  of rank  $2n$  such that  $\tilde{\Gamma} \supset \Gamma$  and  $C^n/\tilde{\Gamma}$  is an abelian variety. Hence the following proposition holds.

**PROPOSITION 7** (Gherardelli and Andreotti [2]). *If  $C^n/\Gamma$  is a quasi-abelian variety, then it is a covering space on an  $n$ -dimensional abelian variety.*

### § 5. $(H, C)$ -groups with positive line bundles

Let  $X$  be a complex manifold.  $X$  is called a weakly 1-complete manifold if there exists a  $C^\infty$  plurisubharmonic exhaustion function on  $X$  (Nakano [11]). It is well-known that an  $(H, C)$ -group  $C^n/\Gamma$  is weakly 1-complete (cf. Kazama [4]).

Let  $X$  be an  $n$ -dimensional weakly 1-complete manifold and  $\psi$  be its  $C^\infty$  plurisubharmonic exhaustion function. We set  $X_c = \{x \in X; \psi(x) < c\}$  for all  $c \in \mathbf{R}$ . Let  $E \rightarrow X$  be a holomorphic vector bundle over  $X$ . We denote by  $\Omega^p(E)$  the sheaf of germs of all  $E$ -valued holomorphic  $p$ -forms. We need the following theorems.

**THEOREM A** (Kazama [3]). *Let  $X$  be an  $n$ -dimensional weakly 1-complete manifold and  $E \rightarrow X$  be a holomorphic vector bundle over  $X$  which is positive in the sense of Nakano [10]. Then for any  $c \in \mathbf{R}$ , the restriction map*

$$\rho: H^0(X, \Omega^n(E)) \rightarrow H^0(X_c, \Omega^n(E))$$

*has a dense image with respect to the topology of uniform convergence on all compact sets in  $X_c$ .*

**THEOREM B** (Hironaka, cf. Fujiki [1]). *Let  $X$  be a weakly 1-complete manifold and  $B \rightarrow X$  be a positive line bundle over  $X$ . Then, for any  $c \in \mathbf{R}$  there exist natural numbers  $m, N$  and  $\varphi^{(0)}, \dots, \varphi^{(N)} \in H^0(X_c, \mathcal{O}(B^m))$  for  $d > c$  such that  $\Phi = (\varphi^{(0)}: \dots: \varphi^{(N)})$  embeds  $X_c$  into  $\mathbf{P}^N$  as a locally closed submanifold and  $\Phi^*[e] = B^m$ , where  $\mathbf{P}^N$  is the  $N$ -dimensional complex projective space and  $[e]$  is the hyperplane bundle of  $\mathbf{P}^N$ .*

**PROPOSITION 8.** *Let  $X = C^n/\Gamma$  be an  $(H, C)$ -group. Suppose that  $X$  has a positive theta bundle  $L \rightarrow X$  with theta factor  $\rho$ . Then, for any  $x, y \in C^n$  with  $x \not\equiv y \pmod{\Gamma}$  there exist a natural number  $m$  and theta functions,  $\theta_1, \theta_2$  with theta factor  $\rho^m$  such that  $f(x) \neq f(y)$ , where  $f = \theta_1/\theta_2$ .*

*Proof.* Let  $\pi: C^n \rightarrow X$  be the projection. For any  $x, y \in C^n$  with  $\pi(x)$

$\neq \pi(y)$ , there exists a real number  $c$  such that  $X_c \ni \pi(x), \pi(y)$ . Take  $c' > c$ . By Theorem B there exist natural numbers  $m, N$  and an embedding map  $\Phi: X_{c'} \rightarrow \mathbf{P}^N$ , where  $\Phi = (\varphi^{(0)}: \dots: \varphi^{(N)})$  and  $\varphi^{(j)} \in H^0(X_d, \mathcal{O}(L^m))$ ,  $d > c'$ . Since the canonical bundle  $K$  of  $X$  is analytically trivial, we have  $\mathcal{O}(L^m) \cong \Omega^n(K^{-1} \otimes L^m)$  and  $K^{-1} \otimes L^m$  is positive. Applying Theorem A to  $\Omega^n(K^{-1} \otimes L^m)$ , we can approximate any element in  $H^0(X_d, \mathcal{O}(L^m))$  by elements in  $H^0(X, \mathcal{O}(L^m))$  uniformly on  $\bar{X}_{c'}$ . Therefore there exist  $\tilde{\varphi}^{(0)}, \dots, \tilde{\varphi}^{(N)} \in H^0(X, \mathcal{O}(L^m))$  such that a holomorphic map  $\tilde{\Phi} = (\tilde{\varphi}^{(0)}: \dots: \tilde{\varphi}^{(N)}): X_c \rightarrow \mathbf{P}^N$  separates points  $\pi(x)$  and  $\pi(y)$ . There exist hyperplanes  $H_1$  and  $H_2$  of  $\mathbf{P}^N$  such that  $H_1 \ni \tilde{\Phi}(\pi(x)), H_1 \not\ni \tilde{\Phi}(\pi(y))$  and  $H_2 \ni \tilde{\Phi}(\pi(x)), H_2 \not\ni \tilde{\Phi}(\pi(y))$ . Let  $\ell_j = 0$  be the homogeneous equation of  $H_j$  for  $j = 1, 2$ . We set

$$f := \frac{\ell_1(\tilde{\varphi}^{(0)}, \dots, \tilde{\varphi}^{(N)})}{\ell_2(\tilde{\varphi}^{(0)}, \dots, \tilde{\varphi}^{(N)})}.$$

Then  $f$  is a meromorphic function on  $X$  and  $f(\pi(x)) \neq f(\pi(y))$ . Each section  $\ell_j(\tilde{\varphi}^{(0)}, \dots, \tilde{\varphi}^{(N)})$  corresponds to a theta function  $\theta_j$  with theta factor  $\rho^m$  for  $j = 1, 2$ . Then we have  $f \circ \pi = \theta_1/\theta_2$ .

**PROPOSITION 9.** *Let  $X = \mathbf{C}^n/\Gamma$  be an (H, C)-group. If  $X$  has a positive line bundle, then it is a quasi-abelian variety.*

*Proof.* Let  $L \rightarrow X$  be a positive line bundle over  $X$ . By Proposition 3, we have  $L \cong L_0 \otimes L_1$ , where  $L_0$  is a topologically trivial line bundle and  $L_1$  is a theta bundle. Under our assumption, every topologically trivial line bundle over  $X$  is given by a representation of  $\Gamma$  ([14]). Hence we may assume that  $L$  is a positive theta bundle. Suppose that  $L$  is given by a theta factor  $\rho$  of type  $(\mathcal{H}, \psi, \mathcal{L}, \mathcal{L})$ .

It suffices to show that  $\text{Ker}(\mathcal{H}) = \{0\}$ . If  $\text{Ker}(\mathcal{H}) \neq \{0\}$ , then there exist  $x, y \in \text{Ker}(\mathcal{H})$  such that  $x \neq y \pmod{\Gamma}$ . By Proposition 8 there exist a natural number  $m$  and theta functions  $\theta_1, \theta_2$  with theta factor  $\rho^m$  such that  $f(x) \neq f(y)$ , where  $f = \theta_1/\theta_2$ . We may assume that  $\theta_1$  and  $\theta_2$  are reduced theta functions of type  $(m\mathcal{H}, \psi^m, 0, 0)$ . Since  $\text{Ker}(m\mathcal{H}) = \text{Ker}(\mathcal{H})$ ,  $\theta_1$  and  $\theta_2$  must be constant on  $\text{Ker}(\mathcal{H})$  (Proposition 5). It is a contradiction.

From Propositions 7 and 9 the following theorem holds.

**THEOREM 1.** *Let  $X = \mathbf{C}^n/\Gamma$  be an (H, C)-group. Then the following statements (1), (2) and (3) are equivalent.*

- (1)  $X$  has a positive line bundle.

- (2)  $X$  is a quasi-abelian variety.  
 (3)  $X$  is a covering space on an  $n$ -dimensional abelian variety.

## §6. Some remarks

H. Kazama proposed the following problem of the structure of weakly 1-complete manifolds (in *Sūgaku* 32 (1980), Iwanami Shoten): Let  $X$  be a weakly 1-complete manifold. Is it possible to explain  $X$  by Stein manifolds and projective algebraic compact manifolds (for example, as a fibre space) when there exists a positive line bundle over  $X$  and  $H^0(X, \mathcal{O}) = \mathbb{C}$ ?

In this section we shall give the affirmative answer to the above problem in the case of  $(H, C)$ -groups.

Let  $X = \mathbb{C}^n/\Gamma$  be an  $(H, C)$ -group with rank  $\Gamma = n + q$ . If  $X$  is a quasi-abelian variety, there exists a hermitian form  $\mathcal{H}$  on  $\mathbb{C}^n \times \mathbb{C}^n$  such that

- (a)  $\text{Im } \mathcal{H}$  is  $\mathbb{Z}$ -valued on  $\Gamma \times \Gamma$ ,  
 (b)  $\mathcal{H}$  is positive definite on  $\mathbb{C}_\Gamma^p \times \mathbb{C}_\Gamma^q$ .

Let  $\mathcal{A} = (\text{Im } \mathcal{H})|_{\mathbb{R}_\Gamma^{n+q} \times \mathbb{R}_\Gamma^{n+q}}$ . Since  $\mathcal{A}$  is an alternating form, rank  $\mathcal{A}$  is an even number. We set rank  $\mathcal{A} = 2r$ . Then  $2q \leq 2r \leq n + q$ . The following definition is due to Gherardelli and Andreotti [2].

**DEFINITION.** When rank  $\mathcal{A} = 2(q + p)$ , we say that a quasi-abelian variety  $X = \mathbb{C}^n/\Gamma$  is of kind  $p$ .

Using the proof of Proposition 6 and a result of period basis of abelian variety, we obtain the following theorem.

**THEOREM 2** (Gherardelli and Andreotti [2]). *Let  $X = \mathbb{C}^n/\Gamma$  be a quasi-abelian variety of kind  $p$  with rank  $\Gamma = n + q$ . Then  $X$  is a fibre bundle over a  $(q + p)$ -dimensional abelian variety with fibres  $\mathbb{C}^p \times (\mathbb{C}^*)^{n-q-2p}$ .*

By Theorems 1 and 2, the problem given the beginning in this section is affirmative in the case of  $(H, C)$ -groups.

We do not know whether a weakly 1-complete manifold is globally embeddable in a complex projective space or not if it has a positive line bundle. But an  $(H, C)$ -group with positive line bundle is embedded in a complex projective space by Theorems 1 and 2.

## REFERENCES

- [ 1 ] Fujiki, A., On the blowing down of analytic spaces, Publ. RIMS Kyoto Univ., **10** (1975), 473–507.
- [ 2 ] Gherardelli, F. and Andreotti, A., Some remarks on quasi-abelian manifolds, Global analysis and its applications, Vol. II, 203–206, Intern. Atomic Energy Agency, Vienna, 1974.
- [ 3 ] Kazama, H., Approximation theorem and application to Nakano's vanishing theorem for weakly 1-complete manifolds, Mem. Fac. Sci. Kyushu Univ., **27** (1973), 221–240.
- [ 4 ] Kazama, H., On pseudoconvexity of complex abelian Lie groups, J. Math. Soc. Japan, **25** (1973), 329–333.
- [ 5 ] Kazama, H.,  $\bar{\partial}$  cohomology of  $(H, C)$ -groups, Publ. RIMS Kyoto Univ., **20** (1984), 297–317.
- [ 6 ] Kazama, H. and Umeno, T., Complex abelian Lie groups with finite-dimensional cohomology groups, J. Math. Soc. Japan, **36** (1984), 91–106.
- [ 7 ] Kopfermann, K., Maximal Untergruppen abelscher komplexer Liescher Gruppen, Schr. Math. Inst. Univ. Münster No. 29, 1964.
- [ 8 ] Morimoto, A., Non-compact complex Lie groups without non-constant holomorphic functions, Proc. of Conf. on Complex Analysis (Minneapolis 1964), Springer-Verlag, 1965, 256–272.
- [ 9 ] Morimoto, A., On the classification of noncompact complex abelian Lie groups, Trans. Amer. Math. Soc., **123** (1966), 200–228.
- [10] Nakano, S., On complex analytic vector bundles, J. Math. Soc. Japan, **7** (1955), 1–12.
- [11] Nakano, S., Vanishing theorems for weakly 1-complete manifolds, Number theory, algebraic geometry and commutative algebra, Kinokuniya, Tokyo, 1973.
- [12] Pothering, G., Meromorphic function fields of non-compact  $C^n/G$  Thesis, Univ. of Notre Dame, 1977.
- [13] Vogt, Ch., Geradenbündel auf toroiden Gruppen, Dissertation, Düsseldorf, 1981.
- [14] Vogt, Ch., Line bundles on toroidal groups, J. reine angew. Math., **335** (1982), 197–215.
- [15] Vogt, Ch., Two remarks concerning toroidal groups, Manuscripta Math., **41** (1983), 217–232.

*Department of Mathematics  
Toyama University  
Gofuku, Toyama, Japan*