

Advancements and developments in the detection and control of invasive weeds: a global review of the current challenges and future opportunities

Review

Cite this article: Roberts J and Florentine S (2024) Advancements and developments in the detection and control of invasive weeds: a global review of the current challenges and future opportunities. *Weed Sci.* **72**: 205–215. doi: [10.1017/wsc.2024.13](https://doi.org/10.1017/wsc.2024.13)

Received: 20 November 2023
Revised: 4 February 2024
Accepted: 15 February 2024
First published online: 29 February 2024

Associate Editor:

William Vencill, University of Georgia

Keywords:

Agricultural development; artificial intelligence; climate change; herbicide resistance; integrated systems; invasive species; weed control

Corresponding author:

Singarayer Florentine;
Email: s.florentine@federation.edu.au

Jason Roberts¹ and Singarayer Florentine² 

¹Research Assistant, Future Regions Research Centre, Institute of Innovation, Science and Sustainability, Federation University Australia, Ballarat, VIC, Australia and ²Professor, Future Regions Research Centre, Institute of Innovation, Science and Sustainability, Federation University Australia, Ballarat, VIC, Australia; School of Science, RMIT University, Melbourne, VIC, Australia

Abstract

Weed invasion has become increasingly recognized as a major threat to the practice of sustainable agriculture and the maintenance of natural ecosystems around the world. Without effective and ongoing management strategies, many weed species have the aggressive capacity to alter ecosystem functions and reduce the economic potential of the land into which they have been introduced. Although traditional weed management strategies can be useful in eliminating certain weeds, these approaches can be costly, economically damaging, and laborious and can result in variable long-term success. To further add to these challenges, several weed species have now developed resistance to a range of herbicide modes of action, which, to date, have been the major mechanism of weed control. As a result, it is anticipated that the use of emerging technology will help to provide a solution for the economical and environmentally sustainable management of various weeds. Of particular interest, emerging technology in the areas of weed detection and control (chemical, mechanical, electrical, laser, and thermal) has shown promising signs of improving long-term weed management strategies. These methods can also be assisted by, or integrated alongside, other technology, such as artificial intelligence or computer vision techniques for improved efficiency. To provide an overview of this topic, this review evaluates a range of emerging technology used for the detection and control of various weeds and explores the challenges and opportunities of their application within the field.

Introduction

The intrusion of weeds into agricultural and natural ecosystems is considered as a major driver toward agricultural production loss and biodiversity decline around the world (Kumar Rai 2022; Storkey et al. 2021). Weeds have the capacity to compete against and displace native or desirable species, and without deliberate and ongoing management interventions, they will continue to economically and environmentally degrade the land they have invaded (Kumar Rai 2022; Kumar Rai and Singh 2020). Exacerbating the urgency for efficient weed management, it is anticipated that the global population will reach 10 billion by 2050, and as a result, the global demand for agricultural products is expected to increase by more than 56% during this time (van Dijk et al. 2021). For the agricultural industry to meet such challenges, careful consideration regarding the most economical and environmentally sustainable production methods are required (Westwood et al. 2017). It has also been noted that the influence of climate change, which is likely to result in elevated atmospheric CO₂ levels, higher temperatures, and more variable weather events, will further add to the challenges, particularly in the area of weed management (Clements and Jones 2021; Malhi et al. 2021; Varanasi et al. 2016). These changes will likely impose stronger selection pressures on existing flora and further increase the expansion and impact of several weed species into regions where they may not have previously occurred (Beaury et al. 2020; Clements and Jones 2021; Ziska et al. 2019).

Although conventional weed control methods using herbicides or mechanical devices can provide some level of control, they often show variable success and require repeated, ongoing modifications to provide long-term success (Langmaier and Lapin 2020; Maqsood et al. 2020; Shahzad et al. 2021). A confounding issue is that, given the current reliance on chemical approaches for managing weeds, the repeated use of some herbicides can contribute to the development of herbicide resistance—there are now 269 species reported to have evolved resistance to 21 of the 31 known herbicide modes of action across 72 countries (Benbrook 2016; Heap 2023). Not only is the widespread use of chemicals adding to the potential development of herbicide resistance, but they can also contaminate and pollute the surrounding environment

© The Author(s), 2024. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



and waterways if they are not appropriately applied to the actively growing plant (Creech et al. 2017; Sankhla et al. 2016). To combat these concerns, innovations in weed management approaches are urgently required to assist land managers with more economical and environmentally sustainable methods of weed control. To address these concerns, it is suggested that the use of a range of emerging technologies will contribute to more sustainable weed management practices. In this regard, this review explores the use of emerging technology in the areas of weed detection and control (chemical, mechanical, electrical, laser, and thermal). It also identifies both the challenges and opportunities of where this technology can be applied in the field or integrated with other techniques such as artificial intelligence or computer vision methods. This information will be of value in identifying future directions and research opportunities in the field of emerging weed technology.

Artificial Intelligence in Weed Management

It is encouraging to recognize that the use of artificial intelligence has shown significant global potential in assisting the agricultural industry, particularly in the field of weed management (Amend et al. 2019; Costello et al. 2022; Fernandez-Quintanilla et al. 2018; Ghatrehsamani et al. 2023). Artificial intelligence can be described as an advanced machine learning system that can emulate the actions of humans, providing alternatives to constant and costly human intervention (Amend et al. 2019; Fernandez-Quintanilla et al. 2018; Ghatrehsamani et al. 2023; Partel et al. 2019). This form of technology has already provided several benefits in the area of weed management and is capable of further assisting, or being integrated with, machine learning systems and robotic devices for improved weed detection and control (Partel et al. 2019). As this area of research is increasingly developing and its full potential has yet to be discovered, the following sections of this review will highlight where artificial intelligence has the capacity to be used alongside a range of emerging technology to assist with improved weed management.

Weed Detection Methods

It is widely agreed that the early detection and control of a weed is a critical step within a weed management program to help reduce its long-term impact on the surrounding environment (Roslim et al. 2021). To assist in this task, recent developments have identified several data and imagery detection methods that can improve the time taken to identify and control a weed within the field (Esposito et al. 2021; Mohidem et al. 2021; Singh et al. 2020). Of the identified weed detection methods within the literature, the most promising that have been recently developed or that are undergoing further research for improvement include the use of (1) unmanned aerial vehicles, (2) all-terrain vehicles, (3) field robotics, (4) remote sensing, (5) proximal sensing, (6) plant signaling methods, and (7) crop genetic modification (Table 1). A common challenge when using certain detection methods such as unmanned aerial vehicles, all-terrain vehicles, and field robotics is their unintentional movement when capturing data or imagery. This movement, often caused by the wind or vehicle motion, increases the risk of motion blur, which can limit the quality of data and imagery and limit the success of deep learning models analyzing the data (Genze et al. 2023). To address this issue, research by Genze et al. (2023) has proposed the use of a deblurring segmentation model known as DeBlurWeedSeg, which has shown to successfully mitigate motion

blur when detecting weeds such as common lambsquarters (*Chenopodium album* L.) in grain sorghum [*Sorghum bicolor* (L.) Moench]. Although this model shows promising signs of reducing motion blur and increasing detection, further factors such as crop and weed height, the type of weed species, and environmental conditions all need to be considered due to their potential influence (Genze et al. 2023). If these factors can be accounted for, or if additional models, data, or imagery can be integrated, then it may be possible to mitigate the influence of motion blur, although further investigation into this combination across a range of crop-weed scenarios would be required.

A common aspect of these detection methods is their ability to capture high-quality data and imagery within the field (Coleman et al. 2023; Esposito et al. 2021; Lati et al. 2014; Mohidem et al. 2021; Pallottino et al. 2019; Su 2020; Sujaritha et al. 2017; Weiss et al. 2020). To achieve this, the use of specialized cameras systems or sensors are required, which will often include the use of hyperspectral imagery, multispectral imagery, red-green-blue or VIS (visible) imagery, satellite imagery, thermal imagery, and 3D stereo imagery (Esposito et al. 2021; Rosle et al. 2022; Su 2020; Xia et al. 2015; Table 2). The most commonly used system is the red-green-blue imagery system, as it is a low-cost operational option that can calculate different vegetation indices to distinguish between crop and weed species (Su 2020; Xia et al. 2015). Although this system has the capacity to effectively identify various weed species, it requires significant geometric plant data abilities to confidently recognize a range of species with high-level accuracy (Su 2020; Xia et al. 2015; Xu et al. 2021). To improve its efficiency in situations where crop and weed species appear geometrically similar, it is suggested that the use of multimodal information that uses red-green-blue imagery with depth information will help to improve weed detection accuracy (Xu et al. 2021, 2024). This approach has been referred to as the WeedsNet system, and Xu et al. (2024) demonstrate that the use of multimodal information has the potential to complement red-green-blue imagery in accurately detecting several grass and broadleaf weeds in a wheat (*Triticum aestivum* L.) crop. In this regard, such technology may have the potential to complement red-green-blue imagery systems in other agricultural settings, although this developing technology would require further detailed investigation in a range of other crop situations to determine its applicability and effectiveness in the field.

Another scheme known as the multispectral system has proven to be a powerful alternative to the red-green-blue imagery system, as it can capture and calculate a wider range of vegetation indices and spectral band wavelengths (Esposito et al. 2021; Lara et al. 2020; Lu et al. 2020). In a similar way, hyperspectral sensor camera systems are also powerful, being able to record thousands of radiometric narrow-band images from hundreds of spectral band wavelengths (Esposito et al. 2021; Su 2020). Although hyperspectral imagery has the potential to provide high accuracy in identifying several weeds, research has identified a range of inconsistencies, including insufficient feature extractions and calibration issues that limit its repeated accuracy (Diao et al. 2023; Peleg et al. 2005). It has also been noted that multispectral and hyperspectral systems often take a long time to calculate and collect data accumulation, as well as requiring specified [tailored] algorithms for analysis. As a consequence, there is an increase in the challenges and time taken to produce detailed precision weed maps (Zou et al. 2021). To attend to some of these issues, it has been suggested that the use of specialized models or machine learning techniques can provide further enhancement and

Table 1. Benefits, challenges and limitations of various weed detection methods

Detection method	What is it?	Benefits	Challenges/limitations	Reference
Unmanned aerial vehicles	The use of aerial devices to collect data and imagery, such as an aircraft or drone	<ul style="list-style-type: none"> • Can be used in difficult to access terrain • Can be automated with GPS or sensory system • Can be used in variable weather conditions • Quick method to cover large areas 	<ul style="list-style-type: none"> • Limited battery life • Prone to motion blur • Requires a skilled operator or GPS navigation system • Influenced by wind • May not be practical in all agricultural settings (e.g., orchards) 	Anderegg et al. 2023; Esposito et al. 2021; Mohidem et al. 2021
All-terrain vehicles	The use of a ground vehicle to collect data and imagery	<ul style="list-style-type: none"> • Data and imagery collected closer to the plant. • Can be used in variable weather conditions • Can be automated with GPS or sensory system 	<ul style="list-style-type: none"> • Limited access to certain areas • Slow process to cover a large area • Prone to motion blur • Limited movability around sensitive areas 	Coleman et al. 2023; Laursen et al. 2017
Field robotics	The use of robotic devices such as small autonomous vehicles	<ul style="list-style-type: none"> • Flexible movability (between crops or sensitive areas) • Can provide high-quality data and images (closer to the plant) • Can be used in variable weather conditions • Can be used alongside some control methods 	<ul style="list-style-type: none"> • Large devices have limited access to certain areas. • Limited battery life if not integrated with another power source • Some devices are bulky and expensive. • May diverge from a planned path if GPS systems or pathways are not accurate 	Francoeur-Leblond 2006; Sujaritha et al. 2017
Remote sensing	Classification of a plant/weed using distinct spectral signatures from satellite imagery or imagery data from a distance	<ul style="list-style-type: none"> • Provides accurate crop and weed density maps • Data can be obtained instantaneously (in the field of view). • Nondestructive method to monitor vegetation 	<ul style="list-style-type: none"> • Requires high-resolution data and satellite imagery • Timing of data collection is critical (may be influenced by weather patterns). 	Lamb and Brown 2001; Sishodai et al. 2020; Weiss et al. 2020; Xue and Su 2017
Proximal sensing	Classification of a plant/weed using field-based sensors close to the plant	<ul style="list-style-type: none"> • Data can be obtained almost instantaneously. • Can provide high spatial resolution • Provides data and imagery from different angles rather than just from above the plant 	<ul style="list-style-type: none"> • Variable lighting or weather conditions may influence results. • Spatial resolution will depend on the distance from and proximity of the sensor to the plant. 	Pallottino et al. 2019; Rañçon et al. 2023
Plant signaling methods	Uses exogenous fluorescent signals on crop plants to help differentiate them from other species	<ul style="list-style-type: none"> • Generally not influenced by environmental factors • Fluorescent markers naturally break down • Differentiates between crop plants and weeds 	<ul style="list-style-type: none"> • If a plant does not display high enough fluorescent markers, it may be identified as a weed. • Markers may break down before certain weeds emerge. 	Andújar et al. 2018; Longchamps et al. 2010; Su 2020
Crop Genetic Modification (e.g., crop coloring)	Genetically modifies a crop species, allowing it to express certain characteristics that can be detected more easily by imagery	<ul style="list-style-type: none"> • Increases imagery detection and classification between a crop and weed species • Consistent accuracy in detection 	<ul style="list-style-type: none"> • Limited research in this field for a range of crop species • Genetic modification of a crop may be restricted due to regulations. • If plants do not have the modification, they may be detected as a weed, resulting in crop loss. 	Lati et al. 2014

detection efficiency (Murad et al. 2023). In this respect, a range of techniques have been developed to assist in this area, with some examples including the use of (1) artificial intelligence-based image analysis (Aitkenhead et al. 2003; Haq et al. 2023); (2) deep learning systems and algorithms (such as artificial neural networks, convolutional neural networks, deep neural networks) (Hasan et al. 2021; Murad et al. 2023); (3) image processing techniques (including clustering, generative adversarial networks, Hilbert transformation, histograms of gradients, linear iterative, local binary patterns) (Murad et al. 2023; Nixon and Aguado 2019); and (4) machine learning systems and algorithms (such as adaptive

boosting, artificial neural networks, decision trees, k-nearest neighbor, and support vector machines) (Murad et al. 2023).

Chemical Control

Recent developments in the field of chemical control have shown promising signs in improving the management of a range of weed species around the world (Amend et al. 2019; Ghatrehsamani et al. 2023; Roslim et al. 2021; Table 3). One area in particular that has been shown to be of assistance in this field is the use of autonomous chemical control, which allows for a weed to be

Table 2. Benefits, challenges and limitations of various imagery and sensor systems for weed detection

Imagery system	What is it?	Benefits	Challenges/limitations	Reference
Red-green-blue or VIS (visible) imagery	Uses a digital camera that is equipped with a red, green, and blue filter with an image sensor	<ul style="list-style-type: none"> • Low operational cost • Easy to use and often readily available • Often a quick method of weed detection • Can be integrated with machine learning for improved accuracy 	<ul style="list-style-type: none"> • Requires significant geometric plant variations for accurate detection • May not be suitable for all agricultural settings (e.g., when crops appear similar to a weed species) 	Park et al. 2016; Su 2020; Xu et al. 2021
Hyperspectral imagery	Captures data and imagery from a series of signals with continuous channels of narrow spectral bands	<ul style="list-style-type: none"> • Rapid classification of a plant species • Can also be used for other purposes (e.g., mapping of soil conditions) • Records hundreds to thousands of radiometric bands • Can be integrated with machine learning systems 	<ul style="list-style-type: none"> • Can have complexities with radiometric calibration • Expensive to use • Often a much larger system that may require specialized devices for field application • Can only detect certain bands at a given time 	Che'Ya et al. 2021; Esposito et al. 2021; Lu et al. 2020; Peleg et al. 2005; Scherrer et al. 2019; Su 2020
Multispectral imagery	Captures data and imagery from discrete bands across a broad spectral signal	<ul style="list-style-type: none"> • Powerful operation that can calculate a wide range of vegetation indices • More powerful than red-green-blue imagery systems • Can be integrated with machine learning systems 	<ul style="list-style-type: none"> • Expensive to use • Often a much larger system that may require specialized devices for field application • Sometimes has a lower resolution than red-green-blue systems, therefore needs to be used closer to the ground surface (e.g., lower altitude using UAVs) 	Esposito et al. 2021; Lu et al. 2020; Mazzia et al. 2020
Satellite imagery	Captures images above the plants using satellites	<ul style="list-style-type: none"> • Quick method to collect data and imagery 	<ul style="list-style-type: none"> • Influenced by cloud or weather patterns • Does not provide the most detailed imagery or data 	Weiss et al. 2020
Thermal imagery	Captures imagery based on the surface temperature of a plant	<ul style="list-style-type: none"> • Can also be used for other purposes (plant stress, moisture stress, herbicide resistance) • Can be integrated with other methods such as remote sensing for improved accuracy 	<ul style="list-style-type: none"> • Not always consistent • Can be influenced by environmental factors • May only be suitable for certain situations 	Eide et al. 2021; Xu et al. 2023
3D stereo imagery	Uses a range of imagery (red-green-blue, LiDAR, spectroscopy, thermal, and ultrasound) to detect and weed species in three dimensions	<ul style="list-style-type: none"> • Capacity to create 3D models of plant characteristics • Can provide high-quality data and imagery with high resolution (can be expensive) 	<ul style="list-style-type: none"> • Relies on a range of imagery systems • Time and cost of this method needs further investigation. • Highly complex due to different plant structures, therefore further automatic programs needed 	Andújar et al. 2011, 2018

selectively targeted and sprayed, reducing off-target damage and the need for large-scale or widespread applications (Partel et al. 2019). These benefits can reduce excess herbicides leaching into the soil or into surrounding waterways, while also having the benefit of reducing the cost of materials and labor needed to treat a field each season (Balafoutis et al. 2017; Partel et al. 2019). While a range of autonomous chemical control options are currently available or under development for the control of various weed species (Table 1), it has been noted that most of these chemical devices have been tested or designed to be used within cropping systems, all of which generally have heterogeneous weed distributions that occur at a range of levels (Allmendinger et al. 2022). In this regard, this technology may be difficult to use within natural ecosystems and is problematic in difficult-to-access terrain or regions with varying levels of weed infestations. As such, the use of a range of weed detection methods and imagery or sensor systems discussed

earlier in this review may have the potential to assist in this area. Of particular interest, integrating such methods may enable more accurate detection and subsequent control of a weed in areas where it may appear geometrically similar to a crop or native species (Ghatrehsamani et al. 2023; Partel et al. 2019), although this area of research would require specific investigation relating to each scenario.

Herbicide-Resistance Management

It has been claimed that the use of artificial intelligence and specialized camera systems have the potential to assist in the field of herbicide-resistance management in agroecosystems (Amend et al. 2019; Ghatrehsamani et al. 2023; Picoli et al. 2017; Roslim et al. 2021). With the growing concern of more frequent occurrences of herbicide resistance, it is important to quickly

Table 3. Examples of autonomous chemical weed control devices reported within the global literature.^a

Product name	Brand/company	How does it work?
ARA	Ecorobotix	Uses Convolutional neural network (CNN)-based detection with multi-camera vision to spot spray weeds
Avirtech-MIMO AutoWeed	Avirtech AutoWeed	Uses 4D radar imaging and unmanned aerial vehicles to map and patch spray weeds Uses deep learning machine technology fitted with high-resolution cameras to detect and spot spray weeds
Bilberry	Bilberry	Uses red-green-blue imagery systems with an artificial intelligence-based detection system for spot spraying
EcoPatch FD20	Dimensions Agri Technologies Farmdroid	Uses red-green-blue imagery with an artificial intelligence-based detection system for spot spraying Uses Real time kinematics - Global Positioning System (RTK-GPS) systems to record the precise location of crop seeds and spot sprays weeds
Greeneye H-sensor	Greeneye Technology Agricon	Uses red-green-blue imagery with an artificial intelligence-based detection system for spot spraying Uses a bi-spectral camera with artificial intelligence-based detection to identify weeds
Kilter AX-1	Kilter Systems	Uses RTK-GPS to detect and selectively spray weeds
Ladybird	University of Sydney	Uses a hyperspectral and thermal camera to detect and sport spray weeds
Robotti	Agrointelli	Uses deep learning systems and RTK-GPS autonomous systems and a LiDAR camera
See and Spray	Blue River Technology	Uses red-green-blue imagery with a CNN-based detection method to spot spray weeds.
Smart Spraying	BASF, Bosch, Amazone	Uses bi-spectral cameras to identify weeds and spot sprays them
Weed-It	Weed-It	Uses a blue LED spectrometer to detect green vegetation
Weedseeker	Trimble Agriculture	Uses infrared sensor technology with a high-resolution spectrometer

^aSources: Allmendinger et al. (2022); Amend et al. (2019); Utstumo et al. (2018); Wu et al. (2020); Zhang et al. (2022).

identify and manage invasive populations before they reproduce to form a new generation of resistant plants. One method that has shown promise in the identification of herbicide-resistant plants is the use of thermal imagery assisted by machine learning systems (Eide et al. 2021; Picoli et al. 2017). Research has indicated that thermal imagery can assist in the detection of glyphosate-resistant plants, as treated plants often experience stress and an inhibition of stomatal conductance that leads to reduced photosynthetic rates and increased surface temperatures (Eide et al. 2021; Picoli et al. 2017; Shirzadifar et al. 2020). As a consequence, thermal imagery, assisted by machine learning techniques have the capability of identifying these plants within a field, which can then be controlled by other mechanisms (Eide et al. 2021; Shirzadifar et al. 2020). On the other hand, research by Eide et al. (2021) suggests that when thermal imagery is used in isolation, it is not a completely reliable indicator to predict herbicide resistance, particularly when dealing with herbicides with different modes of action. In this regard, further research into combining thermal imagery with other machine learning techniques may assist in identifying herbicide-resistant weed populations before they can reproduce. It has also been noted that integrating artificial intelligence in real-time image processing may help to assist in identifying herbicide-resistant and susceptible plants by identifying a range of plant and soil characteristics, although this area of research requires further investigation among different herbicides and crop-weed scenarios (Ghatrehsamani et al. 2023). It has also been noted that due to the limited number of new or emerging herbicide modes of action, the integrated use of existing bioherbicides could be another option to increase control efficiency and help reduce potential herbicide resistance in some weed species (Roberts et al. 2022).

Mechanical Control

Robotics and machine learning capabilities have shown promising signs as an emerging method to mechanically control a range of invasive weeds (Fennimore et al. 2016; Oliveira et al. 2021). For example, a study by Bakker et al. (2010) indicated that the use of intelligent autonomous weederers with real-time kinematics global positioning systems can be used to complete interrow hoeing within corn (*Zea mays* L.) crops, with minimal to no damage to the

surrounding crop plants. Although this method seems promising, the speeds of such devices are often much slower than traditional methods that often operate at a minimum speed of 4 km h⁻¹ or up to 12 km h⁻¹ for methods such as harrowing (Bakker et al. 2010; Bowman 2002). Due to these slower rates of speed, some robotic devices may not be able to completely bury or uproot certain weed species in a given time, ultimately allowing them to reestablish (Bakker et al. 2010). This clearly indicates that careful consideration of both the type of weed species and the speed of the mechanical device is needed for the implementation of these methods. Another study by Nørremark et al. (2012) showed 91% success rates in treating various interrow weeds with a robotic cycloid hoe, while research by Kunz et al. (2015) showed that the use of a camera-guided interrow hoeing device (Kult Robocrop®) reduced weed density by 89% in soybean [*Glycine max* (L.) Merr.] crops and 87% in sugar beet (*Beta vulgaris* L.) crops. It appears that camera-guided devices, fitted with machine learning capabilities, can be utilized to help guide the robotic device in targeting specific weeds within these areas, which can markedly improve these methods (Kunz et al. 2015). Such technology may also help to identify weeds that are growing very close to a crop plant and specifically target it or identify and map it for an alternative treatment to minimize potential crop damage.

A study by Van Evert et al. (2011) was associated with the development and testing of an autonomous robotic device to control broadleaf dock (*Rumex obtusifolius* L.) on a commercial farm. The device navigated using global navigation satellite system technology and included a downward-facing camera for plant detection and a mechanical tool with rotating blades that was lowered and activated once the plant was identified (Van Evert et al. 2011). This method was reported to have a 93% detection rate and a 75% success rate in the control of the species. It is anticipated that this method could be used to help decrease the quantity of herbicide needed to treat a field and reduce the costs associated with on-farm labor. Robotic weed control devices can also work in conjunction with other devices, with research by Noguchi et al. (2004) creating a primary and secondary system allowing for several devices to work together.

Another device that has been developed as an autonomous mechanical and chemical robot within the agricultural industry is

Table 4. Common electrical devices commercially developed around the world for weed control

Name	Brand/company	Areas used	Mode of action and use	Reference
Lightening Weeder	Lasco Inc.	United States	Uses spark discharge methods from electrodes mounted onto machinery, generally positioned slightly above the crop plants to target taller weeds	Vigneault et al. 1990
Weed Zapper™ (Annihilator and Terminator Series)	Old School Manufacturing	Canada, United States	Uses several electrodes mounted onto a flexible boom arm with horizontal electrodes above the crop canopy; allows models can be modified and configured for specific use	Schreier et al. 2022
XPower	Zasso™ (XPU and XPS)	Europe, South America	Uses continuous plant electrode contact and can be flexible with its use in a wide range of areas (e.g., under trees, vines or along roadsides)	Slaven et al. 2023
Rootwave™ (PRO and Top Fruit)	Rootwave™	Europe	Uses continuous plant electrode contact with the use of a handheld electric weeder for small spot-spray applications of weeds	Feys et al. 2023

the AgBotII. This device is capable of accurately identifying 90% of the selected species and has shown success in controlling various weeds such as wild oats (*Avena fatua* L.) and common sowthistle (*Sonchus oleraceus* L.). It is a powerful alternative, as it can provide both mechanical and chemical control, allowing an integrated control method, and alternately addressing control mechanisms with each pass (Fennimore et al. 2016; Oliveira et al. 2021). Another device that has been developed for the use of robotic weed control is the BoniRob platform produced by Bosch Deepfield Robotics, which has shown up to 94% success in controlling various weeds (Langsenkamp et al. 2014). This device is capable of identifying and selectively targeting a weed by means of mechanical control (Langsenkamp et al. 2014). Although promising, this technique can be time-consuming and may not be applicable for deeper or heavier clay-type soils, as it is more suitable for sandy light soils (Langsenkamp et al. 2014). In this regard, further investigation on a range of autonomous mechanical devices will need to consider more localized conditions to allow adjustment for greater efficacy in control options. Further investigation will also need to consider (1) which weed species can be mechanically controlled and which will require follow up or integrated methods of control; (2) the accuracy of control to limit potential off-target damage or unwanted soil disturbance, particularly when used alongside planted crops or native vegetation; (3) any potential spread of weeds and their propagules into other areas of the field if the devices are not cleaned or treated appropriately; and (4) the likely change to crop patterns or rows to help facilitate automated cultivation in two directions for improved efficiency (Fennimore et al. 2016; Sharma et al. 2017).

Electrical Weed Control

A range of research studies have shown that electrical weed control has the potential to successfully control various weeds, with several products already being commercially developed for use around the world (Table 4). This method delivers an electrical current to the targeted plant and can be applied by two main methods: (1) the spark-discharge method or (2) a continuous electrode-plant contact method (Slesarev 1972; Wilson and Anderson 1981). The spark-discharge method transfers a brief high-voltage current directly into a plant, while the continuous electrode-plant contact requires ongoing contact between the electrodes and the plant in order to allow a lethal dose of electricity to pass through the foliage, stems, and roots of the plant (Slesarev 1972; Wilson and Anderson

1981). These methods often cause damage to the plant's cells and structures by increasing temperature and vaporizing volatile liquids, ultimately damaging cell membranes (Slesarev 1972; Wilson and Anderson 1981). In some cases, electrical treatment may not completely kill a plant and may only cause damage in some areas, allowing the plant to regenerate over time (Slesarev 1972; Wilson and Anderson 1981). To achieve successful control, several factors need to be considered, such as (1) the type of device used and the amount of energy output for a lethal dose; (2) contact time with the plant; (3) surrounding vegetation, to ensure targeted plants are not shielded by other vegetation; (4) surrounding environmental conditions; and (5) the potential risk of fire (Bloomer et al. 2024; Landers et al. 2016; Lehnhoff et al. 2022; Slesarev 1972; Vigneault et al. 1990; Wilson and Anderson 1981). Of particular interest, research by Schreier et al. (2022) found a strong correlation between the increase of plant moisture content and the decreased level of weed control using electricity on several weed species such as barnyard grass [*Echinochloa crus-galli* (L.) P. Beauv.], common ragweed (*Ambrosia artemisiifolia* L.), giant foxtail (*Setaria faberi* Herrm.), giant ragweed (*Ambrosia trifida* L.), horseweed [*Coryza canadensis* (L.) Cronquist], waterhemp [*Amaranthus tuberculatus* (Moq.) Sauer], and yellow foxtail [*Setaria pumila* (Poir.) Roem. & Schult.], in a soybean-cropping system. In this regard, the use of electrical weed control needs to carefully consider any potential influence from the surrounding environment, as this is likely to influence the success of this method (Lati et al. 2021; Schreier et al. 2022).

It is clear that electric weed control can be suitable for a range of cropping systems, with research suggesting that it can be successfully used to control a diverse range of species (Bloomer et al. 2024; Landers et al. 2016). Research by Landers et al. (2016) in Brazil and Paraguay used a plant-electrode contact machine (16.6 km h⁻¹) and obtained between 94% to 100% control after 28 d for weeds such as high mallow (*Malva sylvestris* L.), smallflower galinsoga (*Galinsoga parviflora* Cav.), *S. oleraceus*, and wild poinsettia (*Euphorbia heterophylla* L.). On the other hand, only up to 75% success rates were realized when trying to control garden spurge [*Chamaesyce hirta* (L.) Millsp.], thus showing that certain weed species may respond differently to each treatment (Landers et al. 2016). Electrical weed control can also be used to reduce seed development, with an example of the Weed Zapper™ 6R30 (spark-discharge) achieving 54% to 80% reduction in *A. artemisiifolia*, *A. tuberculatus*, *A. trifida*, *S. faberi*, *S. pumila*, and common cocklebur (*Xanthium strumarium* L.) (Schreier et al. 2022).

Although there are several benefits of using electrical weed control methods, there are also several risks associated with its use (Bloomer et al. 2024). One drawback of using such methods is the low rate of speed and the considerable time needed to treat large fields (Llewellyn et al. 2016). A potential solution to this challenge would be to integrate this technology with artificial intelligence or machine learning devices to help identify and then treat weeds autonomously (Llewellyn et al. 2016; Machleb et al. 2020). It is also important to consider any potential resistance to electrical weed control (Beckie et al. 2020; Somerville et al. 2017). In particular, some species with higher levels of cellulose or lignin within their cell walls may have a higher resistance to bursting or becoming vaporized; likewise, plants with a hairy, thicker, or waxy epidermis may also be more protected from electrical control (Bauer et al. 2020; Vigneault et al. 1990). Plants that are not completely controlled may regenerate at a later time from their root systems; therefore, careful postcontrol monitoring may be required (Bloomer et al. 2024; Lehnhoff et al. 2022).

Research has suggested that the use of electrical weed control can also impact soil biota, as it may travel through a plant's root system and into the soil or adjacent water, thus potentially affecting the cellular constituents of surrounding organisms (Ruf et al. 2023). The effect of electrical weed control on the surrounding organism will depend on the voltage and length of the application (Ruf et al. 2023). For example, research by Lati et al. (2021) used an application of 0.05 Wh, which resulted in an increase of more than 40 °C in the shoots and roots of black nightshade (*Solanum nigrum* L.) and redroot pigweed (*Amaranthus retroflexus* L.), which is likely to influence the surrounding soil conditions. Ruf et al. (2023) also found a reduction in earthworm biomass within the top 25 cm of the soil profile after using the Zasso™ XPower XP300 applicator at 3 km h⁻¹ for 2 wk compared with an uncontrolled area. Research has also shown several physical changes to the earthworms from areas that have been treated with electrical weed control, such as a change in skin color, the presence of necrotic tissue, and damage to other cells (Ruf et al. 2023). On the other hand, contrasting results have shown that some macro- and mesofauna can survive in areas treated by electrical weed control and maintain more stable populations compared with other methods such as mechanical control where there are large soil disturbance events (Löbmann et al. 2022). Based on the differences in these findings, it is suggested that localized populations, environmental conditions, or soil types may play an important role in the response of soil biota, and therefore would require localized investigations to determine any long-term effects of this method.

Laser Weed Control

The combination of lasers and various weed detection systems has been shown to be a promising method in weed control, particularly in the early life-cycle stage (Heisel et al. 2002; Rakhmatulin et al. 2021; Wang et al. 2019). These devices have the capacity to be integrated with machine learning devices, including classification algorithms and automated devices for efficient weed control within agroecosystems (Ghatrehsamani et al. 2023; Rakhmatulin et al. 2021). The most commonly used lasers include carbon dioxide lasers, diode lasers, and fiber laser devices (Coleman et al. 2021; Gates et al. 1965; Heisel et al. 2002; Wöltjen et al. 2008). These devices work by emitting an infrared beam that is absorbed by the plant's cells, consequently burning them (Gates et al. 1965; Heisel et al. 2002). Several studies have shown that the use of lasers can be effective in controlling various weeds at different rates, with some

examples including *E. crus-galli* at 54 J per plant (Marx et al. 2012) and rigid ryegrass (*Lolium rigidum* Gaudin) at 76.4 J (Coleman et al. 2021). Several other species that have also been severely damaged or controlled using laser weed control include *A. fatua* (Bayramian et al. 1992), cereal rye (*Secale cereale* L.) (Bayramian et al. 1992), tobacco (*Nicotiana tabacum* L.) (Wöltjen et al. 2008), and water hyacinth [*Eichhornia crassipes* (Mart.) Solms; syn.: *Pontederia crassipes* (Mart.) Solms] (Couch and Gangstad 1974). A common trend within the literature shows that each weed species often requires a different contact time or energy dose to be controlled or partially damaged by the use of lasers (Marx et al. 2012; Rakhmatulin and Andreassen 2020). Research has also shown that many of these robotic devices, such as the commercially available LaserWeeder by Carbon Robotics, are often very expensive and very slow, only reaching speeds of 1.6 km h⁻¹ (Vijayakumar et al. 2023). For such applications to become more widely available, a time–cost–value analysis may be useful to determine whether this is the most appropriate method for a specific weed situation. Another important consideration regarding laser weed control is its influence on the surrounding soil biota, with long-term data on a range of environments required to examine this influence at a broader range (Khan et al. 2020).

Thermal Weed Control

The use of thermal weed control has been successfully integrated within the agricultural industry as a pre-crop emergence technique for weed control (Seaman 2016). Thermal weed control works by emitting quantities of intense heat directly to a targeted plant, which can increase its temperature and thus physically disrupt its cells (Brodie et al. 2019; Seaman 2016). Thermal weed control can often be in the form of a flame, hot oil, steam, or radiation (Brodie et al. 2019). Such technology has been implemented across various agroecosystems for the control of various weeds such as bermudagrass [*Cynodon dactylon* (L.) Pers.], *E. crus-galli*, hairy beggarticks (*Bidens pilosa* L.), ragweed parthenium (*Parthenium hysterophorus* L.), and several *Amaranthus* species (Mutch et al. 2008; Ulloa et al. 2012). Further research has also been used to trial the use of hot foam as a thermal weed control, which has shown success in controlling more than 75% of various species such as little mallow (*Malva parviflora* L.) and wild mustard (*Sinapis arvensis* L.) (Antonopoulos et al. 2023). The impact of thermal weed control is claimed to be more successful on annual species, with the effect on perennial species being more variable due to their more developed and structured root systems or underground rhizomes, which can reshoot if they are not completely damaged (Cisneros and Zandstra 2008). As a consequence, thermal weed control may not be suitable for all weed species and may thus result in surviving weed species creating monospecific stands, all of which can result in the need for further control and resources. For this method to be more widely used with confidence, further investigation on thermal weed control on a range of weed species across different environments would be needed.

The use of radiation-based thermal control, often referred to as microwave radiation, has shown promising results in the control of various weed species (Brodie 2012; Brodie et al. 2019). Under laboratory conditions, Aygun et al. (2017) have shown that microwave radiation energy has the capacity to kill various weed species such as *C. dactylon*, johnsongrass [*Sorghum halepense* (L.) Pers.], *S. nigrum*, and *X. strumarium*. Although this method successful for individual species, each species generally required a different speed and rate of microwave radiation for its effective

control (Aygun et al. 2017). Further research also suggests that different species respond differently to a changing application and rate of microwave radiation (Brodie and Hollins 2015; Kaçan et al. 2018). In particular, a study in Australia by Brodie and Hollins (2015) identified that wild radish (*Raphanus raphanistrum* L.) required at least 60 J/cm⁻² for 100% mortality, while *L. rigidum* required 370 J/cm⁻². Despite these successes, the use of microwave radiation energy for weed control often requires at least 10 times more energy than traditional methods such as chemical control, which therefore has currently limited its commercial use (Brodie 2012). One solution that could improve this method is its potential use alongside artificial intelligence and automated weed control devices. This could allow for plant detection and specific plant control using microwave energy to limit the need for its use across a large scale, although this method would require further investigation and an economic benefit analysis before it could be confidently used as a key weed control method.

Conclusion and Future Management Considerations

It is clear that the use of emerging technology is helping to shape the future of weed management and control around the world and has provided improved and sustainable alternatives for the detection and control of various weed species. Not only can these advancements in technology improve the way various weeds are controlled, but they can provide long-term economic and environmental benefits compared with traditional methods of weed control. However, despite their current success in agroecosystems, several areas of consideration need to be taken into account for their widespread and long-term application. Regarding the use of weed detection systems, some methods will need to be carefully chosen to ensure they are suitable to the specific environments where they are intended to be used. For example, unmanned aerial vehicles fitted with red-green-blue imagery may provide coverage across a large area, but they might not be suitable to all agricultural settings; for example, orchards or areas where crop and weed patterns appear geometrically similar would not be candidates for their use. In this case, the integration of other methods or imagery and sensor systems will need to be considered for improved accuracy. Artificial intelligence could also provide an additional level of support in classifying imagery. Regarding the use of chemical control and herbicide-resistance management, emerging technology in the detection of weeds has allowed for improved detection rates and subsequent control of a species before it has the ability to produce seeds and further impact the environment. Autonomous mechanical control has also been shown to be a feasible option in controlling various weeds as a nonchemical method of control. Despite promising signs, further investigation will need to consider the cost-benefit of using such devices, as they are often much slower and more expensive than conventional methods, which is also a consideration for electrical, laser, and thermal methods. It is suggested that this approach will become more practical and provide greater confidence in sound weed control strategies with the aid of machine learning systems and the integration of artificial intelligence systems.

Acknowledgments. This research received no specific grant from any funding agency or the commercial or not-for-profit sectors. Authors would like to acknowledge Peter Vamplew (Federation University) and Manzur Murshed (Deakin University) for the feedback on an earlier version of this article. The authors declare no competing interests.

References

- Aitkenhead M, Dalgetty I, Mullins C, McDonald A, Strachan N (2003) Weed and crop discrimination using image analysis and artificial intelligence methods. *Comput Electron Agric* 39:157–171
- Allmendinger A, Spaeth M, Saile M, Peteinatos GG, Gerhards R (2022) Precision chemical weed management strategies: a review and a design of a new CNN-based modular spot sprayer. *Agronomy* 12:1620–1641
- Amend S, Brandt D, Di Marco D, Dipper T, Gässler G, Höferlin M, Gohlke M, Kesenheimer K, Lindner P, Leidenfrost R, Michaels A, Mugele T, Müller A, Riffel T, Sampangi Y, Winkler J (2019) Weed management of the future. *Künstliche Intelligenz* 33:411–415
- Anderegg J, Tschurr F, Kirchgeßner N, Treier S, Schmucki M, Streir B, Walter A (2023) On-farm evaluation of UAV-based aerial imagery for season-long weed monitoring under contrasting management and pedoclimate conditions in wheat. *Comput Electron Agric* 204:107558–107571
- Andújar D, Calle M, Fernández-Quintanilla C, Ribeiro Á, Dorado J (2018) Three-dimensional modelling of weed plants using low-cost photogrammetry. *J Sens* 18:1077–1088
- Andújar D, Escolà A, Dorado J, Fernández-Quintanilla C (2011) Weed discrimination using ultrasonic sensors. *Weed Res* 51:543–547
- Antonopoulos N, Kanatas P, Gazoulis I, Tataridas A, Ntovakos D, Ntaoulis V-N, Zavra S-M, Travlos I (2023) Hot foam: evaluation of a new, non-chemical weed control option in perennial crops. *Smart Agric Technol* 3:100063–100072
- Aygun I, Cakir E, Kacan K (2017) Determination of possibilities of microwave application for killing weeds. *Int J Adv Sci Eng Technol* 5:33–36
- Bakker T, Asselt K, Bontsema J, Müller J, Straten G (2010) Systematic design of an autonomous platform for robotic weeding. *J Terramech* 47:63–73
- Balafoutis A, Beck B, Fountas S, Vangeyte J, Wal T, Soto I, Gómez-Barbero M, Barnes A, Eory V (2017) Precision agriculture technologies positively contributing to GHG emissions mitigation, farm productivity and economics. *Sustainability* 9:1339–1367
- Bauer M, Marx C, Bauer F, Flury D, Ripken T, Streit B (2020) Thermal weed control technologies for conservation agriculture—a review. *Weed Res* 60:241–250
- Bayramian AJ, Fay PK, Dyer WE (1992) Weed control using carbon dioxide lasers. *Proceedings of the Western Society of Weed Science* 45:55–56
- Beaury EM, Fusco EJ, Jackson MR, Lahinhad BB, Morelli TL, Allen JM, Pasquarella V, Bradley B (2020) Incorporating climate change into invasive species management: insights from managers. *Biol Invasions* 22:233–252
- Beckie HJ, Flower KC, Ashworth MB (2020) Farming without glyphosate?. *Plants* 9:1–15
- Benbrook CM (2016) Trends in glyphosate herbicide use in the United States and globally. *Environ Sci Eur* 28:1–15
- Bloomer DJ, Harrington KC, Ghanizadeh H, James TK (2024) Robotics and shocks: emerging non-herbicide weed control for vegetable and arable cropping. *NZ J Agric Res* 67:81–103
- Bowman G, ed (2002) *Steel in the Field: A Farmer's Guide to Weed Management Tools*. Sustainable Agriculture Network Handbook 2. Beltsville, MD: Sustainable Agriculture Network. Pp 128 p
- Brodie G (2012) Applications of microwave heating in agricultural and forestry-related industries. The development and applications of microwave heating. *IntechOpen* 1:45–78
- Brodie G, Hollins E (2015) The effect of microwave treatment on ryegrass and wild radish plants and seeds. *Glob J Agric Innov Res Dev* 2:16–24
- Brodie G, Khan MJ, Gupta D (2019) Microwave soil treatment and plant growth in sustainable crop production. Pages 1–18 in *Sustainable Crop Production*. IntechOpen
- Che'Ya NN, Dunwoody E, Gupta M (2021) Assessment of weed classification using hyperspectral reflectance and optimal multispectral UAV imagery. *J Agron* 11:1435–1451
- Cisneros JJ, Zandstra BH (2008) Flame weeding effects on several weed species. *Weed Technol* 22:290–295
- Clements DR, Jones VL (2021) Rapid evolution of invasive weeds under climate change: present evidence and future research needs. *Front Agron* 3:1–16

- Coleman GRY, Macintyre A, Walsh MJ, Salter WT (2023) Investigating image-based fallow weed detection performance on *Raphanus sativus* and *Avena sativa* at speeds up to 30 km h⁻¹. *Comput Electron Agric* 215: 108419–108430
- Coleman M, Kristiansen P, Sindel B, Fyfe C (2021) Wild Radish (*Raphanus raphanistrum*): Weed Management Guide for Australian Vegetable Production. Armidale, NSW: School of Environmental and Rural Science, University of New England. Pp 12 p
- Costello B, Osunkoya OO, Sandino J, Marinic W, Trotter P, Shi B, Gonzalez F, Dhileepan K (2022) Detection of parthenium weed (*Parthenium hysterophorus* L.) and its growth stages using artificial intelligence. *J Agric* 12:1838–1861
- Couch R, Gangstad E (1974) Response of water hyacinth to laser radiation. *Weed Sci* 22:450–453
- Creech CF, Henry RS, Werle R, Sandell LD (2017) Performance of postemergence herbicides applied at different carrier volume rates. *Weed Technol* 29:611–624
- Diao Z, Guo P, Zhang B, Yan J, He Z, Zhao S, Zhao C, Zhang J (2023) Spatial-spectral attention-enhanced Res-3D-OctConv for corn and weed identification utilizing hyperspectral imaging and deep learning. *Comput Electron Agric* 212:108092–108121
- Eide A, Koparan C, Zhang Y, Ostlie M, Howatt K, Sun X (2021) UAV-assisted thermal infrared and multispectral imaging of weed canopies for glyphosate resistance detection. *J Remote Sens* 13:4606–4624
- Espósito M, Crimaldi M, Cirillo V, Sarghini F, Maggio A (2021) Drone and sensor technology for sustainable weed management: a review. *Chem Bio Technol* 8:1–11
- Fennimore SA, Slaughter DC, Siemens MC, Leon RG, Saber MN (2016) Technology for automation of weed control in specialty crops. *Weed Technol* 30:823–837
- Fernandez-Quintanilla C, Pena J, Andujar D, Dorado J, Ribeiro A, Lopez-Granados F (2018) Is the current state of the art of weed monitoring suitable for site specific weed management in arable crops? *Weed Res* 58:259–272
- Feys J, De Cauwer B, Reheul D, Sciffer C, Clercx S, Palmans S (2023) Impact of electrocution on shoot and tuber vitality of yellow nutsedge (*Cyperus esculentus*). *J Agric* 13:696–712
- Francoeur-Leblond N (2006) Design of a Field Robot for Weeding Purposes. Ste-Anne-de-Bellevue, QC, Canada: Bioresource Engineering, McGill University. Pp 44 p
- Gates DM, Keegan HJ, Schleiter JC, Weidner VR (1965) Spectral properties of plants. *Appl Opt* 4:11–20
- Genze N, Wirth M, Schreiner C, Ajekwe R, Grieb M, Grimm DG (2023) Improved weed segmentation in UAV imagery of sorghum fields with combined deblurring segmentation model. *Plant Methods* 19:1–12
- Ghatrehsamani S, Jha G, Dutta W, Molaei F, Nazrul F, Fortin M, Bansal S, Debangshi U, Neupane J (2023) Artificial intelligence tools and techniques to combat herbicide resistant weeds—a review. *Sustainability* 15:1843–1861
- Haq SI, Tahir MN, Lan Y (2023) Weed detection in wheat crops using image analysis and artificial intelligence (AI). *Appl Sci* 13:8840–8863
- Hasan ASMM, Sohail F, Diepeveen D, Laga H, Jones MGK (2021) A survey of deep learning techniques for weed detection from images. *Comput Electron Agric* 184:106067–106118
- Heap I (2023) The International Herbicide-Resistant Weed Database. <https://www.weedscience.org/Home.aspx>. Accessed: October 10, 2023
- Heisel T, Andreasen C, Christensen S (2002) Sugar beet yield response to competition from *Sinapis arvensis* or *Lolium perenne* growing at three different distances from the beet and removed at various times during early growth. *Weed Res* 42:406–413
- Kaçan K, Çakir E, Aygün İ (2018) Determination of possibilities of microwave application for weed control. *Int J Agric Biol* 20:996–974
- Khan MJ, Jurburg SD, He J, Brody G, Gupta D (2020) Impact of microwave disinfection treatments on the bacterial communities of no-till agricultural soils. *Eur J Soil Sci* 71:1006–1017
- Kumar Rai P (2022) Environmental degradation by invasive alien plants in the Anthropocene: challenges and prospects for sustainable restoration. *Anthr Sci* 1:5–28
- Kumar Rai P, Singh JS (2020) Invasive alien plant species: their impact on environment, ecosystem services and human health. *Ecol Indic* 111: 106020–106040
- Kunz C, Weber J, Gerhards R (2015) Benefits of precision farming technologies for mechanical weed control in soybean and sugar beet—comparison of precision hoeing with conventional mechanical weed control. *Agronomy* 5:130–142
- Lamb DW, Brown RB (2001) PA—precision agriculture: remote-sensing and mapping of weeds in crops. *J Agric Eng Res* 78:117–125
- Landers JN, Challiol M, Vilela L, Lanz S (2016) Case study of zero-tillage organic soybean production in Brazil. *Soil Res* 54:166–172
- Langmaier M, Lapin K (2020) A systematic review of the impact of invasive alien plants on forest regeneration in European temperate forests. *Front Plant Sci* 11:1–15
- Langsenkamp F, Sellmann F, Kohlbrecher M, Kielhorn A, Strothmann W, Michaels A, Ruckelshausen A, Trautz D (2014) Tube Stamp for mechanical intra-row individual plant weed control. Pages 16–19 in *Proceedings of the 18th World Congress of CIGR, Beijing, China*
- Lara AEP, Pedraza C, Jamaica-Tenjo DA (2020) Weed estimation on lettuce crops using histograms of oriented gradients and multispectral images, pattern recognition applications in engineering. *IGI Global* 1: 204–228
- Lati RN, Filin S, Aly R, Lande T, Levin I, Eizenberg H (2014) Using genetically modified tomato crop plants with purple leaves for absolute weed/crop classification. *Pest Manag Sci* 70:1059–1065
- Lati RN, Rosenfeld L, David IB, Bechar A (2021) Power on! Low-energy electrophysical treatment is an effective new weed control approach. *Pest Manag Sci* 77:4138–4147
- Laursen MS, Jørgensen RN, Dyrmann M, Poulsen RN (2017) RoboWeedSupport—sub millimetre weed image acquisition in cereal crops with speeds up till 50 km/h. In *J Biol Biomol Agri Food Biotechnol Eng* 11:317–321
- Lehnhoff EA, Neher P, Indacochea A, Beck L (2022) Electricity as an effective weed control tool in non-crop areas. *Weed Res* 62:149–159
- Llewellyn R, Ronning D, Clarke M, Mayfield A, Walker S, Ouzman J (2016) Impact of Weeds on Australian Grain Production: The Cost of Weeds to Australian Grain Growers and the Adoption of Weed Management and Tillage Practices. Canberra, ACT: Grains Research and Development Corporation. Pp 112 p
- Löbmann A, Klauk B, Lang C, Petgen M, Petersen J (2022) Electrical weed control and its effect on soil organisms. Pages 277–282 in *Proceedings of the 30th German Workshop on Questions of Weed Biology and Control, Julius-Kühn-Archiv, Germany*
- Longchamps L, Panneton B, Samson G, Leroux GD, Thériault R (2010) Discrimination of corn, grasses and dicot weeds by their UV-Curr Robot Rep induced fluorescence spectral signature. *Precis Agric* 11:181–97
- Lu B, Dao PD, Liu J, He Y, Shang J (2020) Recent advances of hyperspectral imaging technology and applications in agriculture. *Remote Sens* 12: 2659–2703
- Machleb J, Peteinatos GG, Kollenda BL, Andujar D, Gerhards R (2020) Sensor-based mechanical weed control: Present state and prospects. *Comput Electron Agric* 176:105638–105650
- Malhi GS, Kaur M, Kaushik P (2021) Impact of climate change on agriculture and its mitigation strategies: a review. *Sustainability* 13:1318–1339
- Maqsood Q, Abbas RN, Iqbal MA, Serap KA, Iqbal A, El Sabagh A (2020) Overviewing of weed management practices to reduce weed seed bank and to increase maize yield. *Planta Daninha* 38:1–10
- Marx C, Barcikowski S, Hustedt M, Haferkamp H, Rath T (2012) Design and application of a weed damage model for laser-based weed control. *Biosyst Eng* 113:148–157
- Mazzia V, Comba L, Khaliq A, Chiaberge M, Gay P (2020) UAV and machine learning based refinement of a satellite-driven vegetation index for precision agriculture. *J Sens* 20:2530–2546
- Mohidem NA, Che'Ya NN, Juraimi AS, Illahi WFF, Roslim MHM, Sulaiman N, Saberioon M, Noor NM (2021) How can unmanned aerial vehicles be used for detecting weeds in agricultural fields? *J Agric* 11: 1004–1031

- Murad NY, Mahmood T, Forkan ARM, Morshed A, Jayaraman PP, Siddiqui MS (2023) Weed detection using deep learning: a systematic literature review. *J Sens* 23:3670–3715
- Mutch DR, Thalmann SA, Martin TE, Baas DG (2008) Flaming as a Method of Weed Control in Organic Farming Systems. East Lansing: Michigan State University Extension Bulletin E3038. Pp 4 p
- Nixon M, Aguado A (2019) Feature Extraction and Image Processing for Computer Vision. 4th ed. Cambridge, MA: Academic Press. Pp 360 p
- Noguchi N, Will J, Reid J, Zhang Q (2004) Development of a master-slave robot system for farm operations. *Comput Electron Agric* 44:1–19
- Norremark M, Griepentrog HW, Nielsen J, Søgaard HT (2012) Evaluation of an autonomous GPS-based system for intra-row weed control by assessing the tilled area. *Precis Agric* 13:149–162
- Oliveira LFP, Moreira AP, Silva MF (2021) Advances in agriculture robotics: a state-of-the-art review and challenges ahead. *J Robot Syst* 10:52–83
- Pallottino F, Antonucci F, Costa C, Bisaglia C, Figorilli S, Menesatti P (2019) Optoelectronic proximal sensing vehicle-mounted technologies in precision agriculture: a review. *Comput Electron Agric* 162:859–873.
- Park W, Pak S, Shim H, Lé HAN, Im M, Chang S, Yu J (2016) Photometric transformation from RGB Bayer filter system to Johnson-Cousins BVR filter system. *Adv Sci Res* 57:509–518
- Partel V, Charan Kakarla S, Ampatzidis Y (2019) Development and evaluation of a low-cost and smart technology for precision weed management utilizing artificial intelligence. *Comput Electron Agric* 157:339–350
- Peleg K, Anderson GL, Yang C (2005) Repeatability of hyperspectral imaging systems—quantification and improvement. *Int J Remote Sens* 26:115–139
- Picoli GJ, Carbonari CA, Matos AK, Rodrigues LF, Velini ED (2017) Influence of glyphosate on susceptible and resistant ryegrass populations to herbicide. *Planta Daninha* 35:1–12
- Rakhmatulin I, Andreassen C (2020) A concept of a compact and inexpensive device for controlling weeds with laser beams. *J Agron* 10:1616–1634
- Rakhmatulin I, Kamilaris A, Andreassen C (2021) Deep neural networks to detect and classify weeds from crops in agricultural environments in real-time: a review. *J Remote Sens* 13:4486–4510
- Rançon F, Keresztes B, Deshayes A, Tardif M, Abdelghafour F, Fontaine G, Da Costa JP, Germain C (2023) Designing a proximal sensing camera acquisition system for vineyard applications: results and feedback on 8 years of experiments. *J Sens* 23:847–105
- Roberts J, Florentine S, Fernando WGD, Tennakoon KU (2022) Achievements, developments and future challenges in the field of bioherbicides for weed control: a global review. *Plants* 11:2242–2260
- Rosle R, Sulaiman N, Che'Ya NN, Radzi MFM, Omar MH, Berahim Z, IAhi WFF, Shah JA, Ismail MR (2022) Weed detection in rice fields using UAV and multispectral aerial imagery. *Chem Proc* 10:44–55
- Roslim MHM, Juraimi AS, Che'Ya NN, Sulaiman N, Manaf MNHA, Ramli Z, Motmainna M (2021) Using remote sensing and an unmanned aerial system for weed management in agricultural crops: a review. *J Agron* 11:1809–1823
- Ruf T, Oluwaroye M, Leimbrock L, Emmerling C (2023) Field fodder conversion using electricity – Negative effects on earthworms and changes in labile carbon fractions. *Soil Tillage Res* 232:105746–105761
- Sankhla MS, Kumari M, Nandan M, Kumar R, Agrawal P (2016) Heavy metals contamination in water and their hazardous effect on human health—a review. *Int J Curr Microbiol Appl Sci* 5:759–766
- Scherrer B, Sheppard J, Jha P, Shaw JA (2019) Hyperspectral imaging and neural networks to classify herbicide-resistant weeds. *J Appl Remote Sens* 13:1–16
- Schreiber H, Bish M, Bradley K (2022) The impact of electrocution treatments on weed control and weed seed viability in soybean. *Weed Technol* 36:481–489
- Seaman A (2016) Production Guide for Organic Carrots for Processing. Geneva, NY: New York State Agricultural Experiment Station. Pp 51 p
- Shahzad M, Jabran K, Hussain M, Raza MAS, Wijaya L, El-Sheikh MA, Alyemeni MN (2021) The impact of different weed management strategies on weed flora of wheat-based cropping systems. *PLoS ONE*, 16:1–15
- Sharma DK, Tomar S, Chakraborty D (2017) Role of earthworm in improving soil structure and functioning. *Curr Sci* 113:1064–1071
- Shirzadifar A, Bajwa S, Nowatzki J, Bazrafkan A (2020) Field identification of weed species and glyphosate-resistant weeds using high resolution imagery in early growing season. *Biosyst Eng* 200:200–214
- Singh V, Rana A, Bishop M, Filippi AM, Cope D, Rajan N, Bagavathiannan M (2020) Chapter three—unmanned aircraft systems for precision weed detection and management: prospects and challenges. *Adv Agron* 159: 93–134
- Sishodia RP, Ray RL, Singh SK (2020) Applications of remote sensing in precision agriculture: a review. *J Remote Sens* 12:3136–3167
- Slaven MJ, Koch M, Borger CPD (2023) Exploring the potential of electric weed control: a review. *Weed Sci* 1–19
- Slesarev V (1972) Use of electric spark discharges in agriculture. *Zemledelie* 9:56–57
- Somerville GJ, Powles SB, Walsh MJ, Renton M (2017) Why was resistance to shorter-acting pre-emergence herbicides slower to evolve? *Pest Manag Sci* 73:844–851
- Storkey J, Mead A, Addy J, MacDonald AJ (2021) Agricultural intensification and climate change have increased the threat from weeds. *Glob Change Biol* 27:2416–2425
- Su W-H (2020) Crop plant signalling for real-time plant identification in smart farm: A systematic review and new concept in artificial intelligence for automated weed control. *Artif Intell Agric* 4:271–281
- Sujaritha M, Annadurai S, Satheshkumar J, Sharan SK, Mahesh L (2017) Weed detecting robot in sugarcane fields using fuzzy real time classifier. *Comput Electron Agric* 134:160–171
- Ulloa SM, Datta A, Bruening C, Gogos G, Arkebauer TJ, Knezevic SZ (2012) Weed control and crop tolerance to propane flaming as influenced by the time of day. *Crop Prot* 31:1–7
- Utstumo T, Urdal F, Brevik A, Dørum J, Netland J, Overskeid Ø, Berge TW, Gravdahl JT (2018) Robotic in-row weed control in vegetables. *Comput Electron Agric* 154:36–45
- Van Dijk M, Morley T, Rau ML, Saghai Y (2021) A meta-analysis of projected global food demand and population at risk of hunger for the period 2010–2050. *Nat Food* 2:494–501
- Van Evert FK, Samsom J, Polder G, Vijn M, Van Dooren HJ, Lamaker A, Van Der Heijden GWAM, Kempenaar C, Van Der Zalm T, Lotz LAP (2011) A robot to detect and control broad-leaved dock (*Rumex obtusifolius* L.) in grassland. *J Field Rob* 28:264–277
- Varanasi A, Prasad PVV, Jugulam M (2016) Chapter three—impact of climate change factors on weeds and herbicide efficacy. *Adv Agron* 135:107–146
- Vigneault C, Benoit DL, McLaughlin NB (1990) Energy aspect of weed electrocution. *Rev Weed Sci* 5:15–26
- Vijayakumar V, Ampatzidis Y, Schueller JK, Burks T (2023) Smart spraying technologies for precision weed management: a review. *Smart Agric Technol* 6:1–16
- Wang A, Zhang W, Wei X (2019) A review on weed detection using ground-based machine vision and image processing techniques. *Comput Electron Agric* 158:226–240
- Weiss M, Jacob F, Duveiller G (2020) Remote sensing for agricultural applications: a meta-review. *Remote Sens Environ* 236:111402–111421
- Westwood JH, Charudattan R, Duke SO, Fennimore SA, Marrone P, Slaughter DC, Swanton C, Zollinger R (2017) Weed management in 2050: perspectives on the future of weed science. *Weed Sci* 66:275–285
- Wilson RG, Anderson FN (1981) Control of three weed species in sugarbeets (*Beta vulgaris*) with an electrical discharge system. *Weed Sci* 29:93–98
- Wöltjen C, Haferkamp H, Rath T, Herzog D (2008) Plant growth depression by selective irradiation with CO₂ and diode laser. *Biosyst Eng* 110:316–324
- Wu X, Aravecchia S, Lottes P, Stachniss C, Pradalier C (2020) Robotic weed control using automated weed and crop classification. *J Field Rob* 37:322–40
- Xia C, Wang L, Chung B-K, Lee J-M (2015) In situ 3D segmentation of individual plant leaves using a RGB-D camera for agricultural automation. *J Sens* 15:20463–20479
- Xu B, Meng R, Chen G, Liang L, Lv Z, Zhou L, Sun Z, Zhao F, Yang W (2023) Improved weed mapping in corn fields by combining UAV-based spectral, textural and thermal measurements. *Pest Manag Sci* 79:2591–2602
- Xu K, Yuen P, Xie Q, Zhu Y, Cao W, Ni J (2024) WeedsNet: a dual attention network with RGB-D image for weed detection in natural wheat field. *Precis Agric* 25:460–485

- Xu K, Zhu Y, Cao W, Jiang X, Jiang Z, Li S, Ni J (2021) Multi-modal deep learning for weeds detection in wheat field based on RGB-F images. *Front Plant Sci* 12:1–10
- Xue J, Su B (2017) Significant remote sensing vegetation indices: a review of developments and applications. *J Sens* 2017:1–18
- Zhang W, Miao Z, Li N, He C, Sun T (2022). Review of current robotic approaches for precision weed management. *Curr Rob Rep* 3:139–151
- Ziska LH, Blumenthal DM, Franks SJ (2019) Understanding the nexus of rising CO₂, climate change, and evolution in weed biology. *Invasive Plant Sci Manag* 13:79–88
- Zou K, Chen X, Xiang F, Zhou H, Zhang CA (2021) A field weed density evaluation method based on UAV imaging and modified U-Net. *J Remote Sens* 13:310–329