

Research Article

Cite this article: Wawer J, Andrukajtis M, Karczewski J, Zielińska-Jurek A and Piątek R (2023). Fibrillar aggregates in powdered milk. *Journal of Dairy Research* **90**, 70–74. <https://doi.org/10.1017/S002202992300002X>

Received: 2 November 2021
Revised: 15 December 2022
Accepted: 18 December 2022
First published online: 25 January 2023

Keywords:

Amyloids; food safety; powdered milk; protein aggregation

Author for correspondence:

Jarosław Wawer, Email: jarwawer@pg.edu.pl

¹Department of Physical Chemistry, Faculty of Chemistry, Gdańsk University of Technology, Narutowicza Str. 11/12, 80-233 Gdańsk, Poland; ²Faculty of Applied Physics and Mathematics, Institute of Nanotechnology and Materials Science, Gdańsk University of Technology, Narutowicza Str. 11/12, 80-233 Gdańsk, Poland; ³Department of Process Engineering and Chemical Technology, Faculty of Chemistry, Gdańsk University of Technology, Narutowicza Str. 11/12, 80-233 Gdańsk, Poland; ⁴Department of Molecular Biotechnology and Microbiology, Faculty of Chemistry, Gdańsk University of Technology, Narutowicza Str. 11/12, 80-233 Gdańsk, Poland and ⁵BioTechMed Center, Gdańsk University of Technology, Narutowicza Str. 11/12, 80-233 Gdańsk, Poland

Abstract

This research paper addresses the hypothesis that powdered milk may contain amyloid fibrils. Amyloids are fibrillar aggregates of proteins. Up to this time, research on the presence of amyloids in food products are scarce. To check the hypothesis we performed thioflavin T fluorescence assay, X-ray powder diffraction, atomic force microscopy and fluorescence microscopy imaging. Our preliminary results show that commercially available milks contain fibrils that have features characteristic to amyloids. The obtained results can be interpreted in two opposite ways. The presence of amyloids could be considered as a hazard due to the fact that food products may induce amyloid related diseases. On the other hand, the presence of amyloids in traditionally consumed foodstuffs could serve as proof that fibrils of food proteins do not pose a threat for consumers.

Under unfavourable conditions, proteins can fold and aggregate abnormally, which can lead to the formation of amyloids. The monomeric protein macromolecules in amyloid fibrils are non-covalently bonded (Arnaudov and De Vries, 2005) and the resulting structures are rich in β -sheets (Sunde *et al.*, 1997). The presence of amyloids in the human body is associated with serious diseases including Alzheimer's disease, Parkinson's disease and spongiform encephalopathies (Chiti and Dobson, 2006). Amyloidogenesis appears to be universal process which results from an intrinsic property of the polypeptide chain (Chiti *et al.*, 1999).

Food processing conditions can destabilise the native conformation of proteins and thus may promote protein fibrillation (Jansens *et al.*, 2019; Lambrecht *et al.*, 2019; Monge-Morera *et al.*, 2020). So far, the presence of amyloids in common foodstuffs has been confirmed for only a few products. Probably, the most widely cited work in this matter is by Solomon *et al.* (2007). It provides the proof that *foie gras* contains amyloids and has amyloidogenic potential. Recently, it has been shown that amyloids can be formed during the boiling of egg white (Monge-Morera *et al.*, 2020). According to a study by Yoshida *et al.* (2009), renal and muscular tissues of aged cattle may also contain amyloids. The reports on detection of amyloids in food products are limited and thus more research are needed.

Although studies on the presence of amyloid in foods are scarce, efforts to obtain and test these aggregates from food proteins for potential use in the food industry are extensive (Mohammadian and Madadlou, 2018). Research such as this aims to improve the properties of the product or to give them new features (Cao and Mezzenga, 2019). Exceptionally large number of studies were performed for milk proteins. The comprehensive description of these works can be found in the review of Lambrecht *et al.* (2019). On the basis of experimental evidence, it is apparent that milk proteins readily form amyloids in-vitro (Al-Shabib *et al.*, 2019).

Taking all this into consideration, it would be interesting to address the hypothesis that amyloids are present in highly processed dairy products. In this work, we conducted tests to detect amyloids in powdered milks. For this purpose, we performed thioflavin T fluorescence assay, X-ray powder diffraction, atomic force microscopy and fluorescence microscopy imaging. The obtained results suggest that reconstituted powdered milk may contain amyloid-like fibrillar objects.

Materials and methods*Preparation of solutions*

Three brands of powdered milk were selected for the research. The composition of these products is presented in Table S1 in the online Supplementary File. The products were purchased

© The Author(s), 2023. Published by Cambridge University Press on behalf of Hannah Dairy Research Foundation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.






Fig. 1. Plots of the ThT fluorescence intensity against wavelength for samples of (a) milk with buffer and ThT (b) milk in buffer without ThT (c) buffer with ThT without milk. The fluorescence spectra are shown for Milk 1, Milk 2 and Milk 3 (counted from the left side).

from a local supermarket. Two brands of milk were intended for feeding infants after birth (milk 1 and 2) and one for feeding older babies over 12 months of age (milk 3). These products are legally classified as ‘foodstuffs for particular nutritional uses’. Milks 1 and 2 were, therefore, infant formulae, originally labelled by the producer as ‘infant milk’. Milk 3 was follow-on formula, originally labelled by the producer as ‘modified milk’. We chose these particular products for several reasons. This type of milk must be of the highest quality. Its composition must be well-balanced to meet the nutritional needs of young babies, and above all, the product must be safe to consume. It is obvious that infants are a very special and fragile group of individuals, and our work can help identify potential hazards, should they exist.

All powdered milks were produced by Europe-wide known manufacturers. The milks were reconstituted in accordance with the manufacturer’s instructions and further diluted for analysis. The deionised water used for the solutions preparation was filtered using 0.1 μm syringe filters (Minisart Cat. No. 16553).

Thioflavin T fluorescence assay (ThT assay)

The solution of the milk was diluted in the filtered phosphate buffer (50 mM, pH = 6.7) to a final concentration of 0.8 mg of protein per ml. The 2.5 mM Thioflavin T solution (Sigma-Aldrich Cat. No. T3516) was introduced to milk solution to obtain 25 μM ThT solution. The samples were analysed in a Jasco FP 8300 spectrofluorometer. The excitation wavelength was 440 nm and the emission spectra were collected from 460 to 550 nm. The excitation bandwidth was equal to 5 nm and the emission bandwidth was set to 10 nm. The scan speed was 200 nm/min, the data interval was equal 0.5 nm and the response time was set to 50 ms. The voltage of the photomultiplier was set to 370 V. Three spectra were averaged for each sample.

Atomic force microscopy (AFM)

The sample of milk was diluted with filtered water to a concentration of 0.05 mg of protein per ml. Fifty microlitres of this solution was placed on the freshly cleaved surface of the mica plate. After 5 min of incubation the plate was washed twice with 50 μl of filtered water and then dried and stored in desiccator. The AFM scans were collected using Nanosurf Easyscan 2 microscope (contact

mode). The speed of scan was equal 1.5 s per line, and the images with resolution 512 \times 512 px were recorded. The applied force was equal 20 nN. The data were analysed using Gwyddion 2.50 software.

Fluorescence microscopy (FM)

Fluorescence was recorded with the Olympus IX73 inverted fluorescence microscope equipped with the UCPlanFL N 20x/0.70 or LUCPlanFL N 40x/0.60 objectives, a Olympus U-HGLGPS fluorescence light source and a Hamamatsu Orcaflash 2.8 CMOS digital camera. Blue fluorescence of milk protein aggregates was generated using exciter-emitter filter set dedicated to DAPI (Semrock, IDEX Corporation). Images were recorded using the Olympus cellSens Dimension 1.18 software. The milk solutions were prepared as for the Thioflavin T fluorescence assay.

X-ray powder diffraction (XRD)

The structure of the samples and phase composition were investigated by X-ray powder diffraction (Cu K- α radiation, Rigaku MiniFlex 600 X-Ray diffractometer). The sample of powdered milk was deposited in the glass holder at room temperature, and crystal structures were determined from an XRD pattern measured in the range of $2\theta = 20\text{--}80^\circ$. The analysis and Rietveld refinements were performed with the HighScorePlus software package (Malvern Panalytical, Malvern UK) and the ICDD database.

Results and discussion

The measurement of the thioflavin T fluorescence (ThT assay) is considered as a gold standard for the detection of amyloids in the sample (Biancalana and Koide, 2010). The recorded emission spectra for the analysed solutions of the powdered milks are presented in Figure 1. The fluorescence intensities for the samples containing ThT and milk were significantly higher than intensities recorded for the milk in the buffer or the solution of ThT in the absence of milk. This increase is typical for the systems containing amyloid fibrils (Frare *et al.*, 2004). The positive results of the ThT assay have been previously reported for a raw milk (Lencki, 2007). The authors of the work cited above analysed the casein micelles using the transmission electron microscope.




Fig. 2. Atomic force microscopy height images of the analysed samples (a) milk 1 (b) milk 2 (c) milk 3. Scale bar 1 μm .

However, they did not show that the tested samples contained mature amyloids.

In the current work we analysed the structure of aggregates with atomic force microscopy (AFM). The obtained height images are presented in Figure 2. We observed fibrillar objects in all samples. The morphology of the fibrils was similar to amyloids formed from β -lactoglobulin (Bolisetty *et al.*, 2011) and α -lactalbumin (Antosova *et al.*, 2019) but different than for

κ -casein amyloids (Lee *et al.*, 2019). The objects found in samples of milk 1 and 2 were 0.5–2.5 μm long and 5–7 nm high. There were fewer fibrils in the sample of milk 3 and their morphology was different. These fibrils were longer (1.2–2.8 μm) and somewhat thicker (5–13 nm). Moreover, they did not form clusters and were rather straight. In all samples, small spherical objects were adsorbed to the fibrils. The aggregates reported by us have different morphology than objects presented in earlier research (Lencki, 2007; Qi, 2007) in which authors analysed the structure of the casein micelles.

The aggregates in samples of milks were visualised by fluorescence microscopy. We performed the analysis using Thioflavin T as a molecular probe and the samples were prepared as for the fluorescence spectroscopy. The obtained results are shown in Figure 3. In all samples small spherical objects were observed with size around 1–2 μm . Larger clusters with the diameter ranging from 10 μm to more than 20 μm were also present. These aggregates had a similar granular morphology and substructure composed of smaller objects. Unfortunately, the fluorescence microscopy does not have sufficient resolution to show the amyloids directly. However, the fact that the detected aggregates were fluorescent indicates that they may contain amyloid fibrils.

Additionally, the powdered milks were analysed using X-ray powder diffraction (see online Supplementary File materials and methods). The sharp diffraction peaks indicate that the samples contained well-ordered crystalline compounds. The recorded pattern can be interpreted as evidence of the presence of amyloids or may result from the presence of crystalline forms of low molecular weight compounds.

As mentioned in the introduction, the hypothesis that food products may contain amyloids has been presented before (Jansens *et al.*, 2019; Lambrecht *et al.*, 2019; Monge-Morera *et al.*, 2020). Our study confirms these predictions. It is possible that the fibrillar nano-structures discovered in the powdered milks were generated during food processing (Broersen, 2020) or were intentionally added by manufacturers. It also cannot be ruled out that the amyloids were introduced through the starting material. Therefore, it would be advisable to conduct additional research, preferably in the form of comparative studies. The raw milk and the powdered milk synthesised from it could be analysed to search for amyloid fibrils and verify this hypothesis.

The use of amyloids in the food industry raises concerns about their impact on consumers' health (Chaudhry *et al.*, 2008). Risk may result from several reasons:

- The amyloids of food proteins may be cytotoxic (Mocanu *et al.*, 2014)
- Amyloid seeds introduced to solution of protein monomers eliminate induction time of amyloidogenesis and thus the fibrillation may be significantly faster (Dubey *et al.*, 2014).
- Cross-seeding experiments showed that the introduction of pre-formed fibrils to various systems increased the risk of transformation of unlike proteins into amyloids (Dubey *et al.*, 2014).
- Amyloids from food proteins may have amyloidogenic potential. This risk was highlighted in *in-vivo* experiments (in mouse models) using amyloids isolated from tissues of aged cattle (Yoshida *et al.*, 2009) and *foie gras* (Solomon *et al.*, 2007). The other examples of the oral transmission of amyloid related diseases in animals can be found in published research by Murakami *et al.* (2014).
- Ingestion of products contaminated with amyloids may induce neurodegenerative diseases in humans.




Fig. 3. Fluorescence microscopy visualisation of free floating milk protein aggregates labelled by Thioflavin T: (a) milk 1, (c) milk 2 and (e) milk 3. The panels (b, d, f) represent phase-contrast images of corresponding fluorescence images. Scale bars correspond to 20 μm .

The transmissible spongiform encephalopathies are the most widely known amyloid-related diseases that might potentially be caused by consumption of contaminated food products. The family of these fatal disorders includes Creutzfeldt–Jakob disease and Kuru disease, the latter transmitted during cannibalistic rituals (Sikorska and Liberski, 2012). Worryingly, both these diseases have a long incubation period, years or even decades (Ghani *et al.*, 1998; Collinge *et al.*, 2008). This fact potentially makes it more difficult to take preventive action and limit damage to the population caused by infectious agents. The suggestion that diet may play an important role in development of protein misfolding diseases was formulated over 20 years ago (Cathcart and Elliott-Bryant, 1999) but currently remains a hypothesis. Our research can lead to the identification of additional dietary guidelines for people from at-risk groups.

The results obtained by us can be interpreted in two opposite ways. The presence of amyloids could be considered as a hazard

due to the fact that food products may induce amyloid related diseases. On the other hand, prevalence of amyloids in traditionally consumed foodstuffs could serve as a non-direct proof that fibrils of food proteins are safe and do not pose a threat for consumers. To resolve this dilemma more studies are needed (Cao and Mezzenga, 2019; Jansens *et al.*, 2019).

Another challenge is to determine how contaminants and legally added substances influence the amyloidogenesis of food proteins. For instance, milk and milk-based products may contain various additives (Huang *et al.*, 2014), each of which may alter the course of amyloid generation. Our previous works (Wawer *et al.*, 2019, 2021) show that fibrillation can be strongly affected by these types of compounds.

In conclusion, our data point to the presence of amyloid structures in powdered milks. The origin of these amyloids and the significance (if any) of their presence to consumers is still to be elucidated.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S002202992300002X>

Acknowledgement. Support from the Faculty of Chemistry of the Gdańsk University of Technology is acknowledged.

References

- Al-Shabib NA, Khan JM, Malik A, Sen P, Alsenaidy MA, Husain FM, Alsenaid AM, Khan RH, Choudhry H, Zamzami MA, Khan MI and Shahzad SA (2019) A quercetin-based flavanoid (rutin) reverses amyloid fibrillation in β -lactoglobulin at pH 2.0 and 358 K. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* **214**, 40–48.
- Antosova A, Bednarikova Z, Koneracka M, Antal I, Zavisova V, Kubovcikova M, Wu JW, Wang S-S and Gazova Z (2019) Destroying activity of glycine coated magnetic nanoparticles on lysozyme, α -lactalbumin, insulin and α -crystallin amyloid fibrils. *Journal of Magnetism and Magnetic Materials* **471**, 169–176.
- Arnaudov LN and De Vries R (2005) Thermally induced fibrillar aggregation of hen egg white lysozyme. *Biophysical Journal* **88**, 515–526.
- Biancalana M and Koide S (2010) Molecular mechanism of Thioflavin-T binding to amyloid fibrils. *Biochimica et Biophysica Acta – Proteins and Proteomics* **1804**, 1405–1412.
- Bolisetty S, Adamcik J and Mezzenga R (2011) Snapshots of fibrillation and aggregation kinetics in multistranded amyloid β -lactoglobulin fibrils. *Soft Matter* **7**, 493–499.
- Broersen K (2020) Milk processing affects structure, bioavailability and immunogenicity of β -lactoglobulin. *Foods (Basel, Switzerland)* **9**, 874.
- Cao Y and Mezzenga R (2019) Food protein amyloid fibrils: origin, structure, formation, characterization, applications and health implications. *Advances in Colloid and Interface Science* **269**, 334–356.
- Cathcart ES and Elliott-Bryant R (1999) Diet, amyloid enhancing factor (AEF) and amyloidogenesis: an hypothesis. *Amyloid: The International Journal of Experimental and Clinical Investigation: The Official Journal of the International Society of Amyloidosis* **6**, 107–113.
- Chaudhry Q, Scotter M, Blackburn J, Ross B, Boxall A, Castle L, Aitken R and Watkins R (2008) Applications and implications of nanotechnologies for the food sector. *Food Additives and Contaminants – Part A Chemistry, Analysis, Control, Exposure and Risk Assessment* **25**, 241–258.
- Chiti F and Dobson CM (2006) Protein misfolding, functional amyloid, and human disease. *Annual Review of Biochemistry* **75**, 333–366.
- Chiti F, Webster P, Taddei N, Clark A, Stefani M, Ramponi G and Dobson CM (1999) Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 3590–3594.
- Collinge J, Whitfield J, McKintosh E, Frosh A, Mead S, Hill AF, Brandner S, Thomas D and Alpers MP (2008) A clinical study of Kuru patients with long incubation periods at the end of the epidemic in Papua New Guinea. *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 3725–3739.
- Dubey K, Anand BG, Temgire MK and Kar K (2014) Evidence of rapid coaggregation of globular proteins during amyloid formation. *Biochemistry* **53**, 8001–8004.
- Frare E, Polverino De Lauro P, Zurdo J, Dobson CM and Fontana A (2004) A highly amyloidogenic region of hen lysozyme. *Journal of Molecular Biology* **340**, 1153–1165.
- Ghani AC, Ferguson NM, Donnelly CA, Hagenaars TJ and Anderson RM (1998) Epidemiological determinants of the pattern and magnitude of the vCJD epidemic in Great Britain. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **265**, 2443–2452.
- Huang Y, Min S, Duan J, Wu L and Li Q (2014) Identification of additive components in powdered milk by NIR imaging methods. *Food Chemistry* **145**, 278–283.
- Jansens KJ, Rombouts I, Grootaert C, Brijs K, Van Camp J, Van der Meeren P, Rousseau F, Schymkowitz J and Delcour JA (2019) Rational design of amyloid-like fibrillary structures for tailoring food protein techno-functionality and their potential health implications. *Comprehensive Reviews in Food Science and Food Safety* **18**, 84–105.
- Lambrecht MA, Jansens KJ, Rombouts I, Brijs K, Rousseau F, Schymkowitz J and Delcour JA (2019) Conditions governing food protein amyloid fibril formation. Part II: milk and legume proteins. *Comprehensive Reviews in Food Science and Food Safety* **18**, 1277–1291.
- Lee J, Lee JH, Paik SR, Yeom B and Char K (2019) Thermally triggered self-assembly of κ -casein amyloid nanofibrils and their nanomechanical properties. *Polymer* **179**, 121626.
- Lencki RW (2007) Evidence for fibril-like structure in bovine casein micelles. *Journal of Dairy Science* **90**, 75–89.
- Mocanu MM, Ganea C, Siposova K, Filippi A, Demjen E, Marek J, Bednarikova Z, Antosova A, Baran I and Gazova Z (2014) Polymorphism of hen egg white lysozyme amyloid fibrils influences the cytotoxicity in LLC-PK1 epithelial kidney cells. *International Journal of Biological Macromolecules* **65**, 176–187.
- Mohammadian M and Madadlou A (2018) Technological functionality and biological properties of food protein nanofibrils formed by heating at acidic condition. *Trends in Food Science and Technology* **75**, 115–128.
- Monge-Morera M, Lambrecht MA, Deleu LJ, Gallardo R, Louros NN, De Vleeschouwer M, Rousseau F, Schymkowitz J and Delcour JA (2020) Processing induced changes in food proteins: amyloid formation during boiling of hen egg white. *Biomacromolecules* **21**, 2218–2228.
- Murakami T, Ishiguro N and Higuchi K (2014) Transmission of systemic AA amyloidosis in animals. *Veterinary Pathology* **51**, 363–371.
- Qi PX (2007) Studies of casein micelle structure: the past and the present. *Le Lait* **87**, 363–383.
- Sikorska B and Liberski PP (2012) Human prion diseases: from Kuru to variant Creutzfeldt–Jakob disease. In Harris JR (Ed), *Protein Aggregation and Fibrillogenesis in Cerebral and Systemic Amyloid Disease*. Dordrecht Heidelberg, London, New York: Springer, pp. 457–496.
- Solomon A, Richey T, Murphy CL, Weiss DT, Wall JS, Westermark GT and Westermark P (2007) Amyloidogenic potential of foie gras. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 10998–11001.
- Sunde M, Serpell LC, Bartlam M, Fraser PE, Pepys MB and Blake CC (1997) Common core structure of amyloid fibrils by synchrotron X-ray diffraction. *Journal of Molecular Biology* **273**, 729–739.
- Wawer J, Szociński M, Olszewski M, Piątek R, Naczk M and Krakowiak J (2019) Influence of the ionic strength on the amyloid fibrillogenesis of hen egg white lysozyme. *International Journal of Biological Macromolecules* **121**, 63–70.
- Wawer J, Kaczowska E, Karczewski J, Augustin-Nowacka D and Krakowiak J (2021) Influence of stabilizing osmolytes on hen egg white lysozyme fibrillation. *Journal of Biomolecular Structure and Dynamics*. **40**, 13346–13353.
- Yoshida T, Zhang P, Fu X, Higuchi K and Ikeda SI (2009) Slaughtered aged cattle might be one dietary source exhibiting amyloid enhancing factor activity. *Amyloid: The International Journal of Experimental and Clinical Investigation: The Official Journal of the International Society of Amyloidosis* **16**, 25–31.