

AN EXAMPLE OF NORMAL LOCAL RING WHICH IS ANALYTICALLY RAMIFIED

MASAYOSHI NAGATA

Previously the following question was offered by Zariski [6]:

*Is any normal Noetherian local ring analytically irreducible?*¹⁾

In the present note, we will construct a counter-example against the question.

TERMINOLOGY. A ring (integrity domain) means always a commutative ring (integrity domain) with identity. A normal ring is an integrity domain which is integrally closed in its field of quotients. When \mathfrak{o} is an integrity domain, the integral closure of \mathfrak{o} in its field of quotients is called the derived normal ring of \mathfrak{o} .

In our treatment, some basic notions and results on general commutative rings and Noetherian local rings are assumed to be well known (see, for example, [5] and one of [1] or [2]). In particular, some results on regular local rings and completions of local rings are used freely (without references). On the other hand, we will make use of an example constructed in [3, §1] without proof.

§ 1. The construction of an example

Let \mathbf{k}_0 be a perfect field of characteristic 2 and let $u_0, v_0, \dots, u_n, v_n, \dots$ (infinitely many) be algebraically independent elements over \mathbf{k}_0 . Set $\mathbf{k} = \mathbf{k}_0(u_0, v_0, \dots, u_n, v_n, \dots)$. Further let x and y be indeterminates and set $\mathbf{r} = \mathbf{k}\{x, y\}$ (formal power series ring), $\mathfrak{o} = \mathbf{k}^2\{x, y\}[\mathbf{k}]$ and $c = \sum_{i=0}^{\infty} (u_i x^i + v_i y^i)$. Then we set $\mathfrak{s} = \mathfrak{o}[c]$.

PROPOSITION. \mathfrak{s} is a normal Noetherian local ring and the completion of \mathfrak{s} contains non-zero nilpotent elements (that is, \mathfrak{s} is analytically ramified).

Received January 10, 1955.

¹⁾ It was conjectured that the answer is negative by [4] and the present paper answers the conjecture affirmatively.

§ 2. Some preliminary results

LEMMA 1. *The ring \mathfrak{o} is a regular local ring with a regular system of parameters x, y . \mathfrak{r} is the completion of \mathfrak{o} .*

For the proof, see [3, § 1].

LEMMA 2. *An element $\sum a_{ij}x^i y^j$ ($a_{ij} \in \mathbf{k}$) is in \mathfrak{o} if and only if $[\mathbf{k}^2(a_{00}, a_{01}, a_{10}, \dots) : \mathbf{k}^2]$ is finite.*

Proof. $b = \sum a_{ij}x^i y^j$ is in \mathfrak{o} if and only if b is in $\mathbf{k}^2\{x, y\}[u_0, v_0, \dots, u_n, v_n]$ for some n . Therefore we see our assertion easily.

LEMMA 3. *Set $d_n = \sum_{i=0}^{\infty} u_{n+i}x^i$, $e_n = \sum_{i=0}^{\infty} v_{n+i}y^i$ ($n = 0, 1, \dots$). Then $\mathfrak{t} = \mathfrak{o}[d_0, e_0, \dots, d_n, e_n, \dots]$ is a normal ring.²⁾*

Proof. Let f be any element of the derived normal ring of \mathfrak{t} . Since f is in the field of quotients of \mathfrak{t} , f is expressed in the form $(p + qd_n + re_n + se_n d_n)/t$ ($p, q, r, s, t \in \mathfrak{o}$, $t \neq 0$) (because $\mathfrak{o}[d_0, e_0, \dots, d_n, e_n] = \mathfrak{o}[d_n, e_n]$ by the construction). Since p, q, r, s and t are in \mathfrak{o} , there exists an integer N which is not less than n such that the coefficients of them (as the power series in x and y) are in $\mathbf{k}^2(u_0, v_0, \dots, u_{N-1}, v_{N-1})$. Then since $d_n = u_n + u_{n+1}x + \dots + u_{N-1}x^{N-n-1} + x^{N-n}d_N$ and $e_n = v_n + \dots + v_{N-1}y^{N-n-1} + y^{N-n}e_N$, f is in the derived normal ring of $\mathfrak{o}^*[d_N, e_N]$, where $\mathfrak{o}^* = \mathbf{k}^2\{x, y\}[u_0, v_0, \dots, u_{N-1}, v_{N-1}]$ (because p, q, r, s and t are in \mathfrak{o}^* by our assumption and because the square of f is in $\mathbf{k}^2\{x, y\} \subseteq \mathfrak{o}^*$). Since the maximal ideal of \mathfrak{o}^* is generated by x and y , as is easily seen, \mathfrak{o}^* is a (complete) regular local ring. Since the residue class field of \mathfrak{o}^* is represented by $\mathbf{k}^2(u_0, v_0, \dots, u_{N-1}, v_{N-1})$ and since the leading forms of d_N and e_N are u_N and v_N (respectively), the maximal ideal of $\mathfrak{o}^*[d_N, e_N]$ is generated by x and y . Therefore $\mathfrak{o}^*[d_N, e_N]$ is a regular local ring and is a normal ring. Therefore f is in $\mathfrak{o}^*[d_N, e_N]$ and therefore f is in \mathfrak{t} . Thus we see that \mathfrak{t} is normal.

LEMMA 4. *$x\mathfrak{s}$ and $y\mathfrak{s}$ are prime ideals.*

Proof. $\mathfrak{s}/x\mathfrak{s}$ is isomorphic to $\mathbf{k}^2\{y\}[\mathbf{k}][e_0]$, which is an integrity domain. Therefore $x\mathfrak{s}$ is a prime ideal. That $y\mathfrak{s}$ is prime follows similarly.

LEMMA 5. *Let \mathfrak{s}' be the derived normal ring of \mathfrak{s} and let f be an element*

²⁾ By virtue of this result, we see easily that \mathfrak{t} is a regular local ring.

of \mathfrak{s}' . If xyf is in \mathfrak{s} , then f is in \mathfrak{s} .

Proof. Since \mathfrak{o} is Noetherian and since $\mathfrak{s} = \mathfrak{o}[c]$, \mathfrak{s} is Noetherian. Therefore if f is not in \mathfrak{s} , then one of the following must hold (see [5, § 8]): 1) $xy\mathfrak{s}$ has an imbedded prime divisor; 2) there exists at least one minimal prime divisor \mathfrak{p} of $xy\mathfrak{s}$ such that $\mathfrak{s}_{\mathfrak{p}}$ is not normal. Both are impossible because $x\mathfrak{s}$ and $y\mathfrak{s}$ are prime ideals by Lemma 4. Thus we see that f is in \mathfrak{s} .

§ 3. Proof of the proposition

As was noted above, \mathfrak{s} is Noetherian. Since \mathfrak{s} is isomorphic to $\mathfrak{o}[X]/g(X)\mathfrak{o}[X]$, where $g(X) = X^2 - c^2$, the completion of \mathfrak{s} is isomorphic to $\mathfrak{r}[X]/g(X)\mathfrak{r}[X]$ (because \mathfrak{r} is the completion of \mathfrak{o}). The residue class of $X - c$ is not zero and is nilpotent. Therefore the completion of \mathfrak{s} contains non-zero nilpotent elements. Now we will show that \mathfrak{s} is normal. Let f be any element of \mathfrak{s}' . Since \mathfrak{s} is contained in \mathfrak{t} (because $c = d_0 + e_0$) and since \mathfrak{t} is normal by Lemma 3, f is in \mathfrak{t} . Therefore f is of the form $\mathfrak{p} + qd_n + re_n + sd_n e_n$ ($\mathfrak{p}, q, r, s \in \mathfrak{o}$). Then $x^n y^n f$ is in $\mathfrak{o}[d_0, e_0]$. In order to show that f is in \mathfrak{s} , we have only to show that $x^n y^n f$ is in \mathfrak{s} by Lemma 5. Therefore we may assume that $n = 0$. Since f is in the field of quotients of \mathfrak{s} , f is of the form $(t + uc)/v$ ($t, u, v \in \mathfrak{o}$). Since $c = d_0 + e_0$, we see that $(t/v) + (u/v)d_0 + (u/v)e_0 = \mathfrak{p} + qd_0 + re_0 + sd_0 e_0$. Since 1, $d_0, e_0, d_0 e_0$ are linearly independent over \mathfrak{o} , we have $t/v = \mathfrak{p}, u/v = q (= r, s = 0)$. Therefore $f = \mathfrak{p} + qc$, which is in \mathfrak{s} . Therefore \mathfrak{s} is a normal ring.

BIBLIOGRAPHY

- [1] C. Chevalley, On the theory of local rings, *Ann. of Math.*, vol. **44** (1943), pp. 690-708.
- [2] I. S. Cohen, On the structure and ideal theory of complete local rings, *Trans. Amer. Math. Soc.*, vol. **59** (1946), pp. 54-106.
- [3] M. Nagata, Note on integral closures of Noetherian domains, *Memoirs Kyoto, Ser. A.*, vol. **28** (1953), pp. 121-124.
- [4] M. Nagata, On the theory of Henselian rings, II, *Nagoya Math. Journ.*, vol. **7** (1954), pp. 1-19.
- [5] M. Nagata, Basic theorems on general commutative rings, to appear in *Memoirs Kyoto, Ser. A.*
- [6] O. Zariski, Analytical irreducibility of normal varieties, *Ann. of Math.*, vol. **49** (1948), pp. 352-361.

*Mathematical Institute,
Kyoto University*