

with about a litre of the dry sand in which they had been living. We cannot determine if the bombyliid larvae at that time were in the sand or inside the antlion larvae. However, the antlion larvae looked and reacted normally, and were nearly fully grown when collected.

We thank Jack C. Hall, University of California, Riverside, and Norman D. Penny, California Academy of Sciences, San Francisco, for the identifications of the fly and the antlion, respectively. Graham C.D. Griffiths, University of Alberta, Paul H. Arnaud, Jr., California Academy of Sciences, and Hugh B. and Frances O. Leech are thanked for providing other pertinent information.

Cole, F.R., and E.I. Schlinger. 1969. *The Flies of Western North America*. University of California Press, Berkeley. xi + 693 pp., 360 figs.

Smith, R.C. 1934. Notes on the Neuroptera and Mecoptera of Kansas, with keys for the identification of species. *J. Kans. ent. Soc.* 7(4): 120–145, 10 figs.

(Date received: 7 April 1989; date accepted: 27 April 1989)

ERRATUM

Salom, S.M., and J.A. McLean. 1989. Influence of wind on the spring flight of *Trypodendron lineatum* (Oliver) (Coleoptera: Scolytidae) in a second-growth coniferous forest. *Can. ent.* 121: 109–119.

Corrections to Table 2, p. 115:

- (1) Expt. 1, Rep. 4, Distance (5 m): Flight direction is 105 ± 25 , not ± 2 .
- (2) Expt. 2, Rep. 1: Wind direction is NS instead of the blank space.
- (3) Expt. 3, Reps. 1, 2, and 3: Wind directions are 94, 117, and 95, respectively, instead of 274, 297, and 275, respectively.