

---

## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology,  
CAB International Institute of Parasitology,  
395A Hatfield Road,  
St Albans, Herts AL4 0XQ.  
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of Parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

# Contents

	Pages
<i>Dipetalonema viteae</i> and <i>Brugia pahangi</i> transplant infections in gerbils for use in antifilarial screening. J. P. COURT, J. N. STABLES, G. M. LEES, M. R. MARTIN-SHORT and R. RANKIN	1-9
Determination of the minimum time of praziquantel therapy required for the <i>in vitro</i> treatment of protoscoleces of <i>Echinococcus granulosus</i> . D. L. MORRIS, D. H. TAYLOR, D. DANIELS, E. M. RILEY and K. S. RICHARDS	10-14
<i>Fasciola hepatica</i> : development of monoclonal antibodies against somatic antigens and their characterization by ultrastructural localization of antibody binding. R. E. B. HANNA, A. G. TRUDGETT and A. ANDERSON	15-28
Transmission of <i>Schistosoma bovis</i> in Mkulwe (Mbozi District, Mbeya Region, Southern Highlands of Tanzania). J. A. MWAMBUNGU	29-32
<i>Schistosoma curassoni</i> Brumpt, 1931 and <i>S. bovis</i> (Sonsino, 1876) in cattle in northern Nigeria. G. T. NDIFON, C. BETTERTON and D. ROLLINSON (Research Note)	33-34
Comparative efficacy of flubendazole and mebendazole on encysted larvae of <i>Trichinella spiralis</i> (USA strain) in the diaphragm of mice and rats. J. MAKI and T. YANAGISAWA	35-39
Efficacy of various anthelmintics against third-stage larvae of <i>Ancylostoma caninum</i> in the brain of mice. G. M. BHOPALE and B. S. BHATNAGAR	40-44
A survey on the prevalence of gastrointestinal parasites of dogs in the area of Thessaloniki, Greece. S. T. HARALABIDIS, M. G. PAPAACHARIADOU, A. F. KOUTINAS and T. S. RALLIS (Research Note)	45-49
Eosinophilia in <i>Ascaris suum</i> -reinfected mice. K. SUGANE	51-57
Immunization against experimental rabbit cysticercosis using liposome-associated antigen preparations. P. S. CRAIG and O. ZUMBUEHL	58-62
Morphological changes in <i>Schistosoma mansoni</i> caused by the mutagen ethyl methane sulphonate. G. C. COLES and B. M. ZUCKERMAN	63-68
The tissue reactions of mice to infection with <i>Heligmosomoides polygyrus</i> . R. H. CYPESS, H. L. LUCIA, H. A. DUNSFORD and F. J. ENRIQUEZ	69-76
Cyclic and non-cyclic dynamics in populations of the helminth parasites of wood mice, <i>Apodemus sylvaticus</i> . S. S. J. MONTGOMERY and W. I. MONTGOMERY	78-90
A simple method for the oral transfer of <i>Moniliformis moniliformis</i> (Acanthocephala) between rats. R. C. STODDART and D. W. T. CROMPTON	91-94
Announcements	50, 77