

EPV1921

Regular Exercise Influences Sleep Patterns in Depressed Individuals

J. Kim^{1,1*}, S. Jaka², T. Shah², S. Gunturu¹ and O. El Shahawy²¹BronxCare Health System, Bronx and ²New York University Grossman School of Medicine, New York, United States

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2332


Introduction: Physical activity has been associated with improved sleep health, yet the specific impact on individuals with depressive disorders remains underexplored. This study aims to investigate the relationship between regular physical exercise and sleep duration in individuals diagnosed with a depressive disorder, utilizing data from the Behavioral Risk Factor Surveillance System (BRFSS). Understanding this relationship could inform integrative approaches to managing depressive symptoms and improving overall sleep health in this population.

Objectives:

- To analyze the impact of regular physical exercise on sleep duration among individuals with a depressive disorder.
- To assess the potential role of physical exercise in mitigating inadequate and prolonged sleep patterns within this population.

Methods: Data from the BRFSS for the years 2013, 2014, 2016, 2018, 2020, and 2022 were analyzed to explore the relationship between physical exercise and sleep duration in individuals with depressive disorders. The study included 518,214 participants who reported having a depressive disorder. Of these, 342,276 (weighted 68.5%) reported engaging in physical exercise within the past 30 days. Sleep duration was categorized, and regression analysis was used to assess the association between recent physical exercise and sleep duration.

Results: The analysis indicated that individuals with a depressive disorder who engaged in physical exercise in the past 30 days were less likely to experience inadequate sleep (6-7 hours, Odds Ratio [OR] = 0.83, $p < 0.05$) and more likely to achieve adequate sleep (7-9 hours, OR = 1.18, $p < 0.001$) compared to those who did not exercise. They were slightly less likely to have prolonged sleep (9-12 hours, OR = 0.9, $p < 0.05$). No significant associations were found for very inadequate sleep (< 6 hours, OR = 1.34) and very prolonged sleep (> 12 hours, OR = 0.6) with physical exercise ($p > 0.05$).

Image 1:

Conclusions: Regular physical exercise appears to be associated with better sleep outcomes in individuals with depressive disorders, particularly in promoting adequate sleep duration (7-9 hours). Exercise was linked to a reduced likelihood of inadequate sleep (6-7 hours) and a slight decrease in prolonged sleep (9-12 hours). However, no significant associations were found for very short (< 6 hours) or very prolonged (> 12 hours) sleep durations. These findings highlight the potential role of physical exercise in managing sleep health within this population. Incorporating regular physical activity into treatment plans for depression may improve sleep quality and contribute to better overall health outcomes. Further research is needed to understand the mechanisms and long-term effects of exercise on sleep patterns in those with depressive disorders.

Disclosure of Interest: None Declared

EPV1922

Better Sleep, Better Care: Streamlining Obstructive Sleep Apnea Screening in Psychiatric Outpatients

J. Kim^{1*}, A. Ramu¹, D. Khorasani¹, T. Kainth¹, A. Javadi¹ and S. Gunturu¹¹BronxCare Health System, Bronx, United States

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2333

Introduction: Obstructive Sleep Apnea (OSA) significantly complicates psychiatric conditions, yet its systematic screening within psychiatric settings is not common. We present a unique collaborative initiative within a tertiary care psychiatric hospital in South Bronx, USA, with approximately 3000 outpatients. We aim to bridge this gap by implementing a comprehensive OSA screening and referral process to improve patient outcomes.

Objectives:

- Implement a standardized Obstructive Sleep Apnea screening in psychiatric outpatient care.
- Enhance collaboration between psychiatry and sleep medicine for integrated care.
- Address compliance barriers by offering at-home sleep diagnostic options.

Methods: Starting in April 2024, patients in the Adult Outpatient Psychiatry Department are being screened for Obstructive Sleep Apnea using the STOP-Bang questionnaire, with this process ending in October 2024. High-risk patients identified through screening will receive WatchPAT Home Sleep Apnea Testing devices, with distribution and testing to be completed by December 2024. Data collected will include the number of patients screened, proportion identified as high-risk, HSAT completion rates, diagnostic outcomes, and subsequent referrals to sleep medicine services.

Results: The STOP-Bang questionnaire has been successfully integrated into routine clinical assessments for psychiatric outpatients, and the screening phase will conclude in October 2024. Preliminary data shows that a substantial number of patients were identified as high risk for Obstructive Sleep Apnea. The distribution of WatchPAT Home Sleep Apnea Testing devices to these high-risk individuals is underway and will be completed by December 2024. Initial results indicate effective triaging of patients and a high rate of compliance with at-home sleep testing. Detailed findings, including the exact number of patients screened, high-risk identification rates, HSAT completion rates, and diagnostic outcomes, will be presented upon project completion.