

**Demonstration of different  
mutational sites controlling rhamnose fermentation in FIRN  
and non-FIRN  $rha^-$  strains of *Salmonella typhimurium*:  
an essay in bacterial archaeology**

BY A. MORGENROTH\* AND J. P. DUGUID

*Bacteriology Department, University of Dundee*

(Received 27 October 1967)

I. INTRODUCTION

Strains of *Salmonella typhimurium* isolated from natural sources have been differentiated into a number of fermentation types, or 'biotypes', according to their ability to ferment rhamnose, inositol, xylose, glycerol, *d*-tartrate, *i*-tartrate, *l*-tartrate and citrate (Kristensen, Bojlén & Faarup, 1937; Harhoff, 1948; Kauffmann, 1954; Kallings & Laurell, 1957). Another differential character is fimbriation, i.e. the ability to form the non-flagellar filamentous appendages, fimbriae, described by Houwink & Van Iterson (1950), Brinton, Buzzell & Laufer (1954) and Duguid, Smith, Dempster & Edmunds (1955).

In a series of 775 wild-type strains of *S. typhimurium*, Duguid, Anderson & Campbell (1966) found a correlation between the characters of fimbriation and rhamnose fermentation. Out of 669 strains that were  $rha^+$ , i.e. fermented rhamnose in peptone water within 24 h at 37 °C, 663 were also  $fim^+$ , i.e. genotypically fimbriate; 335 of the  $rha^+$  strains, including four that were  $fim^-$ , were also  $inl^+$ , i.e. fermented inositol. On the other hand, 101 out of 106  $rha^-$  strains were  $fim^-$ ; 100 of these were  $fim^- inl^- rha^-$ , and such strains are designated 'FIRN' (fimbriation, inositol and rhamnose negative). Although they are thus correlated in their occurrence, the  $fim^+$  and  $rha^+$  characters are transferred independently in phage transductions (Duguid, Old & Hume, 1962; Old, 1963) and col-factor-mediated conjugations (Subbaiah & Stocker, 1962; Stocker, Smith & Subbaiah, 1963, cited from Sanderson & Demerec, 1965) and map in different positions on the chromosome.

Considerable epidemiological interest attaches to the question of whether the sites of a mutation in a phenotypically homogeneous group of strains isolated from natural sources are identical or different. If they are identical, the strains probably have a common origin and have descended from a single ancestral bacterium in which the mutation took place. If the sites are not identical, the strains probably have descended from different ancestral bacteria that underwent independent mutations, and so are unlikely to be closely related to one another. Evidence indicating whether the sites of a mutation in two phenotypically similar strains are different or identical may be obtained by the observation of whether or not the

\* Present address: Institut für medizinische Mikrobiologie und Immunologie der Universität, Bonn, Germany.

strains recombine to yield the wild type when they are crossed by generalized transduction with phage. Demerec *et al.* (1954) and Hartman (1956), in studies of galactose-negative or auxotrophic mutants of *S. typhimurium* strains LT2 and LT7, found that most pairs of strains with mutations in the same gene locus yielded galactose-positive or prototrophic recombinants when crossed by transduction. Their results showed that independently produced mutations generally give rise to non-identical alleles and that non-identical alleles are generally distinguishable by their ability to recombine in transductions.

The transduction method was first used to study the relationships between strains with mutational deficiencies acquired in the wild state, by Stocker & Edgar (1959). All of fifteen wild-type nicotinamide-requiring ( $\text{nic}^-$ ) strains of *S. typhimurium* phage-type 2, isolated in Britain, Australia, Italy and South Africa, were found by these workers to produce  $\text{nic}^+$  recombinants when transduced with phage A 3 grown on a  $\text{nic}^-$  laboratory mutant of *S. typhimurium* strain LT2; none of the fifteen wild-type strains produced  $\text{nic}^+$  recombinants when transduced with phage propagated on any of seven of these fifteen strains. The results suggested that the 15  $\text{nic}^-$  strains had mutations at identical sites and therefore that they had descended either from a common ancestral  $\text{nic}^-$  mutant bacterium or from different  $\text{nic}^-$  bacteria produced by independent mutations occurring at a single site specially liable to mutation ('hot spot').

Later, Duguid *et al.* (1962) and Old (1963) used the transduction method in a study of wild-type  $\text{fim}^-$  and  $\text{rha}^-$  strains of *S. typhimurium*. They obtained  $\text{fim}^+$  recombinants in crosses between FIRN and non-FIRN  $\text{fim}^-$  strains, but not in any cross between different FIRN strains. They concluded that the site of the  $\text{fim}^-$  mutation was probably identical in all strains of the FIRN group. Old obtained similar findings for the site of the  $\text{rha}^-$  mutation in the same series of FIRN strains. The combined results suggested that all FIRN strains have descended from a common ancestral FIRN bacterium.

The present paper describes the results of attempts to obtain  $\text{rha}^+$  recombinants by transduction in 23 FIRN ( $\text{fim}^- \text{inl}^- \text{rha}^-$ ) and 14 non-FIRN  $\text{rha}^-$  ( $\text{fim}^+ \text{inl}^+ \text{rha}^-$ ) wild-type strains of *S. typhimurium*. Most of the strains, twenty-one FIRN and nine non-FIRN  $\text{rha}^-$ , were different from those examined by Duguid *et al.* and Old. The purpose of the work was to determine whether the non-FIRN  $\text{rha}^-$  group of strains resembles the FIRN group in being homogeneous in respect of the sites of their  $\text{rha}^-$  mutations, and thus whether the non-FIRN  $\text{rha}^-$  strains, which were isolated in Britain, America and Australia, are likely to have descended from a common ancestor. Observations were also made on the patterns of  $\text{rha}^+$  mutation in the two groups.

## 2. MATERIALS AND METHODS

### (i) Strains

We studied twenty-three FIRN and fourteen non-FIRN  $\text{rha}^-$  strains of *Salmonella typhimurium* isolated from different sources. They were selected from a series under examination for biotype and phage-type by one of us (J.P.D.) and

Dr E. S. Anderson. Their particulars are shown in Table 1. Boyd's (1956) strain Q 1 (our SL 375) was used both as phage indicator and as wild-type rha<sup>+</sup> donor strain. We also examined three weakly rhamnase-fermenting (rha<sup>w</sup>) mutant strains derived in different experiments from FIRN strain 2306.

Table 1. *Strains of Salmonella typhimurium used as donors and recipients in the transduction experiments*

Group of strains	Strain no.	Source	Place of origin	Date of isolation	Phage-type	Biotype	
Non-FIRN rha <sup>-</sup> (fim <sup>+</sup> inl <sup>+</sup> rha <sup>-</sup> )	1179	Man	Australia	1960	U 128	19Xd	
	1180	Man	Australia	1960	27	19Xd	
	1541	Man	U.S.A.	1961	26	19Xd	
	1992	Man	Australia	1963	26	19Xd	
	2002	Man	Australia	1963	26	19Xd	
	2037	Man	Australia	1963	26	19Xd	
	2212	Man	England	1962	26	19Xd	
	2213	Man	England	1962	26	19Xd	
	2241	Pig	Wales	1962	25	19Xd	
	2323	Man	England	1963	23	19Xd	
	2368	Mutton	Australia	1964	26	19Xd	
	2382	Foodstuff	England	1964	U 38	19Xd	
	6768	Man	England	1958	31	19Xd	
	6899	Man	England	1958	U 58	19Xd	
	FIRN (fim <sup>-</sup> inl <sup>-</sup> rha <sup>-</sup> )	706	Guinea-pig	Sweden	1959	U 45	18
		735	Unknown	Unknown	.	1	17
1289		Man	England	1962	U 41	17	
1294		Man	England	1962	14a	17	
1723		Man	England	1963	14	17	
1804		Unknown	Netherlands	1960	26	17	
1877		Man	England	1964	NT	17	
1879		Man	England	1964	14	17	
1881		Man	England	1964	13	17	
2025		Man	Australia	1963	Un	15	
2205		Fowl	England	1962	14c	18	
2235		Man	England	1962	14b	17	
2306		Horse	England	1963	U 86	16	
2309		Canary	England	1963	29	17	
2321		Coypu	England	1963	26	17	
2322		Turkey	England	1963	U 86	17	
2333		Sheep	England	1963	U 86	17	
2540		Calf	Scotland	1965	U 128	17	
2555		Fowl	England	1965	U 131	17	
2565		Man	England	1965	U 84	16	
2571	Guinea-pig	England	1961	U 18	16		
2578	Man	England	1963	U 97	17		
2581	Bird	England	1964	U 151	17		
Rhamnase-positive (fim <sup>+</sup> inl <sup>+</sup> rha <sup>+</sup> )	SL 375	(Boyd's strain Q 1)	.	.	1	6	
Rhamnase-weak (fim <sup>-</sup> inl <sup>-</sup> rha <sup>w</sup> )	2306 rha <sup>w</sup> -1	(Spontaneous mutants derived from FIRN strain 2306)	.	.	.	.	
	2306 rha <sup>w</sup> -2		.	.	.	.	
	2306 rha <sup>w</sup> -3		.	.	.	.	

Phage-type NT = new type; Un = untypable. Biotype 19Xd is a new biotype; see text.

The phage-types of the strains were determined at the Enteric Reference Laboratory, Colindale, London. They included twelve of the types 1–31 described by Callow (1959) and eleven types (U 18–U 151) more recently distinguished in the ERL.

The biotypes were determined according to Harhoff's scheme as described by Kallings & Laurell (1957). The FIRN strains belonged to the established biotypes 15, 16, 17 and 18. These four types differ from the others in not fermenting either inositol or rhamnose, and they differ from one another in their reactions with *d*-tartrate, *i*-tartrate, citrate and glycerol. The non-FIRN rha<sup>-</sup> strains all belonged to a single biotype that is different from the twenty-one established ones. This new biotype is provisionally designated '19Xd', because it resembles biotype 19 in all characters except that it gives a positive Bitter xylose (X) and a negative *d*-tartrate (d) reaction. The Bitter fermentation test is done by culture in a liquid salts-ammonia medium with a very small content of peptone; a positive reaction is given by strains that ferment the sugar strongly and rapidly, and a negative reaction by weakly fermenting and non-fermenting strains. All the strains and biotypes used were prototrophic and capable of growth on an ammonia-salts minimal medium containing a utilizable source of carbon and energy.

The *transducing phage* was salmonella phage P 22 (PLT 22, Zinder & Lederberg, 1952), which is known to be a general transducing phage.

#### (ii) Culture media

*Nutrient broth* was Bacto heart infusion broth (Difco) adjusted to pH 7.2. Nutrient agar was this broth solidified with 1.5% Oxoid agar. In semi-solid agar the concentration of agar was 0.3%.

*Rhamnose minimal medium*. The basic minimal medium (Davis & Mingioli, 1950) contained: K<sub>2</sub>HPO<sub>4</sub>, 7 g; KH<sub>2</sub>PO<sub>4</sub>, 3 g; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 g; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.1 g; Oxoid Ionagar No. 2, 15 g; de-ionized water, 1000 ml. A separately sterilized solution of L(+)-rhamnose (British Drug Houses) was added to the basic medium to give a final concentration of 3 g rhamnose per 1000 ml medium.

*Enriched rhamnose minimal medium* was rhamnose minimal medium plus dehydrated nutrient broth, 0.1 g per 1000 ml.

*Rhamnose peptone water* used for fermentation tests was Oxoid peptone water plus L(+)-rhamnose, 10 g per 1000 ml., and bromocresol purple as indicator of pH (range 6.8–5.2). The rhamnose was sterilized separately and added to the sterile peptone water. The medium (pH 7.2) was dispensed in 3 ml volumes in 6 ml screw-capped bottles.

#### (iii) Culture methods

Incubation was aerobic and at 37 °C. Plates were plastic Petri dishes 9 cm in diameter containing 20 ml agar medium. Viable counts were done by spreading measured volumes of diluted culture on plates of nutrient agar. The diluting solution was 0.85% (w/v) NaCl to which 3% (v/v) of nutrient broth was added. Overnight (c. 18 h) broth cultures for experiments were inoculated from stock

cultures on Dorset egg slopes. Transduction and mutation plates were inoculated with 0.2 ml of phage-treated or untreated broth culture, which was spread as uniformly as possible over the whole surface of the plate and allowed to dry in before the start of incubation.

(iv) *Preparation of donor phages*

Donor phages were propagated by the soft agar layer method of Swanstrom & Adams (1951). About  $10^6$  phages and  $3 \times 10^8$  bacteria of the donor strain were mixed in 20 ml melted semi-solid agar at *c.* 46 °C and poured on to five well-dried nutrient agar plates. The plates were incubated at 37 °C and as soon as semi-confluent lysis was visible the five soft agar layers were scraped off and mixed together in 15 ml nutrient broth. The phage was freed from bacteria and agar debris by repeated centrifugation and then sterilized by the addition of 0.4 ml chloroform per 10 ml; the chloroform was removed after 15 min. To avoid errors arising from the persistence of small numbers of unadsorbed wild-type phage virions in the preparations of 'donor phage', the phage was passed a second time through the chosen experimental donor strain of bacteria and only this second lysate was used in transductions as donor phage.

The donor-phage preparations were titrated on lawns of the indicator bacterium SL 375 and each was diluted to a concentration of  $2 \times 10^{10}$  plaque-forming units per ml. Tests for sterility were made at the time of preparation and every 6 weeks thereafter by inoculation into broth and on to enriched rhamnose minimal medium. At the end of the series of experiments the titre of each phage preparation was found to be nearly at its original value.

(v) *Transduction experiments*

An overnight broth culture of the bacterial strain to be tested as recipient was diluted 1 in 50 in fresh broth and then incubated for 3½ h at 37 °C, so that it yielded about  $3.5 \times 10^8$  log.-phase bacteria per ml. One ml of this recipient culture was mixed with 0.1 ml of the donor phage preparation and the mixture was held for 12 min at 37 °C in a water bath to allow adsorption of the phage. The input ratio of phage virions to bacteria was about 6 to 1. Two 0.2 ml samples of the mixture, which together contained about  $1.3 \times 10^8$  phage-infected bacteria, were then spread over two plates of enriched rhamnose minimal medium (20 ml/plate). Similar samples of the recipient bacteria, grown in the same way but without the addition of phage, were inoculated on two other plates as controls, to demonstrate the frequency of mutation to *rha*<sup>+</sup>. The plates were incubated for 65 h at 37 °C and then scored for colonies of *rha*<sup>+</sup> recombinant or mutant bacteria. The long period of incubation was required because *rha*<sup>+</sup> bacteria grow relatively slowly on enriched rhamnose minimal medium and take about 48 h to form moderate-sized colonies.

*Confirmation of the identity of the rha*<sup>+</sup> *bacteria.* About 10% of all *rha*<sup>+</sup> colonies were subcultured on rhamnose minimal medium to confirm the genetic stability

of their acquired rha<sup>+</sup> character. A smaller proportion were tested for fermentation of inositol and agglutination with salmonella O 4 serum. Except for a few of them, whose appearance suggested that they were airborne contaminants, all showed the serological character of *S. typhimurium* and the inositol reaction of the recipient strain under test.

*Expression of results.* The average number of rha<sup>+</sup> colonies obtained per two plates in several repeated experiments with the same pair of donor and recipient strains is recorded as the result for the cross of that pair of strains. The value represents the number of rha<sup>+</sup> recombinant (or mutant) bacteria obtained per  $1.3 \times 10^8$  bacteria plated, i.e. the rate of apparent recombination. The mean values for groups of crosses are given as the number of rha<sup>+</sup> bacteria (colonies) obtained per one bacterium plated.

(vi) *Determination of the frequency of mutation to rha<sup>+</sup>*

The frequency of mutation of the rha<sup>-</sup> bacteria to the rha<sup>+</sup> genotype was observed in the control (no-phage) platings of the recipient bacteria in the transduction experiments and in further similar platings of the same bacteria. About  $1.3 \times 10^8$  bacteria from a  $3\frac{1}{2}$  h broth culture were inoculated per two plates of enriched rhamnose minimal medium and the plates were scored for colonies after 65 h. For comparison with the rate of recombination in the transductional platings, the mutant frequency is stated as the number of rha<sup>+</sup> bacteria (colonies) obtained per rha<sup>-</sup> bacterium plated. This value is probably considerably greater than the rate of mutation per bacterium per generation. Viable counts showed that the small amount of enriching broth in the minimal medium enabled the  $0.65 \times 10^8$  rha<sup>-</sup> bacteria inoculated on each plate to multiply about 50-fold in the first 24 h of incubation. Since the plates were scored for rha<sup>+</sup> colonies after 65 h and since rha<sup>+</sup> bacteria can form fair-sized colonies within 48 h on rhamnose minimal medium, it is probable that any rha<sup>+</sup> mutants arising from the bacterial population present at 24 h would have formed recognizable colonies within 65 h. The number of rha<sup>+</sup> colonies obtained per rha<sup>-</sup> bacterium plated at 0 h was therefore divided by 50 to give an estimate of the number of rha<sup>+</sup> mutant bacteria produced per rha<sup>-</sup> bacterium present at 24 h. This latter rate is taken as possibly approximating to the mutation rate per bacterium per generation.

### 3. RESULTS

(i) *Distinction of rha<sup>+</sup> transduction from rha<sup>+</sup> back-mutation*

The occurrence of recombination in transductions between different rha<sup>-</sup> strains of *Salmonella typhimurium* was determined by observation of the development of rha<sup>+</sup> colonies in platings of the transduction mixture on enriched rhamnose minimal medium. After incubation for 65 h the rha<sup>+</sup> colonies were seen in clear contrast against the thin film of growth of parental rha<sup>-</sup> bacteria supported by the small amounts of enriching broth contained in the medium and added with the inoculum. Evidence that the colony-forming rha<sup>+</sup> bacteria had arisen by

recombination, and not by spontaneous back-mutation, was provided by the demonstration that the recipient-strain bacteria either did not mutate to the *rha*<sup>+</sup> genotype in control platings without phage or did so only at a rate too low to account for the number of colonies obtained in the transduction platings. Since all the twenty-three FIRN and fourteen non-FIRN *rha*<sup>-</sup> strains were able to grow well on minimal medium containing glucose instead of, or in addition to, rhamnose, the absence of growth in transduction or control platings on the rhamnose medium could be attributed to absence of recombinant or mutant bacteria capable of utilizing rhamnose (i.e. *rha*<sup>+</sup>).

Table 2. Frequency of *rha*<sup>+</sup> mutants in control (no phage) platings of 14 non-FIRN *rha*<sup>-</sup> and 23 FIRN strains of *Salmonella typhimurium* on enriched rhamnose minimal medium

Strain(s) plated	No. of platings of $1.3 \times 10^8$ bacteria	No. of <i>rha</i> <sup>+</sup> colonies obtained in all platings	Mutant frequency expressed as no. of <i>rha</i> <sup>+</sup> colonies per bacterium plated
<b>Non-FIRN <i>rha</i><sup>-</sup></b>			
1179	60	2	$2.6 \times 10^{-10}$
1180	24	1	$3.2 \times 10^{-10}$
1541	20	2	$7.7 \times 10^{-10}$
1992	20	2	$7.7 \times 10^{-10}$
2002	28	1	$2.8 \times 10^{-10}$
2037	60	0	$< 1.3 \times 10^{-10}$
2212	20	1	$3.8 \times 10^{-10}$
2213	36	1	$2.1 \times 10^{-10}$
2241	45	4	$6.8 \times 10^{-10}$
2323	24	2	$6.4 \times 10^{-10}$
2368	58	1	$1.3 \times 10^{-10}$
2382	20	1	$3.8 \times 10^{-10}$
6768	34	2	$4.5 \times 10^{-10}$
6899	20	2	$7.7 \times 10^{-10}$
All 14 strains	469	22	$3.6 \times 10^{-10}$
<b>FIRN</b>			
2306	38	0	$< 2.0 \times 10^{-10}$
2309	30	0	$< 2.6 \times 10^{-10}$
Each other strain	20	0	$< 3.8 \times 10^{-10}$
All 23 strains	488	0	$< 1.6 \times 10^{-11}$

(ii) Frequency of mutation to *rha*<sup>+</sup>

*Non-FIRN rha*<sup>-</sup> strains. This group of fourteen strains, all of which were *fim*<sup>+</sup> *int*<sup>+</sup> *rha*<sup>-</sup> and biotype 19Xd, were tested in a total of 469 double-platings of  $1.3 \times 10^8$  bacteria on enriched rhamnose minimal medium, each strain in 20–60 such platings. All but one strain, no. 2037, produced at least one colony of *rha*<sup>+</sup> mutant bacteria, and a total of twenty-two such mutants were obtained from the total of *c.*  $6.1 \times 10^{10}$  *rha*<sup>-</sup> bacteria plated (Table 2). In individual strains, excluding no. 2037, the number of mutants produced ranged from  $1.3 \times 10^{-10}$  to  $7.7 \times 10^{-10}$

per bacterium plated, and the average rate for the 14 strains was  $3.6 \times 10^{-10}$  per bacterium plated. Since the number of bacteria present on the plate after the first 24 h of incubation was about 50 times the number plated, the average mutant frequency per bacterium present at 24 h was about  $7.2 \times 10^{-12}$ ; this value may approximate to the mutation rate per bacterium per generation (see Materials and Methods).

Table 3. *Transduction of the rha<sup>+</sup> character to non-FIRN rha<sup>-</sup> and FIRN strains of Salmonella typhimurium with phage propagated on the rha<sup>+</sup> strain SL375 as donor*

Recipient strain, group and no.	No. of rha <sup>+</sup> colonies obtained per $1.3 \times 10^8$ bacteria plated*	Mean transduction rate for group, as no. of rha <sup>+</sup> recombinants:	
		Per bacterium plated	Per donor phage virion
<b>Non-FIRN rha<sup>-</sup></b>			
1992	17	$1.4 \times 10^{-7}$	$2.6 \times 10^{-8}$
2002	22		
2037	22		
2212	15		
2213	17		
2241	19		
2323	14		
2368	17		
2382	17		
<b>FIRN</b>			
1879	104	$1.5 \times 10^{-6}$	$2.8 \times 10^{-7}$
1881	97		
2306	221		
2309	231		
2322	314		

\* Number of rha<sup>+</sup> colonies obtained on two plates of enriched rhamnose minimal medium plated with  $1.3 \times 10^8$  bacteria infected with  $7 \times 10^8$  phage virions. Results for strains 2037, 2241 and 2322 are means for two double-platings.

*FIRN strains.* This group of twenty-three *fim<sup>-</sup> inl<sup>-</sup> rha<sup>-</sup>* strains were tested in 488 double-platings, each strain in at least twenty double-platings. The mutational behaviour of the group was clearly different from that of the non-FIRN rha<sup>-</sup> group. Not a single rha<sup>+</sup> mutant was obtained from any strain (Table 2). The aggregate results show that the frequency of rha<sup>+</sup> mutants was less than  $1.6 \times 10^{-11}$  per bacterium plated, i.e. less than  $3.2 \times 10^{-13}$  per bacterium present at 24 h. The absence, or extremely low frequency, of rha<sup>+</sup> mutation in the FIRN group was confirmed by our failure to obtain any rha<sup>+</sup> mutants in similar platings, incubated for 5-7 days, of another 150 FIRN strains isolated from different sources.

(iii) *Transduction from a wild-type rha<sup>+</sup> strain as donor*

The efficiency of the transducing system was tested in experiments in which nine non-FIRN rha<sup>-</sup> and five FIRN strains were transduced with phage that had

been propagated on the  $inl^+ rha^+$  strain SL 375 as donor.  $Rha^+$  recombinants were obtained from each of the recipients, the rate being consistently about ten-fold higher in the FIRN than in the non-FIRN  $rha^-$  strains (Table 3). The rate of production of  $rha^+$  bacteria per phage-infected bacterium plated, even in the strain (2323) with the lowest rate ( $1.1 \times 10^{-7}$ ), was much higher than the highest rate of production of  $rha^+$  mutants observed in the control platings of that or any other strain ( $7.7 \times 10^{-10}$  in strains 1541, 1992 and 6899 Table 2). It is concluded therefore, that most of the  $rha^+$  bacteria produced in the platings with phage were transductional recombinants and not spontaneous mutants.

A number of the  $rha^+$  transductant colonies were picked from the platings of the FIRN recipients and pure subcultures of all of them were found to have the  $inl^-$  character of the recipient strain together with the  $rha^+$  character from the donor. This finding shows that the  $inl^+$  and  $rha^+$  characters of the donor were not transduced jointly and that the  $rha^+$  colonies did not arise from donor-strain bacteria that had survived the treatment of the phage preparation with chloroform.

(iv) *Transduction from  $rha^-$  donors to  $rha^-$  recipients*

Phages propagated on 12 non-FIRN  $rha^-$  and 7 FIRN strains as donors were tested with 14 non-FIRN  $rha^-$  and 23 FIRN strains as recipients in a total of 1075 transduction experiments. The results, as the mean number of  $rha^+$  colonies obtained in the different tests of each cross, are shown in Table 4. The means of the rates of apparent transduction per plated bacterium observed for the different pairs of strains in each of the four classes of cross, non-FIRN  $rha^-$  to non-FIRN  $rha^-$ , FIRN to non-FIRN  $rha^-$ , FIRN to FIRN, and non-FIRN  $rha^-$  to FIRN, are shown in Table 5.

*Crosses of non-FIRN  $rha^-$  with FIRN.* Every strain in the non-FIRN  $rha^-$  and FIRN groups produced  $rha^+$  bacteria, presumed to be recombinants, in at least some of its crossings with members of the other group; most strains did so in a majority of such crossings (Table 4). Each of the 14 non-FIRN  $rha^-$  strains tested as recipients gave  $rha^+$  bacteria in crosses with at least 2 FIRN strains as donors, and each of the 23 FIRN strains tested as recipients gave  $rha^+$  bacteria in crosses with at least 1 non-FIRN  $rha^-$  strain as donor. Conversely, each of 12 non-FIRN  $rha^-$  strains tested as donors gave  $rha^+$  bacteria from at least 1 FIRN recipient and each of 7 FIRN strains tested as donors gave  $rha^+$  bacteria from at least 1 non-FIRN  $rha^-$  recipient.

It is thought that most or all of the  $rha^+$  bacteria obtained in these crosses were recombinants, and not mutants, because the average rate of their production in the transduction platings of each recipient strain was considerably greater than the average rate of production of  $rha^+$  mutants in the control platings of the strain (cf. Tables 2 and 4). The difference between the transduction rate and the mutant frequency was much greater in the crosses in which the recipient was FIRN than in those in which it was non-FIRN  $rha^-$ . The FIRN recipients gave the higher rates of transduction (every strain  $> 7.0 \times 10^{-9}$  per bacterium plated, most strains *c.*  $1.0 \times 10^{-7}$ ) and the lower frequencies of mutation (every strain  $< 3.8 \times 10^{-10}$  per

Table 4. Production of rha<sup>+</sup> recombinants in transductions made to different rha<sup>-</sup> strains of Salmonella typhimurium as recipients with phage propagated on different rha<sup>-</sup> strains of S. typhimurium as donors

Recipient strain	Mean no. of rha <sup>+</sup> colonies obtained per 1.3 × 10 <sup>8</sup> recipient bacteria plated when the donor strain was:																			
	non-FIRN rha <sup>-</sup>								FIRN											
	1179	1180	1541	1992	2002	2037	2212	2241	2323	2368	6768	6899	706	735	2306	2309	2571	2578	2581	
Non-FIRN rha <sup>-</sup>																				
1179	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1180	0	0	0	0	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0
1541	0	0	0	0	0	0	1	0	0.5	0	0	0	0	0	0	0	0	0	0	0
1992	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2002	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2037	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2212	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2213	0	0	0	0	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0
2241	0.5	0.5	1	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0
2323	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2368	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2382	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6768	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6899	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
FIRN																				
706	6	0.5	13	10	9	10	18	7	2	7	3	2	0	0	0	0	0	0	0	0
735	21	0	11	16	17	32	20	61	2	6	17	34	.	.	.	.	.	.	.	.
1289	30	0	3	12	22	16	16	47	2	4	6	31	.	.	.	.	.	.	.	.
1294	2	0	3	5	8	6	13	16	1	6	4	1	0	0	0	0	0	0	0	0
1723	.	1	.	.	14	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1804	6	0	8	26	36	39	45	18	5	26	8	3	0	0	0	0	0	0	0	0
1877	.	4	.	.	19	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1879	6	0.5	5	12	26	23	33	13	5	10	9	4	.	.	.	.	.	.	.	.
1881	5	0.3	6	31	29	37	27	19	5	17	13	2	.	.	.	.	.	.	.	.
2025	.	0	.	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
2305	12	1	2	21	43	31	41	26	9	21	21	6	0	0	0	0	0	0	0	0
2235	2	0	2	5	13	8	14	15	0	15	6	5	0	0	0	0	0	0	0	0
2306	24	0.3	8	15	28	23	29	90	0.2	6	16	38	0	0	0	0	0	0	0	0
2309	6	0.5	2	15	36	23	27	28	3	15	12	6	.	.	.	.	.	.	.	.
2321	.	0	.	.	39	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
2322	25	1	9	16	30	13	33	61	.	7	13	18	.	.	.	.	.	.	.	.
2333	12	0.5	6	14	17	21	13	78	2	4	19	30	0	0	0	0	0	0	0	0
2540	22	0.5	5	17	23	28	20	62	3	12	17	29	0	0	0	0	0	0	0	0
2555	8	0	6	23	21	28	26	4	4	13	10	4	0	0	0	0	0	0	0	0
2565	2	1	0.5	1	5	0	4	4	4	1	1	0	0	0	0	0	0	0	0	0
2571	1	0.5	0	0	3	0	0.5	4	2	0	0.5	.	.	.	.	.	.	.	.	.
2578	3	0	2	7	17	8	13	9	3	13	8	2	.	.	.	.	.	.	.	.
2581	7	1	2	5	18	24	17	15	4	10	4	5	0	0	0	0	0	0	0	0

The fractional results 0.5, 0.3 and 0.2 are average values in crosses in which one rha<sup>+</sup> recombinant was obtained in, respectively, 2, 3 or 4 platings of 1.3 × 10<sup>8</sup> bacteria. Average values greater than one are rounded off to the nearest whole number.

bacterium plated). Most of the non-FIRN  $rha^-$  recipients gave a transduction rate of  $c. 4.0 \times 10^{-9}$  and a mutant frequency of  $c. 4.0 \times 10^{-10}$ , and so showed a difference of only about ten-fold between the two rates. The lowest inter-group transduction rate obtained with any recipient strain was  $2.2 \times 10^{-9}$  (non-FIRN  $rha^-$  strain 1992), yet even this minimum transduction rate was substantially higher than the highest mutant frequency shown by any strain ( $7.7 \times 10^{-10}$ , strains 1541, 1992 and 6899). If the conclusion is accepted that most of the  $rha^+$  bacteria produced in each cross were recombinants, the findings are evidence that any non-FIRN  $rha^-$  strain is capable of giving  $rha^+$  recombinants when crossed with any FIRN strain and, therefore, that the sites of the  $rha^-$  mutations in the non-FIRN  $rha^-$  strains are non-identical and non-overlapping with the sites of the  $rha^-$  mutations in the FIRN strains.

Table 5. Rates of apparent transduction of  $rha^+$  in four classes of crosses of FIRN and non-FIRN  $rha^-$  strains compared with the rates of production of  $rha^+$  mutants in control platings of the recipient strains

(The transduction values are calculated from the aggregated results for the different pairs of strains in each class of cross (Table 4) and the mutation values from the aggregated results for the different recipient strains (Table 2).)

Class of cross		Mean no. of $rha^+$ bacteria obtained	
Donor strain	Recipient strain	Per bacterium plated	Per donor phage virion
Non-FIRN $rha^-$	FIRN	$1.0 \times 10^{-7}$	$1.7 \times 10^{-8}$
FIRN	FIRN	$< 9.9 \times 10^{-11}$	$< 1.7 \times 10^{-11}$
None (control)	FIRN	$< 1.6 \times 10^{-11}$	.
FIRN	Non-FIRN $rha^-$	$8.9 \times 10^{-9}$	$1.6 \times 10^{-9}$
Non-FIRN $rha^-$	Non-FIRN $rha^-$	$5.5 \times 10^{-10}$	$1.0 \times 10^{-10}$
None (control)	Non-FIRN $rha^-$	$3.6 \times 10^{-10}$	.

The rate of production of  $rha^+$  transductants in the FIRN  $\times$  non-FIRN  $rha^-$  crosses was only about a fifteenth as great as that in the crosses of the same FIRN and non-FIRN  $rha^-$  strains as recipients with the  $rha^+$  strain SL 375 as donor (cf. Tables 3, 4). A likely explanation of this difference is that in 14 out of 15 occasions of transduction in the FIRN  $\times$  non-FIRN  $rha^-$  crosses the  $rha^-$  site (or sites) of the donor was incorporated into the recipient's genome along with the positive allele corresponding to the  $rha^-$  site of the recipient. We have no explanation of why the  $rha^+$  transduction rates were generally about ten-fold higher in the crosses with FIRN recipients, than in those with non-FIRN  $rha^-$  ones. In the class of cross, non-FIRN  $rha^-$  donor  $\times$  FIRN recipient, there was a marked variation in the yield of  $rha^+$  recombinants from the different pairs of strains (Table 4). This variation may be attributable in part to some of the strains being lysogenic for phages homologous or related to the transducing phage, and to differences between donor and recipient bacteria in respect of the restriction-modification character.

*Crosses of FIRN with FIRN.*  $Rha^+$  recombinants were not obtained in any cross between 78 different pairs of strains selected from 7 FIRN strains as donors and 23

FIRN strains as recipients (Table 4). This uniformly negative finding, which confirms that of Old (1963), suggests that the *rha*<sup>-</sup> mutation is located at the same site or at overlapping sites in all FIRN strains.

*Crosses of non-FIRN rha*<sup>-</sup> with non-FIRN *rha*<sup>-</sup>. Single *rha*<sup>+</sup> bacteria were obtained in a few (17 out of 168) of the crosses made between 12 non-FIRN *rha*<sup>-</sup> strains as donors and 14 non-FIRN *rha*<sup>-</sup> strains as recipients (Table 4). These *rha*<sup>+</sup> bacteria probably arose by spontaneous mutation in the recipient-strain bacteria, not by recombination, because the frequency of their production (average for group:  $5.5 \times 10^{-10}$  per bacterium plated) was not significantly greater than the frequency of production of *rha*<sup>+</sup> mutant bacteria in the control platings of the recipient strains (average  $3.6 \times 10^{-10}$ ). Although the non-FIRN *rha*<sup>-</sup> strains were less competent as recipients of *rha*<sup>+</sup> than the FIRN strains, they gave *rha*<sup>+</sup> recombinants at a fairly high rate ( $1.1-1.7 \times 10^{-7}$ ) in crosses with the donor strain SL 375 (Table 3). The level of their general competence as recipients therefore does not appear to be low enough to account for the extreme infrequency of *rha*<sup>+</sup> bacteria in their crosses with non-FIRN *rha*<sup>-</sup> donors. It may be concluded, rather, that *rha*<sup>+</sup> recombination is impossible in crosses between any two strains in the non-FIRN *rha*<sup>-</sup> group because the *rha*<sup>-</sup> mutation is located at the same site or at overlapping sites in all members of the group.

(v) *Rhamnose-weak (rha<sup>w</sup>) mutation*

*FIRN rha<sup>w</sup> mutants.* Although FIRN bacteria do not give rise to colonies of *rha*<sup>+</sup> mutants when plated on enriched or unenriched rhamnose minimal medium, they give rise to 'weakly' rhamnose-fermenting (*rha<sup>w</sup>*) mutants which grow out selectively during prolonged culture in rhamnose peptone water. The FIRN strains invariably failed to ferment rhamnose in peptone water within 24 h at 37 °C.

Table 6. *Times of acid production in cultures of FIRN strain 2306 and non-FIRN rha*<sup>-</sup> strain 2213 grown from inocula of 1 or 2 bacteria in 3 ml volumes of rhamnose peptone water at 37 °C

Time of incubation (days)	No. of cultures producing acid within stated time of incubation		Time of incubation (days)	No. of cultures producing acid within stated time of incubation	
	Strain 2306 (40 cultures tested)	Strain 2213 (60 cultures tested)		Strain 2306 (40 cultures tested)	Strain 2213 (60 cultures tested)
1	0	0	7	29	9
1.5	0	1	8	33	12
2	0	1	9	34	14
2.5	0	2	10	34	16
3	1	2	12	35	16
3.5	6	3	14	36	16
4	12	3	16	38	16
4.5	14	3	18	.	20
5	19	5	20	.	20
6	27	7			

Every strain, however, gave 'late' fermentation when incubation was prolonged for a sufficient period, which varied from 3 to 40 days. The time of fermentation differed widely even in repeated tests of the same strain. For example, 40 cultures of FIRN strain 2306 were grown in bottles of 3 ml rhamnose peptone water from inocula of one or two bacteria taken from the same  $10^7$ -fold diluted 20 h broth culture and were observed daily for 16 days. Acid was produced by one of the 40 cultures in 3 days, by 19 within 5 days and by 38 within 16 days (Table 6).

The late fermentation of rhamnose in peptone water was due to the production and selective outgrowth of  $rha^w$  mutant bacteria. Examination of the colonies obtained in platings on nutrient agar showed that  $rha^w$  bacteria made up about 50 % of the population in the fermented cultures, but were absent from the unfermented ones of the same age. The  $rha^w$  colonies were recognized because pure sub-cultures from them gave fermentation in rhamnose peptone water within 24 h. In tests inoculated with about  $10^7$   $rha^w$  bacteria the times taken by the mutants of different FIRN strains to produce acid varied from 10 to 24 h. In similar tests, different wild-type and mutant  $rha^+$  strains produced acid in periods of from 4 to 9 h. The time taken for fermentation by  $rha^w$  and  $rha^+$  bacteria also varied with the size of the inoculum (Table 7).

Table 7. *Times of acid production in cultures of  $rha^+$  strain SL375 and rhamnose-weak strain 2306  $rha^w$ -1 grown from different-sized inocula in 3 ml volumes of rhamnose peptone water at 37 °C*

No. of viable bacteria inoculated	Time (h) of acid production by	
	SL375 ( $rha^+$ )	2306 $rha^w$ -1
$10^8$	3.5	10
$10^7$	4.5	13
$10^6$	5.5	13
$10^5$	6.5	15.5
$10^4$	8	19.5
$10^3$	10.5	21
$10^2$	11.5	22
$10^1$	13	24-36

If it is assumed that the generally observed failure of a small proportion of the cultures of a FIRN strain in rhamnose peptone water to give late fermentation is due to the failure of these cultures to produce even a single  $rha^w$  mutant bacterium it may be concluded that the rate of mutation of FIRN bacteria to the  $rha^w$  form was not much greater than about  $3 \times 10^{-10}$  per bacterium per generation. This calculation is based on the observation that an unfermented culture of FIRN bacteria in 3 ml rhamnose peptone water yields, in about 48 h, a maximum viable population of about  $10^9$  per ml.

The ability of the  $rha^w$  bacteria to ferment rhamnose in peptone water was associated with their having the ability to utilize rhamnose for growth in this medium. The viable counts and turbidity values of  $rha^w$  cultures grown for 24 h in rhamnose peptone water were 2-3-fold higher than those of cultures grown in the

same medium without rhamnose. This ability to utilize rhamnose as an extra source of energy was presumably the reason why the small number of  $rha^w$  bacteria that originated by mutation in cultures of FIRN bacteria in rhamnose peptone water were able selectively to outgrow the  $rha^-$  parental bacteria. Surprisingly, however, the  $rha^w$  bacteria were unable to utilize rhamnose for growth on minimal agar media. Cultures grown for 24 h on enriched rhamnose minimal medium showed little if any greater multiplication of bacteria than that, *c.* 50-fold, observed in parallel cultures grown on the same medium without rhamnose. As well as thus differing from  $rha^+$  bacteria in giving slower fermentation of rhamnose in peptone water and failing to utilize rhamnose in minimal medium, the  $rha^w$  bacteria also differed from  $rha^+$  bacteria in not giving a positive reaction in Bitter rhamnose medium.

*Rha<sup>w</sup> to rha<sup>+</sup> mutation.* The  $rha^w$  mutants of all FIRN strains were capable of undergoing a further mutation to the strongly rhamnose-fermenting  $rha^+$  genotype. A majority of their platings on enriched rhamnose minimal medium yielded a few  $rha^+$  colonies when incubated for 3 days. The frequency of mutation in a series of platings of strain 2306  $rha^w$ -1 was  $1.6 \times 10^{-7}$  per  $rha^w$  bacterium plated, or  $3.2 \times 10^{-9}$  per  $rha^w$  bacterium present after 24 h. These  $rha^+$  mutants of FIRN strains had the same rhamnose-fermentation properties as wild-type  $rha^+$  strains and  $rha^+$  mutants derived from non-FIRN  $rha^-$  strains. They differed, however, from the latter mutants in being capable of derivation only by two successive mutations and two stages of selection, the first in peptone water (for  $rha^w$ ) and the second on minimal medium (for  $rha^+$ ).

*Non-FIRN rha<sup>w</sup> mutants.* Like the FIRN strains, all the non-FIRN  $rha^-$  strains gave late fermentation in rhamnose peptone water. The time of fermentation varied widely in different tests of the same strain, but usually lay between 2 and 14 days. The results for a series of tests inoculated with 1 or 2 bacteria taken from the same broth culture of strain 2213 are shown in Table 6. Some of the late-fermented cultures of each non-FIRN  $rha^-$  strain were found to contain  $rha^w$  mutant bacteria with similar properties to FIRN  $rha^w$  mutants, but others contained  $rha^+$  mutants.

*Stability of mutants.* The  $rha^w$  and  $rha^+$  mutant genotypes derived from FIRN and non-FIRN  $rha^-$  strains appeared to be fairly stable. The rhamnose-fermentation properties remained unchanged when the mutants were carried through ten or more serial subcultures on nutrient agar not containing rhamnose.

#### (vi) *Transduction to FIRN rha<sup>w</sup> recipients*

Three independently isolated  $rha^w$  mutants of FIRN strain 2306, namely mutants 2306  $rha^w$ -1, 2306  $rha^w$ -2 and 2306  $rha^w$ -3, were tested as recipients in transductions with phages propagated on these three mutants, on wild-type  $rha^+$  strain SL 375, non-FIRN  $rha^-$  strains 2002 and 2241, and FIRN strains 706, 735 and 2306. The numbers of  $rha^+$  colonies obtained in platings of the  $rha^w$  bacteria treated with phages from  $rha^w$  donors and FIRN donors were no greater than the numbers obtained in control platings without phage. These colonies presumably

arose from spontaneous mutations, so that there was no evidence that  $rha^+$  recombination took place in any of the crosses between the different  $rha^w$  strains or between  $rha^w$  recipients and FIRN donors. The crosses of the  $rha^w$  recipients with the non-FIRN  $rha^-$  and wild-type  $rha^+$  donors gave ten times as many  $rha^+$  colonies as the control platings and the majority of these  $rha^+$  bacteria are presumed to have been recombinants.

(vii) *Transduction from FIRN  $rha^w$  donors*

The FIRN ( $rha^-$ ) strain 2306 was tested as recipient in transductions with phages propagated on strain 2306  $rha^w$ -2 as donor. Platings on enriched rhamnose minimal medium did not give colonies, presumably because  $rha^w$  bacteria cannot grow on this medium. The production of  $rha^w$  transductants was demonstrated instead by culture in rhamnose peptone water, which is selective for  $rha^w$  bacteria. An overnight broth culture of the recipient strain was diluted 1 in 50 in rhamnose peptone water and ten 3 ml volumes of the diluted culture were incubated in 6 ml bottles for  $3\frac{1}{2}$  h, to yield *c.*  $3.5 \times 10^8$  bacteria per ml. Donor phage was added to a concentration of  $6 \times 10^9$  virions per 3 ml and the cultures were re-incubated for several days and observed daily for production of acid. Similarly treated cultures of the recipient bacteria without the addition of phage served as negative controls, and cultures of the recipient bacteria to which about 15  $rha^w$  bacteria of the donor strain were added served as positive controls. The negative control cultures of the  $rha^-$  bacteria gave late, mutational fermentation at different times between 70 h, when the first of them fermented, and 9 days, when eight out of ten had fermented. Fermentation took place much earlier in the  $rha^-$  cultures treated with phage from the  $rha^w$  donor, presumably as a result of the early production of  $rha^w$  transductant bacteria, and in the positive controls; all these cultures formed acid within 73 h and seven out of ten of the transduction cultures did so in 48–54 h.  $Rha^w$  bacteria were isolated from the fermented transduced cultures.

The finding that the  $rha^w$  character may be transduced into FIRN  $rha^-$  recipients from a FIRN  $rha^w$  mutant as donor and the  $rha^+$  character into FIRN  $rha^w$  recipients from a wild-type  $rha^+$  donor agrees with the suggestion that FIRN strains bear two mutations at different sites in their rhamnose region. The absence of  $rha^+$  recombinants in the crosses tested between independently derived  $rha^w$  mutants of a FIRN strain is in keeping with the view that these mutants arise by back-mutation at the same  $rha^-$  site in different FIRN strains and not by suppressor mutations at different sites.

#### 4. DISCUSSION

The results of the mutation and transduction studies of the non-rhamnose-fermenting FIRN and non-FIRN  $rha^-$  groups of *Salmonella typhimurium* suggest that the site, or sites, of the  $rha^-$  mutation, or mutations, are identical or overlapping in all strains in either group, but are not identical or overlapping as between the two groups. In the tests of mutation to  $rha^+$ , all the twenty-three

FIRN strains behaved in the same way: all failed to produce colonies of rha<sup>+</sup> mutant bacteria when plated directly on enriched rhamnose minimal medium, all produced weakly fermenting rha<sup>w</sup> mutants in ageing cultures in rhamnose peptone water, in which they thereby caused 'late' (> 2 days) fermentation, and all produced colonies of rha<sup>+</sup> mutants in platings of their rha<sup>w</sup> mutants. The pattern of mutation appeared uniform also among the strains of the non-FIRN rha<sup>-</sup> group, but the pattern in this group differed from that in the FIRN group. All but one of the fourteen non-FIRN rha<sup>-</sup> strains produced one or more colonies of rha<sup>+</sup> mutant bacteria in direct platings on enriched rhamnose minimal medium, and all strains produced rha<sup>+</sup> bacteria in some late-fermenting cultures in rhamnose peptone water and rha<sup>w</sup> bacteria in other such cultures. The fact that restoration of the rha<sup>+</sup> genotype required two successive mutations in FIRN strains (rha<sup>-</sup> to rha<sup>w</sup> to rha<sup>+</sup>) but only one mutation in non-FIRN rha<sup>-</sup> strains (rha<sup>-</sup> to rha<sup>+</sup>) suggests that FIRN strains bear two mutational defects at different sites in the rhamnose region of the chromosome and non-FIRN rha<sup>-</sup> strains one such defect at a third site. In FIRN strains the rha<sup>-</sup> to rha<sup>w</sup> mutation may be a back-mutation at one of the two sites, which partially restores fermenting activity, and the rha<sup>w</sup> to rha<sup>+</sup> mutation may be a back-mutation at the second site, which is expressed only when the first site is also active. In the non-FIRN rha<sup>-</sup> strains the rha<sup>-</sup> to rha<sup>+</sup> mutation may be a back-mutation at the single rha<sup>-</sup> site and the rha<sup>-</sup> to rha<sup>w</sup> mutation may be a partially effective suppressor mutation. Power (1967) has shown that in *Escherichia coli* the fermentation of rhamnose is controlled by a cluster of four genes, three of which appear to be structural genes for L-rhamnose isomerase, L-rhamnulokinase and L-rhamnulose 1-phosphate aldolase, and the fourth an activator-producing regulatory gene. *S. typhimurium* probably possesses a similar rhamnose operon, but we have not determined which of its genes contain the rha<sup>-</sup> sites in FIRN and non-FIRN rha<sup>-</sup> bacteria.

The rha<sup>w</sup> biotype, which permits slow (10–24 h) fermentation of rhamnose in peptone water but not the utilization of rhamnose in minimal medium, has not hitherto been clearly defined. Although it is not included among the twenty-one wild-type biotypes recognized by Kristensen *et al.* (1937), Harhoff (1948), and Kallings & Laurell (1957), it is not confined to laboratory mutants. Dr B. A. D. Stocker identified and sent to us five wild-type rha<sup>w</sup> strains isolated in 1962, and one of us (J. P. D.) and Dr E. S. Anderson have detected several other such strains in a large series of isolates of *S. typhimurium*. All these wild-type rha<sup>w</sup> strains were fim<sup>+</sup> and *d*-tartrate-positive, and so differed in one or other character from the rha<sup>w</sup> mutants derived from FIRN and non-FIRN rha<sup>-</sup> strains. The reason why rha<sup>w</sup> bacteria cannot utilize rhamnose in minimal medium, although they are able to utilize it in peptone water, is unknown.

The finding that rha<sup>+</sup> recombinants were produced in transductional crosses between any one FIRN strain and any one non-FIRN rha<sup>-</sup> strain is clear evidence that the sites of the rha<sup>-</sup> mutations are non-identical and non-overlapping between the two groups. The consistent absence of rha<sup>+</sup> recombinants in crosses between different FIRN strains confirms and extends the findings of Old (1963) and

suggests that the sites of the  $rha^-$  mutations are identical or overlapping in all members of the FIRN group. The significance of the finding in crosses between different strains in the non-FIRN  $rha^-$  group is less clear. These strains produced small numbers of  $rha^+$  bacteria by mutation and it was impossible to determine with certainty whether the few  $rha^+$  bacteria obtained in the transduction platings were mutants or recombinants. Since, however, the rate of production of  $rha^+$  bacteria in the non-FIRN  $rha^- \times$  non-FIRN  $rha^-$  crosses was not significantly higher than the rate of production of  $rha^+$  mutants in the control platings, it is probable that  $rha^+$  recombination did not take place in any of the crosses and, therefore, that the sites of the  $rha^-$  mutations are identical or overlapping in all members of the non-FIRN  $rha^-$  group. This conclusion would be in keeping with the observed similarity of mutational behaviour in the non-FIRN  $rha^-$  strains.

The conclusion drawn from the present findings and those of Old (1963) that all FIRN strains have their  $rha^-$  mutations located at identical sites and the similar conclusion drawn from the findings of Duguid *et al.* (1962) and Old (1963) that all FIRN strains have their  $fim^-$  mutations located at identical sites, suggest strongly that all these strains have a common ancestry. FIRN strains have probably originated from the much commoner  $fim^+ inl^- rha^+$  strains of *S. typhimurium* by successive mutations in the  $fim$  and  $rha$  genes. If different lines of FIRN strains had originated independently at different times and places, they would almost certainly have differed from one another in the sites of their mutations. The apparent identity of the mutational sites in all FIRN strains, including ones isolated in several different countries, argues that all of them have descended from a single ancestral FIRN bacterium, which itself was derived from a  $fim^+ inl^- rha^+$  bacterium. One may suppose that the latter bacterium first acquired one of the two characters,  $fim^-$  and  $rha^-$ , by mutation and that after an unknown number of generations one of its descendants acquired the other character by a second mutation. The progeny of the double-mutant, or FIRN bacterium, must then have spread widely throughout the world and become diversified in biotype and phage-type by mutation and lysogenic conversion. The intermediate, single-mutant class of bacteria, which would have been either  $fim^- inl^- rha^+$  or  $fim^+ inl^- rha^-$ , apparently failed to proliferate widely, since only two strains of the former type and none of the latter were found among 775 wild-type strains from different sources examined by Duguid *et al.* (1966). If, as our mutational studies suggest, a FIRN bacterium bears two  $rha^-$  mutations at different sites, the derivation of the first FIRN bacterium from a  $fim^+ inl^- rha^+$  ancestor must have involved three, not two, mutations and the production of an additional intermediate class of mutant.

For reasons similar to those adduced in regard to the FIRN group, the apparently negative results of the transductional crosses between different strains in the non-FIRN  $rha^-$  group suggest that all strains in this latter group have descended from a single non-FIRN  $rha^-$  bacterium that originated by  $rha^-$  mutation in a bacterium of the common  $fim^+ inl^+ rha^+$  type of *S. typhimurium*. The ancestral bacterium was presumably biotype 6, which, except for its being  $rha^+$ , resembles

in all biotyping characters the biotype 19Xd, to which all the non-FIRN rha<sup>-</sup> strains belong. The identity of the rha<sup>-</sup> sites in the different non-FIRN rha<sup>-</sup> strains is not likely to be due to a special mutability of this site, since there is no obvious reason why such a specially mutable site should be present only in strains of biotype 6. Our fourteen non-FIRN rha<sup>-</sup> strains included isolates from Britain, Australia and the U.S.A., and representatives of eight phage-types, and it is apparent that this race, like the FIRN race, has become widely disseminated from its place of origin and widely diversified in phage-type.

Although the conclusions we have drawn are speculative, our results suggest that transduction studies of wild-type strains with mutational deficiencies may make it possible to elucidate the recent evolutionary history of particular strains and types of bacteria.

#### SUMMARY

Observations of mutation and phage-mediated transduction of rhamnose-fermentation properties were made on two groups of wild-type rhamnose-non-fermenting (rha<sup>-</sup>) strains of *Salmonella typhimurium* isolated in several different countries. The 'FIRN' group included twenty-three strains in the biotypes 15, 16, 17 and 18, and the 'non-FIRN rha<sup>-</sup>' group included fourteen strains in a new biotype, provisionally designated 19Xd. The two groups showed different patterns of mutational behaviour and the pattern was uniform for the different strains in each group. Non-FIRN rha<sup>-</sup> strains gave rise directly to colonies of rha<sup>+</sup> mutant bacteria in platings on rhamnose minimal media. FIRN strains were unable to do this, but produced rha<sup>+</sup> bacteria by two successive mutations, the first of which yielded weakly fermenting (rha<sup>w</sup>) bacteria capable of fermenting rhamnose in peptone water, but not of utilizing it for growth on minimal medium.

Rha<sup>+</sup> recombinant bacteria were produced in transductional crosses between any one FIRN strain and any one non-FIRN rha<sup>-</sup> strain, but not in any cross between one FIRN strain and another FIRN strain or between one non-FIRN rha<sup>-</sup> strain and another non-FIRN rha<sup>-</sup> strain. These findings suggest that although the sites of the rha<sup>-</sup> mutations are different between the FIRN and non-FIRN rha<sup>-</sup> groups, they are located at identical sites on the chromosome in all strains in either group. It is concluded that all FIRN strains have probably descended from a single ancestral strain which underwent mutations at two sites in the rhamnose region of the chromosome and that all non-FIRN rha<sup>-</sup> strains have descended from a single ancestral bacterium which underwent a mutation at a third site in the rhamnose region.

We wish to thank Mrs Ruth Barker, B.Sc., for making some of the isolations of rha<sup>+</sup> mutants and observations of mutation rates, Dr E. S. Anderson for supplying and phage-typing strains, and Dr B. A. D. Stocker for many helpful suggestions and criticisms. One of us (A.M.) is indebted to the British Council for a Scholarship, during tenure of which this work was done.

## REFERENCES

- BOYD, J. S. K. (1956). Immunity of lysogenic bacteria. *Nature, Lond.* **178**, 141.
- BRINTON, C. C. JR, BUZZELL, A. & LAUFFER, M. A. (1954). Electrophoresis and phage susceptibility studies on a filament producing variant of the *E. coli* B bacterium. *Biochim. biophys. Acta* **15**, 533-542.
- CALLOW, BESSIE R. (1959). A new phage-typing scheme for *Salmonella typhimurium*. *J. Hyg., Camb.* **57**, 346-359.
- DAVIS, B. D. & MINGIOLI, E. S. (1950). Mutants of *Escherichia coli* requiring methionin or vitamin B12. *J. Bact.* **60**, 17-28.
- DEMEREK, M., MOSER, H., HEMMERLY, J., BLOMSTRAND, I., DEMEREK, Z. E., FITZGERALD, P. L., GLOVER, S. W., HANSON, J. F., NIELSEN, F. J. & YURA, T. (1954). Bacterial genetics, I. *Yb. Carnegie Instn. Wash.* **53**, 225-241.
- DUGUID, J. P., ANDERSON, E. S. & CAMPBELL, I. (1966). Fimbriae and adhesive properties in salmonellae. *J. Path. Bact.* **91**, 107-138.
- DUGUID, J. P., OLD, D. C. & HUME, V. B. (1962). Transduction of fimbriation and rhamnose fermentation characters in *Salmonella typhimurium*. *Proc. Genet. Soc. Edinb.* p. 138.
- DUGUID, J. P., SMITH, I. W., DEMPSTER, G. & EDMUNDS, P. N. (1955). Non-flagellar filamentous appendages ('fimbriae') and haemagglutinating activity in *Bacterium coli*. *J. Path. Bact.* **70**, 335-348.
- HARHOFF, N. (1948). Gastroenteritisbaciller af Salmonellagruppen i Danmark. *Nyt nordisk Forlag*, pp. 23, 42. Copenhagen.
- HARTMAN, P. E. (1956). Genetic studies with bacteria. *Publ. Carnegie Instn.* no. 612, pp. 35-62.
- HOUWINK, A. L. & VAN ITERSOM, W. (1950). Electron-microscopical observations on bacterial cytology. II. Study on flagellation. *Biochim. biophys. Acta* **5**, 10-44.
- KALLINGS, L. O. & LAURELL, A. B. (1957). Relation between phage types and fermentation types of *Salmonella typhimurium*. *Acta path. microbiol. scand.* **40**, 328-342.
- KAUFFMANN, F. (1954). *Enterobacteriaceae*, 2nd ed., p. 42. Copenhagen: Ejnar Munksgaard.
- KRISTENSEN, M., BOJLÉN, K. & FAARUP, C. (1937). Bacteriologisk-epidemiologiske Erfaringer om Infektioner med Gastroenteritisbaciller af Paratyfusgruppen. *Bibltk Laeger* **129**, 310-375.
- OLD, D. C. (1963). Fimbriation in Enterobacteriaceae. Ph.D. thesis, University of Edinburgh.
- POWER, J. (1967). The L-rhamnose genetic system in *Escherichia coli* K-12. *Genetics* **55**, 557-568.
- SANDERSON, K. E. & DEMEREK, M. (1965). The linkage map of *Salmonella typhimurium*. *Genetics* **51**, 897-913.
- STOCKER, B. A. D. & EDGAR, J. B. (1959). *Annual Report of the Board of Governors of the Lister Inst.* p. 13. Also *Proc. Path. Soc. Gt Brit. Ireland*, 8 January 1959 (unpublished).
- SUBBALAH, T. V. & STOCKER, B. A. D. (1962). Mapping of the correlated characters fimbriation and rhamnose-fermentation in *Salmonella typhimurium* using col-agents as fertility factors. *Proc. Genet. Soc. Edinb.* pp. 138.
- SWANSTROM, M. & ADAMS, M. H. (1951). Agar layer method for production of high titer phage stocks. *Proc. Soc. exp. Biol. Med.* **78**, 372-375.
- ZINDER, N. D. & LEDERBERG, J. (1952). Genetic exchange in *Salmonella*. *J. Bact.* **64**, 679-699.