

- 2 Holma KM, Melartin TK, Holma IA, Isometsä ET. Predictors for switch from unipolar major depressive disorder to bipolar disorder type I or II: a 5-year prospective study. *J Clin Psychiatry* 2008; **69**: 1267–75.
- 3 Wada K, Sasaki T, Jitsuiki H, Takaishi Y. One-year outcomes of unipolar depression patients with manic or hypomanic switch during acute antidepressant treatment. *Int J Psychiatry Clin Pract* 2013; **17**: 219–22.
- 4 Jin W, Ma Y, Qiu D, Wang G, Wang X, Fan X, et al. Comparison of switching rate from depression to BPD between different classes of antidepressants [Chinese]. *Chin J Nerv Ment Dis* 2007; **33**: 565–6.
- 5 Li Z, Zhang C, Fan J, Yuan C, Huang J, Chen J, et al. Brain-derived neurotrophic factor levels and bipolar disorder in patients in their first depressive episode: 3-year prospective longitudinal study. *Br J Psychiatry* 2014; **205**: 29–35.
- 6 Forty L, Smith D, Jones L, Jones I, Caesar S, Cooper C, et al. Clinical differences between bipolar and unipolar depression. *Br J Psychiatry* 2008; **192**: 388–9.

ZeZhi Li, MD, PhD, **Chen Zhang**, MD, PhD, **Yiru Fang**, MD, PhD, Division of Mood Disorders, Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai, China. Email: yirufang@aliyun.com

doi: 10.1192/bjp.206.1.79a

Correction

Towards a more nuanced global mental health. *BJP*, 204, 415–417.

Reference 10 should read:

- 10 Wunderink L, Nienhuis FJ, Sytema S, Slooff CJ, Knegtering R, Wiersma D. Guided discontinuation versus maintenance treatment in remitted first-episode psychosis: relapse rates and functional outcome. *J Clin Psychiatry* 2007; **68**: 654–61.

doi: 10.1192/bjp.206.1.80