

## A REMARK ON THE PERMUTATION REPRESENTATIONS AFFORDED BY THE EMBEDDINGS OF $O_{2m}^{\pm}(2^f)$ IN $Sp_{2m}(2^f)$

SIMON GUEST , ANDREA PREVITALI and PABLO SPIGA

(Received 4 April 2013; accepted 1 August 2013; first published online 19 September 2013)

### Abstract

We show that the permutation module over  $\mathbb{C}$  afforded by the action of  $Sp_{2m}(2^f)$  on its natural module is isomorphic to the permutation module over  $\mathbb{C}$  afforded by the action of  $Sp_{2m}(2^f)$  on the union of the right cosets of  $O_{2m}^+(2^f)$  and  $O_{2m}^-(2^f)$ .

2010 *Mathematics subject classification*: primary 20B05; secondary 20H30.

*Keywords and phrases*: permutation groups, permutation character.

### 1. Introduction

That a given finite group can have rather different permutation representations affording the same permutation character was shown by Helmut Wielandt in 1979. For instance, the actions of the projective general linear group  $PGL_d(q)$  on the projective points and on the projective hyperplanes afford the same permutation character, but these actions are not equivalent when  $d \geq 3$ . A more interesting example is offered by the Mathieu group  $M_{23}$ . Here we have two primitive permutation representations of degree 253 affording the same permutation character, but with nonisomorphic point stabilisers (see [3, page 71]).

Establishing which properties are shared by permutation representations of a finite group  $G$  with the same permutation character has been the subject of considerable interest. For instance, it was conjectured by Wielandt [14] that, if  $G$  admits two permutation representations on  $\Omega_1$  and  $\Omega_2$  that afford the same permutation character, and if  $G$  acts primitively on  $\Omega_1$ , then  $G$  acts primitively on  $\Omega_2$ . This conjecture was first reduced to the case where  $G$  is almost simple by Förster and Kovács [5] and then solved (in the negative) by Guralnick and Saxl [7]. Some more recent investigations on primitive permutation representations and their permutation characters can be found in [13].

In this paper we construct two considerably different permutation representations of the symplectic group that afford the same permutation character. We let  $q$  be a power of 2,  $G$  be the finite symplectic group  $Sp_{2m}(q)$ ,  $V$  be the  $2m$ -dimensional natural module

for  $\text{Sp}_{2m}(q)$  over the field  $\mathbb{F}_q$  of  $q$  elements, and  $\pi$  be the complex permutation character for the action (by matrix multiplication) of  $G$  on  $V$ . Since  $q$  is even, the orthogonal groups  $O_{2m}^+(q)$  and  $O_{2m}^-(q)$  are maximal subgroups of  $G$  (see [4]). For  $\varepsilon \in \{+, -\}$ , we let  $\Omega^\varepsilon$  denote the set of right cosets of  $O_{2m}^\varepsilon(q)$  in  $G$ , and we let  $\pi^\varepsilon$  denote the permutation character for the action of  $G$  on  $\Omega^\varepsilon$ .

**THEOREM 1.1.** *The  $\mathbb{C}G$ -modules  $\mathbb{C}V$  and  $\mathbb{C}\Omega^+ \oplus \mathbb{C}\Omega^-$  are isomorphic. That is,  $\pi = \pi^+ + \pi^-$ .*

We find this behaviour quite peculiar considering that the  $G$ -sets  $V$  and  $\Omega^+ \cup \Omega^-$  are rather different. For instance,  $G$  has two orbits of size 1 and  $q^{2m} - 1$  on  $V$ , and has two orbits of size  $q^m(q^m + 1)/2$  and  $q^m(q^m - 1)/2$  on  $\Omega^+ \cup \Omega^-$ . Moreover, the action of  $G$  on both  $\Omega^+$  and  $\Omega^-$  is primitive, but the action of  $G$  on  $V \setminus \{0\}$  is not when  $q > 2$ .

### 2. Proof of Theorem 1.1

Inglis [8, Theorem 1] shows that the orbitals of the two orthogonal subgroups are self-paired, hence the characters  $\pi^+$  and  $\pi^-$  are multiplicity-free (see [1, Section 2.7]). We will use this fact in our proof of Theorem 1.1.

**PROOF OF THEOREM 1.1.** Let 1 denote the principal character of  $G$ . Observe that  $\pi = 1 + \pi_0$ , where  $\pi_0$  is the permutation character for the transitive action of  $G$  on  $V \setminus \{0\}$ . In particular, for  $v \in V \setminus \{0\}$ , we have  $\pi_0 = 1_{G_v}^G$ , where  $G_v$  is the stabiliser of  $v$  in  $G$ . Frobenius reciprocity implies that  $\langle \pi_0, \pi_0 \rangle = \langle \pi_0|_{G_v}, 1 \rangle$ , and this equals the number of orbits of  $G_v$  on  $V \setminus \{0\}$ . We claim that  $G_v$  has  $2q - 1$  orbits on  $V \setminus \{0\}$ . More precisely we show that, given  $w \in v^\perp \setminus \langle v \rangle$  and  $w' \in V \setminus v^\perp$ , the elements  $\lambda v$  (for  $\lambda \in \mathbb{F}_q \setminus \{0\}$ ),  $w$ , and  $\lambda w'$  (for  $\lambda \in \mathbb{F}_q \setminus \{0\}$ ) are representatives for the orbits of  $G_v$  on  $V \setminus \{0\}$ . Since  $G_v$  fixes  $v$  and preserves the bilinear form  $(\cdot, \cdot)$ , these elements are in distinct  $G_v$ -orbits. Let  $u \in V \setminus \{0\}$ . If  $u \in \langle v \rangle$ , then  $u = \lambda v$  for some  $\lambda \neq 0$ , and hence there is nothing to prove. Let  $w_0 = w$  if  $(v, u) = 0$ , and let  $w_0 = ((v, u)/(v, w'))w'$  if  $(v, u) \neq 0$ . By construction, the 2-spaces  $\langle v, u \rangle$  and  $\langle v, w_0 \rangle$  are isometric and admit an isometry  $f$  such that  $v^f = v$  and  $u^f = w_0$ . By Witt’s lemma [9, Proposition 2.1.6],  $f$  extends to an isometry  $g$  of  $V$ . Thus  $g \in G_v$  and  $u^g = w_0$ , which proves our claim. Therefore,

$$\langle \pi_0, \pi_0 \rangle = 2q - 1. \tag{2.1}$$

Next we need to refine the information in (2.1). Let  $P$  be the stabiliser of the 1-subspace  $\langle v \rangle$  in  $G$ . Then  $P$  is a maximal parabolic subgroup of  $G$  and  $P/G_v$  is cyclic of order  $q - 1$ . Write  $\eta = 1_{G_v}^P$  and observe that  $\eta = \sum_{\zeta \in \text{Irr}(P/G_v)} \zeta$ , where by an abuse of terminology we identify the characters of  $P/G_v$  with the characters of  $P$  containing  $G_v$  in the kernel. Thus

$$\pi_0 = 1_{G_v}^G = (1_{G_v}^P)^G = \eta_P^G = \sum_{\zeta \in \text{Irr}(P/G_v)} \zeta_P^G.$$

Since every character of  $G$  is real-valued [6], we must have

$$(\bar{\zeta})_P^G = \overline{\zeta_P^G} = \zeta_P^G,$$

where  $\bar{x}$  denotes the complex conjugate of  $x \in \mathbb{C}$ . Let  $\mathcal{S}$  be a set of representatives, up to complex conjugation, of the nontrivial characters of  $\text{Irr}(P/G_v)$ . Since  $|P/G_v| = q - 1$  is odd, we see that  $|\mathcal{S}| = q/2 - 1$ . We have

$$\pi_0 = 1_P^G + 2 \sum_{\zeta \in \mathcal{S}} \zeta_P^G.$$

If we write  $\pi' = \sum_{\zeta \in \mathcal{S}} \zeta_P^G$ , then  $\pi_0 = 1_P^G + 2\pi'$ .

Since  $1_P^G$  is the permutation character of the rank-three action of  $G$  on the one-dimensional subspaces of  $V$ , we have  $1_P^G = 1 + \chi^+ + \chi^-$  for some distinct nontrivial irreducible characters  $\chi^+$  and  $\chi^-$  of  $G$ . Let  $\Gamma$  be the graph with vertex set the 1-subspaces of  $V$  and edge sets  $\{\langle v \rangle, \langle w \rangle\}$  whenever  $v \perp w$ . Observe that  $\Gamma$  is strongly regular with parameters

$$\left( \frac{q^{2m} - 1}{q - 1}, \frac{q^{2m-1} - q}{q - 1}, \frac{q^{2m-2} - 1}{q - 1} - 2, \frac{q^{2m-2} - 1}{q - 1} \right).$$

Hence the eigenvalues of  $\Gamma$  have multiplicity

$$\frac{1}{2} \left( \frac{q^{2m} - q}{q - 1} - q^m \right) \quad \text{and} \quad \frac{1}{2} \left( \frac{q^{2m} - q}{q - 1} + q^m \right)$$

(see [2, page 27]).

Interchanging the roles of  $\chi^+$  and  $\chi^-$  if necessary, we may assume that  $\chi^-(1) < \chi^+(1)$ . The above direct computation proves that

$$\chi^-(1) = \frac{1}{2} \left( \frac{q^{2m} - q}{q - 1} - q^m \right) \quad \text{and} \quad \chi^+(1) = \frac{1}{2} \left( \frac{q^{2m} - q}{q - 1} + q^m \right) \tag{2.2}$$

(compare [10, Section 1]).

Fix  $\zeta \in \mathcal{S}$ . We claim that  $\zeta_P^G$  is irreducible. From Mackey’s irreducibility criterion [11, Proposition 23, Section 7.3], we need to show that for every  $s \in G \setminus P$ , we have  $\zeta_{sPs^{-1} \cap P} \neq \zeta^s$ , where  $\zeta^s$  is the character of  $sPs^{-1} \cap P$  defined by  $(\zeta^s)(x) = \zeta(s^{-1}xs)$  and, as usual,  $\zeta_{sPs^{-1} \cap P}$  is the restriction of  $\zeta$  to  $sPs^{-1} \cap P$ . Fix a monomorphism  $\psi$  from  $P/P'$  into  $\mathbb{C}^*$ . Since  $\zeta$  is a class function of  $P$ , we need to consider only elements  $s$  in distinct  $(P, P)$ -double cosets. These correspond to the  $P$ -orbits  $\langle v \rangle$ ,  $v^\perp \setminus \langle v \rangle$  and  $V \setminus v^\perp$ . Let  $H = \langle v, u \rangle$  be a hyperbolic plane and choose  $s \in G$  such that

$$vs = u, \quad us = u \quad \text{and} \quad s_{H^\perp} = 1_{H^\perp}.$$

A calculation shows that  $\zeta^s(x) = \psi(\mu^{-1}) = \overline{\zeta(x)}$ , where  $vx = \mu v$ . Since  $q - 1$  is odd, we have  $\zeta(x) \neq \zeta^s(x)$  when  $\mu \neq 1$ . Therefore  $\zeta \neq \zeta^s$ . Finally, choose  $s \in G$  such that  $(v, u, w, z)s = (w, z, v, u)$ , where  $H = \langle v, u \rangle \perp \langle w, z \rangle$  is an orthogonal sum of hyperbolic planes and  $s_{H^\perp} = 1_{H^\perp}$ . Another calculation shows that  $\zeta^s(x) = \psi(\lambda)$  and  $\zeta(x) = \psi(\mu)$ , where  $vx = \mu v$  and  $wx = \lambda$ . If  $\mu \neq \lambda$ , then  $\zeta^s(x) \neq \zeta(x)$  and hence  $\zeta^s \neq \zeta$ . Our claim is now proved.

Write  $\pi' = \sum_{i=1}^{\ell} m_i \chi_i$  as a linear combination of the distinct irreducible constituents of  $\pi'$ . Observe that, by the previous paragraph, each  $\chi_i$  is of the form  $\zeta_p^G$ , for some  $\zeta \in \mathcal{S}$ . Therefore  $\chi_i$  has degree  $|G : P|$  for each  $i$  and, in particular,  $1, \chi^+$  and  $\chi^-$  are not irreducible constituents of  $\pi'$ .

The number of irreducible constituents of  $\pi_0$  is

$$1 + 1 + 1 + 2(m_1 + \dots + m_{\ell}) = 3 + 2|\mathcal{S}| = 3 + 2\left(\frac{q}{2} - 1\right) = q + 1,$$

and by (2.1),

$$3 + 4m_1^2 + \dots + 4m_{\ell}^2 = 2q - 1.$$

Multiplying the first equation by  $-2$  and adding the second equation,

$$-3 + 4m_1(m_1 - 1) + \dots + 4m_{\ell}(m_{\ell} - 1) = -3.$$

It follows that  $m_1 = \dots = m_{\ell} = 1$ , and hence  $\ell = q/2 - 1$ . This shows that  $\pi'$  is multiplicity-free.

Summing up,

$$\pi_0 = 1 + \chi^+ + \chi^- + 2\pi', \quad \langle \pi', \pi' \rangle = \frac{q}{2} - 1, \quad \langle 1 + \chi^+ + \chi^-, \pi' \rangle = 0. \tag{2.3}$$

We now turn our attention to the characters  $\pi^+$  and  $\pi^-$ . By Frobenius reciprocity, or by [8, Theorem 1(i) and (ii)], we see that

$$\langle \pi^+, \pi^+ \rangle = \langle \pi^-, \pi^- \rangle = \frac{q}{2} + 1. \tag{2.4}$$

By [8, Lemma 2(iii) and (iv)], the orbits of  $O_{2m}^-(q)$  in its action on  $\Omega^+$  are in one-to-one correspondence with the elements in  $\{\alpha + \alpha^2 \mid \alpha \in \mathbb{F}_q\}$ . In particular, we have  $\langle \pi^+|_{O_{2m}^-(q)}, 1 \rangle = |\{\alpha + \alpha^2 \mid \alpha \in \mathbb{F}_q\}| = q/2$ . Now Frobenius reciprocity implies that

$$\langle \pi^+, \pi^- \rangle = \frac{q}{2}. \tag{2.5}$$

Next we show that

$$\langle \pi_0, \pi^+ \rangle = \langle \pi_0, \pi^- \rangle = q. \tag{2.6}$$

Using Frobenius reciprocity, it suffices to show that the number of orbits of  $O_{2m}^{\pm}(q)$  on  $V \setminus \{0\}$  is  $q$ . Fix  $\varepsilon \in \{+, -\}$  and let  $Q^{\varepsilon}$  be the quadratic form on  $V$  preserved by  $O_{2m}^{\varepsilon}(q)$ . For  $\lambda \in \mathbb{F}_q$ , we see from [9, Lemma 2.10.5(ii)] that  $\Omega_{2m}^{\varepsilon}(q)$  is transitive on  $V_{\lambda}^{\varepsilon} = \{v \in V \setminus \{0\} \mid Q^{\varepsilon}(v) = \lambda\}$ . In particular,  $\{V_{\lambda}^{\varepsilon} \mid \lambda \in \mathbb{F}_q\}$  is the set of orbits of  $\Omega_{2m}^{\varepsilon}(q)$  on  $V \setminus \{0\}$ . Since  $O_{2m}^{\varepsilon}(q)$  is the isometry group of  $Q^{\varepsilon}$  we see that  $\{V_{\lambda}^{\varepsilon} \mid \lambda \in \mathbb{F}_q\}$  is also the set of orbits of  $O_{2m}^{\varepsilon}(q)$  on  $V \setminus \{0\}$ , and (2.6) is now proved.

Since  $\pi^+$  is multiplicity-free, up to reordering, by (2.3) and (2.6), we may assume that

$$\pi^+ = 1 + a\chi^- + b\chi^+ + \sum_{i=1}^t \chi_i + \rho,$$

where  $a, b \in \{0, 1\}$ ,  $0 \leq t \leq q/2 - 1$  and  $\langle \pi_0, \rho \rangle = 0$ . By (2.6), we have  $q - 2 \geq 2t = q - 1 - a - b \geq q - 3$ . Hence  $2t = q - 2$  and  $\{a, b\} = \{0, 1\}$ . Since

$$\pi^+(1) = |\Omega^+| = \frac{q^m(q^m + 1)}{2}$$

and

$$\pi'(1) = \left(\frac{q}{2} - 1\right) |G : P| = \left(\frac{q}{2} - 1\right) \frac{q^{2m} - 1}{q - 1},$$

it follows by (2.2) that  $a = 0$  and  $b = 1$ . By (2.4), we have  $\pi^+ = 1 + \chi^+ + \pi'$ .

Now (2.3)–(2.6) imply immediately that  $\pi^- = 1 + \chi^- + \pi'$ . This shows that

$$\pi^+ + \pi^- = (1 + \chi^+ + \pi') + (1 + \chi^- + \pi') = 1 + 1 + \chi^+ + \chi^- + 2\pi' = 1 + \pi_0 = \pi,$$

which completes the proof of Theorem 1.1.  $\square$

We note that the ‘ $q = 2$ ’ case of Theorem 1.1 was first proved by Siemons and Zalesskii in [12, Proposition 3.1]. This case is particularly easy to deal with (considering that the action of  $G$  on both  $\Omega^+$  and  $\Omega^-$  is 2-transitive) and its proof depends only on Frobenius reciprocity. However, the general statement (valid for every even  $q$ ) of Theorem 1.1 was undoubtedly inspired by their observation. In fact, since the submission of this paper, Zalesskii has informed the authors that he has obtained a proof of Theorem 1.1 using some relatively sophisticated methods from the character theory of the groups of Lie type.

Finally, we observe that Theorem 1.1 reproduces the following result as an immediate corollary (see [4, Theorem 6]).

**COROLLARY 2.1.** *Every element of  $\mathrm{Sp}_{2m}(q)$  is conjugate to an element of  $\mathrm{O}_{2m}^+(q)$  or of  $\mathrm{O}_{2m}^-(q)$ .*

**PROOF.** Let  $g \in \mathrm{Sp}_{2m}(q)$ . Then

$$\pi(g) = (1 + \pi_0)(g) = 1(g) + \pi_0(g) \geq 1(g) = 1$$

and therefore, since  $\pi = \pi^+ + \pi^-$  by Theorem 1.1, either  $\pi^+(g) \geq 1$  or  $\pi^-(g) \geq 1$ ; that is,  $g$  fixes some point in  $\Omega^+$  or in  $\Omega^-$ . In the first case  $g$  has a conjugate in  $\mathrm{O}_{2m}^+(q)$  and in the second case  $g$  has a conjugate in  $\mathrm{O}_{2m}^-(q)$ .  $\square$

## References

- [1] P. J. Cameron, *Permutation Groups*, London Mathematical Society Student Texts, 45 (Cambridge University Press, Cambridge, 1999).
- [2] P. J. Cameron and J. H. van Lint, *Designs, Graphs, Codes and Their Links*, London Mathematical Society Student Texts, 22 (Cambridge University Press, Cambridge, 1991).
- [3] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups* (Clarendon Press, Oxford, 1985).
- [4] R. H. Dye, ‘Interrelations of symplectic and orthogonal groups in characteristic two’, *J. Algebra* **59** (1979), 202–221.

- [5] P. Förster and L. G. Kovács, 'A problem of Wielandt on finite permutation groups', *J. Lond. Math. Soc. (2)* **41** (1990), 231–243.
- [6] R. Gow, 'Products of two involutions in classical groups of characteristic 2', *J. Algebra* **71** (1981), 583–591.
- [7] R. M. Guralnick and J. Saxl, *Primitive Permutation Characters*, London Mathematical Society Lecture Note Series, 165 (Cambridge University Press, Cambridge, 1992), 364–376.
- [8] N. F. J. Inglis, 'The embedding  $O(2m, 2^k) \leq Sp(2m, 2^k)$ ', *Arch. Math.* **54** (1990), 327–330.
- [9] P. Kleidman and M. Liebeck, *The Subgroup Structure of the Finite Classical Groups*, London Mathematical Society Lecture Notes, 129 (Cambridge University Press, Cambridge, 1990).
- [10] M. W. Liebeck, 'Permutation modules for rank 3 symplectic and orthogonal groups', *J. Algebra* **92** (1985), 9–15.
- [11] J. P. Serre, *Linear Representations of Finite Groups*, Graduate Texts in Mathematics, 42 (Springer, New York, 1977).
- [12] J. Siemons and A. Zalesskii, 'Regular orbits of cyclic subgroups in permutation representations of certain simple groups', *J. Algebra* **256** (2002), 611–625.
- [13] P. Spiga, 'Permutation characters and fixed-point-free elements in permutation groups', *J. Algebra* **299** (2006), 1–7.
- [14] H. Wielandt, 'Problem 6.6, the Kourovka notebook', *Amer. Math. Soc. Transl. Ser. (2)* **121** (1983).

SIMON GUEST, Mathematics, University of Southampton,  
Highfield, Southampton SO17 1BJ, UK  
e-mail: [s.d.guest@soton.ac.uk](mailto:s.d.guest@soton.ac.uk)

ANDREA PREVITALI, Dipartimento di Matematica e Applicazioni,  
University of Milano-Bicocca, Via Cozzi 53, 20125 Milano, Italy  
e-mail: [andrea.previtali@unimib.it](mailto:andrea.previtali@unimib.it)

PABLO SPIGA, Dipartimento di Matematica e Applicazioni,  
University of Milano-Bicocca, Via Cozzi 53, 20125 Milano, Italy  
e-mail: [pablo.spiga@unimib.it](mailto:pablo.spiga@unimib.it)