

Standard Paper

Resolving the phylogenetic relationship between *Parmotrema crinitum* and *Parmotrema perlatum* populations

Ayoub Stelate^{1,2,4} , Ruth Del-Prado², David Alors³ , Hikmat Tahiri⁴, Pradeep K. Divakar²  and Ana Crespo² 

¹Department of Experimental Plant Biology, Charles University Faculty of Science, Viničná 5, 128 44 Prague, Czech Republic; ²Departamento de Farmacología, Farmacognosia y Botánica, Facultad de Farmacia, Universidad Complutense de Madrid, Madrid 28040, Spain; ³Microalgal Biotechnology Laboratory, Jacob Blaustein Institutes for Desert Research, Ben Gurion University of the Negev, Sede-Boker Campus 8499000, Israel and ⁴Department of Biology, Faculty of Sciences, Mohammed V University, Av Ibn Battouta 4, BP 1014 RP. Rabat, Morocco

Abstract

The widespread species *Parmotrema crinitum* (Ach.) M. Choisy and *Parmotrema perlatum* (Huds.) M. Choisy are mainly distinguished by their reproductive strategies. While *P. crinitum* propagates by isidia, *P. perlatum* produces soredia. In this study, we aim to evaluate the phylogenetic relationship between both species and to critically examine their species boundaries. To this purpose, 46 samples belonging to *P. crinitum* and *P. perlatum* were used in our analysis, including 22 for which we studied the morphology and chemistry, before extracting their DNA. We used 35 sequences of the internal transcribed spacer region of nuclear ribosomal DNA (ITS) of *Parmotrema perlatum* from Europe and Africa (20 of which were newly generated), and 11 of *Parmotrema crinitum* from Europe, North America and North Africa (two newly generated). Additionally, 28 sequences of several species from *Parmotrema* were included in the ITS dataset. The ITS data matrix was analyzed using different approaches, such as traditional phylogeny (maximum likelihood and Bayesian analyses), genetic distances, automatic barcode gap discovery (ABGD) and the coalescent-based method poisson tree processes (PTP), in order to test congruence among results. Our results indicate that all samples referred to *P. crinitum* and *P. perlatum* nested in a well-supported monophyletic clade, but phylogenetic relationships among them remain unresolved. Delineations inferred from PTP, ABGD and genetic distance analyses were comparable and suggested that *P. crinitum* and *P. perlatum* belong to the same lineage. Interestingly, two samples of *P. perlatum* separate in a different monophyletic clade, which is supported as a different lineage by all the analyses.

Key words: genetic diversity, ITS, lichen, phylogenetic analyses, species delimitation

(Accepted 25 February 2022)

Introduction

Lichenized fungi form mutualistic relationships with photoautotrophic organisms (photobionts), mainly green algae (*Trebouxiophyceae* and *Ulvophyceae*) and/or cyanobacteria. The lichen symbiosis has been highly successful within fungi, especially *Ascomycota*, with c. 19 400 currently accepted species (Lücking *et al.* 2017) and an estimated diversity of more than 28 000 species (Lücking *et al.* 2009; Leavitt *et al.* 2013). Additionally, lichens are commonly used to assess environmental disturbance, serving as bioindicators of air pollution, forest age and health, and climate change (Nimis *et al.* 2002; Crespo *et al.* 2004; Giordani & Brunialti 2015; Sujetovienė 2015; Sancho *et al.* 2019; Abas 2021).

Recognizing phylogenetic relationships and delimiting species in lichens are crucial for ecological and conservation studies, assessing biotic diversity, and identifying factors driving diversification.

Author for correspondence: Ayoub Stelate. E-mail: Ayoub.stelate@gmail.com

Cite this article: Stelate A, Del-Prado R, Alors D, Tahiri H, Divakar PK and Crespo A (2022) Resolving the phylogenetic relationship between *Parmotrema crinitum* and *Parmotrema perlatum* populations. *Lichenologist* 54, 183–194. <https://doi.org/10.1017/S0024282922000147>

They are also important for future investigations because phylogenetic differences may not be fully reflected in the phenotype.

Traditionally, to infer taxonomic boundaries in lichen-forming fungi, thin-layer chromatography (Culbertson 1972), morphology and the expression of signature secondary metabolites, and isolation and identification of lichen substances have been used (Huneck & Yoshimura 1996; Huneck 1999). However, these characters are highly variable and their homology has proved difficult to assess. For example, it has been shown that apothecial form and spore wall thickness have changed in parallel within *Pertusaria*. The *Pertusaria*-type ascus is plesiomorphic within the *Pertusariaceae* and thus cannot be used to circumscribe *Pertusaria* (Lumbsch & Schmitt 2001). However, with the use of molecular tools, it has been shown that phenotype-based taxonomy may not reflect natural groups, including cases in which morphologically distinct forms formerly recognized as distinct species are shown to represent a polymorphic species (see e.g. Boluda *et al.* 2019), and other cases where several morphologically cryptic species are masked within a single nominal taxon (Del-Prado *et al.* 2016).

Within lichen-forming *Ascomycetes*, *Parmeliaceae* (*Lecanorales*) constitutes one of the largest and best-studied families (Crespo

et al. 2007, 2011; Thell *et al.* 2012; Divakar *et al.* 2015). This family is usually characterized morphologically by a certain type of ascoma ontogeny and the presence of an ascomal structure known as a cupulate exciple (Henssen & Jahns 1974). *Parmeliaceae* also comprises species which are frequently used in biomonitoring studies of atmospheric pollution, such as *Parmelia sulcata*, *Flavoparmelia caperata*, *Parmotrema perlatum* and *Punctelia subrudecta* (Hawksworth & Rose 1970; Crespo *et al.* 2004; De La Cruz *et al.* 2018). The application of phylogenetic analysis based on molecular (DNA) characters to delimit species allows us to determine *a posteriori* which types of phenotypic characters are good predictors of phylogenetic species and demonstrate how these characters evolve in this family and in lichenized fungi in general. These molecular data have led to the recognition of morphologically cryptic species, such as *Parmelia serrana* (Molina *et al.* 2004), *P. barrenoae* (Divakar *et al.* 2005a), *P. encryptata* (Molina *et al.* 2011a) and *Melanelixia californica* (Divakar *et al.* 2010), and conversely also to the union of species traditionally regarded as morphologically distinct (see e.g. Boluda *et al.* 2019). Recently, coalescent-based species delimitation approaches have shown to be well suited to critically evaluate species boundaries in *Parmeliaceae*, as well as lichens in general (Leavitt *et al.* 2015). Furthermore, these methods can accurately display relationships when incomplete lineage sorting and gene tree heterogeneity hide phylogenetic relationships among species (Knowles & Carstens 2007; Camargo *et al.* 2012). Commonly used methods to critically evaluate species boundaries include the poisson tree processes (PTP) model (Zhang *et al.* 2013), the automatic barcode gap discovery (ABGD) (Puillandre *et al.* 2012), the general mixed Yule coalescent model (GMYC) (Pons *et al.* 2006; Monaghan *et al.* 2009), and SpedeSTEM (Ence & Carstens 2011).

Parmotrema (Massalongo 1860) is one of the largest genera in the parmelioid group of the family *Parmeliaceae*. It includes c. 300 described species with an apparent centre of speciation in the Pacific Islands, tropical and subtropical regions of South America (Spielmann & Marcelli 2020). The species of the genus are characterized by a pored epicortex, large thalli with broad lobes, a broad, naked marginal zone on the lower surface, and large, thick-walled, ellipsoid ascospores, sublageniform or filiform conidia (Elix 1993), and (commonly) marginal cilia (Hale 1974). Reproductive strategies vary among *Parmotrema* taxa. Sexual reproduction is restricted to characteristic fungal fruiting bodies (ascmata) producing ascospores. Ascospores are dispersed independently of the photosynthesizing partner (photobiont) and require reacquisition of the appropriate photobiont partner to re-establish the lichenized condition. Species within *Parmotrema* also commonly propagate asexually by means of vegetative diaspores, either isidia or soredia. These specialized vegetative reproductive propagules contain both fungal and algal symbionts, eliminating the need for independent acquisition of the appropriate photobiont.

In a molecular phylogeny of parmotremoid lichens (Blanco *et al.* 2005), a single sample of both *Parmotrema perlatum* (Huds.) M. Choisy and *Parmotrema crinitum* (Ach.) M. Choisy, originally from Portugal, were included and these formed a well-supported monophyletic group, indicating that these specimens could be conspecific. Therefore, a critical evaluation of species boundaries is necessary.

Parmotrema crinitum is characterized by the presence of coralloid branched, apically ciliate isidia or often eciliate isidia (Divakar & Upreti 2005), and the stictic acid complex in the medulla. According to Elix (1994), *P. crinitum* is a cosmopolitan species that is widespread in temperate, tropical regions and even

in high humidity, sub-boreal forests (Elix 1994; Kurokawa & Lai 2001). Many European countries have reported the presence of *P. crinitum* (Jabłońska *et al.* 2009) as have some Asian countries such as Japan (Yoshimura 1974), China (Wei 1991) and Taiwan (Wang-Yang & Lai 1976).

Parmotrema perlatum is a greenish grey foliose lichen, saxicolous or corticolous, loosely adnate, with rounded lobes. It can be recognized by its broad lobes, irregularly branched (7.0 cm), laterally overlapping, with frequent black cilia (0.20–0.50–1.00 × 0.02–0.10) mm, evenly distributed but less common in the lobe apices. Its soralia are marginal and linear, sometimes widely distributed or subcontinuous, concolorous with the thallus, and the medulla is white. The thallus contains atranorin, stictic acid, hypostictic acid, menegazziac acid and norstictic acid (Spielmann & Marcelli 2009). *Parmotrema perlatum* is a widespread species in temperate regions of the Northern and Southern Hemispheres: Asia (Hale 1965; Kurokawa 1991; Kurokawa & Lai 2001), Europe (Hale 1965), Africa (Hale 1965; Swinscow & Krog 1988), North America (Hale 1965; Brodo *et al.* 2001; Nash & Elix 2002), Central America (Hale 1965), South America (Hale 1965), Australia (Kantvilas 2019) and New Zealand (Blanchon 2013).

Morphologically, *P. crinitum* and *P. perlatum* can easily be separated based on the characteristics associated with their different dispersal strategies. *Parmotrema crinitum* has apically ciliate isidia or often eciliate isidia, and *P. perlatum* has marginal soralia. *Parmotrema perlatum* is one of the most common and widely utilized lichens in the Ayurvedic system of medicine and has been overexploited for uses in traditional medicine in India (Kumar & Upreti 2001). In India, this species is currently considered threatened (Divakar & Upreti 2005).

The aim of this study was to evaluate the phylogenetic relationships between *P. crinitum* and *P. perlatum*, and elucidate the possible monophyly between both species. Additionally, PTP, ABGD and genetic distance analyses were included to critically assess species boundaries. The ITS data matrix consisted of a total of 74 sequences of *Parmotrema* species, including 11 of *P. crinitum* and 35 of *P. perlatum*, plus two outgroup sequences of *Crespoa carneopruinata*. Morphological and chemical features of all samples were critically examined.

Material and Methods

Taxon sampling

Sequence data of the ITS locus were analyzed from 76 specimens, of which 11 sequences were of *Parmotrema crinitum* (two individuals were newly sequenced and nine sequences downloaded from GenBank) and 35 of *Parmotrema perlatum* (20 individuals were sequenced in this study and 15 sequences downloaded from GenBank). The new specimens were collected from distant geographical regions throughout the species distributions (Table 1). Additionally, 30 ITS sequences from 30 specimens, belonging to 14 different species (6 were newly sequenced and 24 sequences downloaded from GenBank), are included in this study to test the monophyly of both species within the genus. This total included *Crespoa carneopruinata* which was selected as outgroup since it has previously been shown to be the sister group of *Parmotrema* (Divakar *et al.* 2015). Details of the material studied, including GenBank Accession numbers, are shown in Table 1. Species recognition was based mainly on morphological and chemical characters.

Table 1. *Parmotrema* species used in this study with GenBank Accession numbers of the ITS sequences and voucher information for the specimens. *Crespoa carneopruinata* was included in the phylogenetic analyses as outgroup. * = newly generated sequences from specimens for which morphological and chemical characters are given in Table S1 (available online).

Species	Locality	Voucher specimen	Collector	GenBank Accession number
<i>Crespoa carneopruinata</i> 1	Costa Rica	15171a	R. Lücking	KP888204
<i>C. carneopruinata</i> 2	Costa Rica	15171a	R. Lücking	EF042904
<i>Parmotrema clavuliferum</i>	Gomera, Islas Canarias, Spain	MAF-Lich 20689	A. Crespo, A. Santos	AY586577
<i>P. crinitum</i> 1	North Carolina, USA	44262	J. C. Lendemer	KP943761
<i>P. crinitum</i> 2	La Palma, Islas Canarias, Spain	MAF-Lich 16189	A. Crespo, A. Santos	HM017032
<i>P. crinitum</i> 3	North Carolina, USA	43711	J. C. Lendemer	KP943759
<i>P. crinitum</i> 4	North Carolina, USA	43863	J. C. Lendemer	KP943760
<i>P. crinitum</i> 5	La Palma, Islas Canarias, Spain	MAF-Lich 16188	A. Crespo, A. Santos	HM017033
<i>P. crinitum</i> 6*	Madeira, Portugal	MAF-Lich 20704	P. K. Divakar, M. Talavera	ON312512
<i>P. crinitum</i> 7	Lisboa, Igreja Nova, Portugal	MAF-Lich 6061	P. K. Divakar, M. Talavera	AY586565
<i>P. crinitum</i> 8*	Madeira, Portugal	MAF-Lich 20703	A. Crespo, A. Santos	ON312511
<i>P. crinitum</i> 9	Tenerife, Islas Canarias, Spain	MAF-Lich 16174	A. Crespo, A. Santos	HM017030
<i>P. crinitum</i> 10	Tenerife, Islas Canarias, Spain	MAF-Lich 16170	A. Crespo, A. Santos	HM017028
<i>P. crinitum</i> 11	Tenerife, Islas Canarias, Spain	MAF-Lich 16169	A. Crespo, R. Del-Prado, A. Santos	HM017029
<i>P. dilatatum</i> 1*	Gomera, Islas Canarias, Spain	MAF-Lich 20698	A. R. Burgaz, M. A. Carrasco	ON312500
<i>P. dilatatum</i> 2*	China	MAF-Lich 10164	D. L. Hawksworth	ON312501
<i>P. flavotinctum</i> *	Pichincha, Ecuador	54174	P. v. d. Boom	ON312516
<i>P. grayanum</i>	La Palma, Islas Canarias, Spain	MAF-Lich 16190	A. Crespo, R. Del-Prado, P. K. Divakar	HM017026
<i>P. haitiense</i>	Australia	MAF-Lich 7657	S. H. Louwhoff	AY581055
<i>P. hypoleucinum</i> 1	Spain	MAF-Lich 7637	A. Crespo, P. K. Divakar	AY586567
<i>P. hypoleucinum</i> 2	Morocco	MAF-Lich 16142	A. Crespo, P. K. Divakar, H. Tahiri,	HM017035
<i>P. hypoleucinum</i> 3	Morocco	MAF-Lich 16141	A. Crespo, P. K. Divakar, H. Tahiri	HM017037
<i>P. hypoleucinum</i> 4	Morocco	MAF-Lich 16147	A. Crespo, P. K. Divakar, H. Tahiri	HM017036
<i>P. mellissii</i> *	Pichincha, Ecuador	54312	P. v. d. Boom	ON312515
<i>P. paulense</i> *	Medellin, Antioquia, Colombia	MAF-Lich 20060	P. K. Divakar	ON312513
<i>P. perforatum</i>	USA	7983	M. Cole	AY586568
<i>P. perlatum</i> 1	Medium Atlas, Fes, Morocco	MAF-Lich 16145	A. Crespo, P. K. Divakar, H. Tahiri	HM017044
<i>P. perlatum</i> 2*	Akchour, Chaouen, Morocco	MAF-Lich 20712	A. Stelate, H. Tahiri	ON312519
<i>P. perlatum</i> 3*	Akchour, Chaouen, Morocco	MAF-Lich 20707	A. Stelate, H. Tahiri	ON312518
<i>P. perlatum</i> 4*	Akchour, Chaouen, Morocco	MAF-Lich 20711	A. Stelate, H. Tahiri	ON312521
<i>P. perlatum</i> 5*	Akchour, Chaouen, Morocco	MAF-Lich 20708	A. Stelate, H. Tahiri	ON312523
<i>P. perlatum</i> 6	Chaouen, Morocco	MAF-Lich 16146	A. Crespo, P. K. Divakar, H. Tahiri	HM017046
<i>P. perlatum</i> 7*	Gomera, Islas Canarias, Spain	MAF-Lich 20700	A. Crespo, A. Santos	ON312505
<i>P. perlatum</i> 8	Eregli Vakif, Zonguldak, Turkey	MAF-Lich 16192	P. K. Divakar, R. Del-Prado	HM017041
<i>P. perlatum</i> 9	Tenerife, Islas Canarias, Spain	MAF-Lich 16173	A. Crespo, R. Del-Prado	HM017042
<i>P. perlatum</i> 10*	Chaouen, Morocco	MAF-Lich 20709	A. Stelate, H. Tahiri	ON312520
<i>P. perlatum</i> 11*	Chaouen, Morocco	MAF-Lich 20710	A. Stelate, H. Tahiri	ON312522
<i>P. perlatum</i> 12	Lisboa, Igreja Nova, Portugal	MAF-Lich 6965	P. K. Divakar, M. Talavera	AY586566
<i>P. perlatum</i> 13	La Palma, Islas Canarias, Spain	MAF-Lich 16179	A. Crespo, R. Del-Prado	HM017040
<i>P. perlatum</i> 14	La Palma, Islas Canarias, Spain	MAF-Lich 16187	A. Crespo, R. Del-Prado	HM017043
<i>P. perlatum</i> 15	Chaouen, Morocco	MAF-Lich 16203	A. Crespo, R. Del-Prado	HM017044
<i>P. perlatum</i> 16*	Gomera, Islas Canarias, Spain	MAF-Lich 20699	A. Crespo, A. Santos	ON312506
<i>P. perlatum</i> 17*	Gomera, Islas Canarias, Spain	MAF-Lich 20693	A. Crespo, A. Santos	ON312510
<i>P. perlatum</i> 18*	Gomera, Islas Canarias, Spain	MAF-Lich 20691	A. Crespo, A. Santos	ON312508
<i>P. perlatum</i> 19	La Palma, Islas Canarias, Spain	MAF-Lich 16186	A. Crespo, R. Del-Prado	HM017048
<i>P. perlatum</i> 20	Tenerife, Islas Canarias, Spain	MAF-Lich 16162	A. Crespo, R. Del-Prado	HM017052
<i>P. perlatum</i> 21*	Gomera, Islas Canarias, Spain	MAF-Lich 20702	A. Crespo, A. Santos	ON312503
<i>P. perlatum</i> 22*	Gomera, Islas Canarias, Spain	MAF-Lich 20696	A. Crespo, A. Santos	ON312496
<i>P. perlatum</i> 23	Tenerife, Islas Canarias, Spain	MAF-Lich 16172	A. Crespo, R. Del-Prado	HM017049
<i>P. perlatum</i> 24	Tenerife, Islas Canarias, Spain	MAF-Lich 16166	A. Crespo, R. Del-Prado	HM017051
<i>P. perlatum</i> 25*	Gomera, Islas Canarias, Spain	MAF-Lich 20690	A. Crespo, A. Santos	ON312507
<i>P. perlatum</i> 26*	Gomera, Islas Canarias, Spain	MAF-Lich 20695	A. Crespo, A. Santos	ON312497

(Continued)

Table 1. (Continued)

Species	Locality	Voucher specimen	Collector	GenBank Accession number
<i>P. perlatum</i> 27	Tenerife, Islas Canarias, Spain	MAF-Lich 16168	A. Crespo, R. Del-Prado	HM017050
<i>P. perlatum</i> 28*	Gomera, Islas Canarias, Spain	MAF-Lich 20701	A. Crespo, A. Santos	ON312504
<i>P. perlatum</i> 29	La Palma, Islas Canarias, Spain	MAF-Lich 16191	A. Crespo, R. Del-Prado	HM017039
<i>P. perlatum</i> 30*	Gomera, Islas Canarias, Spain	MAF-Lich 20697	A. Crespo, A. Santos	ON312502
<i>P. perlatum</i> 31*	Gomera, Islas Canarias, Spain	MAF-Lich 20694	A. Crespo, A. Santos	ON312509
<i>P. perlatum</i> 32*	Gomera, Islas Canarias, Spain	MAF-Lich 20688	A. Crespo, A. Santos	ON312498
<i>P. perlatum</i> 33*	Gomera, Islas Canarias, Spain	MAF-Lich 20692	A. Crespo, A. Santos	ON312499
<i>P. perlatum</i> 34*	Lisboa, Igreja Nova, Portugal	MAF-Lich 20706	P. K. Divakar, A. Agudo	ON312517
<i>P. perlatum</i> 35	Tenerife, Islas Canarias, Spain	MAF-Lich 16171	A. Crespo, A. Santos	HM017027
<i>P. pseudoreticulatum</i>	Eastern Cape, South Africa	MAF-Lich 10287	A. Crespo, A. Santos	AY642828
<i>P. reticulatum</i>	Bluf Area, Kenya	MAF-Lich 10121	P. K. Divakar	HM016956
<i>P. robustum</i>	Portugal	MAF-Lich 10166	P. K. Divakar	AY586569
<i>P. subtinctorium</i> 1	India	GPCC02-000696	P. K. Divakar	AY586558
<i>P. subtinctorium</i> 2*	Western, Bluf Area, Kenya	MAF-Lich 20705	A. Crespo, R. Del-Prado	ON312514
<i>P. tinctorum</i> 1	Shizuoka, Japan	TNS-L-Y.O. 5399	Y. Ohmura	AB177404
<i>P. tinctorum</i> 2	Shizuoka, Japan	TNS-L-Y.O. 5375A	Y. Ohmura	AB177401
<i>P. tinctorum</i> 3	Inhaca Island, Mozambique	MAF-Lich 18193	B. Roca-Valiente	KF129447
<i>P. tinctorum</i> 4	Mt Tsukuba, Japan	MAF-Lich 18163	A. Crespo, P. K. Divakar	KF129414
<i>P. tinctorum</i> 5	Chamoli, India	MAF-Lich 18170	P. K. Divakar	KF129422
<i>P. tinctorum</i> 6	Kanagawa, Japan	TNS-L-Y.O. 5947	M. Takeda	KF129464
<i>P. tinctorum</i> 7	Kanagawa, Japan	TNS-L-Y.O. 5486	Y. Ohmura	KF129463
<i>P. tinctorum</i> 8	Nara, Japan	TNS-L-Y.O. 5813	Y. Ohmura	KF129462
<i>P. tinctorum</i> 9	Osaka, Japan	TNS-L-Y.O. 5480	N. Hamada	KF129460
<i>P. tinctorum</i> 10	Inhaca Island, Mozambique	MAF-Lich 18196	B. Roca-Valiente, A. Lumbreras	KF129450

Chemistry and morphology

Thallus morphology of all new specimens of *Parmotrema crinitum* and *P. perlatum* included in the molecular analyses was studied using a Nikon SMZ-1000 stereomicroscope to identify morphological characteristics of the thallus: lobe shape, isidia, soralia, cilia and rhizines. This is because *P. crinitum* and *P. perlatum* are traditionally differentiated based on these features. Photographs were taken with a Nikon 105 mm f/2.8D AF Micro-Nikkor lens coupled to a Nikon D90 camera.

Spot tests were carried out on the medulla with usual chemical reagents such as aqueous potassium hydroxide (K), Steiner's stable paraphenylenediamine (PD) and aqueous calcium hypochlorite (C).

Thin-layer chromatography (TLC) was carried out following Orange *et al.* (2010). We used TLC solvent system C (200 ml toluene/30 ml acetic acid) (Elix & Ernst-Russell 1993), with concentrated acetone extracts at 50 °C spotted onto silica gel 60 F254 aluminum sheets (Merck, Darmstadt). The aluminum sheets were dried for 10 min in an acetic acid atmosphere to maximize resolution.

DNA extraction, PCR and sequencing

Total DNA was extracted from a single, clean (under a dissecting microscope) lichen lobe using the DNeasy Plant Mini Kit (Qiagen, Barcelona) with a slight modification to the manufacturer's instruction (Crespo *et al.* 2001). Genomic DNA (5–25 ng, quantified using a quantitative PCR machine) was used for PCR amplifications of the ITS region. Standard PCR amplifications were conducted in 25 µl reaction volumes containing 12.5 µl of Master Mix (50 units/ml of Taq DNA polymerase supplied in a proprietary reaction buffer (pH 8.5), 400 µM dATP, 400 µM dGTP, 400 µM dCTP, 400 µM dTTP, 3 mM MgCl₂), and 1.5 µl of each primer at 10 µM, 4.5 µl of water (H₂O) and 5 µl of DNA template. Fungal nuclear internal transcribed spacer (ITS) rDNA was amplified using the primer pair ITS1F (5' [CTT GGT CAT TTA GAG GAA GTA A] 3') (Gardes & Bruns 1993)/ LR1 (5' [GGT TGG TTT CTT TTC CT] 3') (Vilgalys & Hester 1990). We also tried primer pair ITS1-LM (5' [GAA CCT GCG GAA GGA TCA TT] 3') (Myllys *et al.* 2001) and ITS2-KL (5' [ATG CTT AAG TTC AGC GGG TA] 3') (Lohtander *et al.* 1998), but we had better results with primer pair ITS1F/LR1. For any failed samples, we tried a nested PCR using the primer pair ITS1F/LR1 for the first PCR, and 5 µl of this PCR product as DNA template for the second PCR using ITS1-LM/ITS2-KL. With this nested PCR, we aimed to amplify a smaller region than with the previous primers. However, it did not yield any additional products.

Amplification was run in an automatic thermocycler (Techne Progene, Jepson Bolton & Co., Watford, Herts, UK) using the following parameters: initial denaturation at 94 °C for 5 min followed by 35 cycles at 94 °C for 1 min, 54 °C (ITS1F/LR1) for 1 min, and 72 °C for 1 min 30 s, with a final extension at 72 °C for 10 min. Amplification products were visualized on 1% agarose gel stained with SYBR green DNA (Life Technologies Corporation, Grand Island, New York, USA), and subsequently purified using ExoSAP-IT (GE Healthcare, Chalfont St Giles, UK) according to the manufacturer's instructions. Sequencing was performed using BigDye Terminator reaction kit (ABI PRISM, Applied Biosystems, Waltham, Massachusetts, USA). Cycle sequencing reactions

were performed with the same sets of primers used in the amplification step. Sequencing reactions were electrophoresed on a 3730 DNA Analyzer (Applied Biosystems) at the Unidad de Genómica (Parque Científico de Madrid).

Sequence alignments and phylogenetic analyses

Sequence fragments generated for this study were assembled and edited using the program SeqMan v. 7 (Lasergene R, DNASTAR, Madison, Wisconsin, USA). Sequence identity was confirmed using the megaBLAST search function in GenBank. For the alignment we prepared a matrix of 480 base pairs (bp). Then we used the program MAFFT v. 7 (Katoh & Standley 2013) with the parameters set to default. If sequences had different lengths, only the part shared by all the sequences was used, and after manually removing ambiguously aligned nucleotide positions, we kept 451 unambiguously aligned base pairs for the final matrix that we used as input for the phylogenetic reconstruction. The alignment was analyzed using maximum likelihood (ML) and a Bayesian Markov chain Monte Carlo (B/MCMC) approach. The ML analysis was performed using an online version of the program RAxML-HPC BlackBox v. 8.2.8 (Stamatakis 2006; Stamatakis *et al.* 2008), implemented on the Cipres science gateway (<https://www.phylo.org/portal2/home.action>; Miller *et al.* 2010). We used a GTRGAMMA model, which includes a parameter (Γ) for rate heterogeneity among sites but chose not to include a parameter to estimate the proportion of invariable sites (Stamatakis 2006; Stamatakis *et al.* 2008). Support values were assessed using the 'rapid bootstrapping' option with 1000 replicates.

For the Bayesian reconstruction, we used MrBayes v. 3.2.1 (Ronquist & Huelsenbeck 2003). The analysis was performed assuming a discrete gamma distribution with six rate categories (GTR + G). The nucleotide-substitution model parameters were selected using the Akaike Information Criterion as implemented in jModelTest (Posada 2008), molecular clock not assumed. A run with 10 million generations, starting with a random tree and employing 12 simultaneous chains, was executed. Trees were saved to a file every 200th generation. The first 2 million generations (i.e. 20 000 trees) were deleted as the 'burn-in' of the chains. We plotted the log-likelihood scores of sample points against generations using Tracer v. 1.5 (Rambaut & Drummond 2007) and determined that stationarity had been achieved when the log-likelihood values of the sample points reached an equilibrium value (Huelsenbeck & Ronquist 2001). The trees obtained before stationarity were discarded. Posterior probabilities (PPs) were obtained from the 50% majority-rule consensus of sampled trees after excluding the initial 20% as burn-in. Only clades that received bootstrap support $\geq 70\%$ in the ML analyses and PP ≥ 0.95 were considered strongly supported. The phylogenetic tree was drawn using FigTree v. 1.2.3 (Rambaut 2009) and modified with CoreDRAW v. 8.

Candidate species identification

In order to establish candidate species limits in the phylogenetic tree, three computational approaches were used:

1) Poisson tree processes (PTP): this method does not require an ultra-metric tree, as the transition point between intra- and inter-specific branching rates is identified directly using the number of nucleotide substitutions (Zhang *et al.* 2013). PTP

incorporates the number of substitutions in the model of speciation and assumes that the probability of a substitution giving rise to a speciation event follows a Poisson distribution. The branch lengths of the input tree are supposed to be generated by two independent classes of Poisson events, one corresponding to speciation and the other to coalescence. The ML phylogeny of the ITS dataset obtained with RAxML was used as the input trees to run PTP species delimitation analysis on the PTP web server (<http://species.h-its.org/ptp/>). We ran the PTP analysis for 100 000 MCMC generations, with a thinning value of 100 and a burn-in of 25%. Outgroup taxa were removed for species delimitation.

- 2) Automatic barcode gap discovery (ABGD): this is an automatic procedure that sorts the sequences into hypothetical species based on the barcode gap. This method automatically finds the distance where the barcode gap is located (Puillandre *et al.* 2012). The ABGD method was carried out on the ITS dataset using the web interface at <http://wwwabi.snv.jussieu.fr/public/abgd/abgdweb.html>. Default parameters were chosen using Kimura 2-parameter (K2P) distances that correct for transition rate bias (relative to transversions) in the substitution process (Kimura 1980). The default for the minimum relative gap width was set to different values between 0.1 and 0.15.
- 3) Genetic distances: pairwise ML distances (given as the number of nucleotide substitutions per site) among the ITS rDNA sequences of *Parmotrema crinitum* and *P. perlatum* were calculated. Genetic distances were calculated with TREE-PUZZLE v. 5.2 (Schmidt *et al.* 2002) using the GTR model of nucleotide substitution, assuming a discrete gamma distribution with six rate categories. The program generates an output file which consists of a triangular matrix with all pairwise distances between all the samples included. This matrix was visualized with Microsoft Excel 2010 and genetic distances between different specimens of *P. crinitum* and *P. perlatum* were manually identified. Candidate species were proposed based on the threshold of 0.016 substitutions per site (s/s) which separates intra- and interspecific distances in parmelioid lichens (Del-Prado *et al.* 2010). Distance values in the matrix ≤ 0.016 s/s have been considered as the values between samples of a single species. By using the Excel filter, we separated values ≤ 0.016 , providing for every specimen included in the analysis a group of specimens that share the values characterizing its species range.

Results

Morphological and chemical analyses

Morphological and chemical characters of the specimens (*P. crinitum* and *P. perlatum*) newly generated and included in the phylogenetic analysis are presented in Supplementary Material Table S1, available online. *Parmotrema crinitum* and *P. perlatum* could not be distinguished based on the colour of the thalli, presence of cilia (Fig. 1A & B) and colour of the lower surface (Fig. 1E & F). However, the reproductive structures allowed unequivocal separation into the two species; *Parmotrema perlatum* has soralia (Fig. 1C) and *P. crinitum* has isidia (Fig. 1D).

The TLC test showed that both species contained the stictic acid complex (stictic acid, menegazzic acid and hypostictic acid) and atranorin. The spot tests of *P. crinitum* were similar to those of *P. perlatum* (Fig. 2, Supplementary Material Table S1).




Fig. 1. Main morphological characters of *P. perlatum* (A, C, E) and *P. crinitum*. (B, D, F), as indicated with arrows. A & B, black, simple cilia. C, marginal soralia concolorous with the thallus. D, isidia with apical cilia. E & F, black lower surface with brown naked zone peripherally. In colour online.

However, the samples *Parmotrema perlatum* 34 and *P. perlatum* 35 lacked menegazzic acid (nos 10 and 12 in Fig. 2). The spot test was congruent with other samples of *P. perlatum*, and the morphology of both samples showed the same characteristics of *P. perlatum*, although the cilia were not as abundant as is usual in this species.

Phylogenetic analysis

The ITS data matrix consisted of a total of 74 sequences of *Parmotrema* species, including 46 of *P. crinitum* and *P. perlatum* (11 sequences of *P. crinitum* and 35 sequences of *P. perlatum*), with two sequences of *Crespoa carneopruinata* were included as

outgroup (Table 1). The final matrix used as input for the phylogenetic reconstruction contained 451 unambiguously aligned base pairs (bp). The tree reconstruction (Fig. 3) comprised the following species of *Parmotrema*, included to test the monophyly of *P. crinitum* and *P. perlatum*: *Parmotrema clavuliferum*, *P. dilatatum*, *P. flavotinctum*, *P. grayanum*, *P. haitiense*, *P. hypoleucinum*, *P. mellissii*, *P. paulense*, *P. perforatum*, *P. pseudoreticulatum*, *P. reticulatum*, *P. robustum*, *P. subtinctorium* and *P. tinctorum*.

The phylogenetic trees estimated from ML and Bayesian methods did not show any well-supported conflict; ML topologies are presented with bootstrap and B/MCMC analysis with posterior probability (ML bootstrap $\geq 70\%$; PP ≥ 0.95 in B/MCMC analysis) (Fig. 3). The LnL value was -1938.068210 for ML, and




Fig. 2. Thin-layer chromatography profile of *Parmotrema* species (included in the DNA analysis) in solvent system C. P is the control, *Pleurosticta acetabulum*. Lanes 1 & 7, *P. dilatatum*. Lane 2, *P. flavotinctum*. Lane 3, *P. mellissii*. Lane 4, *P. paulense*. Lanes 5, 6, 8, 11 & 19, *P. crinitum*. Lane 9, empty. Lanes 10 & 12, *Parmotrema perlatum*. Lanes 13, 14, 15, 16, 17, 18 & 20, *P. perlatum*; these have the same TLC result as *P. crinitum* but were distinguished from each other based on the morphology. a = norstictic acid; b = atranorin; c = protocetraric acid; d = unknown; e = hypostictic acid; f = stictic acid; g = menegazzic acid. In colour online.

–2184.428 for the Bayesian analysis. The phylogenetic analysis showed that all samples of *Parmotrema* included in this analysis formed a monophyletic group. Within this genus the samples of *Parmotrema crinitum* and *P. perlatum* were grouped in one monophyletic group (clade A), with the exception of two samples of *P. perlatum* that were grouped in a separate clade B. While clade A comprised specimens of *P. crinitum* and *P. perlatum* collected from various geographical regions, clade B included only two specimens of *P. perlatum* collected from Tenerife (Canary Islands) and Lisbon (Portugal).

Although four well-supported monophyletic clusters can be recognized in clade A, the phylogenetic relationships among them remained unresolved (Fig. 3).

Cluster A1 (Fig. 3) contained 15 specimens of *Parmotrema perlatum*, most of them collected from Morocco, except four from the Canary Islands and two from Portugal and Turkey.

Cluster A2 included eight specimens of *Parmotrema crinitum* from different geographical regions (Table 1). Cluster A3 grouped 18 samples of *Parmotrema perlatum* from the Canary Islands (Tenerife, Palma, Gomera), and cluster A4 included three samples of *P. crinitum* from the Canary Islands.

Identifying candidate species

We used the same RaxML tree obtained from the phylogenetic analysis to illustrate the delimitation of putative species recognized by the different approaches conducted with the ITS dataset. ABGD, PTP and genetic distance analyses applied to the ITS dataset detected two candidate species that corresponded to the well-supported clades A (including *P. crinitum* + *P. perlatum*) and B (including *P. perlatum*) obtained in the phylogenetic analysis (Fig. 3).




Fig. 3. Phylogenetic relationships between *Parmotrema crinitum* and *P. perlatum*. The green (thickened) line indicates the clade supported by maximum likelihood (ML) and Bayesian analyses; * = clades supported only by ML. ** = clades supported only by Bayesian analysis. Alphanumeric labels indicate clades and clusters. Species delimitation scenarios obtained from PTP, ABGD and genetic distances are indicated in columns to the right. In colour online.

Discussion

Previous molecular phylogenetic studies have proposed the monophyly of *Parmotrema crinitum* and *Parmotrema perlatum* (Blanco *et al.* 2006; Crespo *et al.* 2010). The primary goal of the present investigation was to evaluate the phylogenetic relationship between the two species using ML and Bayesian analyses. Additionally, we used different approaches to assess the species boundary, such as genetic distances based on the threshold of 0.015–0.017 s/s established by Del-Prado *et al.* (2010) to measure intra- and interspecific genetic distances in parmelioid lichens, and separate ranges of intra- and interspecific divergence. This threshold was established using both phylogenetically and morphologically well-delimited species (for details see Del-Prado *et al.* (2010)). We also used the poisson tree processes (PTP) model (Zhang *et al.* 2013) and the automatic barcode gap discovery (ABGD; Puillandre *et al.* 2012). Furthermore, we combined ecological, biogeographical, morphological and chemical data.

Most of the samples belonging to *P. crinitum* and *P. perlatum* were recovered in a well-supported monophyletic clade (clade A, Fig. 3), with the exception of two samples of *P. perlatum*, one from Tenerife (Canary Islands) and one from Lisbon (Portugal), that are separated in the monophyletic clade B, supported by ML and B/MCMC analysis. Morphologically, *P. perlatum* in clade A has more cilia compared to *P. perlatum* in clade B.

Thin-layer chromatography (TLC) revealed that the samples in clade B (nos 10 and 12 in Fig. 2) differ chemically from the samples of *Parmotrema perlatum* grouped in clade A. Several studies have shown that environmental factors, such as light, temperature, pH and culture media, can influence the secondary metabolism in lichens (BeGora & Fahselt 2001). However, we have samples from the same localities and same conditions as the samples in clade A. Furthermore, ABGD, PTP and genetic distance analyses supported clade B as a new sister group to clade A (Fig. 3), suggesting polyphyly, a common phenomenon in *Parmeliaceae* in general (Lumbsch & Leavitt 2011; Leavitt *et al.* 2016) and *Parmotrema* in particular (Divakar *et al.* 2005b; Del-Prado *et al.* 2016, 2019; Widhelm *et al.* 2016).

Clade A was split into 4 groups; however, the phylogenetic relationships among them remained unresolved (Fig. 3). Previous studies have shown that phylogenetic analyses alone are insufficient to explain phylogenetic relationships within *Parmeliaceae*. For example, based only on maximum parsimony and Bayesian analyses the sorediate *P. sulcata* was shown to belong to the same clade as the isidiate *P. squarrosa* (Molina *et al.* 2004), leaving the authors unable to reach any conclusion regarding species boundaries. Similarly, phylogenetic analyses were insufficient to resolve genetic variability among *Parmelia saxatilis* specimens; whereas samples from distant geographical regions formed a monophyletic group, samples from neighbouring localities were separated (Crespo *et al.* 2002; Molina *et al.* 2011b, 2017). A study on *Usnea perpusilla* demonstrated that it was necessary to use a combined approach with molecular and morphological data to assess species boundaries in closely related and morphologically variable species (Wirtz *et al.* 2008). For this reason, coalescent-based species delimitation analyses have been applied with the goal of explaining relationships among clades and delimiting species boundaries (Parnmen *et al.* 2012; Leavitt *et al.* 2013).

PTP, ABGD and genetic distance analyses supported clades A and B as putative distinct species. The values obtained were within the interspecific ranges (genetic distances as defined in

Del-Prado *et al.* (2010)), which would support the two samples of *P. perlatum* in clade B as a separate species, different from clade A. Focusing on clade A, our species delimitation approaches (PTP, ABGD and genetic distance) supported *P. crinitum* and *P. perlatum* as one species. However, based only on one molecular marker (ITS) and the various phylogenetic analyses and species delimitation approaches used, can we consider *Parmotrema perlatum* conspecific with *P. crinitum*?

Similar cases have been reported in the *Parmotrema perforatum* species complex (Widhelm *et al.* 2016), and authors have suggested that the phylogenetic relationships between sexual and asexual populations of this species group could be more complex than previously assumed. They also suggest that traditional tools based on reproductive mode and secondary metabolites (Culberson & Culberson 1973) are no longer the key to identify species such as *P. perforatum* (Widhelm *et al.* 2016).

A previous study on *Umbilicaria* (Ott *et al.* 2004) focused on *Umbilicaria kappenii* and *U. antarctica*, which are distinguished only by their reproductive strategies. *Umbilicaria antarctica* propagates by thalloconidia and *U. kappenii* exhibits a variety of asexual propagules: soredia, adventive lobes and sorediate thallies. To infer phylogenetic relationships between both species, the authors used molecular data from three loci. Results indicated that all samples morphologically referred to *U. antarctica* and *U. kappenii* form a monophyletic group and they proposed placing *U. kappenii* into synonymy with *U. antarctica* (Ott *et al.* 2004). In *Parmelia*, Molina *et al.* (2011b) rejected the previous hypothesis that *P. sulcata* and *P. squarrosa* form a monophyletic group (Molina *et al.* 2004) and based on phylogenetic analyses and species delimitation approaches confirmed that *P. sulcata* is not conspecific with *P. squarrosa*. In addition, *P. squarrosa* is a reproductively isolated lineage and genetic distances clearly separate this from other *Parmelia* species (Del-Prado *et al.* 2010). Within *Usnea*, studies have suggested that *Usnea subfloridana* was a secondary species derived from *U. florida* (Clerc 1984, 1987, 1997; Purvis *et al.* 1992). *Usnea florida* and *U. subfloridana* show many morphological similarities but they have different reproductive strategies. *Usnea florida* always displays many apothecia and produces no specialized asexual propagules. *Usnea subfloridana* has soralia, isidiomorphs and occasionally apothecia. Multilocus phylogenetic analyses based on sequences of the ITS, IGS and LSU regions of the nuclear ribosomal DNA and the gene coding for β -tubulin, *Mcm7*, *RPB1* and *RPB2* showed that specimens of the two morphospecies formed one monophyletic intermixed group, and not two groups corresponding to morphology (Articus *et al.* 2002; Mark *et al.* 2016). This topology was further confirmed with a coalescent-based species delimitation approach (Mark *et al.* 2016). Authors have suggested that they could be conspecific but taxonomic conclusions must await further study. Moreover, a recent study using RADseq data suggests that closely related lichen species may need genome-wide data to test their species boundaries (Grewe *et al.* 2018).

Traditionally it was thought that asexually reproducing species in lichens, and in filamentous fungi in general, was an evolutionary dead end (Normark *et al.* 2003). However, recent molecular studies have demonstrated that lineages with vegetative propagation can also present high genetic diversity (e.g. *Parmelia sulcata* (Molina *et al.* 2011b), *Parmotrema reticulatum* (Del-Prado *et al.* 2016)). However, even in the absence of sexual reproduction, lichens can exchange photobionts and this process could provide opportunities for gene transfer (Piercey-Normore 2006; Nelsen & Gargas 2008).

While our study confirmed the monophyly of an intermixed clade of *Parmotrema crinitum* and *P. perlatum*, the taxonomic conclusion must await additional studies including more markers. The phylogenetic tree lacked ML or B/MCMC support for other widely accepted *Parmotrema* species, such as *P. pseudoreticulatum* and *P. reticulatum*. A more comprehensive taxon sampling and additional molecular markers will therefore be needed before making a formal taxonomic decision on the status of clade B.

Acknowledgements. This study was supported by the Spanish Ministerio de Ciencia e Innovacion (PID2019-105312GB-I00) and the Santander-Universidad Complutense de Madrid (PR87/19-22637 and G/6400100/3000).

Author ORCIDs.  Ayoub Stelate, 0000-0001-8929-3046; David Alors, 0000-0001-7288-9521; Pradeep Kumar Divakar, 0000-0002-0300-0124; Ana Crespo, 0000-0002-5271-0157.

Supplementary Material. To view Supplementary Material for this article, please visit <https://doi.org/10.1017/S0024282922000147>.

References

- Abas A (2021) A systematic review on biomonitoring using lichen as the biological indicator: a decade of practices, progress and challenges. *Ecological Indicators* **121**, 107197.
- Articus K, Mattsson JE, Tibell L, Grube M and Wedin M (2002) Ribosomal DNA and B-tubulin data do not support the separation of the lichens *Usnea florida* and *U. subfloridana* as distinct species. *Mycological Research* **106**, 412–418.
- BeGora MD and Fahselt D (2001) Usnic acid and atranorin concentrations in lichens in relation to bands of UV irradiance. *Bryologist* **104**, 134–140.
- Blanchon DJ (2013) Auckland lichens. *Auckland Botanical Society Journal* **68**, 21–27.
- Blanco O, Crespo A, Divakar PK, Elix JA and Lumbsch HT (2005) Molecular phylogeny of parmotrema-like lichens (*Ascomycota*, *Parmeliaceae*). *Mycologia* **97**, 150–159.
- Blanco O, Crespo A, Ree RH and Lumbsch HT (2006) Major clades of parmelioid lichens (*Parmeliaceae*, *Ascomycota*) and the evolution of their morphological and chemical diversity. *Molecular Phylogenetics and Evolution* **39**, 52–69.
- Boluda CG, Rico VJ, Divakar PK, Nadyeina O, Myllys L, McMullin RT, Zamora JC, Scheidegger C and Hawksworth DL (2019) Evaluating methodologies for species delimitation: the mismatch between phenotypes and genotypes in lichenized fungi (*Bryoria* sect. *Implexae*, *Parmeliaceae*). *Persoonia* **42**, 75–100.
- Brodo IM, Sharnoff SD and Sharnoff S (2001) *Lichens of North America*. New Haven, Connecticut: Yale University Press.
- Camargo A, Morando M, Avila LJ and Sites JW (2012) Species delimitation with ABC and other coalescent-based methods: a test of accuracy with simulations and an empirical example with lizards of the *Liolaemus darwini* complex (Squamata: *Liolaemidae*). *Evolution* **66**, 2834–2849.
- Clerc P (1984) Contribution à la révision de la systématique des Usnées (*Ascomycotina*, *Usnea*) d'Europe. 1. *Usnea florida* (L.) Wigg. emend. Clerc. *Cryptogamie, Bryologie et Lichénologie* **5**, 333–360.
- Clerc P (1987) Systematics of the *Usnea fragilesens* aggregate and its distribution in Scandinavia. *Nordic Journal of Botany* **7**, 479–495.
- Clerc P (1997) Notes on the genus *Usnea* Dill. ex Adanson. *Lichenologist* **29**, 209–215.
- Crespo A, Blanco O and Hawksworth DL (2001) The potential of mitochondrial DNA for establishing phylogeny and stabilizing generic concepts in the parmelioid lichens. *Taxon* **50**, 807–819.
- Crespo A, Molina MC, Blanco O, Schroeter B, Sancho LG and Hawksworth DL (2002) rDNA ITS and β -tubulin gene sequence analyses reveal two monophyletic groups within the cosmopolitan lichen *Parmelia saxatilis*. *Mycological Research* **106**, 788–795.
- Crespo A, Divakar PK, Argüello A, Gasca C and Hawksworth DL (2004) Molecular studies on *Punctelia* species of the Iberian Peninsula, with an emphasis on specimens newly colonizing Madrid. *Lichenologist* **36**, 299–308.
- Crespo A, Lumbsch HT, Mattsson JE, Blanco O, Divakar PK, Articus K, Wiklund E, Bawingan PA and Wedin M (2007) Testing morphology-based hypotheses of phylogenetic relationships in *Parmeliaceae* (*Ascomycota*) using three ribosomal markers and the nuclear *RPB1* gene. *Molecular Phylogenetics and Evolution* **44**, 812–824.
- Crespo A, Kauff F, Divakar PK, del Prado R, Pérez-Ortega S, Amo de Paz G, Ferencova Z, Blanco O, Roca-Valiente B, Núñez-Zapata J, et al. (2010) Phylogenetic generic classification of parmelioid lichens (*Parmeliaceae*, *Ascomycota*) based on molecular, morphological and chemical evidence. *Taxon* **59**, 1735–1753.
- Crespo A, Divakar PK and Hawksworth DL (2011) Generic concepts in parmelioid lichens, and the phylogenetic value of characters used in their circumscription. *Lichenologist* **43**, 511–535.
- Culberson CF (1972) Improved conditions and new data for identification of lichen products by a standardized thin-layer chromatographic method. *Journal of Chromatography A* **72**, 113–125.
- Culberson WL and Culberson CF (1973) Parallel evolution in the lichen-forming fungi. *Science* **180**, 196–198.
- De La Cruz ARH, De La Cruz JKH, Tolentino DA and Gioda A (2018) Trace element biomonitoring in the Peruvian Andes metropolitan region using *Flavoparmelia caperata* lichen. *Chemosphere* **210**, 849–858.
- Del-Prado R, Cubas P, Lumbsch HT, Divakar PK, Blanco O, Amo de Paz G, Molina MC and Crespo A (2010) Genetic distances within and among species in monophyletic lineages of *Parmeliaceae* (*Ascomycota*) as a tool for taxon delimitation. *Molecular Phylogenetics and Evolution* **56**, 125–33.
- Del-Prado R, Divakar PK, Lumbsch HT and Crespo A (2016) Hidden genetic diversity in an asexually reproducing lichen forming fungal group. *PLoS ONE* **11**, e0161031.
- Del-Prado R, Buaruang K, Lumbsch HT, Crespo A and Divakar PK (2019) DNA sequence-based identification and barcoding of a morphologically highly plastic lichen forming fungal genus (*Parmotrema*, *Parmeliaceae*) from the tropics. *Bryologist* **122**, 281–291.
- Divakar PK and Upreti DK (2005) *Parmelioid Lichens in India (A Revisionary Study)*. Dehradun: Bishen Singh Mahendra Pal Singh.
- Divakar PK, Molina MC, Lumbsch HT and Crespo A (2005a) *Parmelia barreniae*, a new lichen species related to *Parmelia sulcata* (*Parmeliaceae*) based on molecular and morphological data. *Lichenologist* **37**, 37–46.
- Divakar PK, Blanco O, Hawksworth DL and Crespo A (2005b) Molecular phylogenetic studies on the *Parmotrema reticulatum* (syn. *Rimelia reticulata*) complex, including the confirmation of *P. pseudoreticulatum*. *Lichenologist* **37**, 55–65.
- Divakar PK, Figueras G, Hladun N and Crespo A (2010) Molecular phylogenetic studies reveal an undescribed species within the North American concept of *Melanelixia glabra* (*Parmeliaceae*). *Fungal Diversity* **42**, 47–55.
- Divakar PK, Crespo A, Wedin M, Leavitt SD, Hawksworth DL, Myllys L, McCune B, Randlane T, Bjerke JW, Ohmura Y, et al. (2015) Evolution of complex symbiotic relationships in a morphologically derived family of lichen-forming fungi. *New Phytologist* **208**, 1217–1226.
- Elix JA (1993) Progress in the generic delimitation of *Parmelia* sensu lato lichens (*Ascomycotina*: *Parmeliaceae*) and a synoptic key to the *Parmeliaceae*. *Bryologist* **96**, 359–383.
- Elix JA (1994) *Flora of Australia. Volume 55. Lichens: Lecanorales. 2. Parmeliaceae*. Canberra: Australian Biological Resources Study.
- Elix JA and Ernst-Russell KD (1993) *A Catalogue of Standardized Thin Layer Chromatographic Data and Biosynthetic Relationships for Lichen Substances*. 2nd Edn. Canberra: Australian National University.
- Ence DD and Carstens BC (2011) SpedeSTEM: a rapid and accurate method for species delimitation. *Molecular Ecology Resources* **11**, 473–480.
- Gardes M and Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113–118.
- Giordani P and Brunialti G (2015) Sampling and interpreting lichen diversity data for biomonitoring purposes. In Upreti DK, Divakar PK, Shukla V and Bajpai R (eds), *Recent Advances in Lichenology, Vol. 1*. New Delhi: Springer, pp. 19–46.
- Grewe F, Lagostina E, Wu H, Printzen C and Lumbsch HT (2018) Population genomic analyses of RAD sequences resolves the phylogenetic relationship of the lichen-forming fungal species *Usnea antarctica* and *Usnea aurantiacoatra*. *MycKeys* **43**, 91–113.

- Hale ME (1965) A monograph of *Parmelia* subgenus *Amphigymnia*. *Contributions from the United States National Herbarium* **36**, 193–358.
- Hale ME (1974) *Bulbothrix*, *Parmelina*, *Relicina*, and *Xanthoparmelia*, four new genera in the *Parmeliaceae*. *Phytologia* **28**, 479–490.
- Hawksworth DL and Rose L (1970) Qualitative scale of estimating sulfur dioxide air pollution in England and Wales using epiphytic lichens. *Nature* **227**, 145–148.
- Henssen A and Jahns HM (1974) *Lichenes. Eine Einführung in die Flechtenkunde*. Stuttgart: Georg Thieme Verlag.
- Huelsenbeck JP and Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**, 754–755.
- Huneck S (1999) The significance of lichens and their metabolites. *Naturwissenschaften* **86**, 559–570.
- Huneck S and Yoshimura I (1996) *Identification of Lichen Substances*. New York: Springer.
- Jabłońska A, Oset M and Kukwa M (2009) The lichen family *Parmeliaceae* in Poland. I. The genus *Parmotrema*. *Acta Mycologica* **44**, 211–222.
- Kantvilas G (2019) An annotated catalogue of the lichens of Kangaroo Island, South Australia. *Swainsona* **32**, 1–97.
- Katoh K and Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitution through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**, 111–120.
- Knowles LL and Carstens BC (2007) Delimiting species without monophyletic gene trees. *Systematic Biology* **56**, 887–895.
- Kumar K and Upreti DK (2001) *Parmelia* spp. (Lichens) in ancient medicinal plant lore of India. *Economic Botany* **55**, 458–459.
- Kurokawa S (1991) Japanese species and genera of the *Parmeliaceae*. *Journal of Japanese Botany* **66**, 152–159.
- Kurokawa S and Lai MJ (2001) Parmelioid lichen genera and species in Taiwan. *Mycotaxon* **77**, 225–284.
- Leavitt SD, Esslinger TL, Spribille T, Divakar PK and Lumbsch HT (2013) Multilocus phylogeny of the lichen-forming fungal genus *Melanohalea* (*Parmeliaceae*, *Ascomycota*): insights on diversity, distributions, and a comparison of species tree and concatenated topologies. *Molecular Phylogenetics and Evolution* **66**, 138–152.
- Leavitt SD, Moreau CS and Lumbsch HT (2015) The dynamic discipline of species delimitation: progress toward effectively recognizing species boundaries in natural populations. In Upreti DK, Divakar PK, Shukla V and Bajpai R (eds), *Recent Advances in Lichenology*, Vol. 2. New Delhi: Springer, pp. 11–44.
- Leavitt SD, Divakar PK, Crespo A and Lumbsch HT (2016) A matter of time – understanding the limits of the power of molecular data for delimiting species boundaries. *Herzogia* **29**, 479–492.
- Lohtander K, Källersjö M and Tehler A (1998) Dispersal strategies in *Roccellina capensis* (*Arthoniales*). *Lichenologist* **30**, 341–350.
- Lücking R, Rivas Plata E, Chaves JL, Umaña L and Sipman HJM (2009) How many tropical lichens are there... really? *Bibliotheca Lichenologica* **100**, 399–418.
- Lücking R, Hodkinson BP and Leavitt SD (2017) The 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota* – approaching one thousand genera. *Bryologist* **119**, 361–416.
- Lumbsch HT and Leavitt SD (2011) Goodbye morphology? A paradigm shift in the delimitation of species in lichenized fungi. *Fungal Diversity* **50**, 59–72.
- Lumbsch HT and Schmitt I (2001) Molecular data suggest that the lichen genus *Pertusaria* is not monophyletic. *Lichenologist* **33**, 161–170.
- Mark K, Cornejo C, Keller C, Flück D and Scheidegger C (2016) Barcoding lichen-forming fungi using 454 pyrosequencing is challenged by artifactual and biological sequence variation. *Genome* **59**, 685–704.
- Massalongo AB (1860) Esame comparativo di alcuni generi di licheni. *Atti dell' I. R. Istituto Veneto di Scienze, Lettere ed Arti, series 3* **5**, 247–276.
- Miller MA, Pfeiffer W and Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 November 2010, New Orleans, Louisiana, pp. 1–8.
- Molina MC, Crespo A, Blanco O, Lumbsch HT and Hawksworth DL (2004) Phylogenetic relationships and species concepts in *Parmelia* s. str. (*Parmeliaceae*) inferred from nuclear ITS rDNA and β -tubulin sequences. *Lichenologist* **36**, 37–54.
- Molina MC, Divakar PK, Millanes AM, Sanchez E, Del-Prado R, Hawksworth DL and Crespo A (2011a) *Parmelia sulcata* (*Ascomycota: Parmeliaceae*), a sympatric monophyletic species complex. *Lichenologist* **43**, 585–601.
- Molina MC, Del-Prado R, Divakar PK, Sánchez-Mata D and Crespo A (2011b) Another example of cryptic diversity in lichen-forming fungi: the new species *Parmelia mayi* (*Ascomycota: Parmeliaceae*). *Organisms, Diversity and Evolution* **11**, 331–342.
- Molina MC, Divakar PK, Goward T, Millanes AM, Lumbsch HT and Crespo A (2017) Neogene diversification in the temperate lichen-forming fungal genus *Parmelia* (*Parmeliaceae*, *Ascomycota*). *Systematics and Biodiversity* **15**, 166–181.
- Monaghan MT, Wild R, Elliot M, Fujisawa T, Balke M, Inward DJG, Lees DC, Ranaivosolo R, Eggleton P, Barraclough TG, et al. (2009) Accelerated species inventory on Madagascar using coalescent-based models of species delineation. *Systematic Biology* **58**, 298–311.
- Myllys L, Lohtander K and Tehler A (2001) β -tubulin, ITS and group I intron sequences challenge the species pair concept in *Physcia aipolia* and *P. caesia*. *Mycologia* **93**, 335–343.
- Nash TH, III and Elix JA (2002) *Parmotrema*. In Nash TH, III, Ryan BD, Gries C and Bungartz F (eds), *Lichen Flora of the Greater Sonoran Desert Region*, Vol. 1. Tempe, Arizona: Lichens Unlimited, Arizona State University, pp. 318–329.
- Nelsen MP and Gargas A (2008) Dissociation and horizontal transmission of codispersing lichen symbionts in the genus *Lepraria* (*Lecanorales: Stereocaulaceae*). *New Phytologist* **177**, 264–275.
- Nimis PL, Scheidegger C and Wolseley P (2002) *Monitoring with Lichens – Monitoring Lichens*. Dordrecht: Kluwer Academic.
- Normark BB, Judson OP and Moran NA (2003) Genomic signatures of ancient asexual lineages. *Biological Journal of the Linnean Society* **79**, 69–84.
- Orange A, James PW and White FJ (2010) *Microchemical Methods for the Identification of Lichens*. 2nd Edn. London: British Lichen Society.
- Ott S, Brinkmann M, Wirtz N and Lumbsch HT (2004) Mitochondrial and nuclear ribosomal DNA data do not support the separation of the Antarctic lichens *Umbilicaria kappenii* and *Umbilicaria antarctica* as distinct species. *Lichenologist* **36**, 227–234.
- Parnmen S, Rangsiruji A, Mongkolsuk P, Boonpragob K, Nutakki A and Lumbsch HT (2012) Using phylogenetic and coalescent methods to understand the species diversity in the *Cladia aggregata* complex (*Ascomycota, Lecanorales*). *PLoS ONE* **7**, e52245.
- Piercey-Normore MD (2006) The lichen-forming ascomycete *Evernia mesomorpha* associated with multiple genotypes of *Trebouxia jamesii*. *New Phytologist* **169**, 331–344.
- Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD and Vogler AP (2006) Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology* **55**, 595–609.
- Posada D (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**, 1253–1256.
- Puillandre N, Lambert A, Brouillet S and Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology* **21**, 1864–1877.
- Purvis OW, Coppins BJ, Hawksworth DL, James PW and Moore DM (1992) *The Lichen Flora of Great Britain and Ireland*. London: British Lichen Society.
- Rambaut A (2009) *FigTree v. 1.2.3*. [WWW resource] URL <http://tree.bio.ed.ac.uk/software/figtree/> [Accessed 5 November 2010].
- Rambaut A and Drummond J (2007) *Tracer v. 1.5*. [WWW resource] URL <http://beast.bio.ed.ac.uk/Tracer>.
- Ronquist F and Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Sancho LG, Pintado A and Green TGA (2019) Antarctic studies show lichens to be excellent biomonitors of climate change. *Diversity* **11**, 42.
- Schmidt HA, Strimmer K, Vingron M and von Haeseler A (2002) TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics* **18**, 502–504.

- Spielmann AA and Marcelli MP** (2009) *Parmotrema* s.l. (*Parmeliaceae*, lichenized *Ascomycota*) from Serra Geral slopes in central Rio Grande do Sul State, Brazil. *Hoehnea* **36**, 551–595.
- Spielmann AA and Marcelli MP** (2020) Type studies on *Parmotrema* (*Parmeliaceae*, *Ascomycota*) with salazinic acid. *Plant and Fungal Systematics* **65**, 403–508.
- Stamatakis A** (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688–2690.
- Stamatakis A, Hoover P and Rougemont J** (2008) A rapid bootstrap algorithm for the RAxML webservers. *Systematic Biology* **57**, 758–771.
- Sujetovienė G** (2015) Monitoring lichen as indicators of atmospheric quality. In Upreti DK, Divakar PK, Shukla V and Bajpai R (eds), *Recent Advances in Lichenology, Vol. 1*. New Delhi: Springer, pp. 87–118.
- Swinscow TDV and Krog H** (1988) *Macrolichens of East Africa*. London: British Museum of Natural History.
- Thell A, Crespo A, Divakar PK, Kearnefelt I, Leavitt SD, Lumbsch HT and Seaward MRD** (2012) A review of the lichen family *Parmeliaceae* – history, phylogeny and current taxonomy. *Nordic Journal of Botany* **30**, 641–664.
- Vilgalys R and Hester M** (1990) Rapid genetic identification and mapping of enzymatically amplified DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**, 4238–4246.
- Wang-Yang JR and Lai MJ** (1976) Additions and corrections to the lichen flora of Taiwan. *Taiwania* **21**, 226–228.
- Wei JC** (1991) *An Enumeration of Lichens in China*. Beijing: International Academic Publishers.
- Widhelm TJ, Egan RS, Bertoletti FR, Asztalos MJ, Kraichak E, Leavitt SD and Lumbsch HT** (2016) Picking holes in traditional species delimitations: an integrative taxonomic reassessment of the *Parmotrema perforatum* group (*Parmeliaceae*, *Ascomycota*). *Botanical Journal of the Linnean Society* **182**, 868–884.
- Wirtz N, Printzen C and Lumbsch HT** (2008) The delimitation of Antarctic and bipolar species of neuropogonoid *Usnea* (*Ascomycota*, *Lecanorales*): a cohesion approach of species recognition for the *Usnea perpusilla* complex. *Mycological Research* **112**, 472–484.
- Yoshimura I** (1974) *Lichen Flora of Japan in Colour*. Osaka: Hoikusha.
- Zhang J, Kapli P, Pavlidis P and Stamatakis A** (2013) A general species delimitation method with applications to phylogenetic placements. *Bioinformatics* **29**, 2869–2876.