

CONVOLUTION OF RIEMANNIAN MANIFOLDS
AND ITS APPLICATIONS

BANG-YEN CHEN

It is well-known that warped products play some important roles in differential geometry as well as in physics. In this article we extend the notion of warped product to the notion of convolution of Riemannian manifolds. We study the basic properties of convolutions of Riemannian manifolds. We also apply the notion of convolution to establish and characterise the Euclidean version of Segre embedding.

1. CONVOLUTION OF RIEMANNIAN MANIFOLDS

Let (N_1, g_1) and (N_2, g_2) be two Riemannian manifolds and f be a positive differentiable function on N_1 . The well-known notion of *warped product* $N_1 \times_f N_2$ is the product manifold $N_1 \times N_2$ equipped with the warped product metric $g_1 + f^2 g_2$. It is well-known that the notion of warped products plays some important roles in differential geometry as well as in physics (see [7]).

The following notion of convolution of Riemannian manifolds extends the notion of warped products in a natural way. Let (N_1, g_1) and (N_2, g_2) be two Riemannian manifolds and let f and h be two positive differentiable functions on N_1 and N_2 , respectively. Consider the symmetric tensor field ${}_h g_1 *_f g_2$ of type $(0,2)$ on $N_1 \times N_2$ defined by

$$(1.1) \quad {}_h g_1 *_f g_2 = h^2 g_1 + f^2 g_2 + 2fhdf \otimes dh$$

The symmetric tensor field ${}_h g_1 *_f g_2$ is called the *convolution of g_1 and g_2 via h and f* . The product manifold $N_1 \times N_2$ together with ${}_h g_1 *_f g_2$, denoted by ${}_h N_1 \star_f N_2$, is called a *convolution manifold*. When f, h are irrelevant, ${}_h N_1 \star_f N_2$ and ${}_h g_1 *_f g_2$ are simply denoted by $N_1 \star N_2$ and $g_1 *_f g_2$, respectively.

When ${}_h g_1 *_f g_2$ is a positive-definite symmetric tensor, it defines a Riemannian metric on $N_1 \times N_2$. In this case, ${}_h g_1 *_f g_2$ is called a *convolution metric* and the convolution manifold ${}_h N_1 \star_f N_2$ is called a *convolution Riemannian manifold*.

In the first part of this article we show that the notion of convolution of Riemannian manifolds arises naturally. In the second part, we apply the notion of convolution to provide a fundamental study of the differential geometry of the tensor product $C^h \otimes E^p$. In particular, we apply the notion of convolution to establish and characterise the Euclidean version of Segre embedding.

Received 23rd July, 2001

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9727/02 \$A2.00+0.00.

2. PRELIMINARIES

Let N be a Riemannian manifold equipped with a Riemannian metric g . The gradient $\nabla\varphi$ of a function φ on N is defined by $\langle \nabla\varphi, X \rangle = X\varphi$ for vector fields X tangent to N . If N is a submanifold of a Riemannian manifold \widetilde{M} , the formulas of Gauss and Weingarten are given respectively by

$$(2.1) \quad \widetilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y),$$

$$(2.2) \quad \widetilde{\nabla}_X \xi = -A_\xi X + D_X \xi$$

for vector fields X, Y for vector fields X, Y tangent to N and ξ normal to N , where $\widetilde{\nabla}$ denotes the Riemannian connection on \widetilde{M} , σ the second fundamental form, D the normal connection, and A the shape operator of N in \widetilde{M} . The second fundamental form and the shape operator are related by $\langle A_\xi X, Y \rangle = \langle \sigma(X, Y), \xi \rangle$, where $\langle \cdot, \cdot \rangle$ denotes the inner product on M as well as on \widetilde{M} . A submanifold in a Riemannian manifold is called *totally geodesic* if its second fundamental form vanishes identically.

The *equation of Gauss* of N in \widetilde{M} is given by

$$(2.3) \quad \widetilde{R}(X, Y; Z, W) = R(X, Y; Z, W) + \langle \sigma(X, Z), \sigma(Y, W) \rangle - \langle \sigma(X, W), \sigma(Y, Z) \rangle,$$

for X, Y, Z, W tangent to M , where R and \widetilde{R} denote the curvature tensors of N and \widetilde{M} , respectively.

The covariant derivative $\overline{\nabla}\sigma$ of σ with respect to the connection on $TM \oplus T^\perp M$ is defined by

$$(2.4) \quad (\overline{\nabla}_X \sigma)(Y, Z) = D_X(\sigma(Y, Z)) - \sigma(\nabla_X Y, Z) - \sigma(Y, \nabla_X Z).$$

The *equation of Codazzi* is

$$(2.5) \quad (\widetilde{R}(X, Y)Z)^\perp = (\overline{\nabla}_X \sigma)(Y, Z) - (\overline{\nabla}_Y \sigma)(X, Z),$$

where $(\widetilde{R}(X, Y)Z)^\perp$ denotes the normal component of $\widetilde{R}(X, Y)Z$.

Let $\mathbf{C}^m \otimes \mathbf{E}^n$ denote the tensor product of \mathbf{C}^m and \mathbf{E}^n . Then $\mathbf{C}^m \otimes \mathbf{E}^n$ is holomorphically isometric to \mathbf{C}^{mn} . The inner product on $\mathbf{C}^m \otimes \mathbf{E}^n$ is given by

$$(2.6) \quad \langle \alpha \otimes \beta, \gamma \otimes \delta \rangle = \langle \alpha, \gamma \rangle \langle \beta, \delta \rangle,$$

where $\langle \alpha, \gamma \rangle$ is the inner product of $\alpha, \gamma \in \mathbf{C}^m$ and $\langle \beta, \delta \rangle$ the inner product of $\beta, \delta \in \mathbf{E}^n$.

A vector subspace L of complex Euclidean m -space \mathbf{C}^m is called *totally real* if $J(L) \perp L$, where J denotes the complex structure on \mathbf{C}^m . A submanifold M in \mathbf{C}^m is called *totally real* if each tangent space of M is totally real.

A submanifold M in \mathbf{C}^m is called a *CR-submanifold* if there exists on M a differentiable holomorphic distribution \mathcal{H} whose orthogonal complementary distribution \mathcal{H}^\perp is a totally real distribution, that is $J\mathcal{H}^\perp \subset T^\perp M$ (see [1, 2, 3, 4, 5]).

3. SOME NATURAL EXAMPLES OF CONVOLUTION MANIFOLDS

Let C_*^n and E_*^m denote $C^n - \{0\}$ and $E^m - \{0\}$, respectively. Let (z_1, \dots, z_n) denote a complex Euclidean coordinate system of C^n and (x_1, \dots, x_m) a Euclidean coordinate system of E^m .

The following result shows that the notion of convolution of Riemannian manifolds arises naturally.

PROPOSITION 3.1. *For each holomorphic isometric immersion $z : (N_1, g_1) \rightarrow C_*^n$ and each isometric immersion $x : (N_2, g_2) \rightarrow E_*^m$, the map*

$$(3.1) \quad \psi : N_1 \times N_2 \rightarrow C^n \otimes E^m = C^{nm}; (u, v) \mapsto z(u) \otimes x(v), \quad u \in N_1, v \in N_2,$$

gives rise to a convolution manifold $N_1 \star N_2$ equipped with

$$(3.2) \quad {}_\mu g_1 *_\lambda g_2 = \mu^2 g_1 + \lambda^2 g_2 + 2\lambda\mu d\lambda \otimes d\mu,$$

where $\lambda = |z| = \sqrt{\sum_{j=1}^n z_j \bar{z}_j}$ and $\mu = |x| = \sqrt{\sum_{\alpha=1}^m x_\alpha^2}$.

PROOF: For vector fields X, Y tangent to N_1 and Z, W tangent to N_2 , we have

$$(3.3) \quad d\psi(X) = X\psi = X \otimes x, \quad d\psi(Z) = Z\psi = z \otimes Z.$$

Also, it follows from the definition of the gradient of $\mu = |x|$ that

$$(3.4) \quad \mu d\mu(\nabla\mu) = \frac{(\sum_{\alpha=1}^m x_\alpha (e_1 x_\alpha))^2}{|x|} = \langle \nabla\mu, x \rangle,$$

where e_1 is a unit vector parallel to gradient of μ . Similarly, we have

$$(3.5) \quad \lambda d\lambda(\nabla\lambda) = \langle \nabla\lambda, z \rangle.$$

From (2.6), (3.3), (3.4) and (3.5), we obtain Proposition 3.1. □

Proposition 3.1 provides many examples of convolution manifolds.

REMARK 3.1. If $x(N_2)$ is contained in the unit hypersphere of C^m centred at the origin, then the convolution $g_1 * g_2$ on the convolution manifold $N_1 \star N_2$ given in Proposition 3.1 is nothing but the warped product metric $g_1 + |z|^2 g_2$.

4. GEOMETRY OF $C_*^h \otimes E_*^p$

In this section we study the geometry of the tensor product $C^h \otimes E^p$ by applying the notion of convolution.

Assume that $z : C_*^h \rightarrow C^h$ and $x : E_*^p \rightarrow E^p$ are the inclusion maps. Let $\psi_{z,x} = z \otimes x$ be the map from $C_*^h \times E_*^p$ into C^{hp} defined by

$$(4.1) \quad \psi_{z,x} = z \otimes x = (z_1 x_1, \dots, z_1 x_p, \dots, z_h x_1, \dots, z_h x_p)$$

for $z = (z_1, \dots, z_h) \in \mathbb{C}_*^h$ and $x = (x_1, \dots, x_p) \in \mathbb{E}_*^p$.

If we put $z_j = u_j + iv_j, i = \sqrt{-1}$, and $\frac{\partial}{\partial z_j} = (1/2)\left(\frac{\partial}{\partial u_j} - i \frac{\partial}{\partial v_j}\right)$ for $j = 1, \dots, h$, then we obtain from (4.1) that

$$(4.2) \quad d\psi_{z,x} \left(\sum_{j=1}^h z_j \frac{\partial}{\partial z_j} \right) = d\psi_{z,x} \left(\sum_{\alpha=1}^h x_\alpha \frac{\partial}{\partial x_\alpha} \right).$$

Notice that $\sum_{j=1}^h z_j \frac{\partial}{\partial z_j}$ and $\sum_{\alpha=1}^h x_\alpha \frac{\partial}{\partial x_\alpha}$ are nothing but the position vectors of \mathbb{C}_*^h and \mathbb{E}_*^p in \mathbb{C}^h and \mathbb{E}^p , respectively. Equation (4.2) implies that the gradient of $|z| = \sqrt{\sum_{j=1}^h z_j \bar{z}_j}$ and of $|x| = \sqrt{\sum_{\alpha=1}^p x_\alpha^2}$ are mapped to the same vector field under $\psi_{z,x}$.

From (4.1) and (4.2) it follows that $d\psi_{z,x}$ has constant rank $2h + p - 1$. Hence $\psi_{z,x}(\mathbb{C}_*^h \times \mathbb{E}_*^p)$ gives rise to a $(2h + p - 1)$ -manifold, denoted by $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$, which is equipped with a Riemannian metric induced from the canonical metric on $\mathbb{C}^h \otimes \mathbb{E}^p$ via $\psi_{z,x}$. From (4.1) we can verify that $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$ is isometric to the warped product $\mathbb{C}_*^h \times S^{p-1}$ with the warped product metric $g = g_1 + \rho_1 g_0$, where ρ_1 is the length of the position function of \mathbb{C}_*^h and g_0 the metric of the unit hypersphere S^{p-1} .

If we denote the vector field of (4.2) by V , then V is a tangent vector field of $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$ with length $|x||z|$.

Let

$$(4.3) \quad \pi : \mathbb{C}_*^h \times \mathbb{E}_*^p \rightarrow \mathbb{C}_*^h \otimes \mathbb{E}_*^p$$

denote the projection: $\pi(u, v) = \psi_{z,x}(u, v) = \{u\} \otimes \{v\} \in \mathbb{C}_*^h \otimes \mathbb{E}_*^p$. It is easy to see that, for each $u \in \mathbb{C}_*^h$ and $v \in \mathbb{E}_*^p$, $\mathbb{C}_*^h \otimes \{v\} =: \psi_{z,x}(\mathbb{C}_*^h \times \{v\})$ is a complex submanifold of complex dimension h and $\{u\} \otimes \mathbb{E}_*^p =: \psi_{z,x}(\{u\} \times \mathbb{E}_*^p)$ is a totally real submanifold of dimension p in \mathbb{C}^{hp} .

On $\mathbb{C}_*^h \times \mathbb{E}_*^p$, if we put

$$(4.4) \quad \begin{aligned} \mathcal{D} &= T(\mathbb{C}_*^h), \quad \mathcal{D}^\perp = \{Z \in T(\mathbb{E}_*^p) : Z\mu = 0\}, \quad \mu = |x|, \\ \mathcal{F} &= T(\mathbb{E}_*^p), \quad \mathcal{F}^\perp = \{X \in T(\mathbb{C}_*^h) : X\lambda = 0\}, \quad \lambda = |z|, \end{aligned}$$

then $\mathcal{D}, \mathcal{D}^\perp, \mathcal{F}$ and \mathcal{F}^\perp can be regarded as distributions on $\mathbb{C}_*^h \times \mathbb{E}_*^p$ in a natural way. Moreover, if we put

$$(4.5) \quad \widehat{\mathcal{D}} = d\pi(\mathcal{D}), \quad \widehat{\mathcal{D}}^\perp = d\pi(\mathcal{D}^\perp), \quad \widehat{\mathcal{F}} = d\pi(\mathcal{F}), \quad \widehat{\mathcal{F}}^\perp = d\pi(\mathcal{F}^\perp),$$

we have the following orthogonal decompositions of the tangent bundle of $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$:

$$(4.6) \quad T(\mathbb{C}_*^h \otimes \mathbb{E}_*^p) = \widehat{\mathcal{D}} \oplus \widehat{\mathcal{D}}^\perp = \widehat{\mathcal{F}}^\perp \oplus \widehat{\mathcal{F}}.$$

Since the Riemannian metric on $C_*^h \otimes E_*^p$ is induced from the convolution:

$$(4.7) \quad {}_h g_1 * f g_2 = \mu^2 g_1 + \lambda^2 g_2 + 2\lambda\mu d\lambda \otimes d\mu, \quad \lambda = |z|, \quad \mu = |x|,$$

the distributions $\widehat{\mathcal{D}}$ and $\widehat{\mathcal{D}}^\perp$ can be regarded as the tangent and normal bundles of $C_*^h \otimes \{v\}, v \in E_*^p$, in $C_*^h \otimes E_*^p$, respectively. Similarly, the distributions $\widehat{\mathcal{F}}$ and $\widehat{\mathcal{F}}^\perp$ can be regarded as the tangent and normal bundles of $\{u\} \otimes E_*^p, u \in C_*^h$, in $C_*^h \otimes E_*^p$, respectively.

We give the following two lemmas for later use.

LEMMA 4.1. *Let ∇ denote the Riemannian connection of $C_*^h \otimes E_*^p$. Then, for any vector fields X in $\widehat{\mathcal{F}}^\perp$ and Z in $\widehat{\mathcal{D}}^\perp$, we have*

- (a) $\nabla_X Z = \nabla_Z X = 0,$
- (b) $\nabla_V Z = \nabla_Z V = Z,$ and
- (c) $\nabla_V X = \nabla_X V = X,$

where V is the vector field given by (4.2).

PROOF: For each vector field X in $\widehat{\mathcal{F}}^\perp$ and Z in $\widehat{\mathcal{D}}^\perp$, there exist vector fields \check{X} in \mathcal{F}^\perp and \check{Z} in \mathcal{D}^\perp such that $d\pi(\check{X}) = X$ and $d\pi(\check{Z}) = Z$. From (4.1), we have

$$(4.8) \quad \check{X}\check{Z}\psi_{z,x} = \check{X} \otimes \check{Z}.$$

Since $\check{X}\lambda = \check{Z}\mu = 0, \lambda = |z|, \mu = |x|$, the vector field $\check{X} \otimes \check{Z}$ is perpendicular to $d\psi_{z,x}\left(\frac{\partial}{\partial z_1}\right), \dots, d\psi_{z,x}\left(\frac{\partial}{\partial z_h}\right), d\psi_{z,x}\left(\frac{\partial}{\partial x_1}\right), \dots, d\psi_{z,x}\left(\frac{\partial}{\partial x_p}\right)$. Thus, for any vector fields X in $\widehat{\mathcal{F}}^\perp$ and Z in $\widehat{\mathcal{D}}^\perp$, we have (a).

Let $\check{V} = z = \sum_{j=1}^h z_j \partial/\partial z_j \in T(C^h)$ and let \check{Z} be any vector in \mathcal{D}^\perp . Then (4.1) implies

$$(4.9) \quad \check{Z}\check{V}\psi_{z,x} = \check{V} \otimes \check{Z}.$$

Since $z \otimes \check{Z} = \check{V} \otimes \check{Z}$ is tangent to $C_*^h \otimes E_*^p$, (4.9) implies

$$(4.10) \quad \nabla_V Z = \check{V} \otimes \check{Z}.$$

On the other hand, we also have $Z = d\psi_{z,x}(\check{Z}) = \check{Z}\psi_{z,x} = \check{V} \otimes \check{Z}$ from (4.1). By comparing this with (4.10), we obtain (b).

Similarly, let $\tilde{V} = x = \sum_{\alpha=1}^p x_\alpha \partial/\partial x_\alpha$ and let \check{X} be any vector field in \mathcal{F}^\perp , we have

$$(4.11) \quad \check{X}\tilde{V}\psi_{z,x} = \check{X} \otimes x.$$

On the other hand, we also have $X = d\psi_{z,x}(\check{X}) = \check{X}\psi_{z,x} = \check{X} \otimes x$. Comparing this with (4.11) gives (c). □

LEMMA 4.2. For each $u \in C_*^h$ and each $v \in E_*^p$, $\{u\} \otimes E_*^p$ and $C_*^h \otimes \{v\}$ are totally geodesic submanifolds of $C_*^h \otimes E_*^p$.

PROOF: First, we recall that the tangent and normal bundles of $C_*^h \otimes \{v\}$ in $C_*^h \otimes E_*^p$ are given respectively by \widehat{D} and \widehat{D}^\perp . Since the distribution \widehat{D} is spanned by $\widehat{\mathcal{F}}^\perp$ and V , statements (a) and (b) of Lemma 4.1 and the formula of Weingarten imply that $C_*^h \otimes \{v\}$ is totally geodesic in $C_*^h \otimes E_*^p$.

Similarly, from statements (b) and (c) of Lemma 4.1 and the formula of Weingarten, we conclude that each $\{u\} \otimes E_*^p$ is a totally geodesic submanifold of $C_*^h \otimes E_*^p$. □

5. CLASSIFICATION OF NATURAL CR-IMMERSIONS OF $C_*^h \otimes E_*^p$

Since there is a canonical holomorphic distribution \widehat{D} on $C_*^h \otimes E_*^p$, we call an isometric immersion $\psi : U \rightarrow C^m$ of an open portion U of $C_*^h \otimes E_*^p$ into C^m a *natural CR-immersion* if ψ carries \widehat{D} into a holomorphic distribution in C^m and carries the orthogonal complementary distribution \widehat{D}^\perp of \widehat{D} into a totally real distribution. Clearly, (4.1) defines a natural CR-immersion of $C_*^h \otimes E_*^p$ into C^{hp} .

THEOREM 5.1. Let $\phi : U \rightarrow C^m$ be a natural CR-immersion from an open portion U of $C_*^h \otimes E_*^p$ into C^m . Then, up to rigid motions of C^m , $\tilde{\phi} = \phi \circ \pi$ is given by

$$(5.1) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p) z_j,$$

where A^1, \dots, A^h are mutually orthogonal vector functions of length $|x|$ which span a totally real subspace of C^m at each point $x = (x_1, \dots, x_p)$ with $\pi(z, x) \in U$. Moreover, A^1, \dots, A^h satisfy

$$(5.2) \quad \begin{aligned} \langle A^j, A_{x_\alpha}^k \rangle &= x_\alpha \delta_{jk}, \quad \langle A_{x_\alpha}^j, A_{x_\beta}^k \rangle = \delta_{jk} \delta_{\alpha\beta}, \\ \langle A^j, iA_{x_\alpha}^k \rangle &= \langle A_{x_\alpha}^j, iA_{x_\beta}^k \rangle = 0 \end{aligned}$$

for $j, k = 1 \dots, h$; $\alpha, \beta = 1, \dots, p$, where $A_{x_\alpha}^j = \partial A^j / \partial x_\alpha$.

Conversely, if A^1, \dots, A^h are h mutually orthogonal E^m -valued functions of length $|x|$ satisfying (5.2), then (5.1) defines a natural CR-immersion of $C_*^h \otimes E_*^p$ into C^m .

PROOF: Suppose that $\phi : U \rightarrow C^m$ is a natural CR-immersion from an open portion of $C_*^h \otimes E_*^p$ into C^m . For each $v \in E_*^p$, Lemma 4.2 implies that $C_*^h \otimes \{v\}$ is a totally geodesic submanifold of $C_*^h \otimes E_*^p$.

Because the restriction of ϕ to $U \cap (C_*^h \otimes \{v\})$ is a holomorphically isometric immersion of $U \cap (C_*^h \otimes \{v\})$ into C^m and $C_*^h \otimes \{v\}$ is a flat Kaehler manifold, the equation of Gauss implies that the restriction of ϕ to $U \cap (C_*^h \otimes \{v\})$ is a totally geodesic holomorphic immersion. Thus, ϕ immerses $U \cap (C_*^h \otimes \{v\})$ into a complex h -plane in

\mathbb{C}^m . Consequently, $\tilde{\phi}$ immerses $\pi^{-1}(U) \cap (\mathbb{C}_*^h \times \{v\})$ into a complex h -plane in \mathbb{C}^m . Hence, we have

$$(5.3) \quad \tilde{\phi}_{z_j z_k} = \tilde{\phi}_{z_j \bar{z}_k} = \tilde{\phi}_{\bar{z}_j \bar{z}_k} = 0, \quad 1 \leq j, k \leq h.$$

Solving (5.3) yields

$$(5.4) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p) z_j + B(x_1, \dots, x_p)$$

for some \mathbb{C}^m -valued functions A^1, \dots, A^h and B . From (5.4) we find

$$(5.5) \quad \tilde{\phi}_{x_\alpha} = \sum_{j=1}^h A_{x_\alpha}^j z_j + B_{x_\alpha}, \quad \alpha = 1, \dots, p.$$

Thus, by (3.15) and (5.5), we obtain

$$(5.6) \quad \sum_{j=1}^h z_j \bar{z}_j = \left\langle \sum_{j=1}^h A_{x_\alpha}^j z_j, \sum_{k=1}^h A_{x_\alpha}^k z_k \right\rangle + 2 \left\langle \sum_{j=1}^h A_{x_\alpha}^j z_j, B_{x_\alpha} \right\rangle + \langle B_{x_\alpha}, B_{x_\alpha} \rangle.$$

Condition (5.6) implies $B_{x_1} = \dots = B_{x_p} = 0$. Hence, B is a constant vector in \mathbb{C}^m . Without loss of generality, we may choose $B = 0$ by applying a suitable translation on \mathbb{C}^m if necessary. Hence, (5.4) reduces to

$$(5.7) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p) z_j.$$

From (5.7), we obtain

$$(5.8) \quad \tilde{\phi}_{u_j} = A^j, \quad \tilde{\phi}_{v_j} = iA^j, \quad \tilde{\phi}_{x_\alpha} = \sum_{j=1}^h A_{x_\alpha}^j z_j$$

for $j = 1, \dots, h$; $\alpha = 1, \dots, p$, where $z_j = u_j + iv_j$. From (3.15) and (5.8), we find

$$(5.9) \quad |x|^2 \delta_{jk} = \langle \tilde{\phi}_{u_j}, \tilde{\phi}_{u_k} \rangle = \langle A^j, A^k \rangle, \quad \langle A^j, iA^k \rangle = \langle \tilde{\phi}_{u_j}, \tilde{\phi}_{v_k} \rangle = 0,$$

$$(5.10) \quad u_j x_\alpha = \langle \tilde{\phi}_{u_j}, \tilde{\phi}_{x_\alpha} \rangle = \left\langle A^j, \sum_{k=1}^h A_{x_\alpha}^k z_k \right\rangle,$$

$$(5.11) \quad v_j x_\alpha = \langle \tilde{\phi}_{v_j}, \tilde{\phi}_{x_\alpha} \rangle = \left\langle iA^j, \sum_{k=1}^h A_{x_\alpha}^k z_k \right\rangle,$$

$$(5.12) \quad |z|^2 \delta_{\alpha\beta} = \langle \tilde{\phi}_{x_\alpha}, \tilde{\phi}_{x_\beta} \rangle = \left\langle \sum_{j=1}^h A_{x_\alpha}^j z_j, \sum_{k=1}^h A_{x_\beta}^k z_k \right\rangle$$

for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$.

The first equation of (5.9) implies that A^1, \dots, A^h are h mutually orthogonal vector functions of length $|x|$. From equations (5.9), (5.10) and (5.11) we have

$$(5.13) \quad \langle A^j, iA^k \rangle = 0, \quad \langle A^j, A^k_{x_\alpha} \rangle = x_\alpha \delta_{jk}, \quad \langle A^j, iA^k_{x_\alpha} \rangle = 0$$

for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$.

By comparing the coefficients of $u_j u_k$ from (5.12), we find

$$(5.14) \quad \langle A^j_{x_\alpha}, A^k_{x_\beta} \rangle + \langle A^j_{x_\beta}, A^k_{x_\alpha} \rangle = 2\delta_{jk} \delta_{\alpha\beta}$$

for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$. If $j = k$, (5.14) implies

$$(5.15) \quad \langle A^j_{x_\alpha}, A^j_{x_\beta} \rangle = \delta_{\alpha\beta}.$$

If $j \neq k$, then, by taking partial derivative of $\langle A^j, A^k_{x_\alpha} \rangle = 0$ with respect to x_β , we obtain

$$(5.16) \quad \langle A^j_{x_\alpha}, A^k_{x_\beta} \rangle = -\langle A^j, A^k_{x_\beta x_\alpha} \rangle = \langle A^j_{x_\beta}, A^k_{x_\alpha} \rangle, \quad j \neq k.$$

Combining this with (5.14) yields $\langle A^j_{x_\alpha}, A^k_{x_\beta} \rangle = 0$ for $j \neq k$. Hence, we have the second equation of (5.2). By comparing the coefficients of $u_j v_k$ from (5.12), we also find

$$(5.17) \quad \langle A^j_{x_\alpha}, iA^k_{x_\beta} \rangle + \langle iA^k_{x_\alpha}, A^j_{x_\beta} \rangle = 0$$

for $1 \leq j \neq k \leq h$ and $1 \leq \alpha, \beta \leq p$.

On the other hand, by applying $\langle \tilde{\phi}_{x_\alpha}, J\tilde{\phi}_{x_\beta} \rangle = 0$ and (5.8), we find

$$(5.18) \quad \left\langle \sum_{j=1}^h A^j_{x_\alpha} z_j, \sum_{k=1}^h iA^k_{x_\beta} z_k \right\rangle = 0.$$

Comparing the coefficients of $u_j u_k$ from (5.18) yields

$$(5.19) \quad \langle A^j_{x_\alpha}, iA^k_{x_\beta} \rangle + \langle A^k_{x_\alpha}, iA^j_{x_\beta} \rangle = 0$$

for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$.

If $j = k$, (5.19) implies $\langle A^j_{x_\alpha}, iA^j_{x_\beta} \rangle = 0$. If $j \neq k$, then, by combining (5.17) and (5.19), we obtain $\langle A^j_{x_\alpha}, iA^k_{x_\beta} \rangle = 0$. Therefore, we have $\langle A^j_{x_\alpha}, iA^k_{x_\beta} \rangle = 0$ for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$. Consequently, we have (5.2). From (5.2) we know that A^1, \dots, A^h span a totally real subspace of \mathbf{C}^m at each point $x = (x_1, \dots, x_p)$ with $\pi(z, x) \in U$.

Statement (b) can be proved by straightforward computation. □

EXAMPLE 5.1. *Let D be an open portion of Euclidean p -space \mathbf{E}^p which does not contain the origin of \mathbf{E}^p . If $A : D \rightarrow \mathbf{E}^m$ is an isometric immersion of the flat space*

D into \mathbf{E}^m satisfying $|A| = |x|$ at $x = (x_1, \dots, x_p) \in D$, then (5.2) holds automatically. Hence, by Theorem 5.1, we know that

$$(5.20) \quad \tilde{\phi}(z, x) = A(x_1, \dots, x_p)z$$

defines a natural CR-immersion from $\mathbf{C}_*^1 \otimes D$ into \mathbf{C}^m . In particular, if $\gamma(s) = (\gamma_1(s), \dots, \gamma_{m-p+1}(s))$ is a unit speed curve satisfying

$$(5.21) \quad |\gamma(s)|^2 = \sum_{j=1}^{m+p-1} \gamma_j^2(s) = s^2,$$

then $A = (\gamma_1(x_1), \dots, \gamma_{m-p+1}(x_1), x_2, \dots, x_p)$ defines an isometric immersion of an open portion D of \mathbf{E}_*^p into \mathbf{E}^m satisfying $|A| = |x|$. Thus,

$$(5.22) \quad \tilde{\phi}(z, x) = A(x_1, \dots, x_p)z, \quad A = (\gamma_1(x_1), \dots, \gamma_{m-p+1}(x_1), x_2, \dots, x_p)$$

defines a natural CR-immersion from $\mathbf{C}_*^1 \otimes D$ into \mathbf{C}^m .

REMARK 5.1. When $m + p - 1 = 2$, then $\gamma_1 = as, \gamma_2 = bs, a^2 + b^2 = 1$, are the only functions satisfying (5.21). However, if $m + p - 1 \geq 3$, then there are many unit speed curves γ , other than lines, which satisfy (5.21).

EXAMPLE 5.2. Suppose that D is an open portion of Euclidean p -space which does not contain the origin. Let $A^j : D \rightarrow \mathbf{E}^{m_j}, j = 1, \dots, h$, be isometric immersions of the flat space D into \mathbf{E}^{m_j} satisfying $|A^1| = \dots = |A^h| = |x|$ on D , then

$$(5.23) \quad \psi(z, x) = (A^1(x_1, \dots, x_p)z_1, \dots, A^h(x_1, \dots, x_p)z_h)$$

defines a natural CR-immersion from $\mathbf{C}_*^h \otimes D$ into $\mathbf{C}^m, m = m_1 + \dots + m_h$.

6. TWO GEOMETRIC CHARACTERISATIONS OF $\psi_{z,x}$

The following result provides a simple geometric characterisation of $\psi_{z,x} = z \otimes x$.

THEOREM 6.1. If $\phi : U \rightarrow \mathbf{C}^m$ is a natural CR-immersion of an open portion U of $\mathbf{C}_*^h \otimes \mathbf{E}_*^p$ into \mathbf{C}^m , then we have:

- (1) The squared norm of the second fundamental form σ of ϕ satisfies

$$(6.1) \quad \|\sigma\|^2 \geq \frac{(2h-1)(p-1)}{|x|^2|z|^2}.$$

- (2) The equality sign of (6.1) holds identically if and only if, up to rigid motions of \mathbf{C}^m , the composition $\tilde{\phi} = \phi \circ \pi$ is given by

$$(6.2) \quad \tilde{\phi}(z, x) = (z_1x_1, \dots, z_1x_p, \dots, z_hx_1, \dots, z_hx_p, 0, \dots, 0),$$

where $\pi : C_*^h \times E_*^p \rightarrow C_*^h \otimes E_*^p$ is the projection and $z = (z_1, \dots, z_h)$ and $x = (x_1, \dots, x_p)$ are natural coordinate systems of C_*^h and E_*^p , respectively.

We need some lemmas.

LEMMA 6.1. *If $\phi : U \rightarrow C^m$ is a natural CR-immersion from an open portion of $C_*^h \otimes E_*^p$ into C^m , then, for any vector fields X in $\widehat{\mathcal{D}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$ and for Z, W in $\widehat{\mathcal{D}}^\perp$, we have*

$$(6.3) \quad \langle \sigma(X, Z), JW \rangle = 0,$$

$$(6.4) \quad \langle \sigma(JV, Z), JW \rangle = \langle Z, W \rangle,$$

$$(6.5) \quad \langle \sigma(V, Z), JW \rangle = 0.$$

PROOF: Suppose Z, W are vector fields in $\widehat{\mathcal{D}}^\perp$. Then we have

$$(6.6) \quad J\nabla_Z W + J\sigma(Z, W) = -A_{JW}Z + D_Z JW.$$

If X is a vector field in $\widehat{\mathcal{D}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$, then both X and JX belong to $\widehat{\mathcal{F}}^\perp$. Thus, by (6.6), we have

$$(6.7) \quad \langle \sigma(JX, Z), JW \rangle = \langle A_{JW}Z, JX \rangle = -\langle \nabla_Z W, X \rangle = \langle \nabla_Z X, W \rangle = 0$$

for X in \mathcal{D} and Z, W on \mathcal{D}^\perp . This proves (6.3).

From (6.6) and statement (b) of Lemma 4.1, we find, for Z, W in $\widehat{\mathcal{D}}^\perp$, that

$$(6.8) \quad \langle \sigma(JV, Z), JW \rangle = \langle A_{JW}Z, JV \rangle = -\langle \nabla_Z W, V \rangle = \langle W, \nabla_Z V \rangle = \langle W, Z \rangle,$$

which proves (6.4).

For Z, W in $\widehat{\mathcal{D}}^\perp$, we also find from (6.6) and Lemma 4.1 that

$$(6.9) \quad \langle \sigma(V, Z), JW \rangle = \langle A_{JW}Z, V \rangle = \langle \nabla_Z W, JV \rangle = \langle W, \nabla_Z JV \rangle = 0,$$

since $JV \in \widehat{\mathcal{F}}^\perp$. This proves (6.5). □

LEMMA 6.2. *If $\phi : U \rightarrow C^m$ is a natural CR-immersion of an open portion of $C_*^h \otimes E_*^p$ into C^m , then, for $X \in \widehat{\mathcal{F}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$ and $Z \in \widehat{\mathcal{D}}^\perp$, the second fundamental form σ of ϕ satisfies*

$$(6.10) \quad |\sigma(X, Z)| = \frac{|X||Z|}{|x||z|}.$$

PROOF: For $X \in \widehat{\mathcal{D}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$ and $Z \in \mathcal{D}^\perp$, the equation of Codazzi implies

$$(6.11) \quad \langle D_{JX}\sigma(X, Z) - \sigma(\nabla_{JX}X, Z) - \sigma(X, \nabla_{JX}Z), JZ \rangle \\ = \langle D_X\sigma(JX, Z) - \sigma(\nabla_X JX, Z) - \sigma(JX, \nabla_X Z), JZ \rangle.$$

Since $C_*^h \otimes \{v\}$ is a totally geodesic submanifold of $C_*^h \otimes E_p^p$ according to Lemma 4.2, $\nabla_X JX$ and $\nabla_{JX} X$ belong to \widehat{D} . From Lemma 4.1, we also know that $\nabla_X Z$ and $\nabla_{JX} Z$ belong to \widehat{D}^\perp . Hence, by (6.11), Lemma 4.1, and Lemma 6.1, we get

$$(6.12) \quad \langle \sigma(X, Z), D_{JX} JZ \rangle + \langle \sigma(\nabla_{JX} X, Z), JZ \rangle = \langle \sigma(JX, Z), D_X JZ \rangle + \langle \sigma(\nabla_X JX, Z), JZ \rangle.$$

From formulas of Gauss and Weingarten, we have

$$(6.13) \quad J\nabla_X Z + J\sigma(X, Z) = -A_{JZ} X + D_X JZ.$$

Since $\nabla_X Z$ lies in \widehat{D}^\perp , (6.13), Lemma 4.1, and Lemma 6.1 imply

$$(6.14) \quad \langle \sigma(JX, Z), D_X JZ \rangle = \langle \sigma(JX, Z), J\nabla_X Z \rangle + \langle \sigma(JX, Z), J\sigma(X, Z) \rangle = \langle \sigma(JX, Z), J\sigma(X, Z) \rangle.$$

Let ν denote the orthogonal complement of $J\widehat{D}^\perp$ in the normal bundle of $C_*^h \otimes E_p^p$ in C^m . Then ν is invariant under the action of the complex structure J of C^m . We denote by σ_ν the ν -component of the second fundamental form σ .

Let $\widetilde{\nabla}$ be the Riemannian connection of C^m . For each $\xi \in \nu$, $Y \in \widehat{D}$, and each tangent vector U of $C_*^h \otimes E_p^p$, we have

$$\langle A_\xi(JY), U \rangle = \langle \sigma(JY, U), \xi \rangle = \langle J\widetilde{\nabla}_U Y, \xi \rangle = -\langle \sigma(Y, U), J\xi \rangle = -\langle A_{J\xi} Y, U \rangle.$$

Hence, we get

$$(6.15) \quad A_{J\xi} Y = -A_\xi(JY), \quad Y \in \widehat{D}, \quad \xi \in \nu.$$

By applying (6.15), we find

$$(6.16) \quad \langle \sigma(JX, Z), J\sigma(X, Z) \rangle = \langle \sigma(JX, Z), J\sigma_\nu(X, Z) \rangle = \langle A_{J\sigma_\nu(X, Z)} JX, Z \rangle = \langle A_{\sigma_\nu(X, Z)} X, Z \rangle = \langle \sigma(X, Z), \sigma_\nu(X, Z) \rangle = |\sigma_\nu(X, Z)|^2.$$

Combining (6.14) and (6.16), we obtain

$$(6.17) \quad \langle \sigma(JX, Z), D_X JZ \rangle = |\sigma_\nu(X, Z)|^2.$$

Replacing X in (6.17) by JX and applying (6.15) yield

$$(6.18) \quad \langle \sigma(X, Z), D_{JX} JZ \rangle = -|\sigma_\nu(X, Z)|^2.$$

On the other hand, by Lemma 4.1 and Lemma 6.1, we have

$$(6.19) \quad \langle \sigma(\nabla_{JX} X, Z), JZ \rangle = \frac{\langle \nabla_{JX} X, JV \rangle}{|x|^2|z|^2} \langle Z, Z \rangle = \frac{\langle JX, \nabla_{JX} V \rangle}{|x|^2|z|^2} \langle Z, Z \rangle = \frac{|X|^2|Z|^2}{|x|^2|z|^2}.$$

Replacing X in (6.19) by JX yields

$$(6.20) \quad -\langle \sigma(\nabla_X JX, Z), JZ \rangle = \frac{|X|^2|Z|^2}{|x|^2|z|^2}.$$

Combining (6.12) and (6.17)-(6.20) gives

$$(6.21) \quad |\sigma_\nu(X, Z)| = \frac{|X||Z|}{|x||z|}$$

for $Z \in \widehat{\mathcal{D}}^\perp$ and $X \in \widehat{\mathcal{D}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$.

On the other hand, from (6.3) of Lemma 6.1, we have $\sigma(X, Z) = \sigma_\nu(X, Z)$. Hence, we obtain (6.10) from (6.21). This proves Lemma 6.3.

Now, we return to the proof of Theorem 6.1. First, by applying (6.4) of Lemma 6.1, we have

$$(6.22) \quad |\sigma(JV, Z)| \geq \langle Z, Z \rangle,$$

with equality holding if and only if $\sigma(JV, Z) = JZ$.

Since $|JV| = |x||z|$, we obtain inequality (6.1) from (6.21) and (6.22).

Suppose that the equality sign of (6.1) holds. Then, by Lemma 6.2, we have

$$(6.23) \quad \sigma(\widehat{\mathcal{D}}, \widehat{\mathcal{D}}) = 0, \quad \sigma(\widehat{\mathcal{F}}, \widehat{\mathcal{F}}) = 0$$

for $Z \in \widehat{\mathcal{D}}^\perp$ and $X \in \widehat{\mathcal{D}}$ with $\langle X, V \rangle = \langle X, JV \rangle = 0$.

Since $\phi : U \rightarrow \mathbb{C}^m$ is a natural CR -immersion from an open portion of $\mathbb{C}_*^h \oplus \mathbb{E}_*^p$ into \mathbb{C}^m , Theorem 5.1 implies that, up to rigid motions of \mathbb{C}^m , the composition $\tilde{\phi} = \phi \circ \pi$ is given by

$$(6.24) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p) z_j,$$

where $A^1(x_1, \dots, x_p), \dots, A^h(x_1, \dots, x_p)$ are orthogonal vector functions of length $|x|$. On the other hand, from Lemma 4.2 and the second equation of (6.23), we know that, for each $u \in \mathbb{C}_*^h$, ϕ immerses $U \cap (\{u\} \oplus \mathbb{E}_*^p)$ into a totally real p -plane in \mathbb{C}^m . Hence, $\tilde{\phi}$ carries $\pi^{-1}(U) \cap (\{u\} \times \mathbb{E}_*^p)$ into a totally real p -plane. Therefore, by applying formula of Gauss, we obtain

$$(6.25) \quad \tilde{\phi}_{x_\alpha x_\beta} = 0, \quad \alpha, \beta = 1, \dots, p.$$

Hence, after solving (6.25), we obtain from (6.24) that

$$(6.26) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h \sum_{\alpha=1}^p c_\alpha^j x_\alpha z_j + \sum_{j=1}^h b^j z_j,$$

for some constant vectors c_α^j and b^j , $\alpha = 1, \dots, p$; $j = 1, \dots, h$.

Equation (6.26) yields

$$(6.27) \quad \tilde{\phi}_{z_j} = \sum_{\alpha=1}^p c_\alpha^j x_\alpha + b^j, \quad \tilde{\phi}_{x_\alpha} = \sum_{j=1}^h c_\alpha^j z_j$$

for $j, k = 1, \dots, h$; $\alpha, \beta = 1, \dots, p$. Hence, by applying (3.15) and (6.27), we find

$$(6.28) \quad |x|^2 = \langle \tilde{\phi}_{z_j}, \tilde{\phi}_{z_k} \rangle = \left\langle \sum_{\alpha=1}^p c_\alpha^j x_\alpha + b^j, \sum_{\alpha=1}^p c_\alpha^k x_\alpha + b^k \right\rangle$$

which implies $b^1 = \dots = b^h = 0$. Thus, (6.26) becomes

$$(6.29) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p) z_j, \quad A^j = \sum_{\alpha=1}^p c_\alpha^j x_\alpha,$$

which gives

$$(6.30) \quad A_{x_\alpha}^j = c_\alpha^j.$$

On the other hand, Theorem 5.1 implies that A^1, \dots, A^h satisfy

$$(6.31) \quad \langle A_{x_\alpha}^j, A_{x_\beta}^k \rangle = \delta_{jk} \delta_{\alpha\beta}, \quad \langle A_{x_\alpha}^j, iA_{x_\beta}^k \rangle = 0$$

for $j, k = 1, \dots, h$; $\alpha, \beta = 1, \dots, p$. Combining (6.30) and (6.31) give

$$(6.32) \quad \langle c_\beta^j, c_\alpha^k \rangle = \delta_{jk} \delta_{\alpha\beta}, \quad \langle c_\beta^j, ic_\alpha^k \rangle = 0$$

for $j, k = 1, \dots, h$; $\alpha, \beta = 1, \dots, p$. Hence $\{c_\alpha^j, j = 1, \dots, h; \alpha = 1, \dots, p\}$ is an orthonormal set which spans a totally real hp -subspace of \mathbf{C}^m . Without loss of generality, we may choose the complex coordinates z_1, \dots, z_m on \mathbf{C}^m such that

$$(6.33) \quad \begin{aligned} c_1^1 &= (1, 0, \dots, 0), \dots, \\ c_p^1 &= (0, \dots, 0, 1, 0, \dots, 0), \quad (1 \text{ appears } p\text{-th place}), \dots, \\ c_1^h &= (0, \dots, 0, 1, 0, \dots, 0), \quad (1 \text{ appears } (h-1)p+1\text{-th place}), \dots, \\ c_p^h &= (0, \dots, 0, 1, 0, \dots, 0) \quad (1 \text{ appears } hp\text{-th place}). \end{aligned}$$

Combining (6.29) and (6.33) gives (6.2).

Conversely, it is straightforward to verify that (6.2) defines a natural CR -immersion of $\mathbf{C}_*^h \otimes \mathbf{E}_*^p$ into \mathbf{C}^m whose second fundamental form satisfies the equality case of (6.1). \square

The following theorem provides another simple geometric characterisation of $\psi_{z,x} = z \otimes x$.

THEOREM 6.2. *Let $\phi : U \rightarrow \mathbb{C}^m$ be a natural CR-immersion from an open portion U of $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$ into \mathbb{C}^m . We have*

- (1) $m \geq hp$.
- (2) *If $m = hp$, then, up to rigid motions of \mathbb{C}^{hp} , $\tilde{\phi} = \phi \circ \pi$ is given by*

$$(6.34) \quad \tilde{\phi}(z, x) = \psi_{z,x}(z, x) = (z_1x_1, \dots, z_1x_p, \dots, z_hx_1, \dots, z_hx_p).$$

PROOF: Let $\phi : U \rightarrow \mathbb{C}^m$ be a natural CR-immersion from an open portion U of $\mathbb{C}_*^h \otimes \mathbb{E}_*^p$ into \mathbb{C}^m . Then Theorem 5.1 implies that, up to rigid motions of \mathbb{C}^m , the composition $\tilde{\phi} = \phi \circ \pi$ is given by

$$(6.35) \quad \tilde{\phi}(z, x) = \sum_{j=1}^h A^j(x_1, \dots, x_p)z_j,$$

where $A^1(x_1, \dots, x_p), \dots, A^h(x_1, \dots, x_p)$ are mutually orthogonal vector functions of length $|x|$. Moreover, A^1, \dots, A^h satisfy

$$(6.36) \quad \langle A^j, A^k \rangle = |x|^2 \delta_{jk}, \quad \langle A^j, iA^k \rangle = 0,$$

$$(6.37) \quad \langle A^j, A_{x_\alpha}^k \rangle = x_\alpha \delta_{jk}, \quad \langle A_{x_\alpha}^j, A_{x_\beta}^k \rangle = \delta_{jk} \delta_{\alpha\beta},$$

$$(6.38) \quad \langle A^j, iA_{x_\alpha}^k \rangle = 0, \quad \langle A_{x_\alpha}^j, iA_{x_\beta}^k \rangle = 0$$

for $j, k = 1, \dots, h; \alpha, \beta = 1, \dots, p$.

From (6.36), (6.37) and (6.38) we know that $\{A_{x_\alpha}^j, j = 1, \dots, h; \alpha = 1, \dots, p\}$ is an orthonormal set which spans a totally real hp -plane in \mathbb{C}^m . Therefore, $m \geq hp$. This proves statement (1).

If $m = hp$, then (6.37)–(6.38) implies that, for each $j \in \{1, \dots, h\}$, A^j defines an isometric immersion from an open domain, say D , of the Euclidean p -space \mathbb{E}^p into a totally real p -subspace of \mathbb{C}^h . Moreover, each A^j satisfies $|A^j|^2 = \sum_{\alpha=1}^p x_\alpha^2$. Since, up to rigid motions of \mathbb{E}^p , the only isometric immersion from the flat p -space $D \subset \mathbb{E}^p$ into \mathbb{E}^p is the inclusion map, each A^j must be an inclusion map of D into \mathbb{E}^p . Furthermore, because A^1, \dots, A^h are orthogonal vector functions in \mathbb{C}^{hp} which satisfy (6.36)–(6.38), we may choose the complex Euclidean coordinate system $\{z_1, \dots, z_h\}$ on \mathbb{C}^{hp} so that (6.35) takes the following special form of (6.34). This proves statement (2). □

REMARK 6.1. Theorem 6.2 can be regarded as the Euclidean version of the natural characterisation of Segre imbeddings obtained in [5].

REMARK 6.2. Further results on convolutions have been obtained in [6].

REFERENCES

- [1] A. Bejancu, *Geometry of CR-Submanifolds*, Mathematics and its Applications **23** (D. Reidel Publishing Co., Dordrecht, 1986).
- [2] B.Y. Chen, 'Some *CR*-submanifolds of a Kaehler manifold. I', *J. Differential Geom.* **16** (1981), 305-322.
- [3] B.Y. Chen, 'Some *CR*-submanifolds of a Kaehler manifold. II', *J. Differential Geom.* **16** (1981), 493-509.
- [4] B.Y. Chen, 'Geometry of warped product *CR*-submanifolds in Kaehler manifolds', *Monatsh. Math.* **133** (2001), 177-195.
- [5] B.Y. Chen, 'Geometry of warped products *CR*-submanifolds in Kaehler manifolds. II', *Monatsh. Math.* **134** (2001), 103-119.
- [6] B.Y. Chen, 'More on convolution of Riemannian manifolds', *Beiträge Algebra Geom.* (to appear).
- [7] B. O'Neill, *Semi-Riemannian geometry with applications to relativity*, Pure and Applied Mathematics **103** (Academic Press, New York, 1983).

Department of Mathematics
Michigan State University
East Lansing, MI 48824-1027
United States of America
e-mail: bychen@math.msu.edu