



was attributable to the joint effects of omitting prophylaxis and conversion to an open procedure (90% CI 14, 56), compared to a 5 pp increase attributable to omission of prophylaxis (90% CI 3, 7) or a 6 pp increase attributable to conversion to an open procedure (90% CI -5, 18) by themselves. The interaction contrast captured this super-additive 24 pp increase in risk (90% CI -1, 47). The relative excess risk due to interaction was 23.81 (90% CI -5.56, 53.19), suggesting a departure from additivity as well. **Conclusions:** Patients undergoing open cholecystectomies stand to benefit the most from antibacterial prophylaxis compared to patients who have laparoscopic cholecystectomies. ASPs could consider reducing or eliminating surgical prophylaxis in low-risk procedures such as laparoscopic cholecystectomy to alleviate selection pressure for antibacterial resistant organisms and preserve its effectiveness for people undergoing higher risk procedures.

Antimicrobial Stewardship & Healthcare Epidemiology 2024;4(Suppl. S1):s141-s142
doi:10.1017/ash.2024.314

Presentation Type:
Poster Presentation - Poster Presentation
Subject Category: SSI
A Deeper Look at Proposed Surveillance of Superficial Incision Surgical Site Infections (SSISIs)

Jessica Seidelman, Duke University; Brittain Wood, Duke University; Polly Padgett, Duke; Valerie Payne, Duke Infection Control Outreach Network; Linda Crane, Duke University Medical Center; Linda Roach, Duke Ic Outreach Network and Deverick Anderson, Duke Center for Antimicrobial Stewardship and Infection Prevention

Although the National Healthcare Safety Network (NHSN) recommends the reporting of superficial incisional SSIs, the standardized infection ratio (SIR) models used by the NHSN exclude superficial incisional SSIs cases. Yet, some superficial incisional SSIs may lead to serious adverse patient outcomes. We previously proposed a new category of such infections: serious superficial incisional surgical site infections (SSISIs), defined as a superficial incisional SSI that (1) required debridement in an operating room and/or (2) led to a hospital readmission within 30 days of surgery. The objective of our study was to determine the prevalence of SSISIs in a large network of community hospitals and compare hospital rankings of SSI rates of organ/space and deep SSIs (complex SSIs) with and without SSISIs. We performed a retrospective descriptive analysis of prospectively collected data on 35 NHSN surgical categories in 47 community hospitals within the Duke Infection Control Outreach Network (DICON) from 1/1/2013-12/31/2022. All hospitals used standardized surveillance and data collection strategies throughout the study period. The Wilcoxon rank-sum was used to test for differences in performance rankings of hospitals sorted by rates of complex SSIs alone compared to complex and SSISI

SSI Depth	Location Where SSI was Diagnosed			
	Current Admission (N=2618)	Outpatient (N=1581)	Readmission (N=7418)	Total (N=11617)
Deep (% row)	293 (11.2)	151 (5.7)	2182 (83.1)	2626
Organ (% row)	1521 (30.4)	164 (3.3)	3310 (66.3)	4995
SSISI (% row)	112 (5.5)	0 (0)	1926 (94.5)	2038
Superficial (% row)	692 (35.3)	1266 (64.7)	0 (0)	1958
Total	2618 (22.5)	1581 (13.6)	7418 (63.9)	11617

Surgical Procedure Category	SSI Depth				
	Organ N=4995 (% column)	Deep N= 2626 (% column)	SSISI N= 2038 (% column)	Non-SSISI Superficial N=1958 (% column)	Total N= 11617
GI	2815 (65.4)	588 (22.4)	653 (32.0)	1013 (51.7)	5069
Orthopedics	847 (17.0)	1051 (40.0)	489 (24.0)	205 (10.5)	2592
Gynecology	710 (14.2)	145 (5.5)	367 (18.0)	480 (24.5)	1702
Neurology	265 (5.3)	458 (17.4)	245 (12.0)	80 (4.1)	1048
Other	272 (5.4)	275 (10.5)	173 (8.5)	126 (6.4)	846
Cardiology	86 (1.7)	109 (4.1)	111 (5.4)	54 (2.8)	360

Rank	SSI Total Rate	Complex SSI Rate	Complex + SSISI Rate	Complex SSI Rank	Complex + SSISI Rank	Change in Rank from Complex to Complex + SSISI Rank
44	0.20	4.55	5.37	1	1	0
43	3.78	2.44	2.89	2	2	0
38	2.80	1.81	2.06	3	3	0
46	2.79	1.80	2.17	3	3	0
45	1.87	1.10	1.40	4	4	0
45	1.73	1.10	1.39	4	4	0
37	1.72	0.80	1.27	5	5	0
32	1.62	0.83	1.11	11	11	0
34	1.52	0.77	0.89	13	13	0
19	1.47	0.68	1.16	9	10	-1
40	1.47	0.65	0.80	19	10	1
22	1.46	1.07	1.28	10	7	1
3	1.42	0.84	1.18	10	9	1
33	1.38	0.57	0.83	25	15	10
47	1.13	0.59	0.74	22	23	-1
26	1.08	0.73	0.82	14	14	-2
18	1.06	0.62	0.81	15	12	0
27	1.06	0.71	0.82	15	12	-2
7	1.05	0.42	0.76	35	20	13
36	1.03	0.60	0.71	17	17	-6
12	1.02	0.65	0.86	20	14	6
36	0.93	0.67	0.74	17	14	3
23	0.92	0.69	0.85	16	19	-3
41	0.92	0.59	0.76	23	21	2
1	0.84	0.55	0.73	26	25	1
13	0.82	0.43	0.59	33	32	1
24	0.80	0.58	0.72	24	26	-2
9	0.79	0.65	0.77	18	20	-2
5	0.77	0.53	0.69	27	28	-1
17	0.76	0.45	0.59	32	31	1
38	0.75	0.47	0.63	31	29	2
39	0.75	0.49	0.60	30	29	1
2	0.72	0.42	0.58	34	36	-2
11	0.71	0.40	0.58	37	34	3
25	0.69	0.37	0.48	38	38	0
14	0.69	0.41	0.53	36	37	-1
16	0.66	0.40	0.60	39	31	8
39	0.61	0.47	0.57	30	35	-5
6	0.53	0.31	0.40	39	39	0
4	0.51	0.26	0.37	42	42	0
31	0.48	0.31	0.41	41	41	0
18	0.44	0.31	0.43	41	40	1
8	0.42	0.38	0.37	43	43	0
42	0.36	0.22	0.28	44	44	0
20	0.21	0.07	0.14	46	45	1
35	0.14	0.09	0.12	45	46	-1
15	0.05	0.02	0.02	47	47	0

rates. A two-tailed P value of .05 or less was considered significant. Overall, 11,617 SSIs occurred after 1,272,257 surgeries (0.91 SSIs/100 procedures). Out of 3,996 superficial SSIs, 2,038 (17.5% overall, 51.0% of superficial incisional) met criteria for SSISI. 112 (5.5%) were diagnosed during the current admission and required takeback to the OR for infection; 1,926 (94.5%) were diagnosed during a readmission; and 3841 (33.1%) were diagnosed during readmission and returned to the OR. (Table1) The highest proportion of SSISIs was diagnosed in patients who underwent gastrointestinal surgery (32.0%) or orthopedic surgery (24.0%). (Table2) Performance ranking of individual hospitals based on rates of complex SSIs, differed significantly when including SSISIs (p=0.02). (Table3) Discussion Our findings suggest that SSISIs make up a moderate but important proportion of SSIs in community hospitals. SSISIs can be identified through established database surveillance looking at objective measures of returning to the OR for debridement and/or readmission within 30 days. Hospital rankings differed significantly when SSISIs were added to complex SSIs to calculate SSI rates. As such, including SSISIs likely provides a more accurate depiction of SSIs with important outcomes and is not as subjective to surveillance bias. Next steps would be specifically to look at outcomes data for complex SSIs compared to SSISIs to fully evaluate the

Disclosure: Jessica Seidelman: UptoDate- Section Editor; BoneSupport-Consultant

Antimicrobial Stewardship & Healthcare Epidemiology 2024;4(Suppl. S1):s142
doi:10.1017/ash.2024.315