

Regular Article

Adopted children's internalizing symptoms and externalizing problems 8 years post-placement: Developmental trajectories and transitions

Amy L. Paine^{1,†} , Oliver Perra^{2,†} and Katherine H. Shelton¹

¹Cardiff University Centre for Human Developmental Science, School of Psychology, Cardiff University, Cardiff, UK and ²School of Nursing and Midwifery, Queen's University Belfast, Belfast, UK

Abstract

Children adopted from public care are more likely to experience mental health problems associated with their histories of early life adversity and instability, but few studies have investigated children's mental health longitudinally across developmental turning points. We followed a sample of $N = 92$ UK domestically adopted children and their families at six time points over eight years post-placement (children's ages ranged from 2 to 15 years). We used multilevel growth curve analysis to model time-related changes in children's internalizing symptoms and externalizing problems and spline models to investigate patterns of change before and after school entry. Children's internalizing symptoms followed a linear increasing trajectory, and externalizing problems followed a quadratic pattern where problems accelerated in early childhood and decelerated in late childhood. Spline models indicated an elevation in internalizing symptoms and externalizing problems as children started school. Internalizing symptoms continued to increase over time after school started and externalizing problems plateaued after the first years in school. Children adopted close in time to school entry displayed more problems when they started school. The transition to school represents a time of vulnerability for adopted children, especially for those who are adopted close in time to this transition, underscoring the need for ongoing support for their mental health across childhood.

Keywords: Adoption; externalizing; internalizing; longitudinal; school; transitions

Introduction

For children who must be removed from their birth parents, placement with an adoptive family provides stability, safety, and a nurturing environment (Brodzinsky & Palacios, 2023; Palacios et al., 2019). Yet each child adopted from public care has a complex and unique early life history that may increase their risk for developing mental health problems. Most children taken into care are removed from their birth families due to maltreatment, neglect, and/or household dysfunction (Palacios et al., 2019). Additionally, all adopted children experience loss of their birth family and encounter varying degrees of instability within their living arrangements before they are placed for adoption (Newton et al., 2000). These early experiences, that for some children are possibly compounded by exposure to substances in utero and a genetic risk for mental health problems, place them at greater risk of developing enduring mental health problems (Dekker et al., 2017; Paine et al., 2021).

Adopted children's mental health

Adopted children's developmental outcomes can vary widely (Grotevant & McDermott, 2014). Adoptees are more likely to have

difficulties in domains of cognitive functioning, academic attainment, and to experience more socioemotional and mental health problems compared to non-adopted children (Brown et al., 2017). However, they tend to out-perform children who remain in vulnerable families or in institutional care; similar findings are suggestive of the protective role of adoptive families (Askeland et al., 2017; Sonuga-Barke et al., 2017; Van den Dries et al., 2009; Vinnerljung & Hjern, 2011). Yet countries vary in the legislative and regulatory use of adoption and children's outcomes differ according to the nature of the adoption context (Dekker et al., 2017). For example, countries differ according to the prevalence of intercountry versus domestic adoption, of time spent in institutional care, with foster families, or in kinship care, and the extent to which contact between the adoptee and their birth family is allowed (Palacios et al., 2019). Context-specific research is therefore essential for informing policy and practice (Rutter et al., 2010).

In the UK, where domestic adoption is prevalent, adopted children are at greater risk for developing mental health problems than the general population and are over-represented in clinical settings (Anthony et al., 2019; DeJong et al., 2016). Children who are older at the time of their adoption are particularly at risk, being likely to have experienced a correspondingly longer exposure to maltreatment and neglect (DeJong et al., 2016; Neil et al., 2020). Older children are also more likely to have spent longer in foster care, which has different associated risk factors; some children may experience instability via multiple moves in care or poor-quality

Corresponding author: Amy L. Paine; Email: paineal@cardiff.ac.uk

[†]Amy L. Paine and Oliver Perra are joint first authors.

Cite this article: Paine, A. L., Perra, O., & Shelton, K. H. (2025). Adopted children's internalizing symptoms and externalizing problems 8 years post-placement: Developmental trajectories and transitions. *Development and Psychopathology*, 1–12, <https://doi.org/10.1017/S0954579425100175>



foster care (Meakings & Selwyn, 2016; Newton *et al.*, 2000) and, upon adoption, some children may find separation from their foster carer very distressing (Neil *et al.*, 2020). Although UK longitudinal research is limited, findings from our cohort study demonstrated that adopted children's mental health problems remained consistently high 4-years post-adoption (Paine *et al.*, 2021). When children's trajectories of internalizing (emotional) and externalizing (behavioral) problems were modeled according to child age, children showed an accelerated increase in problems into middle childhood that differed as a function of their exposure to pre-adoptive risk factors (age at placement, days in care, and number of adverse childhood experiences) (Paine *et al.*, 2021). However, our ability to chart trajectories into the school years was limited due to the young age of the sample. Therefore, the overarching aim of this study was to model time-related changes in adopted children's internalizing symptoms and externalizing problems from early to later childhood, now that children in the sample have all started school.

School transitions

Ecological perspectives emphasize that children develop within multiple, nested contexts, from the proximal interpersonal environment to the broader societal and cultural milieu (Bronfenbrenner, 1977, 1998). Starting school represents a major life transition for all children (Bronfenbrenner & Morris, 2007; Bronfenbrenner, 1977; Rimm-Kaufman & Pianta, 2000) and is an 'agent of developmental change' (Pianta *et al.*, 1995). Entry to school offers new experiences and challenges. Children must navigate relationships in a wider network of peers and with new adult authority figures. They are also expected to adjust to the demands of the new environment including contending with novel goals, routines, and expectations of increased autonomy. According to the Ecological and Dynamic Model of Transition (Rimm-Kaufman & Pianta, 2000), a child's adjustment in the school transition is determined by child characteristics (e.g., readiness, social and cognitive skills), direct (e.g., class size, child relationships) and indirect (e.g., parent-teacher communication) influences that interact through transactional processes. Evidence demonstrates that a good start at school is associated with positive academic and social outcomes (Dockett *et al.*, 2017).

The degree and types of challenges adopted children may experience in their transition to school will depend on each child's individual pre- and post-adoption experiences (Goldberg & Grotevant, 2023; Soares *et al.*, 2017). Yet many adopted children find the social and academic context of school more challenging than their peers (Brown *et al.*, 2017). Possibly, some of these difficulties may be underpinned by problems with processing social and emotional information (Paine *et al.*, 2023), and with cognitive abilities (Paine *et al.*, 2021; Wretham & Woolgar, 2017) that contribute to success in school settings (Boivin & Bierman, 2013; Sabol & Pianta, 2017). Although numerous studies have documented child, parent, and teacher views about school experiences in the context of adoption (Crowley, 2019; Goldberg *et al.*, 2017; Goldberg & Grotevant, 2023), little is known about adopted children's mental health trajectories through this transition. Moreover, we do not yet know whether age at placement is associated with this transition. Not only do prolonged experiences of early adversity place children at risk for more social, emotional, and cognitive difficulties (Paine *et al.*, 2021; Paine *et al.*, 2023), but according to Life Course Theory, the timing of multiple transitions has consequences for development (Elder *et al.*, 2003; Elder, 1998).

For children adopted in later childhood, experiencing the transition of adoption alongside or in temporal proximity to their transition to school, may have important implications for their mental health.

The present study

To our knowledge, no study has investigated adoptees' trajectories of mental health problems across development as children make the transition to formal education. We harnessed six waves of data from a UK longitudinal cohort study that followed children and their adoptive families over eight years post-placement. Our first aim was to investigate trajectories of adopted children's internalizing symptoms and externalizing problems by tracking age related patterns of change from early childhood to early adolescence. Our second aim was to investigate the impact of child (age at placement, sex, number of moves, and adverse childhood experiences) and family (socioeconomic status) factors on children's trajectories. We hypothesized that age at adoption, number of moves and number of adverse childhood experiences would be associated with increasing and accelerated mental health problems (Nadeem *et al.*, 2017; Paine *et al.*, 2021; Vandivere & McClindon, 2010). Our third aim was to investigate the effect of the transition to school on children's trajectories of internalizing symptoms and externalizing problems. Given the changes and new challenges brought about by the school transition (Rimm-Kaufman & Pianta, 2000) we hypothesized a discontinuity in the rate of change and elevation in internalizing symptoms and externalizing problems after school entry.

Method

Design

We used data from the Wales Adoption Cohort Study; a prospective longitudinal study of the early support needs and experiences adopted children and their families; see (Paine *et al.*, 2021) for more details of the study and (Paine *et al.*, 2021) and (Palacios *et al.*, 2019) for background of adoption in the UK. Local authority adoption teams across Wales were asked to send out letters on behalf of the research team to every family with whom they had placed a child for adoption between July 1, 2014 and July 31, 2015. A strategy of rolling recruitment was used, with invitation letters timed to arrive with the families several weeks after the placement began.

Ninety-six families returned the initial questionnaire at 5 months post-placement and were followed up longitudinally over 6 time points over 8 years post-placement. The present study focuses on the questionnaire follow-ups that took place at approximately 5-, 21-, 36-, 48-, 60- and 96-months post-placement (Waves 1 to 6 [W1 to W6] respectively). Of the 96 families that participated in the study at W1, 81 (84.4%) participated at W2, 73 (76.0%) at W3, 68 (70.8%) at W4, 63 (65.6%) at W5, and 70 (72.9%) participated at W6. The present study focuses on $N = 92$ (95.8%) families who completed information about children's mental health on at least one wave out of the six waves of data collection (see Table 1). Among these families, $n = 28$ (30%) had provided data in all six waves of data collection, and further $n = 28$ provided data at five of the six survey points. Only $n = 4$ (4%) and $n = 9$ (10%) provided information on one and two waves of data collection respectively (see Supplementary Material S1 for more details). In the Plan of Analysis section, we make a case for including all these participants in the analyses to minimize bias. Furthermore, sensitivity analyses indicated that the pattern of results did not change when

Table 1. Means, SDs, and ranges of Strengths and Difficulties Questionnaire (SDQ) internalizing and externalizing by wave of data collection

Internalizing Problems	Observations	Mean	SD	Min	Max
Wave 1	58	5.28	3.35	0	14
Wave 2	77	3.74	2.66	0	13
Wave 3	72	3.81	3.00	0	13
Wave 4	67	4.94	4.27	0	18
Wave 5	63	5.51	4.07	0	16
Wave 6	70	6.33	4.08	0	15
Externalizing Problems	Observations	Mean	SD	Min	Max
Wave 1	58	8.33	3.97	1	16
Wave 2	77	6.82	3.64	1	17
Wave 3	72	7.08	3.59	0	20
Wave 4	67	8.54	4.48	0	20
Wave 5	63	8.41	4.06	0	17
Wave 6	70	9.02	4.35	0	17

Note. The number of the sample with valid data was lower in W1 compared to W2 because children under 2 years of age were too young to have an SDQ completed by a parent.

participants who completed less than three waves were excluded (see Supplementary Material Section 3).

Ethical considerations

Informed consent was obtained from parents and ethical permission for the study was granted by the Research Ethics Committee for the School of Psychology at Cardiff University and permission to access the social work records was obtained from the Welsh Government via consultation with the Heads of Children's Services Group and senior adoption managers across the country.

Participants

Of the 92 children who were reported on by their parents in the longitudinal follow-up questionnaires, 45 (48.9%) were female, and were placed for adoption at a mean age of 2.97 years ($SD = 2.10$, range 0 to 9.04 years). Children spent a mean of 541.80 ($SD = 611.01$, range 0 to 2344) days with their birth parents and a mean of 544.71 ($SD = 285.01$, range 207 to 1401) days in care. Children experienced a median of 1 move ($IQR = 1$ to 3) whilst in care and a median of 3 adverse childhood experiences (ACEs) ($IQR = 0$ to 5) out of 10 categories (Felitti et al., 1998) including childhood abuse (emotional, physical, or sexual), neglect, and household dysfunction (domestic violence, parental separation, substance abuse, alcohol abuse, mental illness, or incarceration) evidenced from social worker records for each child (Paine et al., 2021). Twenty-six children (28.3%) were adopted as part of a sibling group.

Parents were a mean age of 40.61 ($SD = 7.06$, range 22 to 62) years at the time of adoption, and the majority (97.8%, $n = 90$) identified as white British. Most parents were in a heterosexual relationship (81.5%, $n = 75$), 5.4% ($n = 5$) were in a same sex relationship and 13% ($n = 12$) were single adopters. At the W1 assessment, there were a median of 4 ($IQR = 3$ to 4) people living in the household and most informants were in either full-time or part time paid work ($n = 68$, 73.9%). Gross family income and education levels were substantially higher than the UK average

(ONS, 2019), where 13% earned more than £75,000 per year and 38% had postgraduate degrees.

Characteristics of the 92 adopted children in the present study were compared to those of all children placed for adoption in Wales the same time window ($N = 374$), by reviewing Child Adoption Reports for all children adopted between July 2014 and July 2015 in Wales (Anthony et al., 2019). The sample was representative of children placed for adoption in this thirteen-month period regarding child sex, age at adoption, and past experiences of abuse and neglect (all $ps > .05$). Attrition analyses showed no differences in sociodemographic characteristics (child sex and age, parent age, relationship status, education, employment status, income, and ethnicity) between those who participated in W1 and W6 of the study (all $ps > .05$).

Procedure

Questionnaires

At each time point, families completed a questionnaire concerning sociodemographic information, pre- and post-adoption experiences, the child and parent's mental health, and adoptive family relationships. Where groups of siblings were placed together, parents were asked to report on the eldest child in the placement to overcome the problem of dependencies in the data from children within the same family. The eldest child was selected to maximize the likelihood that the focal child would be in age range for measures to be completed for them. Questionnaires were completed by either a mother (88.0% at W1, 87.5% at W2, 97.1% at W3, 92.6% at W4, 93.0% at W5, 88.6% at W6), father or non-binary parent. It was encouraged that the questionnaires should be completed by the same parent at each wave, so all families who provided follow-up questionnaires returned at least one completed by the same informant. A remuneration £20 gift voucher was sent to the family upon receipt of the questionnaire at each time point.

Measures

Child internalizing symptoms and externalizing problems

Adoptive parents completed the Strengths and Difficulties Questionnaire (SDQ; Goodman, 1997) at each wave of the study. We used internalizing symptoms (sum of emotional and peer problem scales) and externalizing problems (sum of conduct and hyperactivity scales). A higher score is indicative of more problems for all subscales (where children could score a maximum of 20 in each of these two subscales). The internalizing symptoms and externalizing problems scales had acceptable to good levels of internal consistency across all time points (α values ranged from .60 to .84).

On average, 4.42 SDQ assessments were completed for the 92 children included in data analyses ($SD = 1.52$): Only 4 children (4%) had a single SDQ completed, while 28 (30%) had five SDQs completed across the six waves of data collection, and the same number, 28, had SDQs completed in each of the six waves of data collection. Children's age at the time of assessment varied depending on the timing of their adoption. The average age of the sample with valid data was lower in W2 compared to W1 because children under 2 years of age were too young to have an SDQ completed by a parent. As we illustrate in the Plan of Analyses, the time variable used to model change is children's age, rather than the wave of data collection.

Socioeconomic characteristics

Adoptive parents' sociodemographic information was collected at W1. A general index family socioeconomic status (SES) was

created using Principal Component Analysis using Stata 13 (StataCorp, 2013). The indicators included were being a couple, working full-time, and level of family gross income, see (Paine *et al.*, 2021) for information.

Plan of analyses

A detailed description of our analyses and formal description of the models and their parameters is presented in the Supplementary Materials. To investigate trajectories of internalizing symptoms and externalizing problems across time, we applied multilevel growth curve models (also known as random effects regression models) using Stata 17 (StataCorp, 2021). The multilevel growth curve approach allowed the modeling of change in the outcomes as a function of participants' age, rather than the scheduled wave assessment. To facilitate data interpretation, we centered age at 3 years, so that the intercept in the models represented the expected average outcome score at 3 years of age. Estimation of parameters for participants with incomplete waves was carried out using Maximum Likelihood methods where data were assumed to be Missing at Random (Carpenter *et al.*, 2023). These methods provide more robust and reliable estimates compared to traditional methods such as listwise deletion or average-value imputation (Widaman, 2006). Trajectories of change over time were thus estimated on the full sample with complete and incomplete waves of data collection. However, we also ran sensitivity analyses that confirmed the pattern of results observed when we excluded participants who had completed fewer than three waves of data collection (see Supplementary Material, Section 3).

We modeled a linear, quadratic, and a cubic growth curve for internalizing symptoms and externalizing problems, respectively, comparing these models using the Likelihood Ratio (LR) test. The selected models were then refined by considering the plausibility of accessory parameters (e.g., covariance between intercept and slope residuals). After the best-fitting growth model was selected, covariates were added, thus fulfilling the second study aim. To allow for an asymmetric distribution of SES component scores, we transformed these scores using their square, then standardized the transformed scores to facilitate parameter interpretation in data analyses. A few cases did not provide information on model covariates (number of moves before placement and total number of ACEs). To avoid excluding these cases from analyses and introduce potential bias, we used multiple imputation (MI). MI methods allow to substitute missing data with plausible imputed values by sampling from a predictive distribution based on the observed data. Furthermore, the sampling process allows to account for uncertainty by creating multiple imputed datasets: parameter estimation accounts for this uncertainty by pooling estimates across the imputed datasets. In the analyses, we created 250 imputed datasets using multiple imputation chained equation methods, then used Rubin's rule to integrate results from these datasets with imputed values. More details concerning the methods are provided in Supplementary Material Sections 1 and 2.

Finally, we tested whether children's internalizing symptoms and externalizing problems followed different trajectories before and after entry to school (i.e., the September following their fourth birthday). For this aim, we used *B*-spline models (where the *B* stands for *basis*). *B*-spline models (henceforth, called spline models) allow modeling complex trajectories of change over time. They do so by splitting the time variable into different intervals. Within these intervals, different parameters and polynomial functions specify interval-specific patterns of change. For example,

the outcome may display a linear increase in the first interval, followed by a quadratic accelerated decrease in the second interval. To describe the pattern of change across the whole study with a single continuous function, these polynomials are constrained to meet at pre-defined fixed points, called *knot-points*. This effectively means that interval-specific parameters are weighted to exert increasing influence as they reach their corresponding knot-point. Each parameter exerts an influence that is localized within a specific period: as time moves towards other periods and knot-points, other parameters are gradually "turned on." In this way, spline models allow describing large variations across a study. This feature is particularly useful in modeling patterns of change that diverge or vary around key events or experiences.

In our study, a key event of interest was children's entry to primary school. Therefore, we split the time variable into two intervals that represented the trajectory of children's SDQ scores *before* and *after* school entry. Therefore, we used spline models to describe patterns of change in internalizing symptoms and externalizing problems across developmental periods characterized by a pivotal event, school entry. In our models the knot point was the exact age when each child started school. A detailed and formal presentation of the models is provided in Supplementary Material Section 4. The spline models only included $n = 62$ children who had provided data before and after starting school: The remaining 30 only provided data after they had already started school and were excluded from these analyses.

Since the $n = 62$ included in the Spline analyses had provided data before the transition into primary school, the participants included were significantly younger than the $n = 30$ excluded from the Spline analyses; average age at the start of the study 2.17 ($SD = 1.03$) and 6.07 years ($SD = 1.10$), respectively. Children in this subsample were also significantly younger at the time of placement compared to those excluded, average 1.18 ($SD = 1.04$) and 5.56 ($SD = 1.13$), respectively, and had been exposed to a significantly fewer number of ACEs, average 1.77 ($SD = 2.56$) and 5.00 ($SD = 2.00$) respectively. However, the participants included in the Spline analyses were not significantly different from those excluded considering number of moves before placement (average 1.94 [$SD = 2.08$] and 2.19 [$SD = 1.50$], *Incidence Risk Ratio* = 0.88, 95% *CI* 0.60 to 1.31) and their SES standardized score (average 0.05 [$SD = 0.94$] and -0.11 [$SD = 1.14$] respectively, $t[90] = -0.70$, $p = 0.48$). Overall, the subsample included in the Spline analyses represented a sample of younger adoptees that had been placed earlier and had been exposed to relatively fewer adverse childhood events.

Results

Descriptive statistics

Descriptive statistics are reported in Table 1. A lowess smoother function to represent internalizing symptoms and externalizing problems by children's actual age is presented in Figure 1.

Trajectories of internalizing symptoms

A multilevel regression indicated a significant linear increase in internalizing symptom scores across age. Adding polynomial slopes (quadratic, and cubic) did not improve model fit. More detailed results are reported in Supplementary Material Section 3. The final model retained inter-individual variability around the sample initial status (intercept) and rate of change (slope), but not associations (covariance) between the two. The parameters of the linear multilevel regression of internalizing symptoms on age are




Figure 1. Scatter plot and lowess smoother function across children's age by Strengths and Difficulties Questionnaire scale. Note. The diamond blue markers indicate the observed outcome scores by age across the $N = 92$ participants. Markers are transparent, therefore increasing opacity indicate increasing clustering of observations. The red lines indicate lowess smoother functions of outcomes by age (bandwidth = 0.80).

Table 2. (a) Parameters of the unadjusted multilevel growth curve of internalizing symptoms

	Fixed Effects	Coefficient	95% CI LB	95% CI UB	z
Intercept Slope	Average at 3 years	3.44	2.91	3.97	12.62***
	Linear trend	0.37	0.24	0.50	5.51***
	Random Effects	Coefficient	95% CI LB	95% CI UB	
Variances	Intercept	2.86	1.59	5.14	
	Slope	0.18	0.10	0.31	
	Within-Individuals	5.79	4.85	6.90	

reported in Table 2a, while Figure 2 reports the predicted scores in the internalizing symptoms according to these parameters. The model provided a significant fit to the data, Wald $\chi^2(1) = 30.33$, $p < .001$, and indicated that internalizing symptom raw scores increased by 0.37 units every year on average (95% CI 0.24 to 0.50).

The results of a subsequent model where we entered the main covariates are reported in Table 2b. The model, run on 250 imputed datasets (see Plan of Analyses), indicated a good fit to the data, $F(10, 919297.0) = 4.26$, $p < .001$. In Supplementary Material Section 3 we also report that the analyses did not indicate differences in the rate of change according to covariates sex, family SES, or age at time of placement. The final adjusted model reported in Table 2b indicates no significant associations between average internalizing symptom scores and sex or SES. While there were no significant associations between the average internalizing symptoms and indicators of previous adversity such as the number of moves and the number of ACEs, there was a significant association between internalizing symptom scores and age at time of placement: the results indicated that, compared to children who were placed within the first year of life, those that went into placement around the age when they started school (age >4 to 5) displayed a significant 1.85-unit increase in average internalizing symptom scores. There was also a marginal difference between

those that went into placement at birth and those that went into placement aged over 5 years, whereby the latter were predicted to show a 1.62-unit increase in internalizing symptom scores, $t = 1.75$, $p = 0.079$, see Table 2b.

The results of the adjusted model (Table 2b) also indicated a relatively high level of internalizing symptom scores variability within individuals. Estimated variance within-individual once we controlled for covariates and change over time was approximately 6 points (5.83, see Table 2b).

Overall, the results indicated a significant linear increase in internalizing symptoms across age, but the results also indicated a considerable amount of inter-individual variation in these trajectories, as well as within-individual variation across time. The only covariate that displayed significant associations with this variable was age at time of placement: compared to children that were in placement within their first year of life, those that went into placement aged between >4 to 5 years displayed higher internalizing symptoms on average.

Trajectories of externalizing problems


The analyses suggested that externalizing problem scores followed a quadratic trajectory: adding a quadratic trend according to

Table 2. (b) Parameters of the adjusted multilevel growth curve of internalizing symptoms

	Fixed Effects	Coefficient	95% CI LB	95% CI UB	t
Intercept Slope	Average at 3 Years	2.17	0.39	3.96	2.38*
	Linear trend	0.33	0.20	0.47	4.97***
Age at Placement	Birth to 1 year	reference			
	>1 to 4 years	0.58	-0.63	1.79	0.94
	>4 to 5 years	1.85	0.30	3.39	2.34*
	>5 years	1.62	-0.19	3.43	1.75 ⁺
Number of Moves	None	reference			
	One	0.32	-1.51	2.15	0.34
	Two or More	-0.13	-2.06	1.79	-0.14
Number of ACEs	None	reference			
	From 1 to 5	0.35	-1.09	1.79	0.48
	From 6 to 9	-0.16	-1.69	1.36	-0.21
SES	Std Component Score	0.10	-0.37	0.57	0.41
Sex at Birth	Female	reference			
	Male	0.70	-0.24	1.64	1.47
Variances	Random Effects	Coefficient	95% CI LB	95% CI UB	
	Intercept	2.26	1.19	4.30	
	Slope	0.16	0.09	0.28	
	Within-Individuals	5.83	4.89	6.94	

Note. 95% CI: 95% Confidence Intervals; LB = Lower Bound; UB = Upper Bound. ⁺ $p < .10$; * $p < .05$; ** $p < .01$; *** $p < .001$.

Figure 2. Strengths and Difficulties Questionnaire internalizing symptoms and externalizing problems fitted trajectories by age. Note. Trajectories based on the parameters of multilevel growth models reported in Table 2a and Table 3a. The blue dots and lines represent the scores predicted by the models. The shaded areas represent the 95% confidence intervals of predicted scores.



participants' age increased model fit significantly, $LR \chi^2(4) = 29.23, p < .001$. Furthermore, results suggested significant inter-individual variation in the quadratic slope. The final model and its parameters are reported in Table 3a, while in Figure 2 we report the

predicted externalizing problem scores by age. Details concerning model selection and comparisons are reported in Supplementary Material Section 3. The final model had a significant fit to the data, $Wald \chi^2(2) = 29.21, p < .001$. The quadratic trend indicated a

Table 3. (a) Parameters of the unadjusted multilevel model of externalizing problems

	Fixed Effects	Coefficient	95% CI LB	95% CI UB	z
Intercept Slope	Average at 3 years	6.53	5.83	7.23	18.37***
	Linear trend	0.72	0.39	1.04	4.33***
	Quadratic trend	-0.05	-0.08	-0.01	-2.85**
	Random Effects	Coefficient	95% CI LB	95% CI UB	
Variances	Intercept (a)	5.73	3.25	10.10	
	Linear Slope (b)	1.07	0.51	2.25	
	Quadratic Slope (c)	0.01	0.002	0.02	
	Covariance (a)(b)	0.10	-1.05	1.25	
	Covariance (a)(c)	-0.03	-0.15	0.09	
	Covariance (b)(c)	-0.09	-0.17	-0.01	
	Within-Individuals	5.00	4.07	6.15	

Table 3. (b) Parameters of the adjusted multilevel model of externalizing problems

	Fixed Effects	Coefficient	95% CI LB	95% CI UB	t
Intercept Slope	Average at 3 Years	6.11	3.65	8.58	4.86***
	Linear trend	0.67	0.34	1.00	4.00***
	Quadratic trend	-0.04	-0.08	-0.01	-2.71**
Age at Placement	Birth to 1 year	reference			
	>1 to 4 years	0.33	-1.28	1.93	0.40
	>4 to 5 years	1.38	-0.74	3.50	1.28
	>5 years	1.50	-0.94	3.93	1.20
Number of Moves	None	reference			
	One	-0.06	-2.64	2.51	-0.05
	Two or More	-0.90	-3.58	1.78	-0.66
Number of ACEs	None	reference			
	From 1 to 5	-0.28	-2.23	1.68	-0.28
	From 6 to 9	-0.48	-2.55	1.59	-0.45
SES	Std Component Score	0.42	-0.21	1.59	1.32
Sex at Birth	Female	reference			
	Male	0.93	-0.31	2.17	1.47
	Random Effects	Coefficient	95% CI LB	95% CI UB	
Variances	Intercept (a)	5.35	2.97	9.63	
	Linear Slope (b)	1.01	0.46	2.20	
	Quadratic Slope (c)	0.01	0.002	0.02	
	Correlation (a) (b)	0.09	-0.43	0.56	
	Correlation (a) (c)	-0.22	-0.75	0.48	
	Correlation (b) (c)	-0.95	-0.99	-0.83	
	Within-Individuals	5.01	4.07	6.18	

Note. 95% CI: 95% Confidence Intervals; LB = Lower Bound; UB = Upper Bound. + $p < .10$; * $p < .05$; ** $p < .01$; *** $p < .001$.

steeper increase in externalizing problem scores that became shallower between 7 and 9 years and reached a peak between 9 and 12 years, to then decrease, see Figure 2.

In a further model run based on 250 imputed datasets, we also included covariates: the final parameter estimates are reported in

Table 3b. Previous model tests indicated the rate of change of externalizing problem scores was not affected by key covariates (sex, age at time of placement, family SES), see Supplementary Material Section 3. Overall, none of the covariates on their own indicated a significant association with the average externalizing

Table 4. (a) Parameters of the final spline model of internalizing symptoms

Fixed Effects	Coefficient	95% CI LB	95% CI UB	z
Intercept	3.33	2.76	3.90	11.47**
Linear trend S1	-0.22	-0.74	0.30	-0.82
Linear trend S2	0.97	0.45	1.49	3.65**
Quadratic trend S2	-0.08	-0.17	-0.003	-2.04*
Random Effects	Coefficient	95% CI LB	95% CI UB	
Intercept Variance	2.08	0.99	4.36	
Slope S1 Variance	1.00	0.33	3.03	
Slope S2 Variance	0.23	0.10	0.55	
Within-Individuals Variance	4.66	3.72	5.85	

Table 4. (b) Parameters of the final spline model of externalizing problems

Fixed Effects	Coefficient	95% CI LB	95% CI UB	z
Intercept	6.31	5.57	7.05	16.73**
Linear trend S1	0.38	-0.24	1.00	1.19
Linear trend S2	0.99	0.47	1.51	3.71**
Quadratic trend S2	-0.11	-0.19	-0.03	-2.62*
Random Effects	Coefficient	95% CI LB	95% CI UB	
Intercept Variance	5.20	3.13	8.63	
Slope S1 Variance	2.26	1.18	4.34	
Slope S2 Variance	0.11	0.03	0.51	
Within-Individuals Variance	4.83	3.78	6.18	

Note. S1: Spline 1 — Before school entry; S2: Spline 2 — After school entry; 95% CI: 95% Confidence Intervals; LB: Lower Bound; UB: Upper Bound. * $p < .01$; ** $p < .001$.

problem scores, as indicated by model parameters in Table 3b. The model with covariates displayed a significant fit to the data, $F(11, 498961.0) = 3.28, p < .001$.

Overall, the results indicated that participants' pattern of change in externalizing problems followed a curvilinear trajectory: externalizing problems increased from age 3 years, but this increase decelerated progressively to reach a peak in late childhood. Similarly to the internalizing scores, the results indicated a considerable amount of variation between participants' externalizing trajectories, and furthermore, there was a large within-child variability in externalizing problem scores from one age to another (see Table 3b). None of the covariates considered displayed a unique association with the externalizing scores while we controlled for other covariates.

Internalizing symptoms and externalizing problems at the transition to schooling

We ran spline models on the $n = 62$ children that were observed before and after entering school. Details about model selection are reported in Supplementary Materials Section 5. The spline model for internalizing symptoms indicated a non-significant linear rate of change in the period before children school entry, while the period following school entry was characterized by a quadratic rate of change: we report the estimated parameters of the final spline model in Table 4a. In Figure 3 we report the estimated predicted

scores based on the spline model parameters: the model predicted that internalizing symptoms remained relatively stable before school entry but increased steadily after school entry. Internalizing symptom scores continued to increase, but a shallower decrease was predicted after age 8 years. Inclusion of covariates in the model confirmed the trends indicated, and, similarly to the results from the multilevel growth curve analyses reported in the previous section, indicated elevated internalizing symptom scores of children who were placed for adoption around the time they were bound to start school (age >3 to 5 years), see Supplementary Materials Section 5.

Spline models run on externalizing problems indicated a non-significant linear increase in these scores before children school entry, while externalizing problems were predicted to follow a quadratic trend after children school entry: the model parameters are reported in Table 4b and the predicted scores based on the model are represented in Figure 3. Externalizing problems were predicted to increase more steeply after children school entry. Compared to internalizing symptoms, however, they reached a peak earlier (approximately by 7 years of age) and, while internalizing symptoms were still expected to increase by 10 years of age (although at a slower rate), externalizing problems were predicted to start decreasing by this age. Inclusion of covariates did not indicate these showed significant associations with externalizing problem trajectories, see Supplementary Material Section 5.

Overall, the results of the spline models run on a sub-sample of participants observed before and after school entry suggested large variations in the patterns of change of mental health problems across different stages of childhood, with patterns diverging in correspondence of the transition into primary school. Internalizing symptoms increased steeply after school entry, while after age 8 years they continued to increase at a lower rate. Externalizing problems also increased steeply after school entry, but this increase decelerated after age 7 to then decrease following age 10 years. The results indicate, however, substantial between-child variation in the fitted trajectories, as well as considerable within-child variation from one assessment to another.

Discussion

To our knowledge, the present study represents the first longitudinal study that has charted the mental health trajectories of UK domestically adopted children from early childhood to early adolescence. Across this period, we identified a linear increase in children's internalizing symptoms and a quadratic pattern for externalizing problems, where problems increased from early to middle childhood and appeared to decelerate and decline from late childhood to adolescence. Age at adoption was associated with children's trajectories of internalizing symptoms. Children who were older at the time of adoption had more internalizing symptoms, but this association weakened when children were adopted after age five. Spline models indicated that entry to school marks a turning point of significant increases in children's internalizing symptoms and externalizing problems in their first years of school. Our findings have important implications for guiding interventions and support for adoptees across childhood and into adolescence.

Trajectories and predictors of children's internalizing symptoms and externalizing problems

We found that children's trajectories appeared to mirror what tends to be a minority of children in the general population who




Figure 3. Predicted Strengths and Difficulties Questionnaire scores by age according to spline models. *Note.* $N = 62$ children who provided data before and after school entry. Spline models divided the observation period into intervals, before school entry and after school entry. The red lines represent predicted score according to spline models for internalizing symptoms and externalizing problems respectively. The markers represent the observed outcome scores for individual children. The blue circles indicate scores reported before school entry, while red square markers are used to indicate scores obtained after school entry. Markers are transparent: greater density of colors indicate more observations obtaining the same score at the same age.

exhibit increasing internalizing symptoms and externalizing problems from early childhood to adolescence. In our sample, children's internalizing symptoms followed a linear increasing trajectory from 3 to 15 years. Population studies typically identify a small proportion of children who follow trajectories characterized by increasing levels of internalizing symptoms from early childhood to adolescence (i.e., 11% low and increasing, 3% high and increasing internalizing symptoms; (Papachristou & Flouri, 2020); see also (Davis et al., 2015; Fanti & Henrich, 2010; Parkes et al., 2016). For externalizing problems, adopted children in this sample tended to follow a trajectory of high and increasing scores, following a similar pattern to what is observed in a small proportion of children in population studies (i.e., 5% high and increasing externalizing problems, (Papachristou & Flouri, 2020); see also (Patalay et al., 2016). Our findings suggest, however, that this increase in externalizing problems was followed by a deceleration and decline from late childhood to early adolescence. Nonetheless, the patterns of increasing internalizing symptoms and externalizing problems across childhood for adoptees is concerning, given increasing trajectory patterns are associated with adverse outcomes, such as negative changes in academic attainment (Patalay et al., 2016) and self-harm and drug use in adolescence (Papachristou & Flouri, 2020).

On the basis of prior literature (Nadeem et al., 2017; Paine et al., 2021; Vandiver & McKlindon, 2010) we expected that number of moves before adoption, ACEs, and age at adoption would be associated with increasing and accelerated mental health problems. Our findings partially supported these expectations. Children who were older at the time of adoption experienced more internalizing symptoms, yet the association weakened when children were adopted after the age of 5 years. Although age at adoption does not fully capture pre-adoptive history, it can represent a proxy for early adversity (Tan & Marfo, 2016). Children adopted from public care later in childhood are more likely to have experienced prolonged adversity (Anthony et al., 2019) and instability (Newton et al., 2000) associated with negative cascades into different domains of development (Cicchetti, 2013). Although these findings provide a

useful indicator for practitioners to monitor children at greater risk of difficulties, more research is needed to disentangle specific risk and protective factors associated with children's trajectories of mental health problems (Lacey & Minnis, 2020). Furthermore, it is noteworthy that we detected great intra-individual variability in internalizing symptoms and externalizing problems that we were not able to explain with our age and time-invariant covariates. A question of interest could be to investigate how adoptees' intra-individual variability compares to that of the general population in terms of (in)stability of mental health problems. Quite possibly, the unexplained variability and intra-individual instability in adoptees may be explained by contextual, age-specific factors that were not included in these analyses, such as onset of puberty.

Commencing formal schooling represents a major life transition and period of rapid change for all children (Rimm-Kaufman & Pianta, 2000). Taking a life course perspective (Elder et al., 2003; Elder, 1998), we hypothesized that the demands of this transition might be particularly challenging for children adopted in a similar timeframe. Our subsequent analyses, therefore, pinpointed the short window of time as adoptees made the transition to school. We used Spline models to investigate variations in the patterns of change before and after transition to school. An advantage of using these models resides in the ability to allow for piecemeal patterns of growth at different stages of development, which, in turn, makes them suitable to model significant variations in correspondence of key events or turning points (Howe et al., 2013). One of the challenges in using Spline models lies in choosing sensible numbers and timings of knot-points that define intervals within a study period: in our study we selected a single knot-point that represented the key transition to school and, furthermore, the actual age when each individual child started school defined this knot-point. Thus, the partition of the study period into two intervals with piecemeal growth patterns was based on an objective criterion that represented a key transition in children's lives.

The results of Spline models indicated discontinuities in the rate of change children's internalizing symptoms and externalizing

problems following transition to school. Both children's internalizing symptoms and externalizing problems showed a significant linear increase after children started school. Children's internalizing symptoms continued to rise across the school years, but externalizing problems plateaued after the first three years of school. Later age of adoption was associated with elevated children's internalizing symptoms. Being adopted around school age (>3 to 5 years) was associated with increased internalizing symptoms after school entry compared to children adopted shortly after birth or in early childhood.

Practical implications

The first years of school are vital for future success (Dockett *et al.*, 2017). Our findings highlight the need for special consideration of children who are experiencing multiple life transitions. Our findings underscore the importance of avoiding unnecessary delays to permanency planning, while making sure that transitions move at a pace that is comfortable for the child and accounts for other life events (Neil *et al.*, 2020). Emphasis should also be placed on building collaborative and cohesive interpersonal systems between those who support the child through these normative and less normative transitions (Cook *et al.*, 2018). Part of this approach should involve shared professional understanding of children's adoption status and pre-placement history, where appropriate, to facilitate and anticipate support needs (Goldberg & Grotevant, 2023).

The escalating trajectories of adoptees' internalizing symptoms and externalizing problems through childhood underscores the importance of early intervention when signs of mental health problems emerge. Some children may experience elevated, yet subthreshold problems, that mean they may not qualify for specialist support from mental health services (DeJong *et al.*, 2016). However, taken together with other research (Copeland *et al.*, 2015) our findings suggest these early difficulties place adoptees at risk for escalating problems and disorder later in life. It is vital that alongside the present findings, practitioners recognize the heterogeneity of adoptees' early experiences and the consequences these have for their neurodevelopmental profiles at an individual level (DeJong *et al.*, 2016; Woolgar & Simmonds, 2019). Our previous work indicates that systematic assessment of adoptees' areas of need is a promising way to identify socioemotional and cognitive transdiagnostic risk factors that may underpin mental health problems (Paine *et al.*, 2021, 2023) that may be targeted with early intervention.

Limitations

We acknowledge some study limitations. First, the relatively small sample size precluded disentangling multiple trajectories that can be achieved via larger population studies (Davis *et al.*, 2015; Fanti & Henrich, 2010; Papachristou & Flouri, 2020; Parkes *et al.*, 2016; Patalay *et al.*, 2016). Second, the spline models were based on a smaller subsample of children who had questionnaires returned by parents before and after school entry; children in these analyses entered the study at a younger age, which also meant that many were in placement at an earlier age and were exposed to fewer adverse events relative to the rest of the sample. Third, our findings relied on single informant reports. Although parent reports are a critical source of information, future larger scale studies should avail of multiple reports of more domains of children's school readiness and social and cognitive development (Pears *et al.*, 2013).

This information could also be complemented by children's self-report about their school experiences.

Finally, it is also important to note that relatively few children had reached secondary school age by the time of our last follow up. This therefore precluded us from investigating the transition to secondary school as a further turning point for mental health problems. As such, the apparent deceleration and decline of externalizing problems later in childhood to adolescence should be taken with caution. Transition to secondary school is a period of heightened worry for young people, often concerning new academic and social pressures, a new physical environment (Moore *et al.*, 2021; Rice *et al.*, 2021) and coinciding with ongoing pubertal development. Evidence suggests that children who have high levels of difficulties at the end of primary school continue to experience difficulties across the transition to secondary (Donaldson *et al.*, 2024). For adoptees, these challenges occur alongside a period where they develop a deeper social and cognitive understanding of adoption and adoptive identity (Brodzinsky & Palacios, 2023). Despite these limitations, the results are based on advanced models that enabled a thorough investigation of complex patterns of change, while accounting for potential limitations including the timings of data collection and missingness.

Conclusion

Adoption provides one of the best alternatives for children to have a secure family for life (Brodzinsky & Palacios, 2023). Despite the quality of parenting and home life offered by adoptive families (Anthony *et al.*, 2019; Paine *et al.*, 2021), our analyses demonstrate that through early childhood to early adolescence, children adopted from public care are at greater risk for experiencing increasing internalizing symptoms and externalizing problems. Entry into school may represent a particular challenge and time of vulnerability for young adoptees, particularly if they experience this alongside or temporally close to their adoption. These findings emphasize the vital importance of continued support for adoptive families. Indeed, efforts to support adopted children through the school transition, particularly those also contending with other major life events, may enable them to flourish through the school years.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0954579425100175>.

Data availability statement. The data that support the findings of this study are available on reasonable request from the corresponding author, ALP. The participants of this study did not give written consent for their data to be shared publicly, so due to the sensitive nature of the research and privacy/ethical restrictions, the supporting data is not publicly available.

Acknowledgments. Our sincere thanks go to the staff from the local authority adoption teams in Wales, who kindly assisted with contacting families, and to our research advisory group for their guidance. We thank the families who took part in this study and Sarah Meakings, Charlotte Robinson, and Janet Whitley for research assistance.

Funding statement. The preparation of this manuscript was funded by the Welsh Government. The Wales Adoption Study was initially funded by Health and Care Research Wales, a Welsh Government body that develops, in consultation with partners, strategy and policy for research in the NHS and social care in Wales (2014–2016, Grant reference: SC-12-04; Principal Investigator: Katherine Shelton, co-investigators: Julie Doughty; Sally Holland; Heather Ottaway). Amy L. Paine was funded by the Waterloo Foundation (Grant reference: 738/3512) and the Economic and Social Research Council (Grant ref: ES/T00049X/1).

Competing interests. The authors have no conflict of interest to declare.

References

- Anthony, R. E., Paine, A. L., & Shelton, K. H. (2019). Adverse childhood experiences of children adopted from care: The importance of adoptive parental warmth for future child adjustment. *International Journal of Environmental Research and Public Health*, 16(12), 2212.
- Askeland, K. G., Hysing, M., La Greca, A. M., Aarø, L. E., Tell, G. S., & Sivertsen, B. (2017). Mental health in internationally adopted adolescents: A meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry*, 56(3), 203–213, e201.
- Boivin, M., & Bierman, K. L. (2013). *Promoting school readiness and early learning: Implications of developmental research for practice*. Guilford Publications.
- Brodzinsky, D., & Palacios, J. (2023). *The adopted child*. Cambridge University Press.
- Bronfenbrenner, U. (1977). Toward an experimental ecology of human development. *American Psychologist*, 32(7), 513–531.
- Bronfenbrenner, U. (1998). The ecology of developmental processes. In *Handbook of child psychology* (p. 1). John Wiley & Sons, Inc.
- Bronfenbrenner, U., & Morris, P. A. (2007). The bioecological model of human development. In *Handbook of child psychology* (p. 1). Wiley.
- Brown, A., Waters, C. S., & Shelton, K. H. (2017). A systematic review of the school performance and behavioural and emotional adjustments of children adopted from care. *Adoption & Fostering*, 41(4), 346–368.
- Carpenter, J. R., Bartlett, J. W., Morris, T. P., Wood, A. M., Quartagno, M., & Kenward, M. G. (2023). *Multiple imputation and its application*. John Wiley & Sons.
- Cicchetti, D. (2013). Annual research review: Resilient functioning in maltreated children—past, present, and future perspectives. *Journal of Child Psychology and Psychiatry*, 54(4), 402–422.
- Cook, K. D., Dearing, E., & Zachrisson, H. D. (2018). Is parent-teacher cooperation in the first year of school associated with children's academic skills and behavioral functioning? *International Journal of Early Childhood*, 50(2), 211–226.
- Copeland, W. E., Wolke, D., Shanahan, L., & Costello, E. J. (2015). Adult functional outcomes of common childhood psychiatric problems: A prospective, longitudinal study. *JAMA Psychiatry*, 72(9), 892–899. <https://doi.org/10.1001/jamapsychiatry.2015.0730>
- Crowley, C. (2019). Exploring the views and perceptions of adopted young people concerning their education and social development: An interpretative phenomenological analysis. *Educational Psychology in Practice*, 35(2), 165–183.
- Davis, S., Votruba-Drzal, E., & Silk, J. S. (2015). Trajectories of internalizing symptoms from early childhood to adolescence: Associations with temperament and parenting. *Social Development*, 24(3), 501–520.
- DeJong, M., Hodges, J., & Malik, O. (2016). Children after adoption: Exploring their psychological needs. *Clinical Child Psychology and Psychiatry*, 21(4), 536–550.
- Dekker, M. C., Tieman, W., Vinke, A. G., van der Ende, J., Verhulst, F. C., & Juffer, F. (2017). Mental health problems of Dutch young adult domestic adoptees compared to non-adopted peers and international adoptees. *International Social Work*, 60(5), 1201–1217.
- Dockett, S., Griebel, W., & Perry, B. (2017). Transition to school: A family affair. In *Families and transition to school* (pp. 1–18). Springer.
- Donaldson, C., Hawkins, J., Rice, F., & Moore, G. (2024). *Trajectories of mental health across the primary to secondary school transition* (p. e12244). JCPP Advances.
- Elder, G. H. (1998). The life course as developmental theory. *Child Development*, 69(1), 1–12.
- Elder, G. H., Johnson, M. K., & Crosnoe, R. (2003). *The emergence and development of life course theory*. Springer.
- Fanti, K. A., & Henrich, C. C. (2010). Trajectories of pure and co-occurring internalizing and externalizing problems from age 2 to age 12: Findings from the national institute of child health and human development study of early child care. *Developmental Psychology*, 46(5), 1159–1175.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The adverse childhood experiences (ACE) study. *American Journal of Preventive Medicine*, 14(4), 245–258.
- Goldberg, A. E., Frost, R. L., & Black, K. A. (2017). There is so much to consider”: School-related decisions and experiences among families who adopt noninfant children. *Families in Society*, 98(3), 191–200.
- Goldberg, A. E., & Grotevant, H. (2023). What do teachers know about adoptive families, and how do they use it to serve adopted children? *Adoption Quarterly*, 26(1), 51–76.
- Goodman, R. (1997). The strengths and difficulties questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38(5), 581–586. <https://doi.org/10.1111/j.1469-7610.1997.tb01545.x>
- Grotevant, H. D., & McDerrott, J. M. (2014). Adoption: Biological and social processes linked to adaptation. *Annual Review of Psychology*, 65(1), 235–265.
- Howe, L. D., Tilling, K., Matijasevich, A., Petherick, E. S., Santos, A. C., Fairley, L., Wright, J., Santos, I. S., Barros, A. J., Martin, R. M. and Kramer, M. S. (2013). Linear spline multilevel models for summarising childhood growth trajectories: A guide to their application using examples from five birth cohorts. *Statistical Methods in Medical Research*, 25(5), 1854–1874. <https://doi.org/10.1177/0962280213503925>
- Lacey, R. E., & Minnis, H. (2020). Practitioner review: Twenty years of research with adverse childhood experience scores—advantages, disadvantages and applications to practice. *Journal of Child Psychology and Psychiatry*, 61(2), 116–130.
- Meakings, S., & Selwyn, J. (2016). She was a foster mother who said she didn't give cuddles’: The adverse early foster care experiences of children who later struggle with adoptive family life. *Clinical Child Psychology and Psychiatry*, 21(4), 509–519.
- Moore, G., Angel, L., Brown, R., Van Godwin, J., Hallingberg, B., & Rice, F. (2021). Socio-economic status, mental health difficulties and feelings about transition to secondary school among 10–11 year olds in Wales: Multi-level analysis of a cross sectional survey. *Child Indicators Research*, 14(4), 1597–1615.
- Nadeem, E., Waterman, J., Foster, J., Paczkowski, E., Belin, T. R., & Miranda, J. (2017). Long-term effects of pre-placement risk factors on children's psychological symptoms and parenting stress among families adopting children from foster care. *Journal of Emotional and Behavioral Disorders*, 25(2), 67–81.
- Neil, E., Morciano, M., Young, J., & Hartley, L. (2020). Exploring links between early adversities and later outcomes for children adopted from care: Implications for planning post adoption support. *Developmental Child Welfare*, 2(1), 52–71.
- Newton, R. R., Litrownik, A. J., & Landsverk, J. A. (2000). Children and youth in foster care: Disentangling the relationship between problem behaviors and number of placements. *Child Abuse & Neglect*, 24(10), 1363–1374.
- Office for National Statistics (2019). Gross household income, UK, financial year ending 2018. Retrieved from <https://www.ons.gov.uk/peoplepopulationandcommunity/personalandhouseholdfinances/incomeandwealth/adhocs/009772grosshouseholdincomeukfinancialyearending2018>
- Paine, A. L., Burley, D., Anthony, R., Van Goozen, S. H., & Shelton, K. H. (2021). The neurocognitive profiles of children adopted from care and their emotional and behavioral problems at home and school. *Child Neuropsychology*, 27(1), 17–36.
- Paine, A. L., Fahey, K., Anthony, R. E., & Shelton, K. H. (2021). Early adversity predicts adoptees' enduring emotional and behavioral problems in childhood. *European Child & Adolescent Psychiatry*, 30(5), 721–732.
- Paine, A. L., Perra, O., Anthony, R., & Shelton, K. H. (2021). Charting the trajectories of adopted children's emotional and behavioral problems: The impact of early adversity and postadoptive parental warmth. *Development and Psychopathology*, 33(3), 922–936.
- Paine, A. L., van Goozen, S. H., Burley, D. T., Anthony, R., & Shelton, K. H. (2023). Facial emotion recognition in adopted children. *European Child & Adolescent Psychiatry*, 32(1), 87–99.

- Palacios, J., Adroher, S., Brodzinsky, D. M., Grotevant, H. D., Johnson, D. E., Juffer, F., Martínez-Mora, L., Muhamedrahimov, R. J., Selwyn, J., & Simmonds, J. (2019). Adoption in the service of child protection: An international interdisciplinary perspective. *Psychology, Public Policy, and Law*, 25(2), 57–72.
- Papachristou, E., & Flouri, E. (2020). Distinct developmental trajectories of internalising and externalising symptoms in childhood: Links with mental health and risky behaviours in early adolescence. *Journal of Affective Disorders*, 276, 1052–1060.
- Parkes, A., Sweeting, H., & Wight, D. (2016). Early childhood precursors and school age correlates of different internalising problem trajectories among young children. *Journal of Abnormal Child Psychology*, 44(7), 1333–1346.
- Patalay, P., Fink, E., Fonagy, P., & Deighton, J. (2016). Unpacking the associations between heterogeneous externalising symptom development and academic attainment in middle childhood. *European Child & Adolescent Psychiatry*, 25(5), 493–500.
- Pears, K. C., Fisher, P. A., Kim, H. K., Bruce, J., Healey, C. V., & Yoerger, K. (2013). Immediate effects of a school readiness intervention for children in foster care. *Early Education & Development*, 24(6), 771–791.
- Pianta, R. C., Steinberg, M. S., & Rollins, K. B. (1995). The first two years of school: Teacher-child relationships and deflections in children's classroom adjustment. *Development and Psychopathology*, 7(2), 295–312.
- Rice, F., Ng-Knight, T., Riglin, L., Powell, V., Moore, G. F., McManus, I. C., Shelton, K. H., & Frederickson, N. (2021). Pupil mental health, concerns and expectations about secondary school as predictors of adjustment across the transition to secondary school: A longitudinal multi-informant study. *School Mental Health*, 13(2), 279–298.
- Rimm-Kaufman, S. E., & Pianta, R. C. (2000). An ecological perspective on the transition to kindergarten: A theoretical framework to guide empirical research. *Journal of Applied Developmental Psychology*, 21(5), 491–511.
- Rutter, M., Sonuga-Barke, E. J., Beckett, C., Castle, J., Kreppner, J., Kumsta, R., Schlotz, W., Stevens, S., Bell, C. A., & Gunnar, M. R. (2010). Deprivation-specific psychological patterns: Effects of institutional deprivation. *Monographs of the society for research in child development*, 75(1), i-253.
- Sabol, T. J., & Pianta, R. C. (2017). The state of young children in the United States: School readiness. In *The Wiley handbook of early childhood development programs, practices, and policies* (pp. 1–17). Wiley.
- Soares, J., Barbosa-Ducharne, M., Palacios, J., & Fonseca, S. (2017). Being adopted in the school context: Individual and interpersonal predictors. *Children and Youth Services Review*, 79, 463–470.
- Sonuga-Barke, E. J., Kennedy, M., Kumsta, R., Knights, N., Golm, D., Rutter, M., Maughan, B., Schlotz, W., & Kreppner, J. (2017). Child-to-adult neurodevelopmental and mental health trajectories after early life deprivation: The young adult follow-up of the longitudinal English and Romanian adoptees study. *The Lancet*, 389(10078), 1539–1548.
- StataCorp (2013). (*Version 13*). [Computer software]. College Station, TX: StataCorp LLC.
- StataCorp (2021). *Stata Statistical Software: Release 17*. College Station, TX: StataCorp LLC.
- Tan, T. X., & Marfo, K. (2016). Pre-adoption adversity and behavior problems in adopted Chinese children: A longitudinal study. *Journal of Applied Developmental Psychology*, 42, 49–57.
- Van den Dries, L., Juffer, F., Van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2009). Fostering security? A meta-analysis of attachment in adopted children. *Children and Youth Services Review*, 31(3), 410–421.
- Vandivere, S., & McKlindon, A. (2010). The well-being of US children adopted from foster care, privately from the United States and internationally. *Adoption Quarterly*, 13(3-4), 157–184.
- Vinnerljung, B., & Hjern, A. (2011). Cognitive, educational and self-support outcomes of long-term foster care versus adoption. A Swedish national cohort study. *Children and Youth Services Review*, 33(10), 1902–1910.
- Widaman, K. F. (2006). Best practices in quantitative methods for developmentalists: III. Missing data: What to do with or without them. *Monographs of the Society for Research in Child Development*, 71(3), 42–64.
- Woolgar, M., & Simmonds, J. (2019). The diverse neurobiological processes and legacies of early adversity: Implications for practice. *Adoption & Fostering*, 43(3), 241–255. <https://doi.org/10.1177/0308575919877012>
- Wretham, A. E., & Woolgar, M. (2017). Do children adopted from British foster care show difficulties in executive functioning and social communication? *Adoption & Fostering*, 41(4), 331–345.