

quality improvement strategy was created using the Plan-Do-Study-Act (PDSA) model of creating change. A series of three interventions were implemented and subsequently analysed using the PDSA model. These included creating an admission checklist and placing this in doctors' on-call rooms across the trust, distributing the checklist to wards across hospital sites and service lines and lastly, sending out reminder emails to doctors at the Trust. A re-audit was completed after each intervention.

Results. The baseline BCAC completion rate was 60.5% (n = 76) in February 2023. This increased to 63.75% (n = 80) following the creation of the admission checklist. Distributing the checklist throughout the Trust resulted in a further improvement to 76.5% (n = 81). In July 2023, the final data collected 6 weeks post reminder email demonstrated a BCAC completion rate of 75% (n = 68).

Conclusion. Through the application of the PDSA model, alongside focused interventions aimed at raising Trust-wide awareness of the BCAC form, we managed to obtain a 14.5% increase in BCAC completion rates between 13 February and 31 July 2023. Whilst not achieving the initial goal of 100% completion, this quality improvement project helped to bring a renewed focus to the timely completion of capacity and consent assessments at the Trust. Moving forward, we hope to build on this improvement by exploring additional strategies to reach a 100% BCAC completion rate, thereby promoting patient safety and autonomy.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Confidence in Adherence to Antidepressant Prescribing Guidelines Among Liaison Psychiatrists

Dr Lois Zac-Williams^{1*} and Dr Gareth Smith²

¹Barking, Havering and Redbridge University Hospitals NHS Trust, London, United Kingdom and ²North East London NHS Foundation Trust, London, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.93

Aims. This project aims to increase confidence among Liaison Psychiatrists (LPs) in North East London Foundation Trust (NELFT) regarding their adherence to the prescribing guidelines for antidepressants by 25% in accordance with the standard set by Psychiatric Liaison Accreditation Network (PLAN).

Background

The prescribing guidelines in this project are based on Standard 21 from 7th Edition Standards as devised by PLAN which states:

"When medication is prescribed, specific treatment goals are set with the patient, the risks (including interactions) and benefits are discussed, a timescale for response is set and patient consent is recorded."

This project focuses on antidepressants because they are one of the widely used medications in psychiatry that doctors of all grades working in Liaison Psychiatry will be familiar with to some extent. Adhering to this validated guideline would promote gaining informed consent and patients' involvement in their care, which studies have shown can increase adherence to treatment.

Methods. Circulated an eight-question survey by email based on Standard 21 of 7th Edition Standards document by PLAN to LPs in NELFT. Conducted two Plan-Do-Study-Act (PDSA) cycles. The first PDSA uses a teaching session as the intervention and

explained the importance of antidepressant guidelines and what areas LPs need to address with patients. The second intervention uses a poster to reinforce the key points. After each intervention a reissued survey assesses the change in responses.

Results. The baseline survey response rate was 10 out of 15 LPs, made up of seven consultants, two registrars and one foundation year doctor. The lowest levels of confidence were reported around providing patients with printed information on their prescribed antidepressant with the majority of consultants reporting the lowest level of confidence. The highest levels of confidence across all medical grades were reported around discussing a specific treatment goal and explaining the benefits of treatment with antidepressant medication.

Conclusion. From the baseline data, it can be concluded that providing patients with printed information on newly prescribed antidepressants is the area that LPs, particularly consultants, are the least confident about regarding their adherence to prescribing guidelines. Future cycles of this quality improvement project can assess how incorporating teaching on antidepressant prescribing guidelines into trust induction sessions impacts LPs confidence in their adherence.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

4 Service Evaluation

Review of Rapid Tranquillisation Guidelines Across NHS Trusts in England

Miss Claudia Chavasse^{1*}, Mr James O'Neill¹, Mr Daniel Romeu² and Mr Alex Graham²

¹University of Leeds, Leeds, United Kingdom and ²Leeds and York Partnership Foundation Trust, Leeds, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.99

Aims. Rapid tranquillisation – the parenteral administration of a sedating psychotropic – is frequently utilised to manage acute behavioural disturbances. Each mental health trust in England utilises independent guidelines for rapid tranquillisation, which vary geographically in both recommendations for therapeutic agents, as well as the format in which this information presented. Audits have identified that there is currently poor adherence to rapid tranquillisation protocol guidelines; this may be due to a lack of guideline clarity allowing for personal interpretation. This service evaluation aims to determine the clarity and uniformity of protocols outlined in mental health trust guidelines, in addition to analysing the outcomes of guideline testing to identify if there is consistency between policies, or whether outcomes varied depending on the trust guidelines used.

Methods. Five reviewers (of differing positions throughout clinical training) utilised 52 guidelines from each mental health trust in England, as well as Maudsley and NICE. These were assessed using the same fictional scenario, which simulated a common presentation in which the use of rapid tranquillisation is required. Reviewers deduced the most appropriate therapeutic agent according to the guideline, rated the clarity of each guideline and were invited to leave comments highlighting the guideline's useability.

Results. Seven different management plans were generated by the majority of respondents from the 52 guidelines. Lorazepam was the most frequently selected therapeutic agent.