



COMPOSITIO MATHEMATICA

Non-rigid quartic 3-folds

Hamid Ahmadinezhad and Anne-Sophie Kaloghiros

Compositio Math. **152** (2016), 955–983.

[doi:10.1112/S0010437X15007769](https://doi.org/10.1112/S0010437X15007769)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Non-rigid quartic 3-folds

Hamid Ahmadinezhad and Anne-Sophie Kaloghiros

ABSTRACT

Let $X \subset \mathbb{P}^4$ be a terminal factorial quartic 3-fold. If X is non-singular, X is *birationally rigid*, i.e. the classical minimal model program on any terminal \mathbb{Q} -factorial projective variety Z birational to X always terminates with X . This no longer holds when X is singular, but very few examples of non-rigid factorial quartics are known. In this article, we first bound the local analytic type of singularities that may occur on a terminal factorial quartic hypersurface $X \subset \mathbb{P}^4$. A singular point on such a hypersurface is of type cA_n ($n \geq 1$), or of type cD_m ($m \geq 4$) or of type cE_6, cE_7 or cE_8 . We first show that if $(P \in X)$ is of type cA_n , n is at most 7 and, if $(P \in X)$ is of type cD_m , m is at most 8. We then construct examples of non-rigid factorial quartic hypersurfaces whose singular loci consist (a) of a single point of type cA_n for $2 \leq n \leq 7$, (b) of a single point of type cD_m for $m = 4$ or 5 and (c) of a single point of type cE_k for $k = 6, 7$ or 8 .

1. Introduction

A classical problem in algebraic geometry is to determine which quartic hypersurfaces in \mathbb{P}^4 are rational. In their seminal paper [IM71], Iskovskikh and Manin proved that a non-singular quartic hypersurface $X_4 \subset \mathbb{P}^4$ is *birationally rigid* (see the precise definition below) and, in particular, is not rational.

The classical minimal model program (MMP) shows that a uniruled projective 3-fold Z with terminal singularities is birational to a *Mori fibre space* X/S . More precisely, there is a small morphism $f : \tilde{Z} \rightarrow Z$, where \tilde{Z} is terminal and \mathbb{Q} -factorial (see [KM98, § 6.3]) and the classical MMP $\psi : \tilde{Z} \dashrightarrow X$ terminates with a Mori fibre space X/S (see [KM98, § 3]). Neither the morphism f nor the birational map ψ is unique in general. Mori fibre spaces are end products of the MMP and hence should be seen as *distinguished representatives* of their classes of birational equivalence. In general, there may be more than one Mori fibre space in a class of birational equivalence. The *pliability* of a uniruled terminal 3-fold Z is the set of distinguished representatives in its class of birational equivalence, that is,

$$\mathcal{P}(Z) = \{X/T \text{ Mori fibre space birational to } Z\} / \sim,$$

where \sim denotes the *square birational equivalence* defined in [Cor95, Definition 5.2]. If X itself is a Mori fibre space, X is called *birationally rigid* if its pliability is $\mathcal{P}(X) = \{X\}$.

Received 9 December 2013, accepted in final form 7 July 2015, published online 22 December 2015.

2010 Mathematics Subject Classification 14E05, 14J30, 14J45 (primary).

Keywords: birational maps, quartic hypersurfaces, birational rigidity.

The first author was partially supported by the Austrian Science Fund (FWF) (grant number P22766-N18) and the second author was supported by the Engineering and Physical Sciences Research Council (grant number EP/H028811/1).

This journal is © [Foundation Compositio Mathematica](http://www.cambridge.org/compositio) 2015. This article is distributed with Open Access under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided that the original work is properly cited.

A quartic hypersurface $X \subset \mathbb{P}^4$ with terminal singularities is a Mori fibre space precisely when it is factorial, that is, when every Weil divisor on X is Cartier. The two conditions on the singularities of a Mori fibre space are quite different: requiring that the singularities of X are terminal is a local analytic condition, while factoriality is a global topological condition. Mella extended Iskovskikh and Manin's result and proved that terminal factorial quartic hypersurfaces with no worse than ordinary double points are birationally rigid (see [Mel04, Theorem 2]). Factoriality is a crucial condition for this to hold. Indeed, general determinantal quartic hypersurfaces are examples of rational nodal quartic hypersurfaces (see the introductions of [Pet98, Mel04]) but are not factorial. Todd discussed several examples of non-factorial rational nodal quartic hypersurfaces: the Burkhardt quartic studied in [Tod36] has 45 nodes (see also [Pet98, § 5.1]); an example with 36 nodes is mentioned in [Tod33] (see also [Pet98, Example 6.4.2]) and two examples with 40 nodes are studied in [Tod35] (see also [Pet98, Examples 6.2.1 and 6.2.2]). In fact, *most* terminal non-factorial quartic hypersurfaces are rational [Kal12].

However, factoriality alone is not sufficient to guarantee birational rigidity. There are several known examples of non-rigid terminal factorial quartic hypersurfaces: an example with a cA_2 point was studied in [CM04] and entry No. 5 in [Ahm12, Table 1] is an example with a cA_3 point. In this paper, we show that these examples are not pathological by constructing many examples of non-rigid terminal factorial quartic 3-folds with a singular point of type cA_n for $n \geq 2$. It is conjectured that a terminal factorial quartic 3-fold with no worse than cA_1 points is rigid; we address this conjecture in forthcoming work.

Terminal 3-fold hypersurfaces have isolated cDV singularities [Rei87, (3.1)]; the local analytic type of a singular point thus belongs to one of two infinite families, cA_n for $n \geq 1$, or cD_m for $m \geq 4$ or is cE_6, cE_7 or cE_8 . The first step in our study is to bound the local analytic type of singularities that can occur on a terminal factorial quartic 3-fold. We use topology and singularity theory to bound the local analytic type in the cA_n and cD_m cases, and we show the following proposition.

PROPOSITION 1.1. *If $(P \in X)$ is a cA_n (respectively cD_m) point on a terminal factorial quartic hypersurface $X \subset \mathbb{P}^4$, then $n \leq 7$ (respectively $m \leq 8$).*

The methods used to prove Proposition 1.1 do not restrict the local analytic type of points of type cE . In fact, all possible local analytic types of cE points are realised: we give examples of terminal factorial quartic hypersurfaces with isolated singular points of type cE_6, cE_7 or cE_8 (see Example 4.14). As is noted in Remark 3.10, the bound on the local analytic type of cA points is sharp, but we do not believe that the bound is optimal in the cD case.

If X is a terminal factorial quartic 3-fold, the Sarkisov program shows that any birational map $X \dashrightarrow X'$ to a Mori fibre space X'/S' is the composition of finitely many Sarkisov links (see § 2 for definitions and precise statements). Thus, X is non-rigid precisely when there exists a link $X \dashrightarrow X'$, where X'/S' is a Mori fibre space. Such a link is initiated by a morphism $f: Z \rightarrow X$, where Z is terminal and \mathbb{Q} -factorial, and f contracts a divisor to a singular point or to a curve passing through a singular point. In general, little is known about the explicit form of the morphism f . When f contracts a divisor to a cA_n point $(P \in X)$, Kawakita showed that the germ of f is a weighted blowup, and classified possible weights according to the local analytic type of $(P \in X)$ [Kaw01, Kaw02, Kaw03].

For each n with $2 \leq n \leq 7$, we write down the equation of a quartic hypersurface X with a morphism $f: Z \rightarrow X$ that contracts a divisor to a cA_n point and initiates a Sarkisov link. After a suitable embedding of X as a complete intersection in a larger weighted projective space

$\mathbb{P} = \mathbb{P}(1^5, \alpha, \beta)$, we recover f as the restriction of a weighted blowup $\mathcal{F} \rightarrow \mathbb{P}$ whose weights are determined by Kawakita's classification. The variety \mathcal{F} is a toric variety of Picard rank two, and therefore it is possible to write down explicitly all contracting rational maps $\mathcal{F} \dashrightarrow U$ to a projective variety U . We then check that the birational geometry of \mathcal{F} induces a Sarkisov link $X \dashrightarrow X'$, where X'/S' is a Mori fibre space.

To our knowledge, our construction is the first use of Kawakita's classification to write down explicit *global* uniruled 3-fold extractions. We prove the following theorem.

THEOREM 1.2. *If $(P \in X)$ is a singular point of type cA_n on a terminal factorial quartic 3-fold, then $n \leq 7$. There are examples of non-rigid terminal factorial quartic 3-folds with a singular point of type cA_n for $2 \leq n \leq 7$.*

We also give examples of non-rigid terminal factorial quartic 3-folds with cD_4 , cD_5 and cE_6 , cE_7 and cE_8 singular points (Examples 4.12–4.14). We make the following general conjecture, which generalises [CM04, § 1.3 and Theorem 1.6].

CONJECTURE 1.3. Let $X \subset \mathbb{P}^4$ be a terminal factorial quartic hypersurface. Then $\mathcal{P}(X)$ is finite and $\mathcal{P}(X) = \{[X]\}$ precisely when X has no worse than cA_1 singularities. In particular, no terminal factorial quartic hypersurface is rational.

Outline of the paper

Section 2 recalls general results on the Sarkisov program, that is, on the study of birational maps between Mori fibre spaces, in dimension three and on the geography of models of Mori dream spaces. When X is a terminal \mathbb{Q} -factorial Fano 3-fold with $\rho(X) = 1$, a Sarkisov link $X \dashrightarrow X'$ is initiated by a morphism $f : Z \rightarrow X$ that contracts a single divisor. Here, we state precise conditions for f to initiate a Sarkisov link. We urge the reader who is mainly interested in explicit examples and in bounds on singularities to skip this section on a first reading and refer back to it as and when needed.

Section 3 collects results on terminal singularities in dimension three. We concentrate on the case of terminal Gorenstein singularities, which are those that appear on hypersurfaces. We use the existence of smoothings of terminal Gorenstein Fano 3-folds to bound the local analytic type of singularities on a terminal quartic hypersurface. Last, we recall Kawakita's classification of the germs of divisorial contractions with centre at a cA_n point in terms of the local analytic type of that point.

Section 4 presents our examples of non-rigid terminal factorial quartic 3-folds. We consider hypersurfaces $X \subset \mathbb{P}^4$ that can be embedded as general complete intersections of type $(2, 2, 4)$ in a weighted projective space $\mathbb{P} = \mathbb{P}(1^5, 2^2)$. For suitable weighted blowups $F : \mathcal{F} \rightarrow \mathbb{P}$, the restriction $f = F|_Z : Z \rightarrow X$ (where Z is the proper transform of X) is a divisorial contraction with centre at a cA_n point, and the birational geometry of \mathcal{F} induces a Sarkisov link $X \dashrightarrow X'$. We give examples of non-rigid quartic hypersurfaces with a cA_n point for all $2 \leq n \leq 7$, and explain our construction in detail in a few cases. We also give examples of non-rigid terminal factorial quartic hypersurfaces with singular points of type cD and cE .

2. Preliminary results

Throughout this paper, we work with normal projective varieties over \mathbb{C} .

General results in birational geometry

Let X be a normal projective variety and $\mathbf{k} \in \{\mathbb{Z}, \mathbb{Q}, \mathbb{R}\}$. We denote by $\text{WDiv}_{\mathbf{k}}(X)$ the group of Weil \mathbf{k} -divisors, by $\text{Div}_{\mathbf{k}}(X)$ the group of \mathbf{k} -Cartier \mathbf{k} -divisors on X and by $\sim_{\mathbf{k}}$ and \equiv the \mathbf{k} -linear and numerical equivalences of \mathbb{R} -divisors. We write $\text{Pic}(X)_{\mathbf{k}} = \text{Div}_{\mathbf{k}}(X)/\sim_{\mathbf{k}}$ and $N^1(X)_{\mathbf{k}} = \text{Div}_{\mathbf{k}}(X)/\equiv$.

The nef, effective and pseudo-effective cones in $N^1(X)_{\mathbb{R}}$ are denoted by $\text{Nef}(X)$, $\text{Eff}(X)$ and $\overline{\text{Eff}}(X)$. The movable cone $\overline{\text{Mov}}(X)$ is the closure of the cone in $N^1(X)_{\mathbb{R}}$ spanned by the numerical classes of divisors whose stable base locus has codimension at least 2. If $\mathcal{C} \subset N^1(X)_{\mathbb{R}}$ is a cone, we always denote its closure by $\overline{\mathcal{C}}$.

We recall the definitions of models of divisors introduced in [BCHM10].

DEFINITION 2.1. Let Z be a normal projective variety and $D \in \text{Div}_{\mathbb{Q}}(Z)$.

(1) A birational map $f : Z \dashrightarrow X$ is *contracting* if f is proper and f^{-1} contracts no divisor. The map f is *small* if both f and f^{-1} are contracting birational maps.

(2) Let $D \in \text{Div}_{\mathbb{Q}}(Z)$ be a \mathbb{Q} -Cartier divisor and let $f : Z \dashrightarrow X$ be a contracting birational map such that f_*D is \mathbb{Q} -Cartier. The map f is *D-non-positive* if for a resolution $(p, q) : W \rightarrow Z \times X$,


$$p^*D = q^*D' + E,$$

where E is effective and q -exceptional. When $\text{Supp } E$ contains the strict transform of all f -exceptional divisors, f is said to be *D-negative*.

(3) Assume that D is effective. The map f is a *semiample model* of D if f is D -non-positive, X is normal and projective and D' is semiample. If $\varphi : X \rightarrow S$ is the semiample fibration defined by D' , the ample model of D is the composition $\varphi \circ f : Z \dashrightarrow X \rightarrow S$.

Notation 2.2. If $D = K_Z$ is a canonical divisor on Z , we say that a birational contraction is *K-non-positive* (respectively *K-negative*) instead of *K_Z-non-positive* (respectively *K_Z-negative*).

DEFINITION 2.3. (1) A birational contraction $X \xrightarrow{\varphi} X'$ between \mathbb{Q} -factorial varieties is *elementary* if φ is either a morphism whose exceptional locus is a prime divisor on X or a small birational map that fits into a diagram



where f and f' are morphisms and the Picard ranks of X, X' and W satisfy

$$\rho(X) = \rho(X') = \rho(W) + 1.$$

(2) Assume that X is \mathbb{Q} -factorial and let D be an effective \mathbb{Q} -divisor on X . A *D-MMP* on X is a composition of D -non-positive elementary contractions $X \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_n = X_D$, where X_D is a semiample model for D .

Geography of models of Mori dream spaces

We first recall the definition of Mori dream spaces and the properties that will be important in this paper.

DEFINITION 2.4 [HK00]. Let Z be a projective \mathbb{Q} -factorial variety with $\text{Pic}(Z)_{\mathbb{Q}} = N_{\mathbb{Q}}^1(Z)$; Z is a Mori dream space if:

- (i) $\text{Nef}(Z)$ is the affine hull of finitely many semiample line bundles;
- (ii) there are finitely many small birational maps $f_i : Z \dashrightarrow Z_i$ to projective \mathbb{Q} -factorial varieties Z_i satisfying (i) such that $\overline{\text{Mov}} Z = \bigcup f_i^*(\text{Nef } Z_i)$.

When Z is a Mori dream space, we may run the D -MMP for every \mathbb{Q} -divisor D . More precisely, there is a finite decomposition [KKL14, § 5]

$$\overline{\text{Eff}} Z = \bigsqcup_{i=1}^N \mathcal{C}_i, \quad \text{where, for all } i:$$

- (a) $\overline{\mathcal{C}}_i$ is a rational polyhedral cone;
- (b) there is a birational contraction to a \mathbb{Q} -factorial normal projective variety $\varphi_i : Z \dashrightarrow Z_i$ that is the ample model of all $D \in \mathcal{C}_i$ and a semiample model of all $D \in \overline{\mathcal{C}}_i$.

A Mori dream space Z with $\rho(Z) = 2$ always has a 2-ray configuration, which is defined as follows. Let M_1, M_2 be \mathbb{Q} -divisors such that

$$\overline{\text{Mov}} Z = \mathbb{R}_+[M_1] + \mathbb{R}_+[M_2].$$

Denote by $\varphi_1 : Z \dashrightarrow Z_1$ (respectively $\varphi_2 : Z \dashrightarrow Z_2$) the ample model of $M_1 + \varepsilon M_2$ (respectively $\varepsilon M_1 + M_2$) for an arbitrarily small positive rational number ε . Let $f_i : Z_i \rightarrow X_i$ be the ample model of $(\varphi_i)_* M_i$. Then the birational map φ_i is small, f_i is a fibration when $[M_i]$ lies on the boundary of $\overline{\text{Eff}} Z$ and φ_i is a birational map that contracts a single exceptional divisor otherwise. These maps fit in a diagram which we call a 2-ray configuration:

$$\begin{array}{ccc} Z_1 & \xleftarrow{\varphi_1} & Z & \xrightarrow{\varphi_2} & Z_2 \\ f_1 \downarrow & & & & \downarrow f_2 \\ X_1 & & & & X_2 \end{array} \tag{1}$$

When $f : Z \rightarrow X$ is a divisorial contraction, we may assume that φ_1 is the identity map and that X is the ample model of M_1 (i.e. that $f = f_1$).

The Sarkisov program

We recall a few notions on birational maps between end products of the classical MMP.

DEFINITION 2.5. Let X be a terminal \mathbb{Q} -factorial variety and $p : X \rightarrow S$ a morphism with positive-dimensional fibres (so that $\dim S < \dim X$). Then X/S is a Mori fibre space if $p_* \mathcal{O}_X = \mathcal{O}_S$, $-K_X$ is p -ample and $\rho(X) = \rho(S) + 1$.


The classical MMP shows that any uniruled terminal \mathbb{Q} -factorial variety Z is birational to a Mori fibre space, so that $\mathcal{P}(Z) \neq \emptyset$. The Sarkisov program decomposes any birational map between Mori fibre spaces $[X/S], [X'/S'] \in \mathcal{P}(Z)$ into a finite number of Sarkisov links [Cor95, HM13]. Next we recall the definition of Sarkisov links.

DEFINITION 2.6. A *divisorial contraction* $f : Z \rightarrow X$ is a morphism between terminal \mathbb{Q} -factorial varieties such that $-K_Z$ is f -ample, $f_*\mathcal{O}_Z = \mathcal{O}_X$ and $\rho(Z) = \rho(X) + 1$.

We sometimes call f an *extraction* when we study properties of f in terms of its target X .


DEFINITION 2.7. Let X/S and X'/S' be two Mori fibre spaces. A *Sarkisov link* is a birational map $\varphi : X \dashrightarrow X'$ of one of the following types.

(I) A *link of type I* is a commutative diagram



where $Z \rightarrow X$ is a divisorial contraction and $Z \dashrightarrow X'$ a sequence of flips, flops and inverse flips between terminal \mathbb{Q} -factorial varieties.


(II) A *link of type II* is a commutative diagram



where $Z \rightarrow X$ and $Z' \rightarrow X'$ are divisorial contractions and $Z \dashrightarrow Z'$ a sequence of flips, flops and inverse flips between terminal \mathbb{Q} -factorial varieties.

(III) A *link of type III* is the inverse of a link of type I.

(IV) A *link of type IV* is a commutative diagram



where $X \dashrightarrow X'$ is a sequence of flips, flops and inverse flips between terminal \mathbb{Q} -factorial varieties.

DEFINITION 2.8. Let X/S be a Mori fibre space and $f : Z \rightarrow X$ an extraction; f *initiates a link* if it fits into an Sarkisov link.

The following lemma is a criterion for a divisorial extraction to initiate a link. It is of little practical use, but we want to highlight some of the subtleties that arise when proving that a 2-ray configuration is indeed a Sarkisov link.

LEMMA 2.9. *Let X be a terminal \mathbb{Q} -factorial Fano variety with $\rho(X) = 1$ and let $f : Z \rightarrow X$ be an extraction. Then f initiates a link if and only if the following hold:*

- (i) Z is a Mori dream space;
- (ii) if $\psi : Z \dashrightarrow Z'$ is a small birational map and Z' is \mathbb{Q} -factorial, then Z' has terminal singularities;
- (iii) $[-K_Z] \in \text{int}(\overline{\text{Mov}} Z)$.

Proof. Assume that $f : Z \rightarrow X$ is a divisorial contraction that initiates a link. Then, as X is a Fano 3-fold with rational singularities, $h^1(X, \mathcal{O}_X) = 0$ and, since $f_*\mathcal{O}_Z = \mathcal{O}_X$, by the Leray spectral sequence, $h^1(Z, \mathcal{O}_Z) = 0$ and we have the equality $\text{Pic}(Z)_{\mathbb{Q}} = N_{\mathbb{Q}}^1(Z)$.

Since $\rho(Z) = 2$, if f initiates a Sarkisov link, then, following the notation of Definition 2.7, there are two distinct birational contractions $Z \rightarrow X$ and $Z \dashrightarrow X'$ that are compositions of finitely many elementary maps (flips, flops, inverse flips and divisorial contractions) between terminal \mathbb{Q} -factorial varieties. Thus, Z has a 2-ray configuration as above, and Z is automatically a Mori dream space. The chambers of the decomposition of $\overline{\text{Eff}} Z$ are indexed by the divisorial contraction $Z \rightarrow X$ and by the elementary maps that decompose $Z \dashrightarrow X'$.

Furthermore, if X' is Fano, then

$$-K_Z \in \mathbb{R}_+[f^*(-K_X)] + \mathbb{R}_+[(g \circ \psi)^*(-K_{X'})] = \overline{\text{Mov}} Z \tag{2}$$

and the class of $-K_Z$ is in the interior of $\overline{\text{Mov}} Z$ because X and X' have terminal singularities and $Z \rightarrow X$ and $Z' \rightarrow X'$ are not isomorphisms.

If X'/S' is a Mori fibre space with $\dim S' \geq 1$, then $K_{X'} = \psi_*K_Z$ and

$$-K_Z \in \mathbb{R}_+[f^*(-K_X)] + \mathbb{R}_+[\psi^*(-K_{X'} + A_{S'})] \subsetneq \mathbb{R}_+[f^*(-K_X)] + \mathbb{R}_+[\psi^*A_{S'}] = \overline{\text{Mov}} Z, \tag{3}$$

where $A_{S'}$ is the pullback of a suitable ample divisor on S' and the class of $-K_Z$ is in the interior of $\overline{\text{Mov}} Z$ because, as before, X is terminal so that $\mathbb{R}_+[-K_Z] \neq \mathbb{R}_+[f^*(-K_X)]$ and $-K_Z$ is big so that $-K_Z \notin \mathbb{R}_+[\psi^*A_{S'}]$.

We have seen that Z is a Mori dream space so that if D is any movable \mathbb{Q} -divisor, the D -MMP terminates with a \mathbb{Q} -factorial semiample model for D , which we denote Z_D . The small birational map $Z \dashrightarrow Z_D$ factors $Z \dashrightarrow X'$; therefore, $Z \dashrightarrow Z_D$ is the composition of finitely many elementary contractions between terminal \mathbb{Q} -factorial varieties and, in particular, Z_D is terminal. Let $Z \dashrightarrow Z'$ be an arbitrary small birational map and assume that Z' is \mathbb{Q} -factorial. Let D be the proper transform of an ample \mathbb{Q} -Cartier divisor on Z' ; then D is mobile because $Z \dashrightarrow Z'$ is small. By construction, $Z' \simeq \text{Proj } R(Z, D)$ is the ample model of D and, if Z_D is the end product of a D -MMP on Z , then $Z_D \rightarrow Z'$ is a morphism and a small map. Since Z' and Z_D are both \mathbb{Q} -factorial, it follows that they are isomorphic, so that Z' has terminal singularities.

Conversely, if Z is a Picard rank 2 Mori dream space, then

$$\overline{\text{Mov}} Z = \mathbb{R}_+[M_1] + \mathbb{R}_+[M_2] \subset \overline{\text{Eff}}(Z) = \mathbb{R}_+[D_1] + \mathbb{R}_+[D_2]$$

for effective \mathbb{Z} -divisors M_1, M_2, D_1 and D_2 . Since $f : Z \rightarrow X$ is a divisorial extraction, f is the ample model of $M_1 \neq D_1$; note that D_2 need not be distinct from M_2 . When $M_2 \neq D_2$, let Z' be the ample model of $M_2 + \varepsilon M_1$ for an arbitrarily small positive rational number ε and let X' be the ample model of M_2 . Then $Z \dashrightarrow Z'$ is a small birational map and Z' has terminal singularities by assumption (ii). The birational map $Z \dashrightarrow Z'$ is small; hence, we may identify divisors on Z and on Z' and, under this identification, $\overline{\text{Mov}} Z$ is equal to $\overline{\text{Mov}} Z'$, so that $[-K_Z]$ is in the

interior of $\overline{\text{Mov}} Z'$ by assumption (iii). It follows that the morphism $Z' \rightarrow X'$ is K -negative and that X' has terminal singularities. When $M_2 = D_2$, let X' be the ample model of $M_2 + \varepsilon M_1$ for an arbitrarily small positive rational number ε and let S' be the ample model of M_2 . Then X' is terminal by assumption (ii) and the fibration $X' \rightarrow S'$ is K -negative by assumption (iii). Since M_2 is not a big divisor, $\dim S' < \dim X'$ and X'/S' is a Mori fibre space. \square

Remark 2.10. Condition (ii) may only fail when the 2-ray configuration on Z involves an antiflip, because flips and flops of terminal varieties are automatically terminal. For example, consider

$$Z = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-2) \oplus \mathcal{O}_{\mathbb{P}^1}(-2));$$

then Z is a Mori fibre space (a \mathbb{P}^2 -bundle over \mathbb{P}^1) and a Mori dream space on which (i), (iii) hold but (ii) fails. It follows that the 2-ray configuration on Z does not produce a Sarkisov link. Example 4.11 is a Sarkisov link involving an antiflip, and we check that condition (ii) holds directly.

Remark 2.11. Note that condition (ii) always holds when M_2 is of the form $K_Z + \Theta$ for Θ a nef divisor. Indeed, in that case, every $D \in \overline{\text{Mov}} Z \cap \text{Big } Z$ is of the form $K_Z + \Theta'$ for Θ' nef, and every D -negative birational contraction is K -non-positive (see [Kal13, 2.10]). In particular, the ample model $\varphi_D : Z \dashrightarrow Z_D$ is a D -negative birational contraction and hence is K_Z -non-positive, so that for any resolution $(p, q) : U \dashrightarrow Z \times Z_D$, we have

$$p^* K_Z = q^* K_{Z_D} + E,$$

where E is an effective q -exceptional divisor. This implies that for any divisor F over Z_D , the discrepancy $a_F(Z_D) \geq a_F(Z)$, and that Z_D has terminal singularities if Z does.

3. Terminal singularities on quartic 3-folds

In this section, we recall some results on the local analytic description of terminal hypersurface singularities in dimension three and we bound the local analytic type of singularities on terminal factorial quartic hypersurfaces in \mathbb{P}^4 .

3.1 Local analytic description and divisorial extractions

We first recall a few results on isolated hypersurface singularities.

Singularity theory. Let $\mathbb{C}[[x_1, \dots, x_n]]$ be the ring of complex formal power series in n variables and $\mathbb{C}\{x_1, \dots, x_n\} \subset \mathbb{C}[[x_1, \dots, x_n]]$ the subring of formal power series with non-zero radius of convergence. For $F \in \mathbb{C}\{x_1, \dots, x_n\}$, $(F = 0)$ is a germ of a complex analytic set and the *singularity* $(F = 0)$ is the scheme $\text{Spec}_{\mathbb{C}} \mathbb{C}[[x_1, \dots, x_n]]/(F)$.

Let $F \in \mathbb{C}[[x_1, \dots, x_n]]$ be a power series and d a positive integer. We denote by F_d the degree d homogeneous part of F and by $F_{\geq d}$ the series $\sum_{k \geq d} F_k$. The *multiplicity* of F is $\text{mult } F = \min\{d \in \mathbb{N} \mid F_d \neq 0\}$. Two power series F, G are *equivalent* if there exist an automorphism $\varphi = (\varphi_1, \dots, \varphi_n) \in \text{Aut}(\mathbb{C}[[x_1, \dots, x_n]])$ and a unit $u \in (\mathbb{C}[[x_1, \dots, x_n]])^*$ such that

$$u(x_1, \dots, x_n)G(x_1, \dots, x_n) = F(\varphi_1, \dots, \varphi_n).$$

In other words, F and G are equivalent if the singularities $(F = 0)$ and $(G = 0)$ are isomorphic. We denote the equivalence of power series by $F \sim G$.

In what follows, as we are only interested in isolated critical points, by [AGV85, I, §6.3] we may (and will) assume that the power series u, F and G all have non-zero radius of convergence and that $\varphi \in \text{Aut}(\mathbb{C}\{x_1, \dots, x_n\})$.

DEFINITION 3.1. (1) The singularity $(h(x, y, z) = 0)$ is *Du Val* if h is equivalent to one of the standard forms

$$\begin{aligned} A_n & x^2 + y^2 + z^{n+1} = 0 \quad \text{for } n \geq 0, \\ D_m & x^2 + y^2z + z^{m-1} = 0 \quad \text{for } m \geq 4, \\ E_6 & x^2 + y^3 + z^4 = 0, \\ E_7 & x^2 + y^3 + yz^3 = 0, \\ E_8 & x^2 + y^3 + z^5 = 0. \end{aligned}$$

(2) A singularity $(F(x, y, z, t) = 0)$ is *compound Du Val* if F is equivalent to

$$h(x, y, z) + tf(x, y, z, t) = 0, \tag{4}$$

where $(h = 0)$ is a Du Val singularity. The singularity $(F = 0)$ is cA_n (respectively cD_m, cE_k) if, in (4), h is of type A_n (respectively D_m, E_k) with n (respectively m, k) minimal.

THEOREM 3.2 [Kol98, Theorem 2.8]. *Let $F(x, y, z, t) \in \mathbb{C}[[x, y, z, t]]$ define a cA singularity; then one of the following holds:*

- cA_0 : $F \sim x$;
- cA_1 : $F \sim x^2 + y^2 + z^2 + t^m$ for $m \geq 2$;
- $cA_n, n \geq 2$: $F \sim x^2 + y^2 + f_{\geq n+1}(z, t)$, where $f_{\geq n+1}(z, t)$ has no multiple factors.

Remark 3.3. Up to change of coordinates on $\mathbb{P}_{z,t}^1$, we may assume that z^{n+1} appears with coefficient 1 in $f_{\geq n+1}(z, t)$. Since $(F = 0)$ is an isolated singularity, $f_{\geq n+1}(z, t)$ has no repeated factor and contains at least one monomial of the form t^N or zt^{N-1} for $N \geq n+1$. When $N > n+1$, as in [AGV85, § 12],

$$F \sim x^2 + y^2 + z^{n+1} + t^N \quad \text{or} \quad F \sim x^2 + y^2 + z^{n+1} + zt^{N-1}$$

for $N \geq n+1$. Similarly, when $N = n+1$, we have

$$F \sim x^2 + y^2 + f_{n+1}(z, t),$$

where f_{n+1} is a homogeneous form with no repeated factor of degree $n+1$.

THEOREM 3.4 [Kol98, Theorem 2.9]. *Let $F(x, y, z, t) \in \mathbb{C}[[x, y, z, t]]$ define a cD singularity; then one of the following holds:*

- cD_4 : $F \sim x^2 + f_{\geq 3}(y, z, t)$, where f_3 is not divisible by the square of a linear form;
- $cD_{>4}$: $F \sim x^2 + y^2z + ayt^r + h_{\geq s}(z, t)$, where $a \in \mathbb{C}, r \geq 3, s \geq 4$ and $h_s \neq 0$. This has type cD_m for $m = \min\{2r, s+1\}$ if $a \neq 0$ and $m = s+1$ otherwise.

DEFINITION 3.5. The *Milnor number* of the singularity $(F = 0)$ is

$$\mu(F = 0) = \dim_{\mathbb{C}} \mathbb{C}[[x_1, \dots, x_n]]/J_F,$$

where $J_F = ((\partial F/\partial x_1), \dots, (\partial F/\partial x_n))$ is the Jacobian ideal of F , i.e. the ideal generated by the partial derivatives of F .

If $F \sim G$, the Milnor numbers of $(F = 0)$ and of $(G = 0)$ are equal. The Milnor number $\mu(F = 0)$ is finite precisely when $(F = 0)$ is an isolated singularity.

LEMMA 3.6. (1) *If $(F = 0)$ is a cA_n singularity with $n \geq 1$, then $\mu(F = 0) \geq n^2$. If $F \sim xy + z^{n+1} + t^N$ or $F \sim xy + z^{n+1} + zt^{N-1}$ for $N > n+1$, then $\mu(F = 0) \geq n(N-1)$.*

(2) *If $(F = 0)$ is a cD_m singularity with $m > 4$, then $\mu(F = 0) \geq m(m-2)$.*

Proof. If $(F = 0)$ is a cA_n singularity with $n \geq 1$, then $F \sim xy + f(z, t)$, where $f(z, t)$ has multiplicity greater than or equal to $n + 1$. Since

$$\mu(F = 0) = \mu(xy + f(z, t) = 0), \quad \text{we have } \mu(F = 0) = \dim_{\mathbb{C}} \mathbb{C}[[z, t]]/J_f.$$

In all cases, since $\mu(F = 0)$ is finite and f has no repeated factor, $\mathbb{C}[[z, t]]/J_f$ has dimension $\deg(\partial f/\partial z) \cdot \deg(\partial f/\partial t) \geq n(N - 1)$.

Now assume that $(F = 0)$ is a cD_m singularity with $m > 4$. Then, as in [AGV85, I, § 12], if F_0 is the quasi-homogeneous part of F , then $\mu(F = 0) = \mu(F_0 = 0)$, and $\mu(F_0 = 0)$ is given by the formula [AGV85, I, Corollary 3, p. 200]. In the notation of Theorem 3.4, using the methods of [AGV85, I, § 12], we obtain that:

- (a) either $F_0 \sim x^2 + y^2z + yt^r + z^s$ or $F_0 \sim x^2 + y^2z + yt^r + z^{s-1}t$ and $\mu(F_0 = 0) \geq m(m - 2)$;
- (b) or $F_0 \sim x^2 + y^2z + h(z, t)$, where $h(z, t)$ is of the form $z^s + t^N, z^{s-1}t + t^N, z^N + t^s$ or $z^N + zt^{s-1}$ for some $N \geq s$. In all cases, $\mu(F_0 = 0) \geq m(m - 2)$. □

3.2 Bounding the local analytic type of singularities on a terminal factorial Fano Mori fibre space

We bound the local analytic type of singularities on a terminal factorial Fano 3-fold with Picard rank 1.

THEOREM 3.7 [Nam97]. *Let X be a Fano 3-fold with terminal Gorenstein singularities. Then X has a smoothing, i.e. there is a one-parameter flat deformation*

$$\begin{array}{ccc} X & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \{0\} & \hookrightarrow & \Delta \end{array}$$

with \mathcal{X}_t smooth when $t \neq 0$. For all $t \in \Delta$, \mathcal{X}_t is Fano, $\rho(X) = \rho(\mathcal{X}_t)$ and $-K_X^3 = -K_{\mathcal{X}_t}^3$.

The existence of a smoothing $X \hookrightarrow \mathcal{X}$ allows us to bound the Milnor numbers of singularities on X .

THEOREM 3.8 [NS95, Theorem 3.2]. *Let X be a normal projective 3-fold with isolated rational hypersurface singularities such that $H^2(X, \mathcal{O}_X) = (0)$. Denote by $b_i(X)$ the i th Betti number for the singular cohomology of X . If there is a smoothing $X \hookrightarrow \mathcal{X}$, then*

$$b_4(X) - b_2(X) = b_3(X) - b_3(\mathcal{X}_t) + \sum_{P \in \text{Sing } X} \mu(X, P)$$

for $t \in \Delta \setminus \{0\}$, where $\mu(X, P)$ is the Milnor number of (X, P) .

When X is a terminal and factorial Fano 3-fold, the second and fourth Betti numbers of X are equal, that is, $b_2(X) = b_4(X)$, so that

$$\sum_{P \in \text{Sing } X} \mu(X, P) = b_3(\mathcal{X}_t) - b_3(X) \leq b_3(\mathcal{X}_t). \tag{5}$$

The third Betti numbers of non-singular Fano 3-folds with Picard rank 1 are known (see [IP99, Table 12.2]), and we obtain a bound on the sum of Milnor numbers of singular points on X that only depends on $-K_X^3$. When $-K_X^3 = 4$, we have the following.

PROPOSITION 3.9. *Let $X \subset \mathbb{P}^4$ be a terminal factorial quartic hypersurface. If $(P \in X)$ is a singular point of type cA_n , then n is at most 7.*

If $(P \in X)$ is a singular point of type cD_m , then m is at most 8.

Proof. Let $X \hookrightarrow \mathcal{X}$ be a smoothing; then, for all $t \neq 0$, \mathcal{X}_t is a non-singular quartic hypersurface and $b_3(\mathcal{X}_t) = 60$ (see [IP99, Table 12.2]). By Theorem 3.8, we have that $\mu(X, P)$ is bounded above by 60, and the result follows immediately from the lower bounds obtained in Lemma 3.6. \square

Remark 3.10. The bound on the local analytic type of cA points is sharp; Example 4.9 is an example of a terminal factorial quartic hypersurface with a cA_7 singular point.

We do not believe that the bound on the local analytic type of cD points is optimal, as we have not been able to write down examples attaining it. We give examples of terminal factorial quartic hypersurfaces with isolated singular points of type cD_4, cD_5 and cE_6, cE_7 and cE_8 in § 4.3.

Remark 3.11. By the classification of non-singular Fano 3-folds, the bounds on the local analytic type of singularities lying on a terminal factorial Fano 3-fold with Picard rank 1 and anticanonical degree $-K^3 > 4$ are even more restrictive than in the case of a quartic hypersurface.

Remark 3.12. We can use the same methods to bound the local analytic type of singularities on an arbitrary terminal Gorenstein Fano 3-fold X with $\rho = 1$. Indeed, by Theorem 3.8,

$$\sum_{P \in \text{Sing } X} \mu(X, P) \leq b_3(\mathcal{X}_t) + \sigma(X),$$

where $\sigma(X) = b_4(X) - 1$ is the defect of X . The defect of terminal Gorenstein Fano 3-folds with $\rho = 1$ is bounded in [Kal11]. For example, if X is a (not necessarily factorial) terminal quartic hypersurface, then

$$\sum_{P \in \text{Sing } X} \mu(X, P) \leq 75,$$

so that if $(P \in X)$ is a cA_n point, then $n \leq 8$.

Our main interest in this article is in non-rigid quartic 3-folds; in particular, we have not tried to write down examples of non-factorial quartic hypersurfaces with a singular point of type cA_8 . We believe that such an example would be found with extra work.

3.3 Divisorial extractions with centre at a cA_n point

Kawakita classified the germs of divisorial extractions $f : Z \rightarrow X$ with centre at a cA_n point. We recall this classification here, as we will use it in § 4.

THEOREM 3.13 [Kaw02, Kaw03]. *Let $(P \in X)$ be a cA_n point and $f : Z \rightarrow X$ a divisorial extraction with $f(\text{Exc } f) = \{P\}$. Then one of the following holds.*

- (1) *General type: under a suitable local analytic identification,*

$$(P \in X) \simeq 0 \in \{xy + g(z, t) = 0\} \subset \mathbb{C}^4$$

and f is the blowup of $\mathbb{C}_{x,y,z,t}^4$ with weights $(r_1, a(n+1) - r_1, a, 1)$, where a is the discrepancy of f , a and r_1 are coprime integers and

$$g(z, t) = z^{n+1} + g_{\geq a(n+1)}(z, t)$$

has weighted degree $a(n + 1)$. In particular,

$$E = \text{Exc } f \simeq \{xy + g_{a(n+1)}(z, t) = 0\} \subseteq \mathbb{P}(r_1 a(n + 1) - r_1, a, 1),$$

and, if we denote $r_2 = a(n + 1) - r_1$, then we have $aE^3 = 1/r_1 + 1/r_2$.

(2) *Exceptional type, $n = 1$: under a suitable local analytic identification,*

$$(P \in X) \simeq 0 \in \{xy + z^2 + t^3 = 0\} \subset \mathbb{C}^4$$

and f is the blowup with weights $(1, 5, 3, 2)$; f has discrepancy $a = 4$.

(3) *Exceptional type, $n = 2$: under a suitable local analytic identification,*

$$(P \in X) \simeq 0 \in \{xy + z^3 + g_{\geq 4}(z, t) = 0\} \subset \mathbb{C}^4,$$

the discrepancy of f is 3 and Z has exactly one non-Gorenstein point

$$(Q \in Z) \simeq \{0\} \in \{x^2 + y^2 + z^2 + t^2 = 0\} \subset \mathbb{C}^4 / \frac{1}{4}(1, 3, 3, 2).$$

Remark 3.14. Let $X \subset \mathbb{P}^4$ be a terminal factorial quartic hypersurface. Then the discrepancies of possible divisorial extractions $f : Z \rightarrow X$ with centre at a cA_n point $(P \in X)$ can be bounded in the same way as in Proposition 3.9. Indeed, by Lemma 3.6, if there is an extraction of general type in Theorem 3.13, the Milnor number $\mu(X, P)$ satisfies

$$\mu(X, P) \geq n(a(n + 1) - 1),$$

but, by Theorem 3.8, $\mu(X, P) \leq 60$. It follows that the discrepancy of f satisfies

n	6, 7	5	4	3	2
a	1	≤ 2	≤ 3	≤ 5	≤ 11

4. Examples of non-rigid terminal quartics

In this section, we present examples of non-rigid terminal factorial quartic hypersurfaces $X \subset \mathbb{P}^4$. Each of these examples has a Sarkisov link initiated by an extraction $f : Z \rightarrow X$ that contracts a divisor to a singular point. In most of our examples, the singular point is of type cA_n , so that the germ of f is a weighted blowup as in Kawakita’s classification (Theorem 3.13). Our examples are obtained by *globalising* these germs: we can write down an explicit description of f in projective coordinates.

Let $X \subset \mathbb{P}^4$ be a terminal factorial quartic hypersurface and assume that $P = (1:0:0:0) \in X$ is a cA_n point. Up to projective change of coordinates, the equation of X can be written

$$X = \{\varphi_4(x_0, \dots, x_4) = x_0^2 x_1 x_2 + x_0 \psi_3(x_1, \dots, x_4) + \theta_4(x_1, \dots, x_4) = 0\},$$

where φ_4 is a homogeneous polynomial of degree four in the variables x_0, \dots, x_4 and ψ_3 and θ_4 are homogeneous polynomials in the variables x_1, \dots, x_4 .

The first step in our construction is to look for examples of hypersurfaces $X \subset \mathbb{P}^4$ that can be embedded in a larger weighted projective space \mathbb{P} in such a way that the restriction of a suitable weighted blowup $F : \mathcal{F} \rightarrow \mathbb{P}$ is a divisorial contraction $f : Z \rightarrow X$. We take X to be a complete intersection of the form

$$X = \begin{cases} \alpha = x_0 x_1 + q(x_3, x_4) \\ \beta = x_0 x_2 + q'(x_3, x_4) \\ \alpha \beta + \varphi'_4(x_0, \dots, x_4) = 0 \end{cases} \subset \mathbb{P}(1^5, 2^2) = \mathbb{P}$$

for homogeneous forms α, β of degree two. The equation of the hypersurface $X \subset \mathbb{P}^4$ is recovered by substituting α, β in the third equation.

Explicitly, we want the germ of $f : Z \rightarrow X$ to be a weighted blowup of general type in the classification of Theorem 3.13. This means that up to local analytic identification, denoting by a the discrepancy of f , we have

$$(P \in X) \simeq 0 \in \{xy + g(z, t) = 0\} \subset \mathbb{C}^4, \quad \text{where } g(z, t) = z^{n+1} + g_{\geq a(n+1)}(z, t) \tag{6}$$

and g has weighted degree $a(n + 1)$.

We choose φ_4 so that for suitable α, β , setting $x_0 = 1$ in the third equation gives

$$\alpha\beta + g(x_3, x_4) + (\text{higher weighted order terms}) = 0.$$

In other words, the restrictions of α, β to $\{x_0 = 1\}$ define the local analytic isomorphisms that bring the equation $X \cap \{x_0 = 1\}$ into the desired form (6). The divisorial contractions $f : Z \rightarrow X$ we construct are restrictions of weighted blowups $F : \mathcal{F} \rightarrow \mathbb{P}(1^5, 2^2)$, where the weights assigned to the variables α, β, x_3, x_4 are as in Theorem 3.13.

The second step in our construction is to show that some of these divisorial contractions initiate Sarkisov links. Since \mathcal{F} is a Mori dream space (it is toric), it has a 2-ray configuration as in (1). We check directly that the 2-ray configuration

$$\begin{array}{ccc} Z \subset \mathcal{F} & \xrightarrow{\Phi} & \mathcal{F}^+ \\ \downarrow F & & \downarrow F^+ \\ X \subset \mathbb{P} & & \mathbb{P}^+ \end{array} \quad \text{restricts to a Sarkisov link} \quad \begin{array}{cc} Z & \xrightarrow{\quad} & Z^+ \\ \downarrow & & \downarrow \\ X & & X^+ \end{array} \tag{7}$$

or that we can find another embedding $Z \subset \mathcal{F}'$ via unprojection such that the 2-ray configuration on \mathcal{F}' restricts to a Sarkisov link. Note that we do not make any assumption on the singularities of \mathcal{F} ; in particular, \mathcal{F} needs not be terminal and \mathbb{Q} -factorial. We check the following.

(1) The map $\Phi|_Z$ is an isomorphism in codimension 1 and Z^+ is terminal. In our examples, $\Phi|_Z$ is the composition of finitely many elementary maps

$$Z = Z_0 \xrightarrow{\varphi_0} Z_1 \xrightarrow{\varphi_1} \dots \xrightarrow{\varphi_n} Z_{n+1} = Z^+$$

that are isomorphisms or antiflips, flops and flips (in that order). If $Z_i \xrightarrow{\varphi_i} Z_{i+1}$ is K -non-positive (e.g. a flip or a flop) and Z_i is terminal and \mathbb{Q} -factorial, then so is Z_{i+1} . We need to check directly that antiflips preserve the terminal condition; we do this by identifying the antiflips as inverses of flips appearing in [Bro99].

(2) The restriction of $\mathcal{F}^+ \rightarrow \mathbb{P}^+$ is the contraction of a K -negative extremal ray $Z^+ \rightarrow X^+$.

We always denote by P the point $(1:0:0:0) \in \mathbb{P}^4$.

Remark 4.1. We do not give details of how to check that each of our examples is factorial. This relies on ad hoc methods and the general scheme is as follows. As $X \subset \mathbb{P}$ is a Picard rank 1 hypersurface, X is (\mathbb{Q} -)factorial precisely when the rank of the divisor class group satisfies

$$\text{rkCl}(X) = \rho(X) = \text{rkCl}(\mathbb{P}).$$

Since $\text{Sing } X$ has codimension 3, we have an isomorphism

$$\text{Cl}(X) \simeq \text{Cl}(X \setminus \text{Sing}(X)).$$

Let $\pi : \tilde{\mathbb{P}} \rightarrow \mathbb{P}$ be a map from a smooth variety $\tilde{\mathbb{P}}$ that restricts to a resolution $\tilde{X} \rightarrow X$ with exceptional locus E_X . We have natural isomorphisms

$$\text{Cl}(X) \simeq \text{Cl}(X \setminus \text{Sing}(X)) \simeq \text{Pic}(X \setminus \text{Sing} X) \simeq \text{Pic}(\tilde{X} \setminus E_X)$$

and similarly

$$\text{Cl}(\mathbb{P}) \simeq \text{Pic}(\tilde{\mathbb{P}} \setminus \text{Exc } \pi).$$

Now X is factorial precisely when $\text{Pic}(\tilde{\mathbb{P}} \setminus \text{Exc } \pi) \simeq \text{Pic}(\tilde{X} \setminus E_{\tilde{X}})$. As the classical Grothendieck–Lefschetz theorem guarantees that $\text{Pic}(\tilde{X}) \simeq \text{Pic}(\tilde{\mathbb{P}})$, the result follows by comparison of the kernels of the surjective maps

$$r_1 : \text{Pic}(\tilde{\mathbb{P}}) \rightarrow \text{Pic}(\tilde{\mathbb{P}} \setminus \text{Exc } \pi) \quad \text{and} \quad r_2 : \text{Pic}(\tilde{X}) \rightarrow \text{Pic}(\tilde{X} \setminus E_X).$$

These kernels are isomorphic to the free abelian groups on irreducible divisorial components of $\text{Exc } \pi$ and of E_X , respectively, and can be worked out in each case.

Note that we have a 2-ray configuration (7) and therefore all varieties in (7) are \mathbb{Q} -factorial if and only if one of them is. These varieties are complete intersections in $\mathbb{P}, \mathcal{F}, \mathcal{F}^+$ and \mathbb{P}^+ and their Picard ranks are 1, 2, 2 and 1 by construction, and the method above can be applied to determine the divisor class group of any of them. In some cases, it can be easier to determine \mathbb{Q} -factoriality of another variety in (7) than X (one needs to keep track of Gorenstein indices).

4.1 A family of examples

In this section, we study some links initiated by divisorial contractions with discrepancy 1. We consider the family of quartic hypersurfaces

$$X_4^{i,j} = \{(x_0x_1 - x_3^2)^2 - (x_0x_2 - x_4^2)^2 + x_0^{4-i-j}(x_1^i x_3^j + x_2^i x_4^j) + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4,$$

where (i, j) satisfy $3 \leq i + j \leq 4$ and $(i, j) \neq (4, 0)$. Then $(P \in X^{i,j})$ is a cA_n point with

TABLE 1. Non-rigid quartic hypersurfaces $X^{i,j} \subset \mathbb{P}^4$ with a cA_n point.

(i, j)	(0, 4)	(1, 3)	(2, 2)	(3, 1)	(0, 3)	(1, 2)	(2, 1)	(3, 0)
n	3	4	5	6	2	3	4	5

Indeed, the restriction of $X^{i,j}$ to the affine chart $U_0 = \{x_0 = 1\}$ is

$$\{F(x, y, z, t) = (x - z^2)^2 - (y - t^2)^2 + x^i z^j + y^i t^j + x^4 + y^4 = 0\} \subset \mathbb{C}^4$$

and the singularity $(F = 0)$ is equivalent to $(G = 0)$, where G defines a cA_n singularity. To see this, let $\underline{\varphi} = (\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in \text{Aut } \mathbb{C}[[x, y, z, t]]$ be defined by

$$\begin{cases} \varphi_1(x, y, z, t) = x + y - (z^2 + t^2), \\ \varphi_2(x, y, z, t) = x - y - (z^2 - t^2), \\ \varphi_3(x, y, z, t) = z, \\ \varphi_4(x, y, z, t) = t \end{cases}$$

and check that $F(x, y, z, t) = G(\varphi_1, \varphi_2, \varphi_3, \varphi_4)$, where

$$G(x, y, z, t) = xy + z^{n+1} + t^{n+1} + G'(x, y, z, t), \tag{8}$$

where n is as in Table 1, and $G'(x, y, z, t)$ is a polynomial of degree strictly greater than one with respect to the weights

$$\underline{w} = (1/2, 1/2, 1/n + 1, 1/n + 1).$$

Since G is a semi-quasi-homogeneous polynomial of degree one with respect to \underline{w} , and since no element of its Jacobian algebra has degree strictly greater than one, by [AGV85, I, §12], $G \sim xy + z^{n+1} + t^{n+1}$ and $(P \in X)$ is a cA_n point. The quartic hypersurfaces $X^{i,j}$ are terminal ($\text{Sing } X^{i,j} = \{P\}$) and factorial and hence are Mori fibre spaces.

Remark 4.2. Taking $(i, j) = (4, 0)$ gives a terminal quartic hypersurface with a cA_7 point

$$X = X^{4,0} = \{(x_0x_1 - x_3^2)^2 - (x_0x_2 - x_4^2)^2 + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4,$$

but $X = \{f = 0\}$ is not factorial as $f = q_1q'_1 + q_2q'_2$, where q_i, q'_i are quadric polynomials.

Notation 4.3. We embed $X^{i,j}$ as a complete intersection in a scroll whose coordinates are those of \mathbb{P}^4 on the one hand, and projectivisations of the (non-linear) components of $\varphi \in \text{Aut } \mathbb{C}[[x, y, z, t]]$, the automorphism we used to transform the equation of $X \cap U_0$ into G . Here, $\varphi_3(x, y, z, t) = z$ and $\varphi_4(x, y, z, t) = t$, so we only need to introduce the coordinates

$$\begin{cases} \alpha = x_0^2\varphi_1\left(\frac{x_1}{x_0}, \frac{x_2}{x_0}, \frac{x_3}{x_0}, \frac{x_4}{x_0}\right) = x_0(x_1 + x_2) - (x_3^2 + x_4^2), \\ \beta = x_0^2\varphi_2\left(\frac{x_1}{x_0}, \frac{x_2}{x_0}, \frac{x_3}{x_0}, \frac{x_4}{x_0}\right) = x_0(x_1 - x_2) - (x_3^2 - x_4^2). \end{cases}$$

By construction, X is the complete intersection:

$$X = \begin{cases} \alpha + \beta = x_0x_1 - x_3^2 \\ \alpha - \beta = x_0x_2 - x_4^2 \\ \alpha\beta + x_0^{4-i-j}(x_1^ix_3^j + x_2^ix_4^j) + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathbb{P}(1^5, 2^2).$$

We set $x_0 = 1$ to study X in the neighbourhood of P , and we use the first two equations to eliminate x_1, x_2 , so that $X \cap \{x_0 = 1\}$ is the hypersurface:

$$\{\alpha\beta + x_3^{n+1} + x_4^{n+1} + (\text{higher order terms}) = 0\} \subset \mathbb{C}_{x_3, x_4, \alpha, \beta}^4.$$

We now construct Sarkisov links initiated by divisorial contractions $f : Z \rightarrow X$ that are restrictions of weighted blowups $F : \mathcal{F} \rightarrow \mathbb{P}(1^5, 2^2)$ with centre at P . We assume that the local analytic coordinates for which the germ of f has the description given in Theorem 3.13 are $(\alpha, \beta, x_3, x_4)$. We use the notation of [Ahm14, §3.1]; \mathcal{F} is the Picard rank 2 toric variety $\text{TV}(I, A)$, where

$$I = (u, x_0) \cap (x_1, \dots, x_4, \alpha, \beta)$$

is the irrelevant ideal in the the coordinates ring $\mathbb{C}[u, x_0, \dots, x_4, \alpha, \beta]$ and, for suitable non-zero integers r_1, r_2, w_1, w_2 , the 2×8 matrix A below (the numerical rows) defines the action of $\mathbb{C}^* \times \mathbb{C}^*$ with weights

$$A = \begin{pmatrix} u & x_0 & x_1 & x_2 & x_3 & x_4 & \alpha & \beta \\ 1 & 0 & -w_1 & -w_2 & -a & -1 & -r_1 & -r_2 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 & 2 \end{pmatrix}. \tag{9}$$

For example, $(\mu, \lambda) \in \mathbb{C}^* \times \mathbb{C}^*$ acts on the variable α by

$$((\mu, \lambda), \alpha) \mapsto \mu^{-r_1}\lambda^2\alpha.$$

With the grading defined in (9), taking the ample model of a divisor whose class is in $\binom{0}{k}$,

$$\mathcal{F} \rightarrow \text{Proj} \bigoplus_{k \in \mathbb{N}} H^0\left(\mathcal{F}, \mathcal{O}_{\mathcal{F}}\binom{0}{k}\right)$$

is the morphism given by

$$(u, x_0, x_1, x_2, x_3, x_4, \alpha, \beta) \mapsto (x_0, u^{\omega_1} x_1, u^{\omega_2} x_2, u^a x_3, u x_4, u^{r_1} \alpha, u^{r_2} \beta).$$

This is precisely the weighted blowup $F : \mathcal{F} \rightarrow \mathbb{P}(1^5, 2^2)$ we are after.

In what follows, we always denote by L the pullback of $\mathcal{O}_{\mathbb{P}}(1)$, so that $L \in \binom{0}{1}$, and by $E = \binom{1}{0}$ the F -exceptional divisor L .

The form of the expression (8) imposes that the discrepancy of f is 1. We have $-K_Z = f^*(-K_X) - E$, where $E = \text{Exc } f$, so that $a = 1$ and $r_1 + r_2 = n + 1$, where $(n, (i, j))$ are as in Table 1. Set $r_1 = r$, $r_2 = n + 1 - r$ and assume as we may that $r \leq n + 1 - r$. We have

$$K_{\mathcal{F}} \in \binom{(n+1) + w_1 + w_2 + 1}{-9}.$$

LEMMA 4.4. *The weighted blowup $F : \mathcal{F} \rightarrow \mathbb{P}$ restricts to a divisorial contraction $f : Z \rightarrow X$ with discrepancy 1 if the weights r, w_1, w_2 of F in (9) are one of:*

- (1) $i = 0$ ($n = 2$ or 3) and $w_1 = w_2 = 1, r = 1$ or 2 ;
- (2) $w_1 = w_2 = 2$ and $r \geq 2$ ($n \geq 3$).

Proof. The 3-fold Z is a general complete intersection of three hypersurfaces of degrees determined by r, w_1 and w_2 . Once these degrees are known, we use adjunction to write the anticanonical class of Z . Since $-K_X \in \mathcal{O}(1)$ and $a = 1$, $-K_Z \sim L - E$ and this yields the possible values for r, w_1 and w_2 . For example, if $w_1, w_2, r \geq 2$, Z is the proper transform of X under F and it is a complete intersection:

$$\begin{cases} u^{r-2}\alpha = x_0(u^{w_1-2}x_1 + u^{w_2-2}x_2) - (x_3^2 + x_4^2), \\ u^{n-1-r}\beta = x_0(u^{w_1-2}x_1 - u^{w_2-2}x_2) - (x_3^2 - x_4^2), \\ \alpha\beta + u^{iw_1+j-2}x_1^i x_3^j + u^{iw_2+j-2}x_2^i x_4^j + u^{4w_1-n-1}x_1^4 + u^{4w_2-n-1}x_2^4 = 0, \end{cases}$$

so that

$$Z = \binom{-2}{2} \cap \binom{-2}{2} \cap \binom{-(n+1)}{4} \quad \text{and} \quad -K_Z \in \binom{3 - (w_1 + w_2)}{1}$$

and this forces $w_1 = w_2 = 2$. Other cases are entirely similar. □

We check that, with one exception labelled ‘bad link’, for all weights in Lemma 4.4, the 2-ray configuration on \mathcal{F} induces a 2-ray configuration on $TV(I, A)$ that induces a Sarkisov link for $X^{i,j}$. In the case of the bad link, the second birational contraction $g : Z \dashrightarrow Y$ has relatively trivial canonical class, so that Y is not terminal and the 2-ray configuration is not a Sarkisov link. Table 2 gives details of the construction of each Sarkisov link.

THEOREM 4.5. *There are examples of non-rigid terminal factorial quartic hypersurfaces in \mathbb{P}^4 with an isolated cA_n point for all $2 \leq n \leq 6$. For each combination $((i, j), n)$ in Table 1, $X^{i,j} \subset \mathbb{P}^4$ is a non-rigid terminal factorial quartic 3-fold with an isolated singular point $(P \in X)$ of type cA_n . Table 2 lists Sarkisov links initiated by a divisorial contraction $f : Z \rightarrow X^{i,j}$ with centre at $(P \in X)$ and discrepancy 1. Each entry specifies the weights of $(\alpha, \beta, x_3, x_4)$ for the germ of f in the notation of Theorem 3.13 and gives the explicit construction of the link.*

TABLE 2. Sarkisov links for the hypersurfaces $X^{i,j}$ of Table 1.

$p \in X$	Blowup weights of f	Decomposition of $\Phi _Z$	Y/T
cA_2	(2, 1, 1, 1)	12 flops	$Y_{3,4} \subset \mathbb{P}(1, 1, 1, 1, 2, 2)$
cA_3	(3, 1, 1, 1) for $X^{1,2}$ (3, 1, 1, 1) for $X^{0,4}$	Eight flops	$X^{1,2} \quad (\star)$ Bad link
cA_4	(2, 2, 1, 1)	Four flops	dP_2 fibration over \mathbb{P}^1
cA_5	(3, 2, 1, 1)	Two flops then flip (3, 1, 1, -1, -1; 2)	dP_3 fibration over \mathbb{P}^1
cA_5	(4, 2, 1, 1)	Two flops	Conic bundle over $\mathbb{P}(1, 1, 2)$
cA_5	(3, 3, 1, 1)	\cong then two flips (3, 1, 1, -1, -1; 2)	dP_4 fibration over \mathbb{P}^1
cA_6	(5, 2, 1, 1)	Two flops	$Y_{6,6} \subset \mathbb{P}(1, 1, 2, 3, 3, 5)$
cA_6	(4, 3, 1, 1)	\cong then flip (3, 1, 1, -1, -1; 2)	Conic bundle over $\mathbb{P}(1, 1, 2)$

Proof. Each case is treated individually. To illustrate the computations involved, we treat the cA_6 case in detail. We then say a few words about the cA_2 case, where we recover the example of a non-rigid quartic constructed in [CM04].

Remark 4.6. The case labelled as (\star) in Table 2 is a quadratic involution in the language of [CPR00]. In particular, the link does not produce a new Mori fibre space; it is just a birational selfmap of $X^{1,2}$ that is not an isomorphism.

Non-rigid quartic with a cA_6 singular point. Consider the terminal, factorial quartic hypersurface

$$X = \{(x_0x_1 - x_3^2)^2 - (x_0x_2 - x_4^2)^2 + x_1^3x_3 + x_2^3x_4 + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4.$$

As above, we embed X as a complete intersection in $\mathbb{P} = \mathbb{P}(1^5, 2^2)$, where the variables of weight 2 are α, β :

$$X = \begin{cases} \alpha + \beta = x_0x_1 - x_3^2 \\ \alpha - \beta = x_0x_2 - x_4^2 \\ \alpha\beta + x_1^3x_3 + x_2^3x_4 + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathbb{P}(1^5, 2^2).$$

We construct Sarkisov links initiated by a divisorial contraction $f : Z \rightarrow X$, which is the restriction of a weighted blowup $F : \mathcal{F} \rightarrow \mathbb{P}$. We assume that the weights assigned to the variables $(\alpha, \beta, x_3, x_4)$ are those in Theorem 3.13. By Theorem 3.13 and Lemma 4.4, \mathcal{F} is the Picard rank 2 toric variety $TV(I, A)$, where A is of the form

$$\begin{pmatrix} u & x_0 & x_1 & x_2 & x_3 & x_4 & \alpha & \beta \\ 1 & 0 & -2 & -2 & -1 & -1 & -r & -(7-r) \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 & 2 \end{pmatrix} \quad \text{for } r = 2 \text{ or } 3$$

and Z is a complete intersection of the form $(2L - 2E, 2L - 2E, 4L - 7E)$ in \mathcal{F} .

Case 1. The germ of f is a blowup with weights (5, 2, 1, 1).

We re-order the coordinates of \mathbb{P} and write the action A as follows:

$$\begin{pmatrix} u & x_0 & x_3 & x_4 & \beta & x_1 & x_2 & \alpha \\ 1 & 0 & -1 & -1 & -2 & -2 & -2 & -5 \\ 0 & 1 & 1 & 1 & 2 & 1 & 1 & 2 \end{pmatrix}.$$

In this case, Z is the complete intersection:

$$Z = \begin{cases} u^3\alpha + \beta = x_0x_1 - x_3^2 \\ u^3\alpha - \beta = x_0x_2 - x_4^2 \\ \alpha\beta + x_1^3x_3 + x_2^3x_4 + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathcal{F}.$$


Taking the difference of the first two equations shows that the variable β is redundant and

$$Z = \begin{cases} u^3\alpha = x_0(x_1 + x_2) - (x_3^2 + x_4^2) \\ \alpha(x_0(x_1 - x_2) - (x_3^2 - x_4^2)) + x_1^3x_3 + x_2^3x_4 + u(x_1^4 + x_2^4) = 0 \end{cases} \subset \mathcal{F},$$

where we now denote by \mathcal{F} the toric variety $TV(I, A)$ for

$$A = \begin{pmatrix} u & x_0 & x_3 & x_4 & x_1 & x_2 & \alpha \\ 1 & 0 & -1 & -1 & -2 & -2 & -5 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 \end{pmatrix}.$$

The 2-ray configuration on \mathcal{F} is



where Φ is a small map that is the ample model for $L - (1 + \varepsilon)E$ and $G : \mathcal{F}^+ \rightarrow \mathbb{P}^+$ is a divisorial contraction, where $\mathbb{P}^+ \simeq \text{Proj}(\mathcal{F}, n(L - 2E))$ for suitable $n \gg 1$.

The only pure monomials in u, x_0, x_3, x_4 in the equation of Z are in the expression $x_3^2 + x_4^2$ in the first equation, so that the restriction of Φ_1 to Z is a flop in two lines (a copy of \mathbb{P}^1_{u,x_0} above each of the two points $\{x_3^2 + x_4^2 = 0\} \subset \mathbb{P}^1_{x_3,x_4}$) and, since α does not divide the equations of Z , G does restrict to a divisorial contraction.

To determine \mathbb{P}^+ , we find a suitable change of basis in which to express the action A . In practice, we look for a matrix M in $Sl_2(\mathbb{Z})$ such that

$$M \cdot \begin{pmatrix} -5 \\ 2 \end{pmatrix} = \begin{pmatrix} * \\ 0 \end{pmatrix}.$$

The matrix $M = \begin{pmatrix} 1 & 3 \\ -2 & -5 \end{pmatrix}$ transforms the action A into

$$\begin{pmatrix} u & x_0 & x_3 & x_4 & x_1 & x_2 & \alpha \\ 1 & 3 & 2 & 2 & 1 & 1 & 1 \\ -2 & -5 & -3 & -3 & -1 & -1 & 0 \end{pmatrix},$$

so that $\mathbb{P}^+ = \mathbb{P}(1^2, 2, 3^2, 5)$, with coordinates x_1, x_2 (degree one), $u\alpha$ (degree two), $x_3\alpha, x_4\alpha$ (degree three) and $x_0\alpha^2$ (degree five). Writing the equations of the proper transform of Z^+ shows that $Y \subset \mathbb{P}^+$ is

$$\begin{cases} (u\alpha)^3 = (x_0\alpha^2)(x_1 + x_2) - ((x_3\alpha)^2 + (x_4\alpha)^2), \\ (x_0\alpha^2)(x_1 - x_2) - ((x_3\alpha)^2 - (x_4\alpha)^2) + x_1^3(x_3\alpha) + x_2^3(x_4\alpha) + (u\alpha)(x_1^4 + x_2^4) = 0, \end{cases}$$

so that Y is the complete intersection of two hypersurfaces of degree six in \mathbb{P}^+ . Since $x_3\alpha$ is a section of $g^*\mathcal{O}_{\mathbb{P}^+}(3)$, the 3-fold Y has Fano index 3 and, by construction of f , its basket consists of a single $[5, 2]$ singular point at P_α ; Y is the Fano variety number 41 920 in [GRD].

Case 2. The germ of f is a blowup with weights $(4, 3, 1, 1)$.


We re-order the coordinates of \mathbb{P} and write the action A as

$$\begin{pmatrix} u & x_0 & x_3 & x_4 & \beta & x_1 & x_2 & \alpha \\ 1 & 0 & -1 & -1 & -3 & -2 & -2 & -4 \\ 0 & 1 & 1 & 1 & 2 & 1 & 1 & 2 \end{pmatrix}.$$

The 3-fold Z is the complete intersection:

$$Z = \begin{cases} u^2\alpha = x_0(x_1 + x_2) - (x_3^2 + x_4^2) \\ u\beta = x_0(x_1 - x_2) - (x_3^2 - x_4^2) \\ \alpha\beta + x_1^3x_3 + x_2^3x_4 + u(x_1^4 + x_2^4) = 0 \end{cases} \subset \mathcal{F}.$$

The 2-ray configuration on \mathcal{F} is



where Φ_1 is a small map and \mathcal{F}_1 the ample model of $L - (1 + \varepsilon)E$, Φ_2 is a small map and \mathcal{F}_2 the ample model of $2L - (3 + \varepsilon)E$ and G is a divisorial contraction to $\mathbb{P}^+ \simeq \text{Proj}(\mathcal{F}, n(L - 2E))$ for suitable $n \gg 1$.

Since x_3^2, x_4^2 appear in two of the equations defining Z , Z does not contain any curve contracted by Φ_1 and $\Phi_{1|Z}$ is an isomorphism. We still denote by Z its image under Φ_1 .

We show that $\Phi_{2|Z}$ is a flip. We study the behaviour of Z near $P_\beta = (0:0:0:0:1:0:0:0)$. The restriction of Z to $U_\beta = \{\beta = 1\}$ is a hypersurface: we may use the second and third equations to eliminate u and α , so that $Z \cap U_\beta$ is the hypersurface defined by the first equation. As above, under the change of coordinates associated to

$$\begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix} \in \text{SL}_2(\mathbb{Z}),$$

the action becomes

$$\begin{pmatrix} u & x_0 & x_3 & x_4 & \beta & x_1 & x_2 & \alpha \\ 1 & 1 & 0 & 0 & -1 & -1 & -1 & -2 \\ 2 & 3 & 1 & 1 & 0 & -1 & -1 & -2 \end{pmatrix},$$

so that once the variables u, α and the second and third equations defining Z have been eliminated, we are left with a flip of the hypersurface defined by the first equation of Z in $\mathbb{C}_{x_0, x_3, x_4, x_1, x_2}$, which is

$$x_0(x_1 + x_2) + x_3^2 + x_4^2 + \dots = 0,$$

that is, in the notation of [Bro99], of the form $(3, 1, 1, -1, -1; 2)$. There are thus two flipped curves and, while Z has a $cA/3$ singularity over P_β , Z_2 is Gorenstein over P_β . The map G is a fibration over $\mathbb{P}(\alpha, x_1, x_2) = \mathbb{P}(1, 1, 2)$, and the equations of Z show that the restriction $Z_2 \rightarrow \mathbb{P}(1, 1, 2)$ is a conic bundle.

Non-rigid quartic with a cA_2 point. Consider the terminal, factorial quartic hypersurface

$$X = \{(x_0x_1 - x_3^2)^2 - (x_0x_2 - x_4^2)^2 + x_0(x_3^3 + x_4^3) + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4.$$

By Lemma 4.4, after re-ordering the coordinates of \mathbb{P} , \mathcal{F} can only be the Picard rank 2 toric variety $TV(I, A)$, where

$$\begin{pmatrix} u & x_0 & \alpha & x_1 & x_2 & x_3 & x_4 & \beta \\ 1 & 0 & -1 & -1 & -1 & -1 & -1 & -2 \\ 0 & 1 & 2 & 1 & 1 & 1 & 1 & 2 \end{pmatrix},$$

and Z is given by the equations

$$\begin{cases} \alpha = x_0(x_1 + x_2) - u(x_3^2 + x_4^2), \\ u\beta = x_0(x_1 - x_2) - u(x_3^2 - x_4^2), \\ \alpha\beta + x_0(x_3^3 + x_4^3) + u(x_1^4 + x_2^4) = 0. \end{cases}$$

The first equation shows that we may eliminate the variable α . The first step of the 2-ray configuration of \mathcal{F} is a small map but introduces a new divisor on Z and hence is not a step in the 2-ray game of Z . Note that the second equation is in the ideal (u, x_0) ; we will re-embed Z by unprojection into a toric variety of Picard rank 2 whose 2-ray configuration restricts to suitable maps on Z . To do so, we introduce an unprojection variable s and replace the second equation with

$$\begin{cases} sx_0 = \beta + x_3^2 - x_4^2, \\ su = x_1 - x_2. \end{cases}$$

From the two equations that define s , we eliminate the variables β and x_1 so that the (isomorphic) image of Z under the unprojection is

$$\begin{aligned} & \{(x_0(2x_1 - su) - u(x_3^2 + x_4^2))(sx_0 - x_3^2 + x_4^2) + x_0(x_3^3 + x_4^3) + u(x_1^4 + (x_1 - su)^4) = 0\} \\ & \subset \begin{pmatrix} u & x_0 & x_1 & x_3 & x_4 & s \\ 1 & 0 & -1 & -1 & -1 & -2 \\ 0 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}. \end{aligned}$$

Again, we see that the equation defining Z is in the ideal (u, x_0) . If we denote by f, g (non-unique) polynomials such that $Z = \{x_0f + ug = 0\}$, so that we have

$$f = x_3^3 + x_4^3 + \dots \quad \text{and} \quad g = x_1^4 + (us)^4 + \dots,$$


and introduce a second unprojection variable

$$t = \frac{f}{u} = \frac{g}{x_0} \in \begin{pmatrix} -4 \\ 3 \end{pmatrix},$$

then we see that Z is the complete intersection:

$$\begin{cases} tu = f \\ tx_0 = g \end{cases} \subset \mathcal{F} = \begin{pmatrix} u & x_0 & x_1 & x_3 & x_4 & t & s \\ 1 & 0 & -1 & -1 & -1 & -4 & -2 \\ 0 & 1 & 1 & 1 & 1 & 3 & 1 \end{pmatrix}.$$

The 2-ray configuration of \mathcal{F} is



where Φ is a small map and \mathcal{F}^+ the ample model of $L - (1 + \varepsilon)E$ and G is a divisorial contraction. The restriction $\Phi|_Z : Z \dashrightarrow Z^+$ is a flop in 12 lines that are copies of \mathbb{P}^1_{u,x_0} lying over the 12 points

$$\{x_1^4 + x_3^4 - x_4^4 = x_2^3 + x_3^3 + \dots = 0\} \subset \mathbb{P}_{x_1,x_3,x_4}.$$

Since $Z \not\subset \{s = 0\}$, the restriction $G|_{Z^+} : Z^+ \rightarrow Y$ is a divisorial contraction. As above, applying the coordinate change for the action associated to

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$$

transforms A into

$$\begin{pmatrix} u & x_0 & x_1 & x_3 & x_4 & t & s \\ 1 & 1 & 0 & 0 & 0 & -1 & -1 \\ 1 & 2 & 1 & 1 & 1 & 2 & 0 \end{pmatrix}$$

and we see that $\mathbb{P}^+ = \mathbb{P}(1^4, 2^2)$, with coordinates su, x_1, x_3, x_4 (degree one) and sx_0, st (degree two). The proper transform of Z is

$$Y = \begin{cases} x_1^4 + \dots = 0 \\ x_2^3 + x_3^3 + \dots = 0 \end{cases} \subset \mathbb{P}^+,$$

i.e. Y is the complete intersection of a cubic and a quartic hypersurface in \mathbb{P}^+ , a Fano 3-fold of codimension 2 and genus 2. The map $Z^+ \rightarrow Y$ is a Kawamata blowup of one of the two $1/2(1, 1, 1)$ points on Y . This link is constructed in [CM04]. □

Remark 4.7. In several cases, one or both of the variables α, β are redundant. This means that $f : Z \rightarrow X$ is the restriction of a weighted blowup of some \mathbb{P}' with $\mathbb{P}^4 \subset \mathbb{P}' \subset \mathbb{P}$: the construction could have been obtained with a ‘smaller embedding’. For example, this is the case in our treatment of a terminal factorial quartic hypersurface with a cA_2 point: Corti and Mella [CM04] constructed the same link without introducing α, β . In a given example, it is usually clear how many (if any) variables need to be introduced. We have chosen to always introduce two variables (and then eliminate redundant ones) in order to present our results in a unified way.

Remark 4.8. It is crucial to understand that we make no claim about the existence of Sarkisov links initiated by divisorial contractions $f : Z \rightarrow X$ whose germs have weights different from those in Lemma 4.4, where X is one of the hypersurfaces $X^{i,j}$. Such divisorial contractions may occur, but they are not restrictions of weighted blowups of $\mathbb{P} = \mathbb{P}(1^5, 2^2)$. We expect that in some cases, one may construct such contractions by considering a different embedding of $X \subset \mathbb{P}'$ and looking at restrictions of weighted blowups of \mathbb{P}' . For instance, we do not know whether $X^{3,1}$ admits a Sarkisov link initiated by a divisorial contraction whose germ is a weighted blowup $(6, 1, 1, 1)$.

4.2 Other examples with cA singularities

In this section, we use similar techniques to give an example of a non-rigid terminal factorial quartic hypersurface with a cA_7 singular point. These can also be used to construct Sarkisov links initiated by divisorial contractions with discrepancy $a > 1$ and centre at a cA_n point for $n \geq 2$, where the possible values a, n are determined in Proposition 3.9. We give an example with $n = 2$ and $a = 2$.

Example 4.9. Let X be the hypersurface

$$X = \{(x_0x_1 - x_3^2)^2 - (x_0x_2 - x_4^2)^2 + x_0x_1^3 - x_1^2x_3^2 + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4.$$

The restriction of X to the affine chart $U_0 = \{x_0 = 1\}$ is

$$F(x, y, z, t) = (x - z^2)^2 - (y - t^2)^2 + x^3 - x^2z^2 + x^4 + y^4$$

and the singularity ($F = 0$) is equivalent to ($G = 0$), where G defines a cA_7 singularity. To see this, let $\underline{\varphi} = (\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in \text{Aut } \mathbb{C}[[x, y, z, t]]$, where

$$\begin{cases} \varphi_1(x, y, z, t) = x + y - (z^2(1 - \frac{1}{2}z^2) + t^2), \\ \varphi_2(x, y, z, t) = x - y - (z^2(1 - \frac{1}{2}z^2) - t^2), \\ \varphi_3(x, y, z, t) = z, \\ \varphi_4(x, y, z, t) = t \end{cases}$$

and check that $F(x, y, z, t) = G(\varphi_1, \varphi_2, \varphi_3, \varphi_4)$, where

$$G(x, y, z, t) = xy + z^8 + t^8 + G'(x, y, z, t), \tag{10}$$

where $G'(x, y, z, t)$ is a polynomial of degree strictly greater than one with respect to the weights

$$\underline{w} = (1/2, 1/2, 1/8, 1/8).$$

Since G is a semi-quasi-homogeneous polynomial of degree one with respect to \underline{w} , and since no element of its Jacobian algebra has degree strictly greater than one with respect to these weights,

$$G \sim xy + z^8 + t^8$$

and $(P \in X)$ is a cA_7 point. The hypersurface $X \subset \mathbb{P}^4$ is a terminal and factorial quartic hypersurface and hence is a Mori fibre space.

As in §4.1, we embed X as the complete intersection:

$$\begin{cases} \alpha = x_0(x_1 + x_2) - (x_3^2 + x_4^2) \\ \beta = x_0(x_1 - x_2) - (x_3^2 - x_4^2) \\ \alpha\beta + 2x_1^2(\alpha + \beta) + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathbb{P} = \mathbb{P}(1^4, 2^2).$$


We consider the weighted blowup $F : \mathcal{F} \rightarrow \mathbb{P}$, where \mathcal{F} is the Picard rank 2 toric variety $TV(I, A)$, where $I = (u, x_0) \cap (x_1, \dots, x_4, \alpha, \beta)$ is the irrelevant ideal and A is the action of $\mathbb{C}^* \times \mathbb{C}^*$ with weights

$$\begin{pmatrix} u & x_0 & x_3 & x_4 & x_1 & x_2 & \alpha & \beta \\ 1 & 0 & -1 & -1 & -2 & -2 & -4 & -4 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 & 2 \end{pmatrix}.$$

The proper transform of X is

$$Z = \begin{cases} u^2\alpha = x_0(x_1 + x_2) - (x_3^2 + x_4^2) \\ u^2\beta = x_0(x_1 - x_2) - (x_3^2 - x_4^2) \\ \alpha\beta + 2x_1^2(\alpha + \beta) + x_1^4 + x_2^4 = 0 \end{cases} = \binom{2}{2} \cap \binom{2}{2} \cap \binom{8}{4} \subset \mathcal{F}$$

and we check that the restriction of F to Z is indeed a divisorial contraction $f : Z \rightarrow X$ with discrepancy $a = 1$. The 2-ray configuration on \mathcal{F} is



where Φ is a small map and \mathcal{F}^+ is the ample model of $L - (1 + \varepsilon)E$ and G is a fibration morphism and $\mathbb{P}^+ = \mathbb{P}(1, 1, 2, 2) = \mathbb{P}_{x_1, x_2, \alpha, \beta}$ is the ample model $\text{Proj}(n(L - 2E))$ for suitable $n \gg 1$.

The restriction of Φ to Z is an isomorphism because the monomials x_3^2, x_4^2 appear in the first two equations; we still denote by Z its image. The restriction of G to Z is a conic bundle over the quartic surface $S_4 \subset \mathbb{P}(1, 1, 2, 2)$ defined by the third equation of Z .

The results of § 4.1 and Example 4.9 thus show the following theorem.

THEOREM 4.10. *There are examples of non-rigid terminal factorial quartic hypersurfaces with a singular point of type cA_n for all possible $n \geq 2$.*

Each of the Sarkisov links we have constructed so far is initiated by a divisorial contraction with centre at a cA_n point and discrepancy $a = 1$. We now construct an example with higher discrepancy.

Example 4.11. We construct a Sarkisov link initiated by a divisorial contraction with centre at a cA_2 point. Unlike the previous examples, which only involved flips and flops, the Sarkisov link in this example involves an antiflip.

Consider the terminal, factorial quartic hypersurface

$$X = \{x_0x_1(x_0x_2 - x_4^2) + x_0(x_2^3 + x_3^3) - x_1^4 + x_2^4 + x_3^4\} \subset \mathbb{P}^4.$$

Then $\text{Sing } X = \{P, P_4\}$ and, setting $x_0 = 1$ in the expression above, we see that

$$(P \in X) \sim 0 \in \{xy + z^3 + t^6 + (\text{higher order terms}) = 0\},$$

so that P is a cA_2 point and a divisorial contraction with centre at $(P \in X)$ has discrepancy 1 or 2 by Theorem 3.13. We embed X as

$$X = \begin{cases} \alpha = x_0x_1 \\ \beta = x_0x_2 - x_4^2 \\ \alpha\beta x_0(x_2^3 + x_3^3) - x_1^4 + x_2^4 + x_3^4 = 0 \end{cases} \subset \mathbb{P}(1^5, 2^2).$$

If we consider a divisorial contraction with discrepancy 1, we obtain a link of the same form as above. We now consider the case when f is a divisorial contraction with discrepancy 2. Then, by Theorem 3.13, the weights of $(\alpha, \beta, x_3, x_4)$ are either $(1, 5, 2, 1)$ or $(3, 3, 2, 1)$. We consider the second case and, as in Lemma 4.4, we show that \mathcal{F} is a toric variety $TV(I, A)$, where

$$A = \begin{pmatrix} u & x_0 & x_1 & x_2 & x_3 & x_4 & \alpha & \beta \\ 1 & 0 & -p & -q & -2 & -1 & -3 & -3 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 & 2 \end{pmatrix} \text{ for } (p, q) = (2, 2), (2, 3) \text{ or } (3, 2).$$

We consider the case when $(p, q) = (2, 2)$, so that, after re-ordering,

$$A = \begin{pmatrix} u & x_0 & x_4 & \alpha & \beta & x_1 & x_2 & x_3 \\ 1 & 0 & -1 & -3 & -3 & -2 & -2 & -2 \\ 0 & 1 & 1 & 2 & 2 & 1 & 1 & 1 \end{pmatrix}$$

and the proper transform of X is given by the equations

$$\begin{cases} u\alpha = x_0x_1, \\ u\beta = x_0x_2 - x_4^2, \\ \alpha\beta x_0(x_2^3 + x_3^3) + u^2(-x_1^4 + x_2^4 + x_4^2) = 0. \end{cases}$$

As in the proof of the cA_2 case in Theorem 4.5, the first equation is in the ideal (u, x_0) ; we re-embed Z so that it follows the ambient 2-ray configuration: we introduce an unprojection variable s such that

$$\begin{cases} su = x_1 \\ sx_0 = \alpha \end{cases} \quad s \in \begin{pmatrix} -3 \\ 1 \end{pmatrix}.$$

We then see that the variables x_1 and α are redundant, so that the expression of Z is

$$\begin{cases} u\beta = x_0x_2 - x_4^2, \\ x_0(s\beta + x_3^3 + x_2^3) + u^2((su)^4 + x_2^4 + x_3^4) = 0. \end{cases}$$

As above, since the second equation is in the ideal (u^2, x_0) , we need to introduce a second unprojection variable η such that

$$\begin{cases} \eta u^2 = s\beta + x_3^3 + x_2^3 \\ \eta x_0 = (su)^4 + x_2^4 + x_3^4 \end{cases} \quad \eta \in \begin{pmatrix} -8 \\ 3 \end{pmatrix}.$$


We now get that Z is the complete intersection:

$$\begin{cases} u\beta = x_0x_2 - x_4^2 \\ \eta u^2 = s\beta + x_3^3 + x_2^3 \\ \eta x_0 = (su)^4 + x_2^4 + x_3^4 \end{cases} \subset \mathcal{F},$$

where \mathcal{F} denotes the toric variety $TV(I, A)$ for

$$A = \begin{pmatrix} u & x_0 & x_4 & \beta & x_2 & x_3 & \eta & s \\ 1 & 0 & -1 & -3 & -2 & -2 & -8 & -3 \\ 0 & 1 & 1 & 2 & 1 & 1 & 3 & 1 \end{pmatrix}.$$

The 2-ray configuration on \mathcal{F} is



where:

- $\Phi_1 : \mathcal{F} \dashrightarrow \mathcal{F}_1$ is a small map, and \mathcal{F}_1 is the ample model of $L - (1 + \varepsilon)E$;
- $\Phi_2 : \mathcal{F}_1 \dashrightarrow \mathcal{F}_2$ is a small map, and \mathcal{F}_2 is the ample model of $2L - (3 + \varepsilon)E$;
- $\Phi_3 : \mathcal{F}_2 \dashrightarrow \mathcal{F}_3$ is a small map, and \mathcal{F}_3 is the ample model of $L - (2 + \varepsilon)E$;
- $G : \mathcal{F}_3 \rightarrow \mathbb{P}^+$ is a divisorial contraction, and \mathbb{P}^+ is the ample model of $8L - 3E$.

We study the restriction of this 2-ray configuration to Z . Since the monomial x_4^2 appears in one of the equations of Z , the restriction $\Phi_{1|Z}$ is an isomorphism. We still denote by Z its image.

We now prove that $\Phi_{2|Z} : Z \dashrightarrow Z_2$ is a small birational map. Since it is a K -positive contraction, we also need to prove that Z_2 has terminal singularities. The exceptional locus of $\Phi_{2|Z}$ is at most one dimensional, as the only pure monomial in u, x_0, x_4 that appears in the equations of Z is x_4^2 . The exceptional locus of $(\Phi_{2|Z})^{-1}$ is also at most one dimensional, as pure monomials in x_2, x_3, η, s appear in two of the three equations defining Z .

In order to study $\Phi_{2|Z}$, we localise near P_β . Setting $\beta = 1$, we use the first two equations to eliminate the variables u and s , so that $X \cap \{\beta = 1\}$ is the hypersurface defined by the third equation:

$$\{\eta x_0 = x_2^4 + x_3^2 + (\eta u^2 - x_3^3 - x_2^3)^4 (x_0 x_2 - x_4^2)^4 = 0\}.$$

As above, under the change of coordinates associated to

$$\begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix} \in \text{SL}_2(\mathbb{Z}),$$

the action becomes

$$\begin{pmatrix} u & x_0 & x_4 & \beta & x_2 & x_3 & \eta & s \\ -1 & -1 & 0 & 1 & 1 & 1 & 5 & 2 \\ 2 & 3 & 1 & 0 & -1 & -1 & -7 & -3 \end{pmatrix},$$

so that once the variables u, s and the second and third equations defining Z have been eliminated, we are left with the inverse of a flip of the hypersurface defined by the first equation of Z in $\mathbb{C}_{\eta, x_2, x_3, x_0, x_4}$, which is, in the notation of [Bro99], of the form $(7, 1, 1, -3, -1; 4)$. The map $\Phi_{2|Z}$ is thus an antiflip between 3-folds with terminal singularities. The exceptional locus of $\Phi_{2|Z}$ is empty because the equation of $X \cap U_\beta$ has no pure monomial in x_0, x_4 ; the exceptional locus of $(\Phi_{2|Z})^{-1}$ consists of four lines

$$\{x_2^4 + x_3^4 = 0\} \subset \mathbb{P}(1, 1, 7) = \mathbb{P}_{x_2, x_3, \eta}.$$

By the construction of f , the basket of Z consists of two $[3, 1]$ singularities, one of which lies over P_β . From [Bro99], the basket of Z_2 consists of one $[7, 1]$ singular point and one $[3, 1]$ singular point.

The restriction $\Phi_{3|Z_2}$ is an isomorphism because $x_3^3 + x_2^3$ appears in the second equation defining Z and $x_3^4 + x_2^4$ in the third. We still denote by Z_2 its image by $\Phi_{3|Z_2}$.

The restriction of G to Z_2 is a morphism because $Z_2 \cap \{s = 0\}$ is a prime divisor. As above, under the change of coordinates associated to

$$\begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} \in \text{SL}_2(\mathbb{Z}),$$

the action becomes

$$\begin{pmatrix} u & x_0 & x_4 & \beta & x_2 & x_3 & \eta & s \\ -1 & -2 & -1 & -1 & 0 & 0 & 2 & 1 \\ 1 & 3 & 2 & 3 & 1 & 1 & 1 & 0 \end{pmatrix},$$

so that $\mathbb{P}^+ = \mathbb{P}(1^4, 2, 3^2)$ with coordinates $\eta, s^3 u, s^2 x_2, s^2 x_3$ (degree one), $s^5 x_4$ (degree two) and $s^7 \beta, s^8 x_0$ (degree three).

The proper transform of Z is

$$Y = \begin{cases} (s^3u)(s^7\beta) = (s^8x_0)(s^2x_2) - (s^5x_4)^2 \\ s^7\beta = (s^3u)^2\eta - (s^2x_3)^3 - (s^2x_2)^3 \\ (s^8x_0)\eta = (s^3u)^4 + (s^2x_2)^4 + (s^2x_3)^4 \end{cases} \subset \mathbb{P}^+,$$

so that the variable $s^7\beta$ and the second equation are redundant. The 3-fold Y is a complete intersection of two quartic hypersurfaces $Y_{4,4} \subset \mathbb{P}(1^4, 2, 3)$. This is a Fano 3-fold of index 1 and is the Fano variety number 16 204 in [GRD].

Setting $\eta = 1$ shows that $Y_{4,4} \subset \mathbb{P}^+$ has a singular point of type cE_7 at P_η , and the contraction $g : Z_2 \rightarrow Y$ is a divisorial contraction with centre at P_η . The contraction g has discrepancy 2, and the basket of Z_2 shows that g is, in Kawakita’s notation [Kaw03], a contraction of type I.

In summary, we have constructed a Sarkisov link from a terminal factorial quartic hypersurface X with a cA_2 point ($P \in X$) to a complete intersection $Y_{4,4} \subset \mathbb{P}(1^4, 2, 3)$ with a cE_7 point ($Q \in Y$), which is of the form

$$\begin{array}{ccc} Z & \xrightarrow{\quad \Phi \quad} & Z^+ \\ f \downarrow & & \downarrow g \\ X & & Y_{4,4} \end{array}$$

where:

- f is a discrepancy 2 divisorial contraction with centre at P ;
- $\varphi = \varphi_3 \circ \varphi_2 \circ \varphi_1$, with φ_1, φ_3 isomorphisms and φ_2 an antiflip;
- g is a discrepancy 2 divisorial contraction with centre at Q .

4.3 Examples with cD and cE singularities

We now give examples of non-rigid factorial quartic hypersurfaces with singular points that are not of type cA . The study of the pliability of quartics with cD_m and $cE_{6,7,8}$ singular points is complicated by the fact that, unlike in the cA_n case, there is no classification of the germs of divisorial extractions $f : Z \rightarrow X$ with centre at a cD or a cE point. We only know the germs of a few explicit divisorial extractions with these centres: those that are weighted blowups with discrepancy 1. The following examples are non-rigid quartic hypersurfaces with a cD or a cE singular point.

Example 4.12. Let

$$X = \{x_0^2x_1^2 - x_0(x_2^3 + x_3^3 + x_4^3) + f_4(x_1, x_2, x_3, x_4) = 0\} \subset \mathbb{P}^4$$

be a quartic hypersurface, where f_4 is a general homogeneous polynomial of degree four such that $\{x_2^3 + x_3^3 + x_4^3 = f_4(0, x_2, x_3, x_4) = 0\} \subset \mathbb{P}^2$ consists of 12 points. By generality of f_4 , X is factorial and $\text{Sing } X$ consists of a single point $(1:0:0:0:0)$ that is of type cD_4 . The restriction of the weighted blowup $\mathcal{F} \rightarrow \mathbb{P}$ with centre at P and weights $(2, 1, 1, 1)$ is a divisorial contraction with discrepancy 1. It initiates a Sarkisov link that consists of a flop in 12 lines over the 12 points

$$\{x_2^3 + x_3^3 + x_4^3 = f_4(0, x_2, x_3, x_4) = 0\} \subset \mathbb{P}_{x_2, x_3, x_4}^2$$

followed by a divisorial contraction to a $1/2(1, 1, 1)$ point on a quasi-smooth intersection of a quartic and a cubic $Y = Y_{3,4} \subset \mathbb{P}(1^4, 2^2)$. Note that the Fano 3-fold Y is not a general $Y_{3,4} \subset \mathbb{P}(1^4, 2^2)$. Denoting by x_1, \dots, x_4 the coordinates of weight 1 and by y_1, y_2 those of weight 2, the equation of Y is of the form

$$Y : \begin{cases} y_1x_1 - (x_2^3 + x_3^3 + x_4^3) = 0 \\ y_1y_2 - y_1^2 - f_4(x_1, \dots, x_4) = 0 \end{cases} \subset \mathbb{P}(1^4, 2^2).$$

Note the similarity of this link to the link between a quartic $X' \subset \mathbb{P}^4$ with a cA_2 point and a general $Y_{3,4} \subset \mathbb{P}(1^4, 2^2)$ studied in [CM04]. In our case, Y is a special quasi-smooth model in its family, and we conjecture that as in [CM04] X is birigid, i.e. $\mathcal{P}(X) = \{[X], [Y]\}$.

Example 4.13. Let

$$X = \{x_0^2x_1^2 + x_0x_2^2x_3 + x_1^4 + x_2^4 + x_3^4 + x_4^4 = 0\} \subset \mathbb{P}^4;$$

$\text{Sing } X$ consists of a single point $(1:0:0:0:0)$ that is of type cD_5 . The restriction of the weighted blowup $\mathcal{F} \rightarrow \mathbb{P}$ with centre at P and weights $(2, 1, 2, 1)$ is a divisorial contraction with discrepancy 1. It initiates a Sarkisov link that consists of a flop in four lines lying above $\{x_2^4 + x_4^4 = 0\} \subset \mathbb{P}_{x_2, x_4}^1$ followed by a del Pezzo fibration of degree two.

Example 4.14 (Singular points of cE type). In the same way, let

$$X = \{x_0^2x_1^2 + x_0x_2^3 + x_1^4 + x_2^4 + x_3^4 + x_4^4 = 0\} \subset \mathbb{P}^4;$$

$\text{Sing } X$ consists of a single point $(1:0:0:0:0)$ that is of type cE_6 . The restriction of the weighted blowup $\mathcal{F} \rightarrow \mathbb{P}$ with centre at P and weights $(2, 2, 1, 1)$ is a divisorial contraction with discrepancy 1. It initiates a Sarkisov link that consists of a flop in four lines lying above $\{x_3^4 + x_4^4 = 0\} \subset \mathbb{P}_{x_3, x_4}^1$ followed by a del Pezzo fibration of degree two.

Similarly, if

$$\begin{aligned} Y &= \{(x_0x_1 - x_4^2)^2 + x_0x_2^3 + x_2x_3^3 + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4 \quad \text{and} \\ Z &= \{(x_0x_1 - x_3^2 - x_4^2)^2 + x_0x_2^3 + x_0x_1^2(x_3 + x_4) + x_1^4 + x_2^4 = 0\} \subset \mathbb{P}^4, \end{aligned}$$

then

$$\text{Sing } Y = \text{Sing } Z = \{P\} = (1:0:0:0:0)$$

and $(P \in Y)$ is of local analytic type cE_7 , while $(P \in Z)$ is of local analytic type cE_8 . The 3-folds Y and Z are non-rigid. Indeed, embed Y and Z as complete intersections

$$Y = \begin{cases} \alpha = x_0x_1 - x_4^2 \\ \alpha^2 + x_0x_2^3 + x_2x_3^3 + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathbb{P}_1 = \mathbb{P}(1^5, 2),$$

where the variable of degree two is α , and

$$Z = \begin{cases} \beta = x_0x_1 - x_3^2 - x_4^2 \\ \beta^2 + x_0x_2^3 + x_0x_1^2(x_3 + x_4) + x_1^4 + x_2^4 = 0 \end{cases} \subset \mathbb{P}_2 = \mathbb{P}(1^5, 2),$$

where the variable of degree two is β . Then consider the rank 2 toric varieties \mathcal{F}_1 and \mathcal{F}_2 associated to

$$\begin{pmatrix} u & x_0 & x_1 & x_2 & x_3 & x_4 & \alpha \\ 1 & 0 & -2 & -2 & -2 & -1 & -3 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} u & x_0 & x_1 & x_2 & x_3 & x_4 & \beta \\ 1 & 0 & -3 & -2 & -1 & -1 & -3 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 \end{pmatrix}.$$

The restrictions of the weighted blowup $\mathcal{F}_i \rightarrow \mathbb{P}_i$ with centre at P are divisorial contractions $f_1 : \tilde{Y} \rightarrow Y$ and $f_2 : \tilde{Z} \rightarrow Z$. These both have discrepancy 1, and globalise the germs of weighted blowups with centre at a cE point classified in [Mar96, Theorem 3.2]. Further, f_1 initiates a Sarkisov link between Y and a conic bundle, while f_2 initiates a link between Z and an index 2 Fano 3-fold $Y_{4,6} \subset \mathbb{P}(1^2, 2^2, 3^2)$, which is the Fano variety number 40 369 in [GRD].

ACKNOWLEDGEMENTS

We would like to thank Ivan Cheltsov, Alessio Corti, Masayuki Kawakita, Yuri Prokhorov, Josef Schicho and Constantin Shramov for useful comments and conversations. We thank the referee for many helpful suggestions.

REFERENCES

- Ahm12 H. Ahmadinezhad, *On del Pezzo fibrations that are not birationally rigid*, J. Lond. Math. Soc. (2) **86** (2012), 36–62; MR 2959294.
- Ahm14 H. Ahmadinezhad, *On pliability of del Pezzo fibrations and Cox rings*, J. reine angew. Math. (2014), doi:10.1515/crelle-2014-0095.
- AGV85 V. I. Arnol'd, S. M. Guseĭn-Zade and A. N. Varchenko, *Singularities of differentiable maps: Vol. I: the classification of critical points, caustics and wave fronts*, Monographs in Mathematics, vol. 82 (Birkhäuser, Boston, MA, 1985); translated from the Russian by Ian Porteous and Mark Reynolds.
- BCHM10 C. Birkar, P. Cascini, C. D. Hacon and J. McKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), 405–468.
- Bro99 G. Brown, *Flips arising as quotients of hypersurfaces*, Math. Proc. Cambridge Philos. Soc. **127** (1999), 13–31.
- Cor95 A. Corti, *Factoring birational maps of threefolds after Sarkisov*, J. Algebraic Geom. **4** (1995), 223–254.
- CM04 A. Corti and M. Mella, *Birational geometry of terminal quartic 3-folds. I*, Amer. J. Math. **126** (2004), 739–761.
- CPR00 A. Corti, A. Pukhlikov and M. Reid, *Fano 3-fold hypersurfaces*, in *Explicit birational geometry of 3-folds*, London Mathematical Society Lecture Note Series, vol. 281 (Cambridge University Press, Cambridge, 2000), 175–258.
- GRD *The graded ring database*, <http://www.grdb.co.uk/>.
- HM13 C. D. Hacon and J. McKernan, *The Sarkisov program*, J. Algebraic Geom. **22** (2013), 389–405.
- HK00 Y. Hu and S. Keel, *Mori dream spaces and GIT*, Michigan Math. J. **48** (2000), 331–348.
- IM71 V. A. Iskovskikh and Ju. I. Manin, *Three-dimensional quartics and counterexamples to the Lüroth problem*, Mat. Sb. (N.S.) **86** (1971), 140–166.
- IP99 V. A. Iskovskikh and Yu. G. Prokhorov, *Fano varieties*, in *Algebraic geometry, V*, Encyclopaedia of Mathematical Sciences, vol. 47 (Springer, Berlin, 1999), 1–247.
- Kal11 A.-S. Kaloghiros, *The defect of Fano 3-folds*, J. Algebraic Geom. **20** (2011), 127–149.
- Kal12 A.-S. Kaloghiros, *A classification of terminal quartic 3-folds and applications to rationality questions*, Math. Ann. **354** (2012), 263–296.
- Kal13 A.-S. Kaloghiros, *Relations in the Sarkisov program*, Compositio Math. **149** (2013), 1685–1709.
- KKL14 A.-S. Kaloghiros, A. Küronya and V. Lazić, *Finite generation and geography of models*, in *Minimal models and extremal rays*, Advanced Studies in Pure Mathematics (Mathematical Society of Japan, Tokyo, 2014), to appear.

- Kaw01 M. Kawakita, *Divisorial contractions in dimension three which contract divisors to smooth points*, Invent. Math. **145** (2001), 105–119.
- Kaw02 M. Kawakita, *Divisorial contractions in dimension three which contract divisors to compound A_1 points*, Compositio Math. **133** (2002), 95–116.
- Kaw03 M. Kawakita, *General elephants of three-fold divisorial contractions*, J. Amer. Math. Soc. **16** (2003), 331–362 (electronic).
- Kol98 J. Kollár, *Real algebraic threefolds. I. Terminal singularities*, Collect. Math. **49** (1998), 335–360; dedicated to the memory of Fernando Serrano.
- KM98 J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134 (Cambridge University Press, Cambridge, 1998).
- Mar96 D. Markushevich, *Minimal discrepancy for a terminal cDV singularity is 1*, J. Math. Sci. Univ. Tokyo **3** (1996), 445–456.
- Mel04 M. Mella, *Birational geometry of quartic 3-folds. II. The importance of being \mathbb{Q} -factorial*, Math. Ann. **330** (2004), 107–126.
- Nam97 Y. Namikawa, *Smoothing Fano 3-folds*, J. Algebraic Geom. **6** (1997), 307–324.
- NS95 Y. Namikawa and J. H. M. Steenbrink, *Global smoothing of Calabi–Yau threefolds*, Invent. Math. **122** (1995), 403–419.
- Pet98 K. F. Pettersen, *On nodal determinantal quartic hypersurfaces in \mathbf{P}^4* , PhD thesis, University of Oslo (1998).
- Rei87 M. Reid, *Young person’s guide to canonical singularities*, in *Algebraic geometry (Bowdoin, Brunswick, Maine, 1985)*, Proceedings of Symposia in Pure Mathematics, vol. 46 (American Mathematical Society, Providence, RI, 1987), 345–414.
- Tod33 J. A. Todd, *Configurations defined by six lines in space of three dimensions*, Math. Proc. Cambridge Philos. Soc. **29** (1933), 52–68.
- Tod35 J. A. Todd, *A note on two special primals in four dimensions*, Q. J. Math. **6** (1935), 129–136.
- Tod36 J. A. Todd, *On a quartic primal with forty-five nodes, in space of four dimensions*, Q. J. Math. **7** (1936), 169–174.

Hamid Ahmadinezhad h.ahmadinezhad@bristol.ac.uk
School of Mathematics, University of Bristol, Bristol BS8 1TW, UK

Anne-Sophie Kaloghiros anne-sophie.kaloghiros@brunel.ac.uk
Imperial College London, 180 Queen’s Gate, London SW7 2AZ, UK

Current address: Brunel University London, Uxbridge UB8 3PH, UK