



A Congruence Modulo Four for Real Schubert Calculus with Isotropic Flags

Nickolas Hein, Frank Sottile, and Igor Zelenko

Abstract. We previously obtained a congruence modulo four for the number of real solutions to many Schubert problems on a square Grassmannian given by osculating flags. Here we consider Schubert problems given by more general isotropic flags, and prove this congruence modulo four for the largest class of Schubert problems that could be expected to exhibit this congruence.

Introduction

The number of real solutions to a system of real equations is congruent to the number of complex solutions modulo two. We established a congruence modulo four for many symmetric Schubert problems given by osculating flags, leaving as a conjecture a stronger form of that result [9]. We prove this conjecture for symmetric Schubert problems given by flags that are isotropic with respect to a symplectic form, giving a simpler proof of a stronger and more basic result than that obtained in [9].

This congruence modulo four follows from a result on the real points in fibers of a map between real varieties equipped with an involution. When the fixed point set of the involution has codimension at least two, the number of real points satisfies a congruence modulo four. There is an involution acting on symmetric Schubert problems given by isotropic flags and we can compute the dimension of the fixed point locus in a universal family of Schubert problems. Our inability to compute this dimension when the flags are osculating was the obstruction to establishing the conjecture in [9].

The congruence modulo four often implies a non-trivial lower bound on the number of real solutions to a symmetric Schubert problem given by isotropic flags. Similar lower bounds and congruences in real algebraic geometry have been of significant interest [1, 3, 6, 7, 10, 11, 17, 18, 20, 24]. Another topological study was recently made of this phenomenon in the Schubert calculus [5], and delicate lower bounds [14] were given by computing the signature of a hermitian matrix arising in the proof of the Shapiro Conjecture [15].

In Section 1 we state our main result, whose proof occupies Section 2.

Received by the editors February 6, 2015; revised June 25, 2015.

Published electronically February 23, 2017.

Research of authors F. S. and N. H. supported in part by NSF grant DMS-1001615. Research of author I. Z. supported by NSF grant DMS-140619.

AMS subject classification: 14N15, 14P99.

Keywords: Lagrangian Grassmannian, Wronski map, Shapiro Conjecture.

1 Symmetric Schubert Problems

Let V be a complex vector space of dimension $2m$ equipped with a nondegenerate alternating form $\langle \cdot, \cdot \rangle: V \otimes V \rightarrow \mathbb{C}$. Write \overline{W} for the complex conjugate of a point, vector, subspace, or variety W . A variety W is *real* if it is defined by real equations; equivalently, if $\overline{W} = W$. Write $W(\mathbb{R})$ for the real points of a real variety W , i.e., those that are fixed by complex conjugation. Write S_a for the symmetric group of permutations of $\{1, \dots, a\}$.

The set of m -dimensional linear subspaces of V forms the Grassmannian $\text{Gr}(m, V)$, which is a manifold of dimension m^2 . A *flag* is a sequence $F_\bullet: F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_{2m} = V$ of linear subspaces of V with $\dim F_i = i$. A *partition* is a weakly decreasing sequence of integers $\lambda: m \geq \lambda_1 \geq \dots \geq \lambda_m \geq 0$. A flag F_\bullet and a partition λ determine a Schubert subvariety of $\text{Gr}(m, V)$,

$$X_\lambda F_\bullet := \{H \in \text{Gr}(m, V) \mid \dim H \cap F_{m+i-\lambda_i} \geq i \text{ for } i = 1, \dots, m\}.$$

This has codimension $|\lambda| := \lambda_1 + \dots + \lambda_m$ in $\text{Gr}(m, V)$.

Let $\lambda = (\lambda^1, \dots, \lambda^s)$ be a list of partitions and $F_\bullet^1, \dots, F_\bullet^s$ be general flags. By Kleiman’s Transversality Theorem [12] the intersection

$$(1.1) \quad X_{\lambda^1} F_\bullet^1 \cap X_{\lambda^2} F_\bullet^2 \cap \dots \cap X_{\lambda^s} F_\bullet^s.$$

is either empty or has dimension $\dim \text{Gr}(m, V) - |\lambda^1| - \dots - |\lambda^s|$. Call λ a *Schubert problem* if this expected dimension is zero, so that (1.1) is either empty or consists of finitely many points. The number of points $d(\lambda)$ in (1.1) is independent of the choice of general flags. We will assume that $d(\lambda) \neq 0$. A choice of flags is an *instance* of the Schubert problem λ ; its *solutions* are the points in (1.1). The instance is *real* if for all i , there is some j with $\overline{F_\bullet^i} = F_\bullet^j$ and $\lambda^i = \lambda^j$, for then (1.1) is stable under complex conjugation.

Remark 1.1 Osculating flags provide a rich source of isotropic flags. As explained in [9, §3], a rational normal curve $\gamma: \mathbb{C} \rightarrow V$ induces a symplectic form on V and a symplectic form on V gives rise to a rational normal curve; we may assume that γ is real in that $\gamma(\bar{t}) = \overline{\gamma(t)}$. If γ is a rational normal curve corresponding to the symplectic form $\langle \cdot, \cdot \rangle$, then every osculating flag is isotropic. (For $t \in \mathbb{C}$, the *osculating flag* $F_\bullet(t)$ is the flag whose i -plane $F_i(t)$ is spanned by $\gamma(t)$ and its derivatives $\gamma'(t), \dots, \gamma^{(i-1)}(t)$.)

The study of real solutions to Schubert problems given by flags osculating at real points in Grassmannians and flag manifolds has been quite rich and fruitful [4, 13, 15, 16, 19, 21].

A partition λ is represented by its Young diagram, which is a left-justified array of boxes with λ_i boxes in row i . We display some partitions with their Young diagrams,

$$(2, 1, 1) \longleftrightarrow \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array}, \quad (2, 2) \longleftrightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \quad \text{and} \quad (3, 2, 1) \longleftrightarrow \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \square & & \\ \hline \end{array}.$$

A partition λ is *symmetric* if it is symmetric about its main diagonal, that is, if $\lambda = \lambda'$, where λ' is the transpose of λ . The partitions $(2, 2)$ and $(3, 2, 1)$ are symmetric while $(2, 1, 1)$ is not. A Schubert problem λ is *symmetric* if every partition in λ is symmetric.

Recall that our vector space V was equipped with a nondegenerate alternating bilinear form $\langle \cdot, \cdot \rangle$. A linear subspace W of V has annihilator $\perp(W)$ under $\langle \cdot, \cdot \rangle$,

$$\perp(W) := \{v \in V \mid \langle v, w \rangle = 0 \text{ for all } w \in W\},$$

and we have $\dim W + \dim \perp(W) = 2m$. This induces a map $H \mapsto \perp(H)$ on $\text{Gr}(m, V)$ called the *Lagrangian involution*. Given a flag F_\bullet , we get the flag $\perp(F_\bullet)$ whose i -plane is $\perp(F_{2m-i})$. A flag F_\bullet is *isotropic* if $\perp(F_\bullet) = F_\bullet$.

The *length*, $\ell(\lambda)$ of a symmetric partition is the number of boxes on its main diagonal, so $\ell(2, 2) = 2$ while $\ell(2, 1) = 1$. We state our main theorem.

Theorem 1.2 *Suppose that $\lambda = (\lambda^1, \dots, \lambda^s)$ is a symmetric Schubert problem on $\text{Gr}(m, V)$ and that $F_\bullet^1, \dots, F_\bullet^s$ are isotropic flags defining a real instance of the Schubert problem λ such that (1.1) is finite. If $\sum_i \ell(\lambda^i) \geq m+4$, then the number (counted with multiplicity) of real points in (1.1) is congruent to the number $d(\lambda)$ of complex points, modulo four.*

Remark 1.3 We show in Remark 2.6 that $\sum_i \ell(\lambda^i) \geq m$ and this sum has the same parity as m , so that the condition in Theorem 1.2 for this congruence modulo four is that $\sum_i \ell(\lambda^i)$ is not equal to m or to $m + 2$, which is very mild.

We use the observation of Remark 1.1 that osculating flags are isotropic to deduce a corollary about Schubert problems given by osculating flags. Fix a real rational normal curve $\gamma: \mathbb{C} \rightarrow V$ with corresponding osculating flags $F_\bullet(t)$ for $t \in \mathbb{C}$, and a symmetric Schubert problem $\lambda = (\lambda^1, \dots, \lambda^s)$ on $\text{Gr}(m, V)$. An *osculating instance* of this Schubert problem is a list of distinct complex numbers t_1, \dots, t_s which give corresponding osculating flags $F_\bullet(t_1), \dots, F_\bullet(t_s)$. This osculating instance is *real* if the set $\{t_1, \dots, t_s\}$ is closed under complex conjugation and if $\bar{t}_i = t_j$ implies that $\lambda_i = \lambda_j$.

We deduce a corollary to Theorem 1.2 that implies [9, Conjecture 21], which was the strongest result one could reasonably expect to hold concerning this congruence modulo four for symmetric Schubert problems. This is strictly stronger than all congruence results obtained in [9].

Corollary 1.4 *Suppose that $\lambda = (\lambda^1, \dots, \lambda^s)$ is a symmetric Schubert problem on $\text{Gr}(m, V)$ and that $F_\bullet(t_1), \dots, F_\bullet(t_s)$ are osculating flags defining a real instance of λ . If $\sum_i \ell(\lambda^i) \geq m + 4$, then the number (counted with multiplicity) of real points in*

$$X_{\lambda^1} F_\bullet(t_1) \cap X_{\lambda^2} F_\bullet(t_2) \cap \dots \cap X_{\lambda^s} F_\bullet(t_s)$$

is congruent to the number $d(\lambda)$ of complex points, modulo four.

We do not need to assert that the intersection consists of finitely many points, for it always does [2].

Remark 1.5 When $d(\lambda)$ is congruent to two modulo four and $\sum_i \ell(\lambda^i) \geq m + 4$, there will always be at least two real solutions to a real instance of a symmetric Schubert problem. Such lower bounds implied by Theorem 1.2 occur frequently. Table 1 gives the total number of symmetric Schubert problems in $\text{Gr}(m, V)$ for small values of m , together with the number of those for which Theorem 1.2 implies a lower

bound of two. The timings are reported in GHz-seconds (s), GHz-hours (h), and GHz-years (y).

Table 1: Numbers of symmetric Schubert problem with a lower bound of two.

m	2	3	4	5	6	7
Symmetric	1	8	81	1037	16933	349844
Have lower bound	0	2	14	199	3289	82753
Percentage	0	25	17.3	19.2	19.4	23.7
Time	0.57 s	0.6 s	1.9 s	158 s	17.1 h	1.15 y

Real instances of nine of the smallest problems from Table 1 involving osculating flags were computed [8]. Few other problems from this table are feasible to compute using symbolic methods. For eight (numbers 20,57, 507, 568, 586, 587, 590, and 595 from [8]) at least one real instance had only two real solutions, showing that the lower bound of two is sharp in these cases.

Real osculating instances of problem number 773 from [8] with 70 solutions in $\text{Gr}(5, 10)$ have a sharp lower bound of six real solutions. This is a member of a family of such symmetric Schubert problems. By results in [7, §-3], when $m = 2k+1$ is odd, a real osculating instance of the symmetric problem on $\text{Gr}(2k+1, V)$ given by one condition $(2k, \dots, 2k, 0)$ and $4k-1$ conditions \square , which has $\binom{4k}{2k}$ complex solutions, will have at least $\binom{2k}{k}$ real solutions, and this lower bound is attained.

2 Proof of Theorem 1.2

We follow the main line of argument for the results of [9]. We observe that the Lagrangian involution $H \mapsto \angle(H)$ permutes the solutions to an instance of a symmetric Schubert problem λ given by isotropic flags, and then construct a family $\mathcal{X}_\lambda \rightarrow \mathcal{Z}_\lambda$ whose base parameterizes instances of the Schubert problem λ given by isotropic flags and whose fibers are the solutions to those instances. We then estimate the codimension of the \angle -fixed point locus of the family $\mathcal{X}_\lambda \rightarrow \mathcal{Z}_\lambda$, which shows that the numerical condition $\sum_i \ell(\lambda^i) \geq m+4$ implies that the fixed points have codimension at least two. Finally, we invoke a key lemma from [9] to complete the proof.

2.1 The Lagrangian Grassmannian

An m -dimensional subspace H of V is *Lagrangian* if $\angle(H) = H$. The set of all Lagrangian subspaces of V forms the *Lagrangian Grassmannian* $\text{LG}(V)$. This is smooth of dimension $\binom{m+1}{2}$ and is a homogeneous space for the symplectic group $\text{Sp}(V)$ of linear transformations of V which preserve $\langle \cdot, \cdot \rangle$.

An isotropic flag F_\bullet and a symmetric partition λ determine a Schubert subvariety $Y_\lambda F_\bullet$ of $\text{LG}(V)$, which is the intersection $X_\lambda F_\bullet \cap \text{LG}(V)$,

$$Y_\lambda F_\bullet := \{H \in \text{LG}(V) \mid \dim H \cap F_{m+i-\lambda_i} \geq i \text{ for } i = 1, \dots, m\}.$$

This has codimension $\|\lambda\| := \frac{1}{2}(|\lambda| + \ell(\lambda))$ in $\text{LG}(V)$.

We need the following result which partially explains why these Lagrangian Schubert varieties are relevant for Theorem 1.2.

Proposition 2.1 ([9, Lemma 9]) *Let λ be a partition and F_\bullet a flag. Then*

$$\angle(X_\lambda F_\bullet) = X_{\lambda'} \angle(F_\bullet).$$

Thus, if λ is symmetric and F_\bullet isotropic, then $\angle(X_\lambda F_\bullet) = X_\lambda F_\bullet$ and $Y_\lambda = (X_\lambda F_\bullet)^\angle$, the set of points of $X_\lambda F_\bullet$ that are fixed by \angle . This has the following consequence.

Corollary 2.2 *The Lagrangian involution permutes the solutions to a symmetric Schubert problem given by isotropic flags.*

2.2 Families Associated with Schubert Problems

Let λ be a symmetric Schubert problem. We construct families whose bases parameterize all instances of λ given by isotropic flags and whose fibers are the solutions to the corresponding instance.

The set $\mathbb{F}\ell$ of isotropic flags in V is a flag manifold for $\text{Sp}(V)$ of dimension m^2 . Define

$$\mathcal{U}_\lambda^* := \{(F_\bullet^1, \dots, F_\bullet^s, H) \mid F_\bullet^i \in \mathbb{F}\ell \text{ and } H \in X_{\lambda^i} F_\bullet^i \text{ for } i = 1, \dots, s\}.$$

We have the two projections $\pi: \mathcal{U}_\lambda^* \rightarrow (\mathbb{F}\ell)^s$ and $\text{pr}: \mathcal{U}_\lambda^* \rightarrow \text{Gr}(m, V)$. For isotropic flags $F_\bullet^1, \dots, F_\bullet^s$, the fiber $\text{pr}(\pi^{-1}(F_\bullet^1, \dots, F_\bullet^s))$ consists of the solutions

$$(2.1) \quad X_{\lambda^1} F_\bullet^1 \cap X_{\lambda^2} F_\bullet^2 \cap \dots \cap X_{\lambda^s} F_\bullet^s$$

to the instance of the Schubert problem λ given by the flags $F_\bullet^1, \dots, F_\bullet^s$.

As $\text{Sp}(V)$ does not act transitively on $\text{Gr}(m, V)$, we cannot use Kleiman’s Theorem [12] to conclude that an intersection (2.1) given by general flags is transverse. Transversality follows instead from the main result of [23]. Consequently, there is a nonempty Zariski open subset $\mathcal{O} \subset (\mathbb{F}\ell)^s$ consisting of s -tuples of isotropic flags for which the intersection (2.1) is transverse and therefore consists of $d(\lambda)$ points.

We seek a family $\mathcal{X} \rightarrow \mathcal{Z}$ of instances of λ where $\dim \mathcal{X} = \dim \mathcal{Z}$ and \mathcal{Z} is irreducible with $\mathcal{Z}(\mathbb{R})$ parameterizing all real instances of λ . Since we cannot easily compute the dimension of \mathcal{U}_λ^* , we replace it by a possibly smaller set. Define \mathcal{U}_λ to be the closure of $\pi^{-1}(\mathcal{O})$ in \mathcal{U}_λ^* . Restricting π to \mathcal{U}_λ gives the dominant map

$$(2.2) \quad \pi: \mathcal{U}_\lambda \rightarrow (\mathbb{F}\ell)^s,$$

where a fiber $\pi^{-1}(F_\bullet^1, \dots, F_\bullet^s)$ is a subset of the intersection (2.1) and is equal to it when the intersection is finite. Thus $\dim \mathcal{U}_\lambda = \dim(\mathbb{F}\ell)^s = s \cdot m^2$.

This family (2.2) has the fault that the real points of its base $(\mathbb{F}\ell)^s$ are s -tuples of real isotropic flags, which are only some of the flags giving real instances of λ .

Let $S_\lambda \subset S_s$ be the group of permutations σ of $\{1, 2, \dots, s\}$ with $\lambda^i = \lambda^{\sigma(i)}$ for all $i = 1, \dots, s$. Then $S_\lambda \simeq S_{a_1} \times \dots \times S_{a_t}$, where λ consists of t distinct partitions μ^1, \dots, μ^t with μ^i occurring a_i times. Then S_λ acts on the families $\mathcal{U}_\lambda^*, \mathcal{U}_\lambda \rightarrow (\mathbb{F}\ell)^s$, preserving

fibers,

$$\text{pr}(\pi^{-1}(F_{\bullet}^1, \dots, F_{\bullet}^s)) = \text{pr}(\pi^{-1}(F_{\bullet}^{\sigma(1)}, \dots, F_{\bullet}^{\sigma(s)})) \quad \text{for all } \sigma \in S_{\lambda}.$$

Define $\pi: \mathcal{X}_{\lambda} \rightarrow \mathcal{Z}_{\lambda}$ to be the quotient of $\mathcal{U}_{\lambda} \rightarrow (\mathbb{F}\ell)^s$ by the group S_{λ} .

2.3 Proof of Theorem 1.2

We defer the proof of the following lemma.

Lemma 2.3 *The map $\pi: \mathcal{X}_{\lambda} \rightarrow \mathcal{Z}_{\lambda}$ is a proper dominant map of real varieties of the same dimension with \mathcal{Z}_{λ} smooth and $\mathcal{Z}_{\lambda}(\mathbb{R})$ connected. The Lagrangian involution preserves fibers of π and the codimension in \mathcal{X}_{λ} of the \angle -fixed points $\mathcal{X}_{\lambda}^{\angle}$ is at least $\frac{1}{2}(\sum_i \ell(\lambda^i) - m)$.*

We recall [9, Lemma 5].

Proposition 2.4 *Let $f: X \rightarrow Z$ be a proper dominant map of real varieties of the same dimension with Z smooth. Suppose that X has an involution \angle preserving the fibers of f such that the image in Z of the set of \angle -fixed points has codimension at least 2.*

If $y, z \in Z(\mathbb{R})$ belong to the same connected component of $Z(\mathbb{R})$, the fibers above them are finite, and at least one contains no \angle -fixed points, then we have

$$\#f^{-1}(y) \cap X(\mathbb{R}) \equiv \#f^{-1}(z) \cap X(\mathbb{R}) \pmod{4}.$$

Remark 2.5 Lemma 5 in [9] requires that there are no \angle -fixed points in either fiber $\pi^{-1}(y)$ or $\pi^{-1}(z)$. This may be relaxed to only one fiber avoiding \angle -fixed points, which may be seen using a limiting argument along the lines of the proof of [9, Corollary 7].

Proof of Theorem 1.2 By Lemma 2.3, the hypotheses of Proposition 2.4 hold as the inequality $\sum_i \ell(\lambda^i) \geq m + 4$ implies that $\text{codim } \pi(\mathcal{X}_{\lambda}^{\angle}) \geq \text{codim}(\mathcal{X}_{\lambda}^{\angle}) \geq 2$. Let $(F_{\bullet}^1, \dots, F_{\bullet}^s)$ be isotropic flags defining a real instance of the Schubert problem λ such that (2.1) is finite.

Since this instance is real, for each $i = 1, \dots, s$ if $\overline{F_{\bullet}^i} = F_{\bullet}^j$, then $\lambda^i = \lambda^j$. Thus there is a permutation $\sigma \in S_{\lambda}$ such that $\overline{F_{\bullet}^i} = F_{\bullet}^{\sigma(i)}$ for $i = 1, \dots, s$, and so the image of $(F_{\bullet}^1, \dots, F_{\bullet}^s)$ in \mathcal{Z}_{λ} is a real point $y \in \mathcal{Z}_{\lambda}(\mathbb{R})$. We complete the proof by exhibiting a point $z \in \mathcal{Z}_{\lambda}(\mathbb{R})$ for which $\pi^{-1}(z)$ consists of $d(\lambda)$ real points, none of which are fixed by \angle .

For distinct $t_1, \dots, t_s \in \mathbb{R}$, the intersection

$$(2.3) \quad X_{\lambda^1} F_{\bullet}(t_1) \cap X_{\lambda^2} F_{\bullet}(t_2) \cap \dots \cap X_{\lambda^s} F_{\bullet}(t_s)$$

is transverse and consists of $d(\lambda)$ real points by the Mukhin–Tarasov–Varchenko Theorem [15]. The osculating flags $F_{\bullet}(t_i)$ are real and isotropic, and we would be done if there were no \angle -fixed points in (2.3). This is equivalent to the intersection of the corresponding Lagrangian Schubert varieties being empty. This is unknown, but expected, as it follows from [22, Conjecture 5.1], which is supported by significant evidence.

Since the intersection (2.3) is transverse, if $(E^1_\bullet, \dots, E^s_\bullet) \in (\mathbb{F}\ell)^s$ are real isotropic flags that are sufficiently close to the osculating flags in (2.3), then the intersection

$$(2.4) \quad X_{\lambda^1} E^1_\bullet \cap X_{\lambda^2} E^2_\bullet \cap \dots \cap X_{\lambda^s} E^s_\bullet$$

is transverse and consists of $d(\lambda)$ real points. By Kleiman’s Theorem [12] we may also assume that $(E^1_\bullet, \dots, E^s_\bullet)$ are general in that the intersection

$$(2.5) \quad Y_{\lambda^1} E^1_\bullet \cap Y_{\lambda^2} E^2_\bullet \cap \dots \cap Y_{\lambda^s} E^s_\bullet$$

of Lagrangian Schubert varieties is either empty or has dimension

$$\begin{aligned} \binom{m+1}{2} - \sum_{i=1}^s \|\lambda^i\| &= \binom{m+1}{2} - \frac{1}{2} \sum_{i=1}^s |\lambda^i| - \frac{1}{2} \sum_{i=1}^s \ell(\lambda^i) \\ &\leq \frac{m^2}{2} + \frac{m}{2} - \frac{m^2}{2} - \frac{m}{2} - 2 = -2. \end{aligned}$$

We conclude that (2.5) is empty and therefore (2.4) contains no Lagrangian subspaces.

If $z \in \mathcal{Z}_\lambda(\mathbb{R})$ is the image of $(E^1_\bullet, \dots, E^s_\bullet) \in (\mathbb{F}\ell)^s$, then the fiber $\pi^{-1}(z)$ (which is (2.4)) consists of $d(\lambda)$ real points, none of which are Lagrangian. This completes the proof. ■

Proof of Lemma 2.3 Consider the quotient of $(\mathbb{F}\ell)^s$ by the group S_λ , which is the product $\mathcal{Z}_\lambda = \text{Sym}_{a_1}(\mathbb{F}\ell) \times \text{Sym}_{a_2}(\mathbb{F}\ell) \times \dots \times \text{Sym}_{a_t}(\mathbb{F}\ell)$, where $\text{Sym}_{a_i}(\mathbb{F}\ell)$ is the quotient $(\mathbb{F}\ell)^{a_i}/S_{a_i}$ and λ consists of t distinct partitions μ^1, \dots, μ^t with μ^i occurring a_i times in λ .

For $F_\bullet \in \mathbb{F}\ell$, let $Z_e^\circ F_\bullet \subset \mathbb{F}\ell$ be those flags in linear general position with respect to F_\bullet . This dense subset of $\mathbb{F}\ell$ is a Schubert variety isomorphic to \mathbb{C}^{m^2} . As F_\bullet varies in $\mathbb{F}\ell$, these form an affine cover of $\mathbb{F}\ell$. Given a finite set $\{F^1_\bullet, \dots, F^a_\bullet\}$ of isotropic flags, there is an isotropic flag F_\bullet that is simultaneously in linear general position with each F^i_\bullet , so that $\{F^1_\bullet, \dots, F^a_\bullet\} \subset Z_e^\circ F_\bullet$. Thus $(\mathbb{F}\ell)^a$ is covered by the S_a -invariant affine varieties $(Z_e^\circ F_\bullet)^a$, each isomorphic to $(\mathbb{C}^{m^2})^a$. By descent, this implies that the quotient $\text{Sym}_a(\mathbb{F}\ell) = (\mathbb{F}\ell)^a/S_a$ is well defined and covered by affine varieties $(Z_e^\circ F_\bullet)^a/S_a$, each isomorphic to $(\mathbb{C}^{m^2})^a/S_a \simeq (\mathbb{C}^{m^2})^a$ as $\mathbb{C}^a/S_a \simeq \mathbb{C}^a$. It follows that $\text{Sym}_a(\mathbb{F}\ell)$ is a smooth irreducible variety whose real points are connected which implies the same for \mathcal{Z}_λ .

The map $\pi: \mathcal{U}_\lambda^* \rightarrow (\mathbb{F}\ell)^s$ is proper, as it comes from a projection along a Grassmanian factor. Its fibers are preserved by the Lagrangian involution and are equal over points in an S_λ -orbit. Both properties hold for $\pi^{-1}(\mathcal{O}) \rightarrow \mathcal{O}$ (as \mathcal{O} is S_λ -stable) and therefore for $\pi: \mathcal{U}_\lambda \rightarrow (\mathbb{F}\ell)^s$. We conclude that π descends to the quotient $\pi: \mathcal{X}_\lambda \rightarrow \mathcal{Z}_\lambda$, where it is a proper dominant map and the Lagrangian involution preserves its fibers.

Since $\dim \mathcal{U}_\lambda = \dim(\mathbb{F}\ell)^s = s \cdot m^2$ and S_λ is a finite group, we conclude that $\dim \mathcal{X}_\lambda = \dim \mathcal{Z}_\lambda = s \cdot m^2$.

We study the \angle -fixed points of \mathcal{U}_λ^* which form the universal family,

$$\mathcal{L}_\lambda := \{(F^1_\bullet, \dots, F^s_\bullet, H) \mid F^i_\bullet \in \mathbb{F}\ell \text{ and } H \in Y_{\lambda^i} F^i_\bullet \text{ for } i = 1, \dots, s\}.$$

Consider the projection $\text{pr}: \mathcal{L}_\lambda \rightarrow \text{LG}(V)$. Let $H \in \text{LG}(V)$. Then

$$\begin{aligned} \text{pr}^{-1}(H) &= \{(F_\bullet^1, \dots, F_\bullet^s, H) \mid H \in Y_{\lambda^i} F_\bullet^i \text{ for } i = 1, \dots, s\} \\ &\simeq \prod_{i=1}^s \{F_\bullet \in \mathbb{F}\ell \mid H \in Y_{\lambda^i} F_\bullet\}. \end{aligned}$$

For λ symmetric and $H \in \text{LG}(V)$, define $Z_\lambda(H) := \{F_\bullet \in \mathbb{F}\ell \mid H \in Y_\lambda F_\bullet\}$. This is a Schubert subvariety of $\mathbb{F}\ell$ of codimension $\|\lambda\|$. Thus

$$\text{pr}^{-1}(H) = Z_{\lambda^1}(H) \times Z_{\lambda^2}(H) \times \dots \times Z_{\lambda^s}(H)$$

has codimension $\sum_i \|\lambda^i\| = \frac{1}{2} \sum_i (|\lambda^i| + \ell(\lambda^i))$ in $(\mathbb{F}\ell)^s$ and is irreducible as each $Z_{\lambda^i}(H)$ is a Schubert variety and is therefore irreducible. Thus $\text{pr}: \mathcal{L}_\lambda \rightarrow \text{LG}(V)$ exhibits \mathcal{L}_λ as a fiber bundle. We compute its dimension:

$$\begin{aligned} \dim \mathcal{L}_\lambda &= \dim \text{LG}(V) + \dim \text{pr}^{-1}(H) = \binom{m+1}{2} + s \cdot m^2 - \sum_{i=1}^s \|\lambda^i\| \\ &= s \cdot m^2 - \frac{1}{2} \left(\sum_{i=1}^s \ell(\lambda^i) - m \right). \end{aligned}$$

Thus $\dim \mathcal{U}_\lambda \cap \mathcal{L}_\lambda \leq s \cdot m^2 - \frac{1}{2} (\sum_i \ell(\lambda^i) - m)$. As $\mathcal{U}_\lambda \cap \mathcal{L}_\lambda$ is the set of \angle -fixed points of \mathcal{U}_λ , $\dim \mathcal{U}_\lambda = m^2$, and \mathcal{X}_λ is the quotient of \mathcal{U}_λ by the finite group S_λ , the \angle -fixed points in \mathcal{X}_λ have codimension at least $\frac{1}{2} (\sum_i \ell(\lambda^i) - m)$. ■

Remark 2.6 If λ is a symmetric Schubert problem, the quantity

$$\sum_{i=1}^s \|\lambda^i\| = \frac{1}{2} \sum_{i=1}^s (|\lambda^i| + \ell(\lambda^i)) = \frac{m^2}{2} + \frac{1}{2} \sum_{i=1}^s \ell(\lambda^i)$$

is an integer, so $\sum_i \ell(\lambda^i)$ has the same parity as m . For generic flags $(E_\bullet^1, \dots, E_\bullet^s)$, the intersection (2.5) of Lagrangian Schubert varieties is a subset of the intersection (2.4) of Schubert varieties. By Kleiman’s Theorem, this gives the inequality

$$\binom{m+1}{2} - \sum_{i=1}^s \|\lambda^i\| \leq m^2 - \sum_{i=1}^s |\lambda^i|,$$

which implies that $m \leq \sum_i \ell(\lambda^i)$. Thus the only possibilities for $\sum_i \ell(\lambda^i)$ for which Theorem 1.2 does not imply a congruence modulo four are m or $m + 2$.

When $\sum_i \ell(\lambda^i) = m$, we have $\binom{m+1}{2} = \sum_i \|\lambda^i\|$ so that λ is a Schubert problem for $\text{LG}(V)$ with $c(\lambda)$ solutions. That is, for general isotropic flags $E_\bullet^1, \dots, E_\bullet^s$, the intersection (2.5) is transverse and consists of $c(\lambda)$ points. When $c(\lambda) \neq 0$, the family $\mathcal{U}_\lambda \rightarrow (\mathbb{F}\ell)^s$ is reducible: \mathcal{L}_λ is one component and $\overline{\mathcal{U}_\lambda} \setminus \mathcal{L}_\lambda$ is the other.

For example, the problem $\square^2 \cdot \square^2 = 8$ on $\text{Gr}(4, 8)$ has $\sum_i \ell(\lambda^i) = 4 = m$. The corresponding problem in $\text{LG}(\mathbb{C}^8)$ has four solutions. Thus four of the eight solutions on $\text{Gr}(4, 8)$ will be isotropic and the other four will not be isotropic. In our experimentation (number 490 on [8]) this problem exhibits a congruence modulo four.

When $\sum_i \ell(\lambda^i) = m + 2$, a general intersection (2.5) of Lagrangian Schubert varieties is empty and $\pi^{-1}(0)$ does not meet \mathcal{L}_λ . There are three possibilities.

- (1) $\mathcal{L}_\lambda \subset \mathcal{U}_\lambda$ and $\pi: \mathcal{L}_\lambda \rightarrow \pi(\mathcal{L}_\lambda)$ generically has finite fibers.

- (2) $\mathcal{L}_\lambda \subset \mathcal{U}_\lambda$ and $\pi: \mathcal{L}_\lambda \rightarrow \pi(\mathcal{L}_\lambda)$ has positive-dimensional fibers.
- (3) $\mathcal{L}_\lambda \not\subset \mathcal{U}_\lambda$.

In case (1), $\pi(\mathcal{L}_\lambda)$ has codimension one as does the image of the set of \angle -fixed points of \mathcal{X}_λ , so Proposition 2.4 does not necessarily imply a congruence modulo four. In cases (2) and (3), $\pi(\mathcal{L}_\lambda)$ has codimension two, and so there will be a congruence modulo four.

We have observed some symmetric Schubert problems with $\sum_i \ell(\lambda^i) = m + 2$ that have a congruence modulo four and some that do not (so that (1) holds). For example, $\sigma_1^2 \cdot \sigma_2 \cdot \sigma_3 = 6$ on $\text{Gr}(4, 8)$ has $\sum_i \ell(\lambda^i) = 6 = m + 2$ (number 495 on [8]) and its numbers of real solutions do not exhibit a congruence modulo four, but $\sigma_1^2 \cdot \sigma_2 \cdot \sigma_3 = 8$ on $\text{Gr}(4, 8)$ has $\sum_i \ell(\lambda^i) = 6 = m + 2$ (number 497 on [8]) and its numbers of real solutions appear to be congruent modulo four. We believe that (3) is unlikely and that (2) holds if and only if there is a congruence modulo four.

References

- [1] B. Anderson and U. Helmke, *Counting critical formations on a line*. SIAM J. Control Optim. 52(2014), no. 1, 219–242. <http://dx.doi.org/10.1137/120890533>
- [2] D. Eisenbud and J. Harris, *Divisors on general curves and cuspidal rational curves*. Invent. Math. 74(1983), 371–418. <http://dx.doi.org/10.1007/BF01394242>
- [3] A. Eremenko and A. Gabrielov, *Degrees of real Wronski maps*. Discrete Comput. Geom. 28(2002), no. 3, 331–347. <http://dx.doi.org/10.1007/s00454-002-0735-x>
- [4] ———, *Rational functions with real critical points and the B. and M. Shapiro conjecture in real enumerative geometry*. Ann. of Math. (2) 155(2002), no. 1, 105–129. <http://dx.doi.org/10.2307/3062151>
- [5] L. M. Fehér and Á. K. Matszangosz, *Real solutions of a problem in enumerative geometry*. Period. Math. Hungar. 73(2016), no. 2, 137–156. <http://dx.doi.org/10.1007/s10998-016-0122-7>
- [6] S. Finashin and V. Kharlamov, *Abundance of real lines on real projective hypersurfaces*. Int. Math. Res. Notices (2012).
- [7] N. Hein, C. J. Hillar, and F. Sottile, *Lower bounds in real Schubert calculus*. São Paulo J. Math. Sci. 7(2013), no. 1, 33–58. <http://dx.doi.org/10.11606/issn.2316-9028.v7i1p33-58>
- [8] N. Hein and F. Sottile, *Beyond the Shapiro Conjecture and Eremenko-Gabrielov lower bounds*, 2013, <http://www.math.tamu.edu/~jssecant/lowerBounds/lowerBounds.php>.
- [9] N. Hein, F. Sottile, and I. Zelenko, *A congruence modulo four in real Schubert calculus*. 2014, J. Reine Angew. Math., to appear.
- [10] I. V. Itenberg, V. M. Kharlamov, and E. I. Shustin, *Welschinger invariant and enumeration of real rational curves*. Int. Math. Res. Not. (2003), no. 49, 2639–2653.
- [11] ———, *Logarithmic equivalence of the Welschinger and the Gromov-Witten invariants*. Uspekhi Mat. Nauk 59(2004), no. 6(360), 85–110. <http://dx.doi.org/10.4213/rm797>
- [12] S. L. Kleiman, *The transversality of a general translate*. Compositio Math. 28(1974), 287–297.
- [13] A. Martín del Campo and F. Sottile, *Experimentation in the Schubert calculus*. In: Schubert Calculus (Osaka 2012), Advanced Studies in Pure Mathematics, 71, Mathematical Society of Japan, 2016, pp. 295–335.
- [14] E. Mukhin and V. Tarasov, *Lower bounds for numbers of real solutions in problems of Schubert calculus*. 2014, [arxiv:1404.7194](https://arxiv.org/abs/1404.7194).
- [15] E. Mukhin, V. Tarasov, and A. Varchenko, *The B. and M. Shapiro conjecture in real algebraic geometry and the Bethe ansatz*. Ann. of Math. (2) 170(2009), no. 2, 863–881. <http://dx.doi.org/10.4007/annals.2009.170.863>
- [16] ———, *Schubert calculus and representations of the general linear group*. J. Amer. Math. Soc. 22(2009), no. 4, 909–940. <http://dx.doi.org/10.1090/S0894-0347-09-00640-7>
- [17] C. Okonek and A. Teleman, *Intrinsic signs and lower bounds in real algebraic geometry* J. Reine Angew. Math. 688(2014), 219–241.
- [18] ———, *A wall-crossing formula for degrees of real central projections*. Int. J. Math. 25(2014), 1450038, 34 pp. <http://dx.doi.org/10.1142/S0129167X14500384>

- [19] K. Purbhoo, *Reality and transversality for Schubert calculus in $OG(n, 2n + 1)$* . *Math. Res. Lett.* 17(2010), no. 6, 1041–1046. <http://dx.doi.org/10.4310/MRL.2010.v17.n6.a3>
- [20] E. Soprunova and F. Sottile, *Lower bounds for real solutions to sparse polynomial systems*. *Adv. Math.* 204(2006), no. 1, 116–151. <http://dx.doi.org/10.1016/j.aim.2005.05.016>
- [21] F. Sottile, *The special Schubert calculus is real*. *Electronic Research Announcements of the AMS* 5(1999), 35–39. <http://dx.doi.org/10.1090/S1079-6762-99-00058-X>
- [22] ———, *Some real and unreal enumerative geometry for flag manifolds*. *Mich. Math. J.* 48(2000), 573–592. <http://dx.doi.org/10.1307/mmj/1030132734>
- [23] ———, *General isotropic flags are general (for Grassmannian Schubert calculus)*. *J. Algebraic Geom.* 19(2010), no. 2, 367–370. <http://dx.doi.org/10.1090/S1056-3911-09-00518-9>
- [24] J.-Y. Welschinger, *Invariants of real rational symplectic 4-manifolds and lower bounds in real enumerative geometry*. *C. R. Math. Acad. Sci. Paris* 336(2003), no. 4, 341–344. [http://dx.doi.org/10.1016/S1631-073X\(03\)00059-1](http://dx.doi.org/10.1016/S1631-073X(03)00059-1)

Department of Mathematics and Computer Science, Benedictine College, Atchison, Kansas 66002, USA
e-mail: nhein@benedictine.edu

Department of Mathematics, Texas A&M University, College Station, Texas 77843, USA
e-mail: sottile@math.tamu.edu

Department of Mathematics, Texas A&M University, College Station, Texas 77843, USA
e-mail: zelenko@math.tamu.edu