

viruses. Compared with CAI, HA-RVI patients were older (mean age 37.1±29.7 vs 21.4±27.5 years, $p<0.001$), had recent hospitalisations (OR 1.5, 95% CI:1.1-2.1, $p=0.007$), underlying bronchial asthma or COPD (OR 2.8, 95% CI:1.5-5.3, $p=0.002$) and were immunosuppressed (OR 7.6, 95% CI:4.3-13.4, $p<0.001$). Interestingly, HA-RVI patients were less likely to have fever (49.8% vs 66.4%, $p<0.001$), cough (42.2% vs 67.3%, $p<0.001$) or shortness of breath (17.5% vs 26.6%, $p=0.02$). Despite fewer symptoms, HA-RVI patients were more likely to have pneumonia with abnormal chest x-rays (33.3% vs 22.7%, RR 1.14, 95% CI:1.02-1.29, $p=0.04$), longer lengths of stay (mean 21.2±36.7 vs 5±14 days, $p<0.001$), higher rates of ICU admission (14.9% vs 8.1%, OR 2.0, 95% CI:1.4-2.9, $p<0.001$), and mortality (4.8% vs 0.6%, OR 8.6, 95% CI:3.6-20.5, $p<0.001$). **Conclusions:** Patients who are older, have pre-existing respiratory disorders or are immunosuppressed face greater HA-RVI risk. HA-RVI patients are less likely to exhibit typical respiratory infection symptoms, potentially delaying diagnosis. This probably contributes to increased morbidity and mortality associated with HA-RVI which underscore the importance of hospital infection prevention even for endemic respiratory viruses.

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s22–s23

doi:10.1017/ash.2025.142

Knowledge and practice of surgical hand hygiene of healthcare worker at University Medical Center, Ho Chi Minh City, 2022

Truong Thi Le Huyen¹, Pham Thi Lan¹, Tran Nguyen Giang Huong¹, Nguyen Thi Minh Khai¹, Le Thanh Truyen¹, Huynh Hoang Hai¹, Trinh Thi Thoa¹, Nguyen Vu Hoang Yen¹ and Huynh Minh Tuan^{1,2}

¹University Medical Center, HCM city and ²University of Medicine and Pharmacy at HCM city

Introduction: Nowadays, surgical site infection is one of the four common types of healthcare-associated infections. There are many preventive measures applied and surgical hand hygiene (SHH) is the most effective and the simplest measure. This study aimed to assess the knowledge, practices of SHH among staff and the relationship between knowledge and practice of SHH in Viet Nam. **Methods:** An analytical cross-sectional study was conducted at the University Medical Center (UMC) in Ho Chi Minh City in 2022. The study employed a set of pre-prepared questions for the knowledge assessment section. For the practical assessment section, the research team conducted indirect observation through cameras and filled out a monitoring checklist. The data were analyzed using Stata 13.2. **Results:** Of the 271 healthcare workers, surgeons had the highest proportion at 48.7%, which was 18.6 times higher than that of anesthesiologists. The majority of healthcare workers received training on SHH, accounting for a rate of 95.6%. Among the participants, the overall compliance of SHH before entering the operative room accounted for 85.6%. The percentage of correct general knowledge reached 73.8%, and there was a relationship between correct knowledge and correct practice with $p < 0.01$. **Conclusion:** Our data suggests that having correct general knowledge of SHH is a crucial factor in accurately practicing SHH. Therefore, providing training to impart accurate knowledge about SHH to healthcare staff is necessary to enhance the overall compliance rate of SHH before entering the operating room.

Keywords: Surgical hand hygiene; Surgical hand hygiene procedure; hand hygiene

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s23

doi:10.1017/ash.2025.143

Results of environmental monitoring at pediatrics isolation ward, tertiary hospital in northern of Thailand

Sutthiphan Thanomphan PhD¹, Mattana Chaiprasert B.N.S² and Nittaya Wungsudthisomsri B.N.S³

¹Department of Nursing Research Development Nakornping, Chiang Mai Province., Thailand, ²Department of Pediatrics Patients Nursing Nakornping Hospital, Chiang Mai Province., Thailand and ³Department of Pediatrics Patients Nursing Nakornping Hospital, Chiang Mai Province., Thailand

Background: Throughout healthcare, the physical environment presents an important source of pathogens that can cause healthcare associated infections (HAIs). To keep patients safe, hospitals must maintain a clean environment and minimize the presence of pathogens. **Objectives:** 1. To identify through environmental monitoring the level of cleanliness in area of pediatrics isolation ward. 2. To assessments of environmental cleaning practice of environmental service (EVS) staff, and healthcare worker. **Methodology:** This retrospective study was done in pediatrics isolation ward, tertiary hospital. Sample sizes were 1 EVS staff, 3 nurses aid, and 102 environmental sites. Data were collected from database of infection control program between February 29, to March 2, 2024 via Infection control assessment and response (ICAR: tool for assessing cleaning practice of EVS staff, personnel by direct performance observations), visual assessment, and monitored the residual bioburden by adenosine tri phosphate (ATP) tests, and swab culture of the surface. Data were analyzed by using descriptive statistics. **Results:** The results of this study revealed that level of cleanliness in area of ward by ATP test found, contaminated spots were highest (61.76%, 21/34) and clean spots were lowest (38.24%, 13/34), while swab culture method found contaminated spots were higher (55.88%, 19/34), clean spots were lower (44.12%, 15/34), and visual monitoring found contaminated and clean spots were 35.29%, 12/5 and 64.71%, 22/34 respectively. The most of contaminated sites were bed rails, toilet sink, Treatment and IV care car, door handle, mop, Light scope & blade (5.88%, 6/102). As the results proved the most organisms were *Acinetobacter spp.*, *Escherichia coli*, and *Pseudomonas spp.*, respectively. **Conclusion:** This study suggests that the environmental cleaning in specialized area must be monitored continuous with standard methods. It is necessary to promote education and training staffs follow update practice guidelines, especially the participation of disciplinary team motivated effective activities in reducing the microbial contamination.

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s23

doi:10.1017/ash.2025.144

Knowledge, practice of medical waste classifications and related factors of health staff at Ho Chi Minh City District 4 Hospital

To Thanh Tam^{1,*} and Ngô Đồng Khanh²

¹Ho Chi Minh city District 4 hospital and ²Ho Chi Minh City University of Medicine and Pharmacy

Corresponding author: MSc To Thanh Tam Tel: 0333415037 Email: tothanhtam123@gmail.com

Objectives: In the face of the complicated developments of the COVID-19 epidemic, the increasing number of cases, accompanied by an increase in the number of personal protective equipment has contributed mainly to the increased amount of medical solid waste. Updating knowledge and practicing the correct classifications of solid medical waste according to regulations is an urgent issue to minimize the risk of pandemic spread, health, and the environment, as well as responding to incidents and

exposures. To determine the proportion of health workers with correct knowledge and practice in classifying solid medical waste and related factors at District 4 Hospital, Ho Chi Minh City. **Methods:** A cross-sectional study was conducted on 149 health workers at District 4 hospital in 2022. Self-administrated questionnaires including personal data, 50 knowledge questions and practice checklists for solid medical waste classification were used. Determine the relationship using the χ^2 test, PR, and the 95% confidence interval. **Results:** Health staff have knowledge account for 87.25%; general practice 53.69%. Knowledge of color coding non-infectious hazardous waste accounts for less than 50%. Waste bin cleaning 9.4%, exposure reporting procedures 30.87%. The age group >30, the subclinical departments, the information sources from radio, and friends have a higher rate of practice correctly than the other group, $p < 0.05$. **Conclusions:** Health staff have correct knowledge account for 87.25%; correct practice account for 53.69%. Health facilities need to maintain training on solid waste classification knowledge, focusing on color coding, symbols, handling and responding to incidents of exposure to medical waste and occupational safety. Fully equipped with different means of communication to instruct, supervision classification, collection and transportation of solid waste to take timely remedial measures.

Keywords: classification; solid medical waste

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s23–s24
doi:10.1017/ash.2025.145

Infection rates, risk factors and microbial etiology of Cerebrospinal Fluid (CSF) shunt infections – a single centre prospective study from India

Dinoop Korol Patambah

Background/Objectives: CSF shunts are widely used in neurosurgery practice for temporary or permanent CSF diversion. Patients on CSF shunts are at risk of device-associated CNS infections particularly ventriculitis or meningitis. The study objectives were to delineate the risk factors and infection rates for various shunt procedures and their microbial etiology. **Methods:** This is a single center prospective cohort study. The study period was 2 years (October 2020- September 2022). Patients were categorised using IDSA criteria as Contamination or Colonisation or Infection. Device days were also collected from the Hospital information system (HIS) for calculation of infection rates. Microbial etiology was identified by culture of CSF and shunt catheter tips. Cox regression model was used to estimate hazard risk for various risk factors. **Results:** During the 2-year study period, 161 shunts were inserted. 133 were ventriculo-peritoneal (VP) shunts, 19 were lumbo-peritoneal (LP) shunts, 6 were subdural-peritoneal (SDP) shunts, 2 were syringo-subarachnoid (SS) shunt and 1 cysto-peritoneal (CP) shunt. Hydrocephalus was the commonest indication for a shunt insertion (71.4 %). There were 8 VP shunt and 1 LP shunt infections during the study period. The average infection rates for VP and LP shunts were 6 and 5.2 per procedure, respectively. Gram negative bacteria caused most of the shunt infections (7/9, 77%). The most common organism causing shunt infection was *Klebsiella pneumoniae* (n=4, 44%), followed by *Staphylococcus aureus* (n=2, 22%). The risk factors which were independently associated with increased risk for shunt infection were Pre-OP ASA score >3 [HR:8.28, $p = 0.013$], presence of associated perioperative systemic [HR:3.89, $p = 0.01$] or scalp infections [HR:3.53, $p = 0.005$]. **Conclusion:** VP and LP shunt infection rates were similar in our study. *Klebsiella pneumoniae* was the commonest causative agent causing shunt infection. High Pre-OP ASA score and associated perioperative scalp or systemic infections were independent risk factors for shunt infection.

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s24
doi:10.1017/ash.2025.146

The impact of hospital wide measures to reduce mupirocin resistance among methicillin-resistant *Staphylococcus aureus* in a Singapore hospital

HML Oh¹ and J Chen¹

¹Department of Infectious Diseases Changi General Hospital, Singapore

Introduction: Methicillin-resistant *Staphylococcus aureus* (MRSA) is a leading cause of healthcare associated infections. Colonization with MRSA increases the risk of subsequent nosocomial infection. The primary concern regarding widespread use of mupirocin is the emergence of mupirocin resistance. A prospective cross-sectional study in Singapore in 2013, found mupirocin resistance to be 31.6% in Changi General Hospital (CGH). Annual usage of mupirocin (g) in CGH was 36870 and hospital-onset MRSA bacteremia was 1.1/10,000 patient-days in 2013. **Objective:** To study the impact of hospital measures to reduce mupirocin resistance among MRSA by detection of mupirocin resistance in screening isolates. **Method:** Changi General Hospital is a 1000 bedded acute care hospital. Hospital wide measures were instituted in CGH to reduce mupirocin resistance in MRSA included a) universal body wash with Octenidine for all hospitalized patients in the wards with MRSA cubicles b) 2% mupirocin ointment removed from formulary (available for nasal decolonization only) A study was conducted on MRSA screening isolates from the Microbiology Laboratory between May and September 2019. These were obtained by swabbing nasal, axilla and groin on all newly admitted patients as part of an active surveillance program since 2013. The swabs were streaked onto MRSA¹selective media plates which were incubated at 35 °C for 20 hours and stored at 4 °C. E-test was performed to determine the susceptibility and minimum inhibitory concentration (MIC) of MRSA isolates to mupirocin, oxacillin and vancomycin, following the CLSI guidelines for *S. aureus*. MPCR (multiplex polymerase chain reaction) assay was used for the simultaneous identification of *ileS-2* (primers MupA and MupB) and *mecA* (primers MecA1 and MecA2). PCR amplification of *ileS-2* gene for high level mupirocin resistance and *MecA* gene was performed on Touch thermal cycler. **Results:** 200 MRSA isolates were tested. E-test revealed 5 isolates were detected to be High Level mupirocin-resistant (2.5%) and 69 isolates were detected to be oxacillin-resistant (74%). The MPCR assay detected *mecA* gene in 100% and *ileS-2* gene in 3 isolates (1.5%). **Conclusion:** Our study indicated the low prevalence of high level mupirocin resistance among MRSA screening isolates in 2019 in CGH. This suggested that the hospital wide measures to reduce mupirocin resistance were effective.

Antimicrobial Stewardship & Healthcare Epidemiology 2025;5(Suppl. S1):s24
doi:10.1017/ash.2025.147

Risk factors for Catheter-Associated Urinary Tract Infection (CAUTI) in sepsis patients at RSPAD GATOT SOEBROTO 2022: a quantitative study

Theresia Leonita¹, Soroy Lardo², Maria Selvester Thadeus³, Marlina Dewi Astuti⁴, Martaviani Budiastuti⁵ and Jonny⁶

¹Faculty of Medicine Universitas Pembangunan Nasional “Veteran” Jakarta, Jakarta, Indonesia, ²Division of Tropical and Infectious Diseases, Department of Internal Medicine Gatot Soebroto Army Hospital, Jakarta, ³Department of Pathology Anatomy Universitas Pembangunan Nasional “Veteran” Jakarta, Jakarta, Indonesia, ⁴Department of Internal Medicine Universitas Pembangunan Nasional “Veteran” Jakarta, Jakarta, Indonesia, ⁵Head of PPI Committee Gatot Soebroto Army Hospital, Jakarta and ⁶Division of Nephrology & Hypertension, Department of Internal Medicine Gatot Soebroto Army Hospital, Jakarta

Correspondence E-mail: theresialeonitaa@gmail.com

Background: Urinary tract infection (UTI) is the most dominant case, around 40% of healthcare-associated infections (HAIs). UTI related to