

WALTHAM SUPPLEMENT

Salmonellosis impacts the proportions of faecal microbial populations in domestic cats fed 1–3-d-old chicks*

K. R. Kerr¹, S. E. Dowd² and K. S. Swanson^{1†}

¹Department of Animal Sciences and Division of Nutritional Sciences, University of Illinois at Urbana-Champaign, Urbana, IL, USA

²MR DNA (Molecular Research LP), Shallowater, TX, USA

(Received 1 November 2013 – Final revision received 19 December 2013 – Accepted 14 January 2014)

Journal of Nutritional Science (2014), vol. 3, e30, page 1 of 5

doi:10.1017/jns.2014.20

Abstract

There has been a recent increase in the feeding of unconventional diets, including whole-prey diets, to domestic pet cats. Our objective was to characterise faecal microbial populations of domestic cats fed whole and ground (6·35 mm grind) raw 1–3-d-old chicks (Rodent Pro). Faecal samples were collected from neutered male domestic cats (mean age = 5·7 years) fed these diet items in a crossover design. Bacterial DNA was isolated from faecal samples and amplicons of the 16S rRNA V4–V6 region were generated and analysed by 454 pyrosequencing. Faecal microbial populations of cats fed whole v. ground chicks did not differ. During the study, three cats presented with symptoms of infection (anorexia or diarrhoea) and tested clinically positive for *Salmonella* using a standard PCR method. The remaining cats tested negative. Data were analysed *post hoc* to test for differences in microbial populations due to clinical status. The predominant genera were *Clostridium* (9–30 %), unidentified Lachnospiraceae (10–28 %), *Blautia* (4–19 %), *Peptococcus* (2–19 %) and *Fusobacterium* (2–14 %). Faeces of cats testing clinically positive for *Salmonella* had higher ($P \leq 0.05$) proportions of the genera *Coprococcus* (5·6 v. 0·4 %) and *Escherichia* (subgenera *Shigella*; 1·1 v. 0·3 %). *Salmonella* was not detected in faecal samples utilising the pyrosequencing method; however, there was a shift in microbial populations due to clinical status. The clinical symptoms reported herein may be not only due to the *Salmonella* itself, but also shifts in other gut microbial populations.

Key words: Feline nutrition; Raw diet; *Salmonella*

There has been a recent increase in feeding unconventional diets, including whole-prey diets, to domestic pet cats. Feeding whole prey mimics the diet of small wild cats, which typically eat small mammals, reptiles, birds and insects⁽¹⁾. Whole-prey items fed raw require minimal processing, including no heat treatment or addition of preservatives. Dietary processing and ingredient and nutrient composition impact faecal microbial populations in cats^(2–8); however, little is known about the microbial population of cats fed a ‘wild-type’^(7,8).

Feeding raw foods may increase risk of exposure to potentially pathogenic bacteria for companion animals and human subjects. Whole-prey diet items containing gut contents make it inevitable that the animals ingesting them will be

exposed to foreign bacteria, which may be pathogenic^(9–11). Infection of cats, dogs and human subjects with pathogenic bacteria have been linked to contaminated pet foods, including raw meat-based foods, commercial dry foods and wild-caught prey^(9,11–14). During infection, pathogenic bacteria compete with and displace the commensal species resulting in dysbiosis of microbial population and gastrointestinal upset⁽¹⁵⁾. The authors are aware of no studies that have utilised pyrosequencing to examine the faecal microbial populations of infected cats fed raw diets.

Our primary objective was to characterise the microbiota of cats fed whole and ground 1–3-d-old chicks. During the study three cats presented with symptoms of infection (anorexia or diarrhoea), and multiple pathogenic bacteria were detected in

† Corresponding author: K. S. Swanson, fax +1 217 333 5044, email ksswanso@illinois.edu

* This article was published as part of the WALTHAM International Nutritional Sciences Symposium Proceedings 2013.



their faeces utilising PCR and culture techniques, including *Salmonella*. Therefore a secondary aim was to examine the alterations in the faecal bacterial populations of cats with symptomatic salmonellosis compared with those which tested negative for *Salmonella*.

Experimental methods

Study design

The animal protocol was approved by the University of Illinois Animal Care and Use Committee. Faecal samples were collected from neutered male domestic cats ($n = 12$; mean age = 5.7 years; body condition score 4.5–5.5 of 9). A crossover design was utilised to test the impacts of two dietary treatments: (1) whole or (2) ground (6.35 mm grind) 1–3-d-old chicks (DM: 25 %, crude protein: 71 % DM basis, fat: 20 % DM basis; Rodent Pro). Fresh water was available *ad libitum*. Cats were adapted to diets for 10 d prior to fresh faecal collection (<15 min from defecation). Faecal samples were stored at -80°C until DNA extraction.

Sample analysis

Faecal bacterial DNA was isolated according to procedures described previously⁽¹⁶⁾ using the MO BIO PowerSoil™ Kit (MO BIO Laboratories, Carlsbad, CA, USA). Amplification of a 600 bp sequence of the V4–V6 variable region of the 16S rRNA gene was done using barcoded primers as previously described⁽¹⁷⁾. PCR amplicons were further purified utilising AMPure XP beads (Beckman-Coulter Inc.). Amplicons were combined in equimolar ratios to create a DNA pool that was used for pyrosequencing. DNA quality of amplicon pools was assessed before pyrosequencing using a 2100 Bioanalyser (Agilent Technologies). Pyrosequencing was performed at the W. M. Keck Center for Biotechnology at the University of Illinois utilising a 454 Genome Sequencer and FLX titanium reagents (Roche Applied Science).

Data analysis

High-quality (quality value > 25) sequence data derived from the sequencing process was processed using a proprietary analysis pipeline (www.mrdnalab.com) and as described previously⁽¹⁸⁾.

Clinical testing

During the study, three cats presented with symptoms of infection (anorexia or diarrhoea). Two diet samples and faecal samples for cats with ($n = 3$) and without ($n = 9$) symptoms were submitted to the Veterinary Diagnostic Laboratory (University of Illinois) for *Salmonella* detection using PCR, *Salmonella* antimicrobial resistance ($n = 3$ positive samples only), *Campylobacter* detection using PCR, aerobic culture and anaerobic culture. Positive detection of *Salmonella* was verified by isolation and culture. The cat that presented with anorexia was removed from the larger study. A faecal sample had already been obtained prior to his symptoms manifesting.

Statistical analysis

Sequence percentages at each taxonomic level were analysed using the Mixed models procedure of SAS (version 9.3; SAS Institute, Cary, NC, USA). The fixed effect of diet was tested. Means were separated for treatments using a Fisher-protected least significant difference with Tukey's adjustment. Results are reported as least-squares means with $P \leq 0.05$ defined as significant and $P \leq 0.10$ as trends for treatment effects. Genera data and select species data were analysed *post hoc* to test for differences in microbial populations due to clinical *Salmonella* status.

Results

Dietary treatment did not impact proportions of bacterial phyla or genera (data not shown). Firmicutes (62–86 % of all sequences) was the predominant bacterial phylum in cat faeces. Proteobacteria (0.6–16 % of all sequences), Fusobacteria (2–14 % of all sequences), Actinobacteria (1–10 % of all sequences), Tenericutes (0.7–9 % of all sequences) and Bacteroidetes (0–0.7 % of all sequences) were also present (data not shown). The predominant genera were *Clostridium* (9–30 % of sequences), unidentified Lachnospiraceae (10–27 % of sequences), *Blautia* (4–19 % of sequences), *Peptococcus* (2–9 % of sequences) and *Fusobacterium* (2–14 % of sequences).

The three cats exhibiting symptoms of infection (anorexia or diarrhoea) tested clinically positive for a group B *Salmonella* species using a standard PCR assay (Table 1). Diet samples and the remaining cats tested negative. The *Salmonella* was resistant to the antibiotics Clindamycin, Erythromycin, Penicillin and Oxacillin, and susceptible to Amoxicillin, Ampicillin and fourteen other antibiotics. Regardless of symptoms, the results were similar for *Campylobacter*, aerobic culture and anaerobic culture. *Salmonella* was not detected in any of the samples utilising pyrosequencing; however, the proportions of select faecal genera were modified in cats with symptomatic salmonellosis (Table 2). Cats with symptomatic salmonellosis had increased ($P \leq 0.05$) proportions of *Coprococcus* and *Escherichia* subgenera *Shigella*, and tended to have decreased ($P \leq 0.09$) proportions of *Oscillobacter* and *Anaerotruncus*. Twenty-eight species from the genera *Campylobacter*, *Clostridium*, *Enterococcus* and *Escherichia* were examined. Cats with symptomatic salmonellosis had increased ($P \leq 0.05$) faecal proportions of *Clostridium perfringens* (6 v. 1 % of sequences) and an unidentified species from *Escherichia* subgenera *Shigella* (0.5 v. < 0.1 % of sequences) (data not shown).

Discussion

The gut of cats are colonised by trillions of bacteria, which exist in a symbiotic relationship with the host⁽¹⁹⁾. The majority of these bacteria are neutral to the host, whereas others confer benefits to the host, including proper development of the immune system, the digestion of food and absorption of nutrients, the production of key vitamins (e.g. vitamin K and biotin) and protection against invading pathogenic organisms. During infection, pathogenic bacteria compete with and displace the commensal species resulting in gastrointestinal



Table 1. Potentially pathogenic bacteria detected by PCR and culture methods in the diet and faeces of domestic cats with (*n* 3) and without (*n* 9) symptoms of infection (expressed as number of samples)

Item	Test result	Diet*	Faecal samples	
			No symptoms	Symptomatic†
<i>Salmonella</i> detection PCR				
	Negative	2	9	0
	Positive	0	0	3
<i>Campylobacter</i> detection PCR				
	Negative	2	9	3
	Positive	0	0	0
Aerobic culture				
<i>Clostridium perfringens</i>				
	None reported	2	0	0
	Heavy	0	7	2
	Very heavy	0	2	1
<i>Clostridium</i> species				
	None reported	2	0	2
	Heavy	0	4	1
	Very heavy	0	5	0
Anaerobic culture				
<i>Escherichia coli</i>				
	Few	0	1	0
	Moderate	2	5	1
	Heavy	0	3	1
	Very heavy	0	0	1
<i>Enterococcus</i> species				
	None reported	0	0	1
	Moderate	1	0	1
	Heavy	1	5	1
	Very heavy	0	4	0

*Whole and ground 1–3-d-old chicks (Rodent Pro).

†Symptomatic = anorexia (*n* 1); diarrhoea (*n* 2).

Table 2. Predominant bacterial genera (expressed as percentage of sequences) in faeces of domestic cats with (*n* 3) and without (*n* 9) symptoms of infection

Phylum	Family	Genus	Salmonella negative	Salmonella positive	SEM	P value
Actinobacteria	Coriobacteriaceae	<i>Collinsella</i>	5.1	3.5	1.6	0.49
		<i>Slackia</i>	0.2	0.1	0.1	0.65
	Firmicutes	<i>Clostridiaceae</i>	15.8	22.3	3.1	0.18
		<i>Enterococcaceae</i>	1.4	1.6	0.7	0.74
		<i>Erysipelotrichaceae</i>	1.2	1.4	0.8	0.83
		Unidentified genera	0.7	0.2	0.3	0.19
		<i>Eubacteriaceae</i>	4.2	3.1	1.7	0.38
		<i>Lachnospiraceae</i>	10.1	12	2.8	0.65
		<i>Coprococcus</i>	0.4	5.6	1.4	< 0.01
		<i>Moryella</i>	0.6	0.1	0.3	0.27
		<i>Psuedobutyryvibrio</i>	3.6	4.3	0.9	0.50
Fusobacteria	Proteobacteria	<i>Roseburia</i>	1.0	0.7	0.5	0.67
		Unidentified genera	16.6	12.3	2.4	0.22
		<i>Oscillibacter</i>	0.4	0.1	0.1	0.06
		<i>Peptococcaceae</i>	10.3	10.9	2.5	0.87
		<i>Peptostreptococcaceae</i>	1.6	0.0	1.4	0.39
		Unidentified genera	0.3	0.4	0.1	0.35
		<i>Ruminococcaceae</i>	2.3	1.4	0.4	0.09
		<i>Anaerotruncus</i>	4.5	4.8	1.6	0.89
		<i>Ruminococcus</i>	2.5	1.1	0.8	0.20
		Unidentified genera	8.6	7.4	2.1	0.71
	<i>Fusobacteriaceae</i>	<i>Fusobacterium</i>	1.6	0.3	1.7	0.61
	<i>Campylobacteraceae</i>	<i>Campylobacter</i>	0.3	1.1	0.3	0.03
	<i>Enterobacteriaceae</i>	<i>Escherichia Shigella</i>	1.8	3.5	1.3	0.41
	<i>Succinivibrionaceae</i>	<i>Anaerobiospirillum</i>	0.8	1.3	0.8	0.73
	<i>Sutterellaceae</i>	<i>Sutterella</i>	1.1	0.7	0.5	0.58



upset, including inflammation, nausea, gas production and bloating, and diarrhoea⁽¹⁵⁾. Herein we identified alterations in the faecal bacterial genera of cats with symptomatic salmonellosis.

Salmonella are rod-shaped Gram-negative enterobacteria. Reported *Salmonella* isolates from cats with pathogenic infection include *S. typhimurium* (most commonly isolated), *S. choleraesuis*, *S. dublin*, *S. newport*, *S. arizonae*, *S. saintpaul*, *S. krefeld*, *S. typhisuis*, *S. enteritidis*, *S. hadar*, *S. manhattan*, *S. infantis* and *S. virchow*⁽¹⁴⁾. *Salmonella* infection in cats can be asymptomatic, or cause gastroenteritis, and/or septicaemia⁽¹⁴⁾. The symptoms noted herein, anorexia and diarrhoea, are typical of salmonellosis. Additional symptoms (not observed herein) include fever, lethargy, vomiting, weight loss and dehydration among others. Cats herein were monitored closely, the infections were self-limiting and resolved without treatment within a few days.

Exposure to pathogenic species of strains by faecal-to-oral transmission, and ingestion of contaminated feed may lead to zoonotic infection (transmissible between species). However, not all host animal species will develop infection or overt symptoms when a pathogenic species of strain is present (i.e. poultry species can act as a reservoir for pathogenic *Salmonella* species without overt signs of infection)⁽²⁰⁾. Although the diet samples tested herein were negative for *Salmonella*, Kerr *et al.*⁽¹⁰⁾ detected *Salmonella* in 1–3-d-old chicks obtained from the same supplier. Whole-prey diet items containing gut contents make it inevitable that the animals ingesting them will be exposed to foreign bacteria. Contamination of poultry species with potentially pathogenic bacteria may occur pre-harvest (e.g. breeding, growth, etc.), during harvest (handling and transport) or after harvest (during slaughter, storage or transport). Because whole prey are typically fed raw, and freezing does not kill *Salmonella*, extra caution should be taken during production processes to reduce contact of whole prey with pathogenic bacteria.

The natural history, digestive physiology (e.g. rapid transit time) and commensal microbiota of healthy adult cats may allow them to tolerate exposure and harbour pathogenic bacteria without overt symptoms of disease⁽¹⁴⁾. However, exposure to pathogenic bacteria should be minimised in cats, and the risk for human subjects may be a concern. Proper handling of raw animal products, bowls and other surfaces that come into contact with the raw items can minimise the risk of exposure. Because cats themselves can harbour pathogens, such as *Salmonella*, without overt signs of infection, handling animals fed whole prey may increase risk of infection for human subjects. These potential risks mean that whole-prey feeding may not be appropriate for all households (e.g. people or animals that are immune-compromised).

Salmonella was not confirmed utilising the pyrosequencing technique in this study. However, the proportions of select faecal genera were modified in cats with symptomatic salmonellosis. These data indicate that pyrosequencing may not be a sensitive enough assay for detecting salmonellosis. Even though the culture techniques gave similar results for symptomatic and asymptomatic cats, salmonellosis may have been secondary to infection by another pathogenic bacteria.

Both the results of the aerobic and anaerobic cultures (detection of other potentially pathogenic bacteria) and pyrosequencing (increased proportions of other potentially pathogenic bacteria) support the role of other pathogenic bacteria (e.g. *Clostridium perfringens* and *Escherichia coli* species) in the progression of gastroenteritis in these cats. We do not have enough data to determine which potentially pathogenic bacteria were responsible for the initial infection. Further investigations to determine the relationship between clinical status and microbial shifts may provide insight into the diagnosis of salmonellosis.

Acknowledgements

The authors have no conflicts of interest to declare. This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. In regards to authorship, K. R. K. and K. S. S. contributed to formulation of research, study design and study execution. K. R. K. and S. E. D. contributed to data analysis. K. R. K. and K. S. S. contributed to manuscript writing. As stated in the Experimental Methods section, the animal protocol for this experiment was approved by the University of Illinois Animal Care and Use Committee.

This paper was published as part of the WALTHAM International Nutritional Sciences Symposium Proceedings 2013, publication of which was supported by an unrestricted educational grant from Mars Incorporated. The papers included in these proceedings were invited by the Guest Editor and have undergone the standard journal formal review process. They may be cited.

References

- Plantinga EA, Bosch G & Hendriks WH (2011) Estimation of the dietary nutrient profile of free-roaming feral cats: possible implications for nutrition of domestic cats. *Br J Nutr* **106**, 35S–38S.
- Bermingham EN, Kittelmann S, Henderson G, *et al.* (2011) Five-week dietary exposure to dry diets alters the faecal bacterial populations in the domestic cat (*Felis catus*). *Br J Nutr* **106**, S49–S52.
- Bermingham EN, Young W, Kittelmann S, *et al.* (2013) Dietary format alters fecal bacterial populations in the domestic cat (*Felis catus*). *Microbial Open* **2**, 173–181.
- Hooda S, Vester Bolter BM, Kerr KR, *et al.* (2013) The gut microbiome of kittens is affected by dietary protein:carbohydrate ratio and associated with blood metabolite and hormone concentrations. *Br J Nutr* **109**, 1637–1646.
- Barry K, Middelbos IS, Vester Bolter BM, *et al.* (2012) Effects of dietary fiber on the feline gastrointestinal metagenome. *J Proteome Res* **11**, 5924–5933.
- Tun HM, Brar MS, Khin N, *et al.* (2012) Gene-centric metagenomics analysis of feline intestinal microbiome using 454 junior pyrosequencing. *J Microbiol Methods* **88**, 369–376.
- Desai AR, Musil KM, Carr AP, *et al.* (2009) Characterization and quantification of feline fecal microbiota using cpn60 sequence-based methods and investigation of animal-to-animal variation in microbial population structure. *Vet Microbiol* **137**, 120–128.
- Kerr KR, Dowd SE & Swanson KS (2014) Faecal microbiota of domestic cats fed raw whole chicks *v.* an extruded chicken-based diet. *J Nutr Sci* **3**, doi:10.1017/jns.2014.21.



9. Crump JA, Griffin PM & Angulo FJ (2002) Bacterial contamination of animal feed and its relationship to human foodborne illness. *Clin Infect Dis* **35**, 859–865.
10. Kerr KR, Vester Boler BM, Morris CL, *et al.* (2012) Apparent total tract energy and macronutrient digestibility and fecal fermentative end-product concentrations of domestic cats fed extruded, raw beef-based, and cooked beef-based diets. *J Anim Sci* **90**, 515–522.
11. Tauni MA & Osterlund A (2000) Outbreak of *Salmonella typhimurium* in cats and humans associated with infection in wild birds. *J Small Anim Pract* **41**, 339–341.
12. Dato V, Moll M, Sandt C, *et al.* (2010) Human salmonella infections linked to contaminated dry dog and cat food, 2006–2008. *Pediatrics* **126**, 477–483.
13. Leonard EK, Pearl DL, Finley RL, *et al.* (2011) Evaluation of pet-related management factors and the risk of *Salmonella* spp. carriage in pet dogs from volunteer households in Ontario (2005–2006). *Zoonoses Public Health* **58**, 140–149.
14. Stiver SL, Frazier KS, Mauel MJ, *et al.* (2003) Septicemic salmonellosis in two cats fed a raw-meat diet. *J Am Anim Hosp Assoc* **39**, 538–542.
15. Suchodolski JS, Markel ME, Garcia-Mazcorro JF, *et al.* (2012) The fecal microbiome in dogs with acute diarrhea and idiopathic inflammatory bowel disease. *PLoS ONE* **7**, e51907.
16. McInnes P & Cutting M (2010) Manual of procedures for human microbiome project: core microbiome sampling, protocol A, HMP protocol no. 07–001, version 11. 2010 http://www.hmpdacc.org/doc/sops_2/manual_of_procedures_v11.pdf (accessed March 2013).
17. Cephas KD, Kim J, Mathai RA, *et al.* (2011) Comparative analysis of salivary bacterial microbiome diversity in edentulous infants and their mothers or primary care givers using pyrosequencing. *PLoS ONE* **6**, e23503.
18. Swanson KS, Dowd SE, Suchodolski JS, *et al.* (2011) Phylogenetic and gene-centric metagenomics of the canine intestinal microbiome reveals similarities with humans and mice. *ISME J* **5**, 639–649.
19. Kerr KR, Beloshapka AN & Swanson KS (2013) 2011 and 2012 Early Careers Achievement Awards: use of genomic biology to study companion animal intestinal microbiota. *J Anim Sci* **91**, 2504–2511.
20. White DG, Sudler R, Ayers S, *et al.* (2001) The isolation of antibiotic-resistant *Salmonella* from retail ground meats. *Mass Med Soc* **345**, 1147–1154.