

Keywords: psychiatry; Psychoanalysis; Medical Body

EPP0723

Setting up a clinical psychology service for reception department with consultative care

N. Semenova* and S. Shport

Psychiatry, Moscow Research Institute of Psychiatry – a branch of V. Serbsky National Medical Research Centre for Psychiatry and Narcology, Moscow, Russia, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1039

Introduction: There is a plenty of literature on providing psychological services for psychiatry inpatients and outpatients. Seeing a psychiatrist for the first time can be stressful.

Objectives: This paper will identify challenges in setting up a clinical psychology service for consultees seen in Reception Department with consultative and primary specialized health care. 'Clinical psychology service' is a project in Moscow Research Institute of Psychiatry providing services within the Reception Department setting. A need to address mental ill health issues (getting a diagnosis and treatment plan) within the broader psychosocial needs of consultees has been identified.

Methods: Clinical psychology established a brief intervention 'service' for consultees. Issues of establishing trust within this population suggested the need to provide a 'named' male or female psychologist. The psychologist accompanied psychiatric consultants' recommendations to familiarize a consultee of the availability of the service. Evaluation of the service, including uptake, client satisfaction, and outcome, is ongoing using quantitative and qualitative methods. Data is presented on key themes identified in providing psychological services to consultees.

Results: Key themes identified included: 1. Service development: establishing trust, ensuring confidentiality, close between consultants working, flexibility, crisis management, safety, establish links with other agencies; 2. Complex psychological and social needs: mental health issues, trauma, substance misuse, domestic and sexual violence.

Conclusions: Consultees present with a range of complex psychosocial needs. While this population may have reservations about accessing 'standard' mental health services, a flexible psychology service working in close liaison with psychiatric consultants may be effective in addressing these needs.

Keywords: Reception Department; Consultees; Brief intervention service; Clinical psychology

EPP0724

Prevalence of adhd in the adult population in the Czech Republic and frequency of medication

M. Vňuková and R. Ptáček*

Psychiatry, First Faculty of Medicine Charles University in Prague and General Teaching Hospital, Prague, Czech Republic

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1040

Introduction: Attention Deficit Hyperactivity Disorder (ADHD) is a common neurodevelopmental disorder often diagnosed between the ages of 7 and 10. The estimated prevalence of ADHD in adults is usually 2-5%, which means that in up to half of people this diagnosis persists into adulthood.

Objectives: The aim of this sub-study was to determine whether there are significant differences in ADHD symptomatology between individuals who have taken or are taking prescription drugs and those who have never taken them.

Methods: Data collection was performed by STEM / MARK in January 2019 through the European National Panel. Respondents completed a demographic questionnaire focusing on the history of ADHD and a standardized ADHD Self-Report Scale (ASRS) questionnaire for the symptomatology of ADHD in adulthood.

Results: Of the 1,518 respondents, 3% reported being diagnosed with ADHD / hyperkinetic disorder during their lifetime. According to the ASRS assessment, 119 respondents were classified as suspected ADHD. Overall, men scored higher symptoms of ADHD. The results also show that the group that states that taking medication has a statistically significantly higher average score in ASRS than unmedicated individuals. 6 individuals are taking medication to this day.

Conclusions: The data collected confirm our main hypothesis that ADHD symptomatology has an impact on the daily functioning of individuals in adulthood. Unlike peers with reduced or no ADHD symptoms, these people are far more likely to have time management issues, need to plan their day carefully, and yet often experience problems such as late arrivals due to a lack of anticipation.

Conflict of interest: Financial support was provided by grants : GA ČR – 18 -112 47 S, Progres Q06 1LF

Keywords: adult ADHD prevalence; ASRS; medication

EPP0726

An outpatient clinic for integrative mental health: Patient and treatment characteristics and health outcomes based on patient reported routine outcome monitoring data

R. Hoenders¹, A. Bartels^{1,2}, S. Castelein^{3,4}, E. Visser² and S. Booi^{1,5*}

¹Center For Integrative Psychiatry, Lentis, Groningen, Netherlands;

²Psychiatry, Rgoc, University of Groningen, University Medical Center Groningen, Groningen, Netherlands; ³Clinical Psychology And

Experimental Psychopathology, University of Groningen, Groningen, Netherlands; ⁴Lentis Research, Lentis, Groningen, Netherlands and

⁵Department Of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1041

Introduction: There is an increasing interest in integrative (mental) health care and a growth in centers offering such services, but a paucity of research on patient characteristics, diagnosis, treatments offered, the effects of those treatments and patient satisfaction.

Objectives: To examine the course of mental health outcomes in the context of the nature and quality of care of outpatients at a center for integrative psychiatry in the Netherlands, as well as relevant sociodemographic, clinical, and treatment-related moderators of this course.