



Magnets used to tune supercooled gallium arsenide semiconductors

Physicists from Rice University and Princeton University have discovered how to use one of the information technology industry's mainstay materials—gallium arsenide semiconductors—as an ultrasensitive microwave detector that could be suitable for next-generation computers. The discovery comes at a time when computer chip engineers are racing both to add nanophotonic devices directly to microchips and to boost processor speeds beyond 10 GHz. Their research has been published in the December 10, 2010 issue of *Physical Review Letters* (DOI: 10.1103/PhysRevLett.105.246802; 246802).

“Tunable photon-detection technology in this microwave range is not well-developed,” said Rice physicist R.-R. Du, the study's lead author. “Single-photon detectors based on superconductors in

the 10-GHz to 100-GHz range are available, but their resonance frequency has been difficult to tune. Our findings suggest that tunable single-photon detection may be within reach with ultrapure gallium arsenide.”

The study is the latest result from a long-term collaboration between Du and Princeton University physicist L. Pfeiffer, whose group produces pure samples of gallium arsenide. For this study, Rice graduate student Y. Dai cooled one of Pfeiffer's ultrapure samples to below 4 K—the temperature of liquid helium. She then bombarded the sample with microwaves while applying a weak magnetic field—approximately the same strength as that of a refrigerator magnet. Du and Dai were surprised to find that microwaves of a specific wavelength resonated strongly with the cooled sample. They also found they could use the magnet to tune this resonance to specific microwave frequencies.

Du said previous experiments have typically measured weak resonance effects

from microwaves. “A signal level of 1% is a common measurement. In our case, the change was a thousand times that much.”

While the team does not yet understand the mechanism that leads to such a sensitive reaction, they are eagerly pursuing follow-up research to try to prove they can use the effect for single-photon measurements.

A photon is the smallest possible unit of light or electromagnetic radiation. By incorporating devices that create, transmit, and measure digital information through photons, rather than with electrons, makers of computer chips hope to produce computers that are both faster and more powerful.

“The clock speed of a new computer right now is about 2 GHz,” Du said. “For the next generation, the industry is shooting for around 100 GHz, which is a microwave device. The phenomenon we've observed is in this region, so we hope it may be useful for them.”

Additional co-authors include K. West of Princeton University. □

MRS Publications Hosted on Cambridge Journals Online

COMING EARLY 2011

Expanded benefits for authors, readers and libraries—advanced search capabilities, reference linking, mobile browser accessibility, and more!

www.mrs.org/publications



MRS Bulletin

- New design and layout
- Now includes Energy Quarterly
- Viewable in page-turning format

PLUS ... Join the Conversation on the Materials for Energy blog
www.materialsforenergy.org

- No author page charges
- 24 issues per year
- Free online color
- Electronic-only subscriptions now available
- 3 special focus issues in 2011

MRS Online PROCEEDINGS LIBRARY

- over 68,000 papers, and counting
- all FREE to MRS members

MRS MATERIALS RESEARCH SOCIETY
Advancing materials. Improving the quality of life.

CAMBRIDGE
UNIVERSITY PRESS