



Maternal iodine intake and adherence to iodine supplement recommendations in a group of Chinese women: the results from the WIN cohort study – CORRIGENDUM

S. Liu^{1,2}, A. Sharp², Z. Lu³ and Z. F. Ma¹

¹*Department of Health and Environmental Sciences, Xi'an Jiaotong-Liverpool University, Suzhou, China,*

²*Harris-Wellbeing Research Centre, Faculty of Health & Life Sciences, University of Liverpool, Liverpool, UK,*

³*Department of Biological Sciences, Xi'an Jiaotong-Liverpool University, Suzhou, China*

⁴*Centre for Public Health and Wellbeing, School of Health and Social Wellbeing, College of Health, Science and Society, University of the West of England, Bristol, UK.*

doi: 10.1017/S0029665123000356, published by Cambridge University Press, 8 March 2023.

In the original publication of this article, the wrong figures had been provided for the mean iodine intake plus iodised salt in pregnant and lactating women. This was given as 159.56 µg/d and 157.34 µg/d, but should have been 232 µg/d and 237 µg/d. The article has since been updated to reflect this.

Reference

Liu, S., Sharp, A., Lu, Z., & Ma, Z. (2023). Maternal iodine intake and adherence to iodine supplement recommendations in a group of Chinese women: The results from the WIN cohort study. *Proceedings of the Nutrition Society*, **82**(OCE1), E27. doi:10.1017/S0029665123000356